



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Health and Behavioural Sciences - Papers
(Archive)

Faculty of Science, Medicine and Health

2006

Health benefits of herbs and spices: the past, the present, the future - Public Health

P. G. Williams

University of Wollongong, peterw@uow.edu.au

Publication Details

This is a preprint of an article originally published as Williams, PG, Health benefits of herbs and spices: Public Health, Medical Journal of Australia, 185(4), 2006, Supplement S17-S18. Copyright Australasian Medical Publishing Company 2006. Original journal available [here](#).

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Health benefits of herbs and spices: the past, the present, the future - Public Health

Abstract

Recommendations for intakes of food in the Australian Guide to Healthy Eating do not yet include suggested intakes of herbs and spices, although several dietary guidelines refer to their benefits. Future consideration should be given to including more explicit recommendations about the place of herbs and spices in a healthy diet

Keywords

herbs, spices, dietary guidelines, public health

Disciplines

Arts and Humanities | Life Sciences | Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

This is a preprint of an article originally published as Williams, PG, Health benefits of herbs and spices: Public Health, Medical Journal of Australia, 185(4), 2006, Supplement S17-S18. Copyright Australasian Medical Publishing Company 2006. Original journal available [here](#).

Consumer trends and public health guidelines for culinary herbs and spices

Assoc Prof Peter Williams PhD FDA

Consumption Trends

It is difficult to estimate the current level of consumption of culinary herbs and spices by Australians. The Australian Bureau of Statistics trend data on apparent consumption of foods do not include herbs and spices¹ and results from the last National Nutrition Survey (NNS) in 1995 provide only limited information on consumption. The median daily intake of herbs, spices, seasonings and stock cubes combined was estimated to be 1.4grams per adult (declining with age from 4.2g in 19-24year olds, to 0.7g in those 65years and older), with only 3.1 percent of persons reported consuming this category of food items on the day of survey². However the separate intakes of herbs and spices alone are not reported. The intake of males appeared to be higher than that of females, but the low values make it difficult to assess the significance of this difference. In New Zealand, the consumption of spices, estimated using import data, was 364g/year, or around 1g per day, a similar value to that reported in the NNS³.

One comparison of spices used in representative vegetable and meat-based recipes from 36 different countries found Australia (with a mean of 3.4 spices per recipe) had a moderate level of usage compared to the international mean of 3.9 (ranging from 1.6 in Norway to 6.9 in Indonesia)⁴. However, increased use of herbs and spices as flavourings in foods is a major trend worldwide⁵ with sales growth of 20-30% over the past five years in both the UK and the US⁶. It has been suggested that this trend is partly driven by demographics; as consumers age their palates can become more adventurous. Promotion can also be important: a recent UK advertisement in which Jamie Oliver encouraged consumers to experiment with nutmeg boosted sales of that spice four-fold⁷.

Based on retail sales data, consumption of herbs and spices in Australia has increased in line with global trends and this is expected to continue. The market for local fresh-cut culinary herbs was estimated to be worth over \$62 million per year in 2004 and continues to grow at 20% per annum⁸. Information from major supermarket sales in 2003 estimates suggest that total retail sales of fresh herbs and spices were valued at \$54 million, and a further \$107 million for dried products. The sales volumes of fresh herbs are shown in Table 1.

Dietary guidelines and recommendations

Although there is increasing interest and research in the health-promoting and protective properties of herbs and spices⁹⁻¹¹, there are few authoritative recommendations about intake in existing national dietary guidelines. The first of the Australian guidelines for adults is: Enjoy a wide variety of nutritious foods¹² and in the guidelines for older Australians a food variety checklist is given in an appendix, that includes the

recommendation to use herbs and spices regularly¹³. The same tool has also been used as the basis of a checklist to assess intakes of phytochemical-dense foods, and herbs and spices make up 11 out of the 64 foods scored in the checklist (basil, oregano, mint, dill/fennel, parsley, pepper, ginger, cumin, tumeric, coriander, rosemary/thyme)¹⁴. The higher the score the more adequate the diet is suggested to be in phytochemicals, however it is acknowledged that such food scores need to be further developed and are not backed by any health outcome studies at this stage.

Two other Australian dietary guidelines refer to herbs and spices. In the background chapter on choosing foods low in salt it is stated that among the recommended substitutes for salt are ingredients such as curry spices, garlic and onion, and herbs¹². Secondly, the Dietary Guidelines for Older Australians note particularly that age-related sensory loss of smell and taste is common in older people, especially in those who take many medications, and can have adverse effects on overall nutrient intake. Experimentation with new flavourings such as herbs and spices is suggested to stimulate appetite and support adequate overall intakes¹³.

A few other countries have made similar recommendations about herbs and spices. In the 2005 revision of the Dietary Guidelines for Americans, the chapter on choosing a diet moderate in salt and sodium recommends flavouring with herbs and spices¹⁵ and the Irish dietary guidelines have similar advice. Perhaps the country with the most direct recommendation about the health benefits for culinary herbs is Greece. Their dietary guidelines not only refer to the usefulness of herbs as salt substitutes but also state: “*oregano, basil, thyme and other herbs grown in Greece are good sources of antioxidant compounds*”¹⁶. This emphasis on the health-promoting properties of herbs is of interest given research in Australia that has found that first generation Greek migrants display 35% lower mortality from cardiovascular and overall mortality than Australian-born controls, despite high prevalence of risk factors such as obesity, smoking and sedentary lifestyles. It has been suggested that one of the dietary factors contributing to this lower mortality could be their high intake of antioxidant rich plant foods, including garlic and herbs¹⁷.

Despite the generally supportive statements in the dietary guidelines, the quantitative recommendations for intakes of food in the Australian Guide to Healthy Eating from the NH&MRC do not yet include suggested intakes of herbs and spices¹⁸. Those recommendations are based primarily on the aim of ensuring adequate intakes of nutrients for which recommended dietary intakes have already been established and it is probably too early for there to be more definitive recommendations about foods based on their content of other phytochemicals. It should also be borne in mind that there are possible adverse effects of some spices (such as chilli and peppers) if consumed in large quantities¹⁰, although this is unlikely to be a significant risk at normal levels of use. Thus the apparent increasing consumption of culinary herbs and spices is certainly a welcome trend that is worthy of closer monitoring, and in the future consideration should be given to inclusion of more explicit recommendations about their place in a healthy diet.

Table 1. Major supermarket sales of fresh herbs and spices in Australia 2003⁸

Herb	Kilograms	Retail sales (\$)
Garlic	1,767,000	7,941,100
Ginger	592,900	7,259,900
Chilli	425,000	3,755,300
Basil	74,200	7,373,200
Coriander	73,900	8,860,600
Parsley	30,500	8,415,000
Mint	18,800	2,151,500
Chives	18,400	2,225,500
Lemon grass	17,400	950,300
Rosemary	15,200	1,169,700
Oregano	9,700	594,700
Dill	6,100	1,074,900
Thyme	3,600	693,300
Other	845,800	9,104,000

References

1. Australian Bureau of Statistics. *Apparent consumption of foodstuffs and nutrients, Australia 1992-3. Cat No 4306.0*. Canberra: AGPS, 1995.
2. McLennan W and Podger A. *National Nutrition Survey. Food Eaten. Australia 1995. ABS Cat No 4804.0*. Canberra: Australian Bureau of Statistics, 1999.
3. Fowles J, Mitchell J, and McGrath H. Assessment of cancer risk from ethylene oxide residues in spices imported into New Zealand. *Food Chem Toxicol*. 2001;39:1055-1062.
4. Sherman P and Hash G. Why vegetable recipes are not very spicy. *Evol Hum Behav*. 2001;22:147-163.
5. Sloan A. Top 10 Global Food Trends. *Food Technol*. 2005;59:20-32.
6. Mintel International Group Ltd, *Seasoning - UK - August 2005*. Available at: www.the-infoshop.com/study/mt32490-seasonings.html.
7. Anon. Sector Insight: Seasonings - Added spice. *Marketing*. 2005;Nov 16:36.
8. Parker J. Culinary Herbs, in *The New Crop Industries Handbook. RIRDC Publication No 04/125*, S. Salvin, M. Bourke, and T. Byrne (eds). Canberra: Rural Industries Research and Development Corporation, 2004. Available at: <http://www.rirdc.gov.au/NewCrops/Contents.html>.
9. Craig W. Health-promoting properties of common herbs. *Am J Clin Nutr*. 1999;70:491S-499S.
10. Lampe J. Spicing up a vegetarian diet: chemoprotective effects of phytochemicals. *Am J Clin Nutr*. 2003;78:579S-583S.
11. Billing J and Sherman P. Antimicrobial functions of spices: why some like it hot. *Quart Rev Biol*. 1998;73:3-49.
12. National Health and Medical Research Council. *Food for Health: Dietary Guidelines for Australian Adults*. Canberra: NH&MRC, 2003.
13. National Health and Medical Research Council. *Dietary Guidelines for Older Australians*. Canberra: Australian Government Publishing Service, 1999.
14. Wahlqvist M, Wattanapenpaiboon N, Kannar D, Dalais F, and Kouris-Blazos A. Phytochemical deficiency disorders: inadequate intake of protective foods. *Curr Therap*. 1998;July:53-59.
15. US Department of Agriculture (USDA) and Department of Health and Human Services(HHS), *Dietary Guidelines for Americans 2005*. Available at: www.healthierus.gov/dietaryguidelines.
16. Ministry of Health and Welfare Supreme Scientific Health Council. Dietary guidelines for adults in Greece. *Arch Hellenic Med*. 1999;16:516-524.
17. Kouris-Blazos A. Morbidity mortality paradox of 1st generation Greek Australians. *Asia Pacific J Clin Nutr*. 2002;11(Suppl):S569-S575.
18. Smith A, Kellett E, and Schmerlaib Y. *The Australian Guide to Healthy Eating. Background information for nutrition educators*. Canberra: Commonwealth Department of Health, 1998.