



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Graduate School of Medicine - Papers (Archive)

Faculty of Science, Medicine and Health

2003

Rain, smog, fog and printed educational material

Judy Mullan

University of Wollongong, jmullan@uow.edu.au

Patrick A. Crookes

University of Wollongong, pcrookes@uow.edu.au

Heather Yeatman

University of Wollongong, hyeatman@uow.edu.au

Publication Details

Mullan, J. R., Crookes, P. A. & Yeatman, H. R. (2003). Rain, smog, fog and printed educational material. *Journal of Pharmacy Practice and Research*, 33 (4), 284-286.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Rain, smog, fog and printed educational material

Abstract

Patient education is central to the practice of all health professionals. Pharmacists often use printed educational material, such as consumer medicine information, as adjuncts to verbal communication and education of patients about their medications. A relatively large proportion of the Australian population have low literacy skills and cannot read or understand written patient educational materials which are typically written at levels equivalent to year 9 and above. By using simple readability assessment tools such as RAIN, SMOG or FOG and the UK's Department of Health 'Toolkit for producing patient information' health professionals can ensure that printed educational materials are simple and easily read by a wide patient population. These measures will ensure that pharmacists can become more effective educators to improve patients' medication knowledge and understanding, which will assist with optimising compliance and overall therapeutic outcomes.

Keywords

material, printed, smog, fog, educational, rain

Disciplines

Medicine and Health Sciences

Publication Details

Mullan, J. R., Crookes, P. A. & Yeatman, H. R. (2003). Rain, smog, fog and printed educational material. *Journal of Pharmacy Practice and Research*, 33 (4), 284-286.



87065801

RAIN, SMOG, FOG and Printed Educational Material

Judy R Mullan, Patrick A Crookes, Heather R Yeatman

ABSTRACT

Patient education is central to the practice of all health professionals. Pharmacists often use printed educational material, such as consumer medicine information, as adjuncts to verbal communication and education of patients about their medications. A relatively large proportion of the Australian population have low literacy skills and cannot read or understand written patient educational materials which are typically written at levels equivalent to year 9 and above. By using simple readability assessment tools such as RAIN, SMOG or FOG and the UK's Department of Health 'Toolkit for producing patient information' health professionals can ensure that printed educational materials are simple and easily read by a wide patient population. These measures will ensure that pharmacists can become more effective educators to improve patients' medication knowledge and understanding, which will assist with optimising compliance and overall therapeutic outcomes.

J Pharm Pract Res 2003; 33: 284-6.

INTRODUCTION

Chronic disease is the principal cause of disability, the major reason for seeking health care, and accounts for the majority of healthcare expenditure in Australia and overseas.^{1,3} Although the ageing population has contributed to these increases, the prevalence of chronic disease has risen in virtually every age group.⁴ For patients with chronic disease there is a growing interest in 'self-management' programs that emphasise the patients' central role in managing their illness. Several studies have concluded that low-cost programs involving health and medication education can help to improve health outcomes,⁵ reduce healthcare costs⁶ and reduce hospitalisation due to adverse drug events.⁷ Printed materials are an essential and economical part of this education.⁸ These materials must be easy to read and understand as patient outcomes may depend on their level of comprehension.⁵ Health professionals need to ensure that written patient educational materials, such as consumer medicines information (CMI), are simple and easily read by a wide patient population (people from non-English speaking backgrounds, low literacy and the elderly).

The aim of this paper is to raise awareness of the availability of simple tools, which can be used to ensure the delivery of easy to understand printed educational material. This, in turn, should help to optimise the quality, cost and outcomes of patient care.

DISCUSSION

Self-Management and Patient Education

Two of the principal components of self-management are

Judy R Mullan, BPharm, FSHP, PhD candidate, Graduate School of Public Health, University of Wollongong. *Patrick A Crookes*, Professor, Head of the Department of Nursing, University of Wollongong. *Heather R Yeatman*, Associate Professor, Graduate School of Public Health, University of Wollongong, New South Wales
Address for correspondence: Judy Mullan, 7-9 Elizabeth St, Mangerton NSW 2500, Australia
E-mail: jmullan@uow.edu.au

compliance with medication and adherence to self-treatment guidelines. Informed patients have a better understanding of their medication and can therefore make educated decisions about their treatment. They have been found to be more willing to accept or continue treatment despite minor or transitory adverse effects.⁹

In this era of healthcare reform with its emphasis on self-management and prevention, education of patients and their families has taken on an even greater role than previously.^{5,10} Patient education is any set of planned, educational activities designed to improve patients' health behaviours and/or health status. The purpose of patient education is to maintain or improve health or, in some cases, to slow deterioration. Patient education is central to the practice of all health professionals.

Printed Educational Materials

Pharmacists typically use verbal communication as part of their patient education intervention. CMI leaflets are often used as adjuncts to these verbal education sessions. This printed educational material is convenient, economical and very useful for providing drug information to patients and/or their carer(s).⁸ Unfortunately, in Australia most printed educational material and the CMI leaflets are pitched beyond the level of comprehension of most of the population including, but not only, people who have English as their second language.^{11,12}

Low Literacy in Australia

Low literacy is a pervasive and under-recognised problem in health care in Australia and overseas.^{11,13} A recent study has shown that patients with inadequate literacy skills have a much greater risk of being hospitalised than patients with adequate literacy skills.¹⁴ Low literacy groups include not only the poorly educated but also the elderly and immigrants who speak English as a second language.¹⁵ A 1996 survey of literacy found that 41 to 46% of people aged 65 to 74 years had very poor literacy skills and that of the 20% of the Australian population who are from non-English speaking backgrounds, 2.5% spoke little or no English.¹⁶ There is evidence to suggest that Australian migrants' need for information about their medications and pharmacy services are not being met.¹⁷

Given that a proportion of the Australian population have low literacy skills, it is logical that printed educational material should be pitched at no greater than the average reading ability of the Australian population (estimated at year 8 level).¹² Presently the reading level for printed educational material in Australia and overseas range from year 9 to 14.^{11,18-20}

Drug-Related Errors and Health Literacy

Drug-related errors are the major cause of adverse events, causing an estimated 10 to 20% of all medically related adverse events.²¹⁻²³ Australian research has shown that between 2.4 to 3.6% of all hospital admissions are drug-

related with an estimated cost of \$350 million per annum.²⁴⁻²⁶ A large proportion of these adverse events could be prevented by improving patient education and communication.²⁷ All health professionals, could help to reduce medically related adverse events by ensuring that all printed educational material and CMI leaflets are written at levels no greater than year 6 to 8, in order to increase the level of patient understanding.¹³

Assessment of Health Literacy

Assessing Literacy Levels

There are two obvious ways in which to deal with the problems associated with health literacy. The first involves carrying out simple tests to establish the patients' literacy level. Wide Range Achievement Test-Revised (WRAT-R),²⁸ Rapid Estimate of Adult Literacy in Medicine (REALM)²⁹ and Slosson Oral Reading Test-Revised (SORT-R)³⁰ are simple literacy tests which are quick and easy to administer. The REALM test, unlike the WRAT-R and the SORT-R tests, is healthcare specific thus making it appropriate for the assessment of health literacy. The REALM test involves the patient reading out aloud sixty-six words typically used in medical literature such as 'nutrition', and 'haemorrhoids'.

Alternatively, the Cloze test³¹ and the Test of Functional Health Literacy in Adults (TOFHLA)³² are much more comprehensive. TOFHLA takes approximately 22 minutes to administer, includes 50 comprehension and 17 numerical ability tests based on tasks often required of patients seeking health care, e.g. reading prescription bottles or appointment slips. Time constraints may make testing impractical especially since some of the health-related literacy tests, are time consuming to administer.³³

Readability Assessment of Written Information

The second way in which to deal with the problems associated with health literacy is to assess the readability levels of printed educational material. There are several well-known literacy tests such as SMOG,³⁴ FOG,³⁵ The Fry Readability Formula³⁶ and the Flesch Reading Ease Index,³⁷ which estimate the reading grade level a patient requires to understand the printed educational materials. These tests use sentence number and word length to estimate the reading (grade) level. Unfortunately, these readability tests are generic rather than health-specific making interpretation of the reading level and the understanding of printed educational materials difficult. The Readability Assessment Instrument (RAIN)³⁸ however, is a tool especially designed to assess health and mental health patient education materials. RAIN estimates readability and patient comprehension in terms of the following variables: structure, pronoun references, audience appropriateness, writing style, illustrations, print size, print style, colour and the highlighting of titles and subtitles.

DISCERN³⁹ is another tool, developed by researchers at the University of Oxford, UK. It can assist both patients and health professionals to quickly analyse the quality of the plethora of written health information available not only as printed educational leaflets or booklets but also on the Internet.

In addition to these instruments, there are several computer programs, which can be used to assess readability levels. Some of these include: The Flesch Reading Ease (FRE) formula, Grammatik for Windows, Right Writer, Que software and Readability Calculations.

Preparation of Printed Educational Materials

Many health professionals are involved with the preparation of printed educational material. When preparing or perhaps rewriting these documents it is important to consider both the literacy levels of the Australian population, and the readability levels of the written health information. An Australian study, investigated factors influencing patient satisfaction with, and the utilisation of cancer information booklets and found that patients preferred information to be written in plain English at a readability level of year 8 or lower.⁴⁰

In addition to vocabulary, sentence length and readability levels, other factors such as format and the use of colour illustrations may influence the patient's understanding and satisfaction with the printed educational materials.^{40,41} This will potentially influence the patient's use of the written information.³³ These factors have been incorporated into the 'Toolkit for producing patient information'⁴² developed by the UK's Department of Health. The toolkit is easy to use and ensures that the health practitioners' printed educational materials are of a high standard and understood by a wide population.

CONCLUSION

Patient education is a central part of the practice of all health professionals. Simple and easy-to-read printed educational materials and CMI leaflets are necessary and useful adjuncts to verbal education sessions with patients. The knowledge and information patients gain from these sessions can empower them to make informed decisions about their treatment, compliance and self-management guidelines. These informed decisions ultimately optimise therapeutic outcomes and minimise possible drug-related adverse events. Simple tests such as RAIN, SMOG and FOG as well as the UK's Department of Health 'Toolkit for producing patient information' can be used to ensure that printed educational materials and CMI leaflets are of a high standard and understood by a wide patient population including those from non-English speaking backgrounds, with low literacy, and the elderly.

Acknowledgment

This research is conducted in partial fulfilment of Judy Mullan's PhD of Public Health at the University of Wollongong entitled 'To develop and trial a new warfarin educational program which will improve warfarin compliance and understanding for a warfarin prescribed population'.

This paper is based on a presentation by Judy Mullan 'Rain, smog, fog it's not always about the weather' at the SHPA NSW Branch Conference, 26 October 2002.

Competing interests: None declared

Reference

1. Hoffman C, Rice D, Sung, H. Persons with chronic conditions: their prevalence and costs. *J Am Med Assoc* 1996; 276: 1473-9.
2. Holman H, Lorig K. Patients as partners in managing chronic disease. *BMJ* 2000; 320: 526-7.
3. Consumers' Health Forum of Australia. Healthy ageing discussion paper. Consumers' Health Forum of Australia; Canberra: 2000.
4. Rothenberg RB, Koplan JP. Chronic disease in the 1990s. *Annu Rev Public Health* 1990; 11: 267-96.
5. Rutledge DN, Donaldson NE. Improving readability of print materials in patient care and health services. *Cinahl Information Systems. The Online Journal of Clinical Innovations* 1998; 1: 1-27.
6. Lorig K, Ritter P, Stewart A, Sobel D, Brown B, Bandura A, et al. Chronic disease self-management program: 2-year health status and health care utilization outcomes. *Medical Care* 2001; 39: 1217-23.
7. Wilson RM, Runciman WB, Gibberd RW, Harrison BT, Newby L, Hamilton JD. The Quality in Australian Health Care Study. *Med J Aust* 1995; 163: 458-71.
8. Bernier MJ, Yasko J. Designing and evaluating printed education materials: model and instrument development. *Patient Educ Couns* 1991; 18: 253-63.
9. Graham JD. What consumers want to know about medicines. *Aust Prescr* 1995; 18: 10-11.

10. Jaarsma T, Halfens R, Abu-Saad H, Dracup K, Diederiks J. Self-care and quality of life in patients with advanced heart failure: The effect of a supportive educational intervention. *Heart Lung* 2000; 29: 319-30.
11. Baker SJ. Who can read consumer product information? *Aust J Hosp Pharm* 1997; 27: 126-31.
12. Buchbinder R, Hall S, Grant G, Mylvaganam A, Patrick M. Readability and content of supplementary written drug information for patients used by Australian rheumatologists. *Med J Aust* 2001; 174: 575-8.
13. Mayeaux Jr E, Murphy P, Arnold C, Davis T, Jackson R, Sentell T. Improving patient education for patients with low literacy skills. *Am Fam Physician* 1996; 53: 205-12.
14. Baker D, Parker R, Williams M, Clark W. Health literacy and the risk of hospital admission. *J Gen Intern Med* 1998; 13: 791-8.
15. Doak C, Doak L, Root J. Teaching patients with low literacy skills. 2nd ed. Philadelphia: Lippincott, Williams and Wilkins; 1996.
16. Australian Bureau of Statistics. Aspects of literacy. Canberra: Australian Bureau of Statistics; 1996.
17. Bajramovic J, Tett S. Problems of pharmacy communication in multicultural Australia. *Aust Pharmacist* 2000; 19: 430-4.
18. Rolland PD. Reading levels of drug information printouts: A barrier to effective communication of patient medication information. *Drug Info J* 2000; 34: 1329-38.
19. Davis T, Crouch M, Wills G, Miller S, Abdehou D. The gap between patient reading comprehension and the readability of patient education materials. *J Fam Pract* 1990; 31: 533-8.
20. Estrada CA, Hryniewicz M, Barnes-Higgs V, Collins C, Byrd J. Anticoagulant patient information material is written at high readability levels. *Stroke* 2000; 31: 2966-71.
21. Leape L, Kabaceni A, Berwick D, Roessner J. Reducing adverse drug events. Boston: Institute of Healthcare Improvement; 1998.
22. Bates DW. Frequency, consequences and prevention of adverse drug events. *J Qual Clin Pract* 1999; 19: 13-7.
23. Dartnell J, Anderson R, Chohan V, Galbraith K, Lyon M, Nestor P, et al. Hospitalization for adverse effects related to drug therapy: incidence, avoidability and costs. *Med J Aust* 1996; 164: 659-62.
24. Roughhead EE. The nature and extent of drug-related hospitalisations in Australia. *J Qual Clin Pract* 1999; 19: 19-22.
25. Roughhead EE, Gilbert AL, Primrose JG, Sansom LN. Drug-related hospital admissions: a review of Australian studies published 1988-1996. *Med J Aust* 1998; 168: 405-8.
26. Australian Council for Safety and Quality in Health Care. Second National Report on Patient Safety. Improving medication safety. Canberra: Safety and Quality Council; 2002. Available from: <www.safetyandquality.org/articles/Publications/med_saf_rept.pdf>.
27. Wilson R, Runciman W, Gibberd R, Harrison B, Newby L, Hamilton J. The quality in Australian health care study. *Med J Aust* 1995; 163: 458-71.
28. Jastak S, Wilkinson G. The Wide Range Achievement Test Revised: administration manual (WRAT-3), 1993.
29. Murphy PW, Davis TC, Long SW, Jackson TH, Decker BC. Rapid estimate of adult literacy in medicine (REALM): a quick testing for patients. *J Reading* 1993; 37: 124-30.
30. Slosson RL. Slosson oral reading test (SORT) revised. East Aurora: Slosson Educational Publications; 1990.
31. Taylor W. Cloze procedure: a new tool for measuring readability. *Journalism Quarterly* 1953; 30: 415-33.
32. Nurrs J, Parker R, Williams M, Baker D. Test of functional health literacy in adults. Centre for the study of adult literacy. Atlanta: Georgia State University; 1995.
33. Koo M, Krass I, Aslani P. Factors influencing consumer use of written drug information. *Ann Pharmacother* 2003; 37: 259-67.
34. McLaughlin GH. SMOG grading: a new readability formula. *J Reading* 1969; 12: 639-46.
35. Gunning R. The technique for clear writing. New York: McGraw-Hill; 1952.
36. Fry EB. A readability formula that saves time. *J Reading* 1968; 11: 513-6.
37. Flesch RF. A new readability yardstick. *J Appl Psychol* 1948; 32: 221-3.
38. Singh J. RAIN: Readability assessment instrument manual. Midlothian: Woodlake Institute for Human Services; 1994.
39. Charnock D, Shepperd S, Needham G, Gann R. DISCERN: an instrument for judging the quality of written consumer health information on treatment choices. *J Epidemiol Community Health* 1999; 53: 105-11.
40. Butow P, Brindle E, McConnell D, Boakes R, Tattersall M. Information booklets about cancer: factors influencing patient satisfaction and utilization. *Patient Educ Couns* 1998; 33: 129-41.
41. Redman KB. The practice of patient education. 9th ed. Detroit: Mosby Inc; 2001.
42. The United Kingdom Department of Health. Toolkit for producing patient information. NHS, United Kingdom; 2002. Available from: <www.doh.gov.uk/nhsidentity/toolkit/toolkit-patientinfo.htm>.

Submitted: January 2003
Accepted after external peer review: May 2003

Copyright of Full Text rests with the original copyright owner and, except as permitted under the Copyright Act 1968, copying this copyright material is prohibited without the permission of the owner or its exclusive licensee or agent or by way of a licence from Copyright Agency Limited. For information about such licences contact Copyright Agency Limited on (02) 93947600 (ph) or (02) 93947601 (fax).

Your work was in the public domain at the 26th Federal Conference in your country!!!

Prizes will be awarded to authors of papers submitted to JPPR from Conference contributed papers.

Each prize will be \$2500 assistance for attendance at an SHPA Clinical Conference or Federal Conference.

Articles must be submitted to JPPR by June 30 2004, requesting consideration for the prize. Please refer to this issue or <www.shpa.org.au/journal/austra.htm> for information for authors. Judging will be undertaken by the Editorial Committee of JPPR.

All of us at JPPR would like to wish you and yours a joyful festive season and a happy New Year.