

2007

Aryl Nitro Reduction with Iron Powder or Stannous Chloride under Ultrasonic Irradiation

A. B. Gamble
University of Wollongong

James A. Garner
University of Wollongong, jgarner@uow.edu.au

Christopher Gordon
cpg02@uow.edu.au

S. M. J. O'Conner
University of Wollongong

Paul A. Keller
University of Wollongong, keller@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/scipapers>

 Part of the [Life Sciences Commons](#), [Organic Chemistry Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Gamble, A. B.; Garner, James A.; Gordon, Christopher; O'Conner, S. M. J.; and Keller, Paul A.: Aryl Nitro Reduction with Iron Powder or Stannous Chloride under Ultrasonic Irradiation 2007.
<https://ro.uow.edu.au/scipapers/148>

Aryl Nitro Reduction with Iron Powder or Stannous Chloride under Ultrasonic Irradiation

Abstract

The selective reduction of aryl nitro compounds in the presence of sensitive functionalities, including halide, carbonyl, nitrile and ester substituents under ultrasonic irradiation at 35 kHz is reported in yields of 39-98%. Iron powder proved superior to stannous chloride with high tolerance of sensitive functional groups and high yields of the desired aryl amines in relatively short reaction times. Simple experimental procedure and purification also make the iron reduction of aryl nitro compounds advantageous over other methods of reduction.

Keywords

Nitro reduction, Iron powder, Ultrasound, CMMB

Disciplines

Life Sciences | Organic Chemistry | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

This article was originally published as Gamble, AB, Garner, J, Gordon, CP, O'Conner, SMJ & Keller, PA, Aryl Nitro Reduction with Iron Powder or Stannous Chloride under Ultrasonic Irradiation, Synthetic Communications, 37, 2007, 2777-2786. Original journal article available [here](#)

Aryl Nitro Reduction with Iron Powder or Stannous Chloride under Ultrasonic Irradiation

Allan B. Gamble, James Garner, Christopher P. Gordon, Sean M. J. O'Conner, Paul A. Keller *

Department of Chemistry, University of Wollongong, Wollongong 2522, Australia

Abstract: The selective reduction of aryl nitro compounds in the presence of sensitive functionalities, including halide, carbonyl, nitrile and ester substituents under ultrasonic irradiation at 35 kHz is reported in yields of 39-98%. Iron powder proved superior to stannous chloride with high tolerance of sensitive functional groups and high yields of the desired aryl amines in relatively short reaction times. Simple experimental procedure and purification also make the iron reduction of aryl nitro compounds advantageous over other methods of reduction.

Keywords: Nitro reduction, Iron powder, Ultrasound


Aryl amines are synthetically important compounds which act as precursors to the synthesis of many interesting molecules, and can be readily synthesized from aryl nitro compounds *via* countless reduction methods. The most general methods involve activated metal catalysis^[1] and transition metal catalyzed hydrogenation,^[2] although the latter often employs harsh reaction conditions affecting other reduction sensitive functionalities such as halides, ketones, aldehydes, esters and nitriles in addition to the nitro substituent^[2]. The selective reduction of aryl nitro compounds using iron powder and dilute acid^[3] or stannous chloride^[4] have been reported as efficient methods for the synthesis of aryl amines in good yields. However notable disadvantages to these methods include high reaction temperatures, relatively long reaction times, potential halogenation reactions, and the incompatibility of acid sensitive functional groups associated with the use of a strong acid such as HCl. The use of milder conditions, for example acetic acid as solvent, has been reported but high temperatures and longer reaction times are still evident^[5].

Ultrasound is reported as an alternative energy source for the initiation of organic reactions,^[6] with the potential to accelerate chemical transformations, affect product distributions, improve yields and increase the catalytic activity of metal particles by factors as high as 10^5 .^[7] Sonication has been employed in many different types of reactions, in particular transition metal catalyzed reduction. Generally reactions involving metal reagents exhibit sluggish reaction times due to the presence of surface impurities, but through continuous cleaning and chemical activation of the metal surface, and the high temperature and pressure produced by acoustic cavitation which potentially results in the formation of hot spots,^[8a] an accelerated reaction rate is often associated with ultrasound.^[7]

* Corresponding author. Tel.: +61 2 4221 4692; fax: +61 2 4221 4287; e-mail: keller@uow.edu.au.

There has been a limited number of aryl nitro reduction procedures *via* sonication reported in the literature,^[8] although a recent study utilizing stannous chloride in the presence of ionic liquids provided aryl amines in good yields.^[8a] Other procedures investigated the reduction of nitrobenzene using elemental iron,^[8b] and reduction of some simple aryl nitro compounds using iron in the presence of multiple additional reagents,^[8c] but the tolerance of sensitive functional groups for iron-catalyzed nitro reduction promoted by ultrasound was not examined. The lack of iron and stannous chloride-catalyzed nitro reduction promoted by ultrasound in the literature prompted the search for more efficient, selective and relatively straightforward procedures.


The reduction of aryl nitro compounds is known to proceed *via* the hydroxylamine, followed by azoxy and azo compounds to its corresponding aryl amine after a prolonged reaction time (Scheme 1).^[9] Therefore, the ability of ultrasonic irradiation to accelerate heterogeneous chemical reactions could potentially reduce the reaction time and reduce the amount of intermediates isolated, increasing the yield of aryl amine. To determine the optimal reaction conditions for the reduction of aryl nitro compounds to their corresponding aryl amines under ultrasound conditions without isolation of the intermediates, a series of model reactions were performed (Table 1).



Scheme 1.

From these reactions the optimal conditions were 5 equivalents of iron powder in a mixture of ethanol, glacial acetic acid and water (2:2:1) exposed to ultrasound for 1 hour (Entry 3). The reaction was repeated under thermal conditions (2.5 h at 60 °C) with the yield of aryl amine **2** determined to be 85%, similar to the optimal conditions in entry 3, but the temperature and time of reaction required for the complete conversion of **1** and its intermediates to **2** were notably higher and longer. This demonstrates that the high energy effects of acoustic cavitation in addition to the continuous cleaning of the iron surface due to ultrasound are responsible for the enhanced reaction rate. The latter probably accounts for the greatest enhancement in reaction rate since under thermal heating, surface impurities such as oxides, hydroxides and carbonates inhibit contact between the aryl nitro and the iron surface.^[7] In addition, cleansing of the iron sweeps reactive intermediates or products from the surface making way for subsequent reactions.^[8b]

Table 1. Investigation of ultrasound promoted reduction of 3-nitroanisole **1**



Entry	Reducing Agent	Solvent	Time (h)	Yield (%) ^a
1	$\text{Fe}_{(s)}$ (1 equiv.)	ethanol: glacial acetic acid: water	2	36
2	$\text{Fe}_{(s)}$ (2 equiv.)	ethanol: glacial acetic acid: water	2	59
3	$\text{Fe}_{(s)}$ (5 equiv.)	ethanol: glacial acetic acid: water	1	89
4	$\text{Fe}_{(s)}$ (5 equiv.)	ethanol: water	2	0
5	$\text{Fe}_{(s)}$ (5 equiv.)	glacial acetic acid: water	1.5	75
6	$\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (1 equiv.)	ethanol	2	26
7	$\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (5 equiv.)	ethanol	2	45
8	$\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (10 equiv.)	ethanol	2	76

^aIsolated yield















Entries 1 and 2 demonstrate that the yield of aryl amine **2** decreases significantly as the molar equivalents of iron powder are reduced, probably due to the presence of the intermediates not converted to the aryl amine (scheme 1), and the yield of **2** was slightly reduced in the absence of ethanol (Entry 5). When glacial acetic acid, which presumably helps to activate the iron powder, is not used in the reaction, no reduction of the aryl nitro was observed (Entry 4).

The ability of stannous chloride to selectively reduce an aryl nitro substituent to an aryl amine was also investigated using ethanol as the solvent. Entry 8 demonstrates that for good conversion to the aryl amine, 10 molar equivalents of stannous chloride are required. In the presence of only one equivalent (Entry 6) and 5 equivalents (Entry 7) of stannous chloride the yields of aryl amine were significantly lower.

The optimal reduction conditions (Entry 3) were applied to further nitro aromatic derivatives, examining the selectivity of the reduction for nitro substituents in the presence of other sensitive functionalities (Table 2). Entries 1 to 3 allow comparison between iron and stannous chloride

catalyzed reduction, but due to the superiority of iron reduction, entries 4-8 were only performed *via* this method.

Table 2. Reduction of aryl nitro compounds with reduction sensitive functional groups

Entry	Reagent	Product ^b	Fe _(s) Rxn Time	Yield using Fe _(s) (%) ^a	SnCl ₂ .2H ₂ O Rxn Time	Yield using SnCl ₂ .2H ₂ O (%) ^a
1			1 h	77	2 h	58
2			1 h	85	2 h	54
3			2 h	65	2 h	35
4			1 h	86	-	-
5			1 h	98 ^c	-	-
6			1 h	85	-	-
7			1 h	82	-	-
8			15 min	78	-	-

^aIsolated yield, 100% conversion, monitored by TLC for completion of the reaction

^bProducts characterised by ¹H, ¹³C and MS and known compounds compared to the literature

^c12 equivalents of iron used

Entries 1 and 2 demonstrate the tolerance of iron reduction for ketone functionalities which under catalytic hydrogenation conditions could potentially be reduced to their corresponding alcohols and

methylene groups.^[10] The tolerance of ketone functionalities to iron-catalyzed nitro reduction is well precedented in the literature^[1a,11] and with yields of 77% and 85% for *p*-aminoacetophenone and *m*-aminoacetophenone under ultrasonic irradiation respectively, our conditions also allow complete selectivity for the nitro substituent over the ketone moiety. Both of these compounds have been reported in the literature with yields of 92%^[1a] and 80%^[12] for the *para*- and *meta*- substituted compounds respectively, and although the former yield is higher than our reported yield, the reaction was done under high pressure at a temperature of 210 °C for 2 h with water as solvent. Therefore the iron-catalyzed reduction conditions promoted by ultrasound at 30 °C for 1 h are advantageous as they are significantly milder, safer and easier to perform.

Aryl halides are known to be susceptible to dehalogenation under harsh reduction conditions, in particular catalytic hydrogenation.^[2] Entries 3-7 examine the selective reduction of the aryl nitro substituent over the aryl halide, with good to excellent yields obtained in all reactions. Entries 3, 4, 5 and 6 demonstrate the tolerance of bromine to the optimized iron reduction conditions and with the exception of entry 3 the yields of desired aryl amine are excellent (86%, 98% and 85% for entries 4, 5 and 6 respectively). Although TLC analysis showed 100% conversion of 2-bromo-4-methoxy-6-nitrophenol to its corresponding aryl amine (Entry 3) with no other intermediates identified, it could only be isolated in 65% yield. During work-up, difficulties in isolation of the desired aryl amine were encountered with the product partitioning between both the organic and aqueous phase, therefore contributing to the lower than expected yield. Entry 5 demonstrates the ability of iron-catalyzed reduction under ultrasound to selectively reduce two nitro substituents of a symmetrical dimer with an almost quantitative yield of 98%. The same compound in entry 5 has reportedly been synthesized *via* iron-catalyzed reduction, in the presence of ferric chloride and concentrated HCl at reflux in ethanol, although could only be isolated in 69% yield,^[13] indicating the superiority of iron-catalyzed reduction promoted by ultrasound. Entry 7 demonstrates the tolerance of chlorine substituents on a pyrimidine ring with a yield of 82% obtained, with previously reported iron-catalyzed reduction reporting a yield of 64%.^[14]

Finally, entry 8 was used to examine the tolerance of both an ester and nitrile functionality on an aliphatic chain. The reaction was complete in just 15 mins as indicated by TLC analysis and provided the desired aryl amine in 78% yield, demonstrating the ability of ultrasound to drastically accelerate the iron-catalyzed reduction of an aryl nitro functionality in the presence of these reduction sensitive groups.

In conclusion, an effective and efficient method for the reduction of aryl nitro compounds to their corresponding aryl amine under ultrasonic irradiation has been reported. Although in some instances, marginally higher yields are reported, the reaction of aryl nitro compounds with iron powder in a solvent mixture of ethanol, acetic acid and water promoted by ultrasonic irradiation provides a much more accessible and simpler procedure. Additionally, the short reaction times at relatively low temperature, the use of environmentally benign solvents and cheap reagents make this an attractive and advantageous method for reduction of aryl nitro compounds in organic synthesis.

Experimental

Reagents and solvents used in the experiments were purchased reagent grade and used without further purification. Sonication was performed in an Elma Transsonic T460 ultrasonic cleaning bath (at a frequency of 35 kHz and a nominal power of 85 W) with all reactions exposed to air in standard glassware or glass sample vials with the temperature of the bath maintained at 30 °C. Melting points were determined in a Gallenkamp (Griffin) melting point apparatus. ¹H NMR spectra were acquired on a Varian Unity-300 or Unity-500 spectrometer at 300 MHz and 500 MHz respectively. Spectra were recorded in deuterated chloroform (CDCl₃), using chloroform (δ 7.26 ppm) as internal standard. ¹³C NMR spectra were acquired with a Varian Unity-300 or Unity-500 spectrometer at 76 MHz and 126 MHz respectively using CDCl₃ as solvent and chloroform (δ 77.16 ppm) as the internal standard. Electron Impact (EI) mass spectra were obtained on a Shimadzu QP-5000 MAT-44 quadrupole spectrometer performed *via* a direct insertion technique, with an electron beam of 70 eV and a source temperature <200 °C and electrospray ionization (ESI) mass spectra were obtained on a VG Quattro-triple quadrupole. All data reported for known compounds were spectroscopically identical to that reported in the literature (see compound references).

General Procedure for Nitro Reduction with Iron

To a suspension of **1** (0.147 g, 0.961 mmol) in a mixture of glacial acetic acid (2 mL), ethanol (2 mL) and water (1 mL) was added reduced iron powder (0.279 g, 5.00 mmol). The resulting suspension was exposed to ultrasonic irradiation for 1 h at 30 °C with TLC analysis monitoring for the completion of the reaction. The reaction mixture was filtered to remove the iron residue which was washed with ethyl acetate (30 mL). The filtrate was partitioned with 2M KOH and the basic layer was further extracted with ethyl acetate (3 x 25 mL). The combined organic extracts were washed with brine (2 x 25 mL) and water (3 x 50 mL), dried (MgSO₄) and concentrated under reduced pressure. The crude residue was then subjected to flash silica gel column chromatography (20% ethyl acetate in hexanes) yielding 3-methoxyaniline **2** (89%).

General Procedure for Nitro Reduction with Stannous Chloride

To a solution of **1** (0.148 g, 0.967 mmol) in ethanol (5 mL) was added SnCl₂·2H₂O (2.26 g, 10.0 mmol). The reaction mixture was exposed to ultrasonic irradiation for 2 h at 30 °C until the reaction was complete as indicated by TLC analysis. The solvent was removed under reduced pressure and the crude residue was partitioned between ethyl acetate and 2M KOH. The aqueous layer was extracted with further portions of ethyl acetate (3 x 25 mL) and the combined organic extracts were washed with brine (2 x 25 mL) and water (3 x 50 mL), dried (MgSO₄) and concentrated under reduced pressure. The crude residue was subjected to flash silica gel column chromatography (20% ethyl acetate in hexanes) yielding 3-methoxyaniline **2** (76%).

3-Methoxyaniline (Table 1)^[15]

Dark yellow oil. ¹H NMR (CDCl₃): δ 7.07 (dd, *J* = 8.1 Hz, 7.8 Hz, 1H); 6.33 (dd, *J* = 8.1 Hz, 2.4 Hz, 1H); 6.30 (dd, *J* = 7.8 Hz, 1.8 Hz, 1H); 6.25 (dd, *J* = 2.4 Hz, 2.4 Hz, 1H); 3.77 (s, 3H); 3.66 (bs, 2H). ¹³C NMR (CDCl₃): δ 160.8, 147.9, 130.2, 108.0, 104.0, 101.1, 55.1. MS (EI) *m/z* 123 (M, 100%).

p-Aminoacetophenone (Table 2, Entry 1)^[16]

Opaque crystalline solid. Mp: 102-103 °C (lit.^[16] 106-107 °C) ¹H NMR (CDCl₃): δ 7.79 (d, *J* = 8.7 Hz, 2H); 6.63 (d, *J* = 9.3 Hz, 2H); 4.20 (bs, 2H); 2.49 (s, 3H). ¹³C NMR (CDCl₃): δ 196.7, 151.3, 130.9, 127.9, 113.8, 26.2. MS (EI) *m/z* 135 (M, 60%), 120 (M-15, 100%).

m-Aminoacetophenone (Table 2, Entry 2)^[12]

Light bronze solid. Mp: 88-89 °C (lit.^[12] 98-99 °C). ¹H NMR (CDCl₃): δ 7.31 (ddd, *J* = 7.5 Hz, 1.8 Hz, 1.2 Hz, 1H); 7.25 (dd, *J* = 1.8 Hz, 1.2 Hz, 1H); 7.21 (dd, *J* = 8.1 Hz, 7.5 Hz, 1H); 6.85 (ddd, *J* = 8.1 Hz, 2.1 Hz, 1.2 Hz, 1H); 3.85 (bs, 2H); 2.54 (s, 3H). ¹³C NMR (CDCl₃): δ 198.6, 146.9, 138.3, 129.5, 119.7, 118.9, 114.1, 26.8. MS (EI) *m/z* 135 (M, 85%), 120 (M-15, 100%).

2-Amino-6-bromo-4-methoxyphenol (Table 2, Entry 3)^[17]

Dark brown semi-solid. ¹H NMR (CDCl₃): δ 6.41 (d, *J* = 2.7 Hz, 1H); 6.26 (d, *J* = 2.7 Hz, 1H); 3.70 (s, 3H). ¹³C NMR (CDCl₃): δ 154.3, 136.2, 134.8, 109.8, 105.3, 102.1, 55.9. MS (EI) *m/z* 217 (M⁷⁹Br, 100%), 219 (M⁸¹Br, 90%).

3-Bromo-2,5-dimethoxyaniline (Table 2, Entry 4)

Brown viscous oil. ^1H NMR (CDCl_3 , 500 MHz): δ 6.45 (d, $J = 3.0$ Hz, 1H); 6.23 (d, $J = 2.5$ Hz, 1H); 3.93 (bs, 2H); 3.78 (s, 3H); 3.71 (s, 3H). ^{13}C NMR (CDCl_3 , 126 MHz): δ 156.9, 141.7, 138.7, 117.1, 107.1, 101.3, 59.9, 55.7. MS (EI) m/z 231 (M^{79}Br , 100%), m/z 233 (M^{81}Br , 90%). HRMS (EI) calculated for $\text{C}_8\text{H}_{10}\text{NO}_2\text{Br}$: 230.9895, found 230.9888.

3,3'-Diamino-5,5'-dibromo-2,2'-dimethoxy-1,1'-biphenyl (Table 2, Entry 5)^[13]

Light brown/red crystalline solid. Mp: 172-175 °C (lit.^[13] 185-186 °C). ^1H NMR (CDCl_3 , 300 MHz): δ 6.89 (d, $J = 2.4$ Hz, 2H); 6.80 (d, $J = 2.1$ Hz, 2H); 3.96 (bs, 4H); 3.44 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz): δ 144.0, 141.5, 132.3, 122.8, 118.1, 116.8, 60.1. MS (EI) m/z 400 ($\text{M}^{79}\text{Br}^{79}\text{Br}$, 50%), 402 ($\text{M}^{81}\text{Br}^{79}\text{Br}$, 100%) 404 ($\text{M}^{81}\text{Br}^{81}\text{Br}$, 50%). HRMS (EI) calculated for $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2^{79}\text{Br}^{79}\text{Br}$: 399.9422, found 399.9419.

2-Bromo-3-methylaniline (Table 2, Entry 6)^[18]

Yellow oil. ^1H NMR (CDCl_3): δ 7.00 (dd, $J = 7.8$ Hz, 7.5 Hz, 1H); 6.64 (m, 2H); 3.95 (bs, 2H); 2.39 (s, 3H). ^{13}C NMR (CDCl_3): δ 144.4, 138.8, 127.5, 120.38, 113.2, 112.3, 23.7. MS (EI) m/z 238 (M^{79}Br , 100%), 240 (M^{81}Br , 90%).

5-Amino-4,6-dichloropyrimidine (Table 2, Entry 7)^[14]

White solid. ^1H NMR (CDCl_3): δ 8.21 (s, 1H); 4.50 (bs, 2H). ^{13}C NMR (CDCl_3) δ 146.0, 144.3, 136.0. MS (ES+) m/z 164 ($[\text{MH}^{35}\text{C}^{35}\text{Cl}]^+$, 100%) 166 ($[\text{MH}^{37}\text{C}^{35}\text{Cl}]^+$, 60%), 168 ($[\text{MH}^{37}\text{C}^{37}\text{Cl}]^+$, 10%).

4-Amino-*N*-(1'-cyanoethyl)-*N*-(acetoethyl)aniline (Table 2, Entry 8)

Light brown oil. ^1H NMR (CDCl_3 , 500 MHz): δ 6.66 (d, $J = 8.9$ Hz, 2H); 6.61 (d, $J = 8.9$ Hz, 2H); 4.12, (t, $J = 5.6$ Hz, 2H); 3.66 (s, 2H); 3.49 (t, $J = 6.1$ Hz, 2H); 3.42 (t, $J = 6.1$ Hz, 2H); 2.46 (t, $J = 6.8$ Hz, 2H); 2.00 (s, 3H). ^{13}C NMR (CDCl_3 , 126 MHz): δ 171.2, 140.2, 139.5, 118.9, 118.2, 116.9, 62.2, 51.9, 49.2, 21.1, 16.5. MS (ES+) m/z 248 ($[\text{MH}]^+$, 100%). HRMS (ES+) calculated for $\text{C}_{13}\text{H}_{18}\text{N}_3\text{O}_2$: 248.1399, found 248.1405.

ACKNOWLEDGMENTS

ABG would like to thank the Australian Government for an Australian Postgraduate Award, AVEXA Limited for financial support and the Institute of Biomolecular Science, UOW for financial support.

REFERENCES

- (a) Wang, L.; Li, P.; Wu, Z.; Yan, J.; Wang, M.; Ding, Y. Reduction of nitroarenes to aromatic amines with nanosized activated metallic iron powder in water. *Synthesis* **2003**, 13, 2001-2004; (b) Herepoulos, G. A.; Georgakopoulos, S.; Steele, B. R. High intensity ultrasound-assisted reduction of sterically demanding nitroaromatics. *Tetrahedron Lett.* **2005**, 46, 2469-2473; (c) Nagaraja, D.; Pasha, M. A. Reduction of aryl nitro compounds with aluminium/NH₄Cl: Effect of ultrasound on the rate of the reaction. *Tetrahedron Lett.* **1999**, 40, 7855-7856; (d) Pyo, S. H.; Han, B. H. Reduction of nitroarenes by activated metals. *Bull. Korean Chem. Soc.* **1995**, 16, 181-183.
- (a) Nomura, K. Transition metal catalyzed hydrogenation or reduction in water. *J. Mol. Catal.* **1998**, 130, 1-28; (b) Downing, R. S.; Kunkeler, P. J.; van Bekkum, H. Catalytic syntheses of aromatic amines. *Catal. Today* **1997**, 37, 121-136; (c) Boix, C.; Poliakov, M. Selective reductions of nitroarenes to anilines using metallic zinc in near-critical water. *J. Chem. Soc. Perkin Trans. 1* **1999**, 1487-1490.
- (a) Liu, Y.; Lu, Y.; Prasad, M.; Repic, O.; Blacklock, T. J. A practical and chemoselective reduction of nitroarenes to anilines using activated iron. *Adv. Synth. Catal.* **2005**, 347, 217-219; (b) Hazlet, S. E.; Dornfeld, C. A. The reduction of aromatic nitro compounds with activated iron. *J. Am. Chem. Soc.* **1944**, 66, 1781-1782.
- Bellamy, F. D.; Ou, K. Selective reduction of aromatic nitro compounds with stannous chloride in non acidic and non aqueous medium. *Tetrahedron Lett.* **1984**, 25, 839-842.
- (a) Guay, V.; Brassard, P. The regiospecific synthesis of the A and B rings of phomazarin. *J. Heterocyclic Chem.* **1987**, 24, 1649-1652; (b) Owsley, D. C.; Bloomfield, J. J. The reduction of nitroarenes with iron/acetic acid. *Synthesis* **1977**, 118-120.
- Nuchter, M.; Ondruschka, B.; Jungnickel, A.; Muller, U. Organic processes initiated by non-classical energy sources. *J. Phys. Org. Chem.* **2000**, 13, 579-586.
- Luche, J. L. *Synthetic Organic Sonochemistry*; Plenum Press: New York, 1998.
- (a) Rai, G.; Jeong, J. M.; Lee, Y. S.; Kim, H. W.; Lee, D. S.; Chung, J. K.; Lee, M. C. Ionic liquid mediated efficient reduction of nitroarenes using stannous chloride under sonication. *Tetrahedron Lett.* **2005**, 46, 3987-3990; (b) Hung, H. M.; Ling, F. H.; Hoffmann, M. R. Kinetics and mechanism of the enhanced reductive degradation of nitrobenzene by elemental iron in the presence of ultrasound. *Environ. Sci. Technol.* **2000**, 34, 1758-1763; (c) Han, B. H.; Shin, D. H.; Cho, S. Y. Organic sonochemistry. Ultrasonic acceleration of the reduction of aromatic nitro compounds with hydrazine-iron in the presence of activated carbon. *Bull. Korean Chem. Soc.* **1985**, 6, 320.

9. Pasha, M. A.; Jayashankara, V. P. Reduction of aryl nitro compounds to azoarenes and/or arylamines by Al/NaOH in methanol under ultrasonic conditions. *Ultrason. Sonochem.* **2005**, *12*, 433-435.
10. Blaser, H-U.; Malan, C.; Pugin, B.; Spindler, F.; Steiner, H.; Studer, M. Selective hydrogenation for fine chemicals: Recent trends and new developments. *Adv. Synth. Catal.* **2003**, *345*, 103-151.
11. Radziejewski, C.; Ghosh, S.; Kaiser, E. T. The synthesis of two strongly electron deficient flavin analogs. *Heterocycles* **1987**, *26*, 1227-1238.
12. Simpson, J. C. E.; Atkinson, C. M.; Schofield, K.; Stephenson, O. o-Amino-ketones of the acetophenone and benzophenone types. *J. Chem. Soc.* **1945**, 646-657.
13. Viswanathan, N.; Patankar, S. J. Synthesis of 5,5'-dibromo-8,8'-dihydroxy-7,7'-biquinoline. *Indian J. Chem.* **1985**, *24B*, 1284-1285.
14. Baidur, N.; Chadha, N.; Player, M. R. Solution-phase synthesis of a library of 3,5,7-trisubstituted 3H-[1,2,3]triazole[4,5-d]pyrimidines. *J. Comb. Chem.* **2003**, *5*, 653-659.
15. (a) Lee, D-Y.; Hartwig, J. F. Zinc trimethylsilylamide as a mild ammonia equivalent and base for the amination of aryl halides and triflates. *Org. Lett.* **2005**, *7*, 1169-1172. (b) Wolter, M.; Nordmann, G.; Job, G. E.; Buchwald, S. L. Copper-catalysed coupling of aryl iodides with aliphatic alcohols. *Org. Lett.* **2002**, *4*, 973-976.
16. Pal, M.; Parasuraman, K.; Yeleswarapu, K. R. Palladium-catalysed cleavage of O/N-propargyl protecting groups in aqueous media under copper-free condition. *Org. Lett.* **2003**, *5*, 349-452.
17. Malamas, M. S.; Manas, E. S.; McDevitt, R. E.; Gunawan, I.; Xu, Z. B.; Collini, M. D.; Miller, C. P.; Dihn, T.; Henderson, R. A.; Keith Jr., J. C.; Harris, H. A. Design and synthesis of aryl diphenolic azoles as potent and selective estrogen receptor- β ligands. *J. Med. Chem.* **2004**, *47*, 5021-5040.
18. Krolski, M. E.; Renaldo, A. F.; Rudisill, D. E.; Stille, J. K. Palladium-catalysed coupling of 2-bromoanilines with vinylstannanes. A regiocontrolled synthesis of substituted indoles. *J. Org. Chem.* **1988**, *53*, 1170-1176.