



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Informatics - Papers (Archive)

Faculty of Engineering and Information Sciences

2005

Values of Minors of Some Infinite Families of Matrices Constructed from Supplementary Difference Sets and Their Application to the Growth Problem

C. Koukouvinos

National Technical University of Athens, Greece

M. Mitrouli

University of Athens, Greece

Jennifer Seberry

University of Wollongong, jennie@uow.edu.au

Publication Details

This article was originally published as Koukouvinos, C, Mitrouli, M and Seberry, J, Values of Minors of Some Infinite Families of Matrices Constructed from Supplementary Difference Sets and Their Application to the Growth Problem, *Linear Algebra and its Applications*, 406, 2005, 218-234. Copyright Elsevier 2005. Original journal available [here](#).

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Values of Minors of Some Infinite Families of Matrices Constructed from Supplementary Difference Sets and Their Application to the Growth Problem

Abstract

We obtain explicit formulae for the values of the $2v - j$ minors, $j = 0, 1, 2$ of $(1, -1)$ matrices of order $2v$, v odd, where the matrix is constructed using two circulant or type 1 incidence matrices of $2 - \{v; k_1, k_2, \lambda\}$ sds. This allows us to obtain information on the growth problem for families of matrices with moderate growth. Some of our theoretical formulae imply growth close to the order $2v$ but experimentation has not yet supported this result. An open problem remains to establish whether the $(1, -1)$ CP incidence matrices of certain SBIBDs, can have growth greater than $2v$.

Keywords

: SBIBD, supplementary difference sets, Gaussian elimination, growth, complete pivoting.

Disciplines

Physical Sciences and Mathematics

Publication Details

This article was originally published as Koukouvinos, C, Mitrouli, M and Seberry, J, Values of Minors of Some Infinite Families of Matrices Constructed from Supplementary Difference Sets and Their Application to the Growth Problem, *Linear Algebra and its Applications*, 406, 2005, 218-234. Copyright Elsevier 2005. Original journal available [here](#).

Values of Minors of Some Infinite Families of Matrices Constructed from Supplementary Difference Sets and Their Application to the Growth Problem

C. Koukouvinos*, M. Mitrouli^{†‡} and Jennifer Seberry[§]

October 23, 2004

Abstract

We obtain explicit formulae for the values of the $2v - j$ minors, $j = 0, 1, 2$ of $(1, -1)$ matrices of order $2v$, v odd, where the matrix is constructed using two circulant or type 1 incidence matrices of $2 - \{v; k_1, k_2, \lambda\}$ sds. This allows us to obtain information on the growth problem for families of matrices with moderate growth. Some of our theoretical formulae imply growth close to the order $2v$ but experimentation has not yet supported this result. An open problem remains to establish whether the $(1, -1)$ CP incidence matrices of certain SBIBDs, can have growth greater than $2v$.

Key Words and Phrases: SBIBD, supplementary difference sets, Gaussian elimination, growth, complete pivoting.

AMS Subject Classification: 05B20, 15A15, 65F05, 65G05.

Short running title: Minors of matrices from difference sets and growth problem.

1 Introduction

A set of k residues $D = \{a_1, \dots, a_k\}$ modulo v is called a (v, k, λ) difference set or cyclic difference set, if for every $d \not\equiv 0 \pmod{v}$ there are exactly λ ordered pairs (a_i, a_j) , $a_i, a_j \in D$ such that $a_i - a_j \equiv d \pmod{v}$.

For the purpose of this paper we will define two supplementary difference sets $2 - \{v; k_1, k_2; \lambda\}$, abbreviated as sds, to be two circulant (or type 1) $v \times v$ matrices B_1 and B_2 , with entries 0 or 1, which have exactly k_i entries +1 and $v - k_i$ entries 0, $i = 1, 2$ respectively, in each row and column and for which the inner product of any pair of rows of $[B_1 \ B_2]$ is λ . The $(1, -1)$ incidence matrices of B_i , are obtained by letting $A_i = 2B_i - J$, $i = 1, 2$, where J is the $v \times v$ matrix of all ones. Then we have for all methods below A is of order $2v$ is given by

*Department of Mathematics, National Technical University of Athens, Zografou 15773, Athens, Greece, e-mail: ckoukouv@math.ntua.gr

[†]Department of Mathematics, University of Athens, Panepistemiopolis 15784, Athens, Greece, e-mail: mmitroul@cc.uoa.gr

[‡]The work of this author was supported by a grant from the University of Athens.

[§]School of Information Technology and Computer Science, University of Wollongong, Wollongong, NSW, 2522, Australia, e-mail: jennie@uow.edu.au

$$A = \begin{bmatrix} A_1 & A_2 \\ A_2^T & -A_1^T \end{bmatrix} \quad (1)$$

$$A_1 A_1^T + A_2 A_2^T = 4(k_1 + k_2 - \lambda)I + 2(v - 2(k_1 + k_2 - \lambda))J = (2v - \Lambda)I + \Lambda J$$

where $\Lambda = 2v - 4(k_1 + k_2 - \lambda)$, and I is the identity matrix of order v .

We can write

$$AA^T = (2v + 2)I_{2v} - 2I_2 \times J_v.$$

It is easy to use the determinant simplification theorem [8] to see that for methods I and II, which follows, we have

$$\det A = 2^v(2v - 1)(v - 1)^{v-1}.$$

In this paper we study the family $2 - \{v; k_1, k_2; \lambda\}$ sds, and using $\lambda(v - 1) = k_1(k_1 - 1) + k_2(k_2 - 1)$ we can conclude that

$$\det A = 2^v(k_1 + k_2 - \lambda)^{v-1}(v^2 - 2v(k_1 + k_2) + 2k_1^2 + 2k_2^2).$$

Method I.

When $v = 4t + 1$ is a prime power, there exists a supplementary difference set with parameters $2 - \{v; \frac{v-1}{2}, \frac{v-1}{2}; \frac{v-3}{2}\} = 2 - \{4t + 1; 2t, 2t; 2t - 1\}$, see [14, 15].

Their incidence $(1, -1)$ matrices A_1 and A_2 satisfying the relation

$$A_1 A_1^T + A_2 A_2^T = (2v + 2)I - 2J.$$

Method II.

As described in Baumert [1, p.119], the best known n^{th} power residue difference sets are the quadratic residue sets of Paley [11].

When $v = 4t - 1$ is a prime power, the quadratic residues modulo v form a difference set with parameters $(v, k, \lambda) = (4t - 1, 2t - 1, t - 1)$. Thus, if we take this difference set twice we form a $2 - \{v; \frac{v-1}{2}, \frac{v-1}{2}; \frac{v-3}{2}\} = 2 - \{4t - 1; 2t - 1, 2t - 1; 2t - 1\}$ sds.

Their incidence $(1, -1)$ matrices are identical in this case, say $P = A_1 = A_2$, and then we have

$$2PP^T = (2v + 2)I - 2J.$$

We can write

$$AA^T = (2v + 2)I_{2v} - 2I_2 \times J_v.$$

It is easy to use the determinant simplification theorem [8] to see that for methods I and II we have

$$\det A = 2^v(2v - 1)(v - 1)^{v-1}.$$

We also examine some other methods of constructing sds given in [18].

Method III.

When $q = 25 + 4b^2 \equiv 13 \pmod{16}$ is a prime power, we consider the following $2 - \{q; \frac{q-1}{4}, \frac{q-1}{2}; \frac{5q-17}{16}\}$ sds. In this case we have

$$A_1 A_1^T + A_2 A_2^T = \left(\frac{7q + 5}{4}\right)I + \left(\frac{q - 5}{4}\right)J.$$

As an example for $q = 29$ we obtain a $2 - \{29; 7, 14; 8\}$ sds, which gives

$$A_1 A_1^T + A_2 A_2^T = 52I + 6J.$$

Method IV.

When $q = 49 + 4b^2 \equiv 5 \pmod{16}$ is a prime power, we consider the following $2 - \{q; \frac{q-1}{2}, \frac{3(q-1)}{4}, \frac{13q-33}{16}\}$ sds. In this case we have

$$A_1 A_1^T + A_2 A_2^T = \left(\frac{7q+17}{4}\right)I + \left(\frac{q-17}{4}\right)J.$$

As an example for $q = 53$ we obtain a $2 - \{53; 26, 39; 41\}$ sds, which gives

$$A_1 A_1^T + A_2 A_2^T = 97I + 9J.$$

Method V.

When $q = 1 + 4b^2 \equiv 5 \pmod{16}$ is a prime power, we consider the following $2 - \{q; \frac{q-1}{2}, \frac{3(q-1)}{4}, \frac{13q-33}{16}\}$ sds. In this case we have

$$A_1 A_1^T + A_2 A_2^T = \left(\frac{7q+17}{4}\right)I + \left(\frac{q-17}{4}\right)J.$$

As an example for $q = 37$ we obtain a $2 - \{37; 18, 27; 28\}$ sds, which gives

$$A_1 A_1^T + A_2 A_2^T = 69I + 5J.$$

In the present paper we get values for the pivots [17] of $2 - \{v; k_1, k_2; \lambda\}$ sds, and $(1, -1)$ matrices of order $2v$ made from them. Calculations have given moderate values of growth for such matrices. An open problem concerning the possibility of finding $(1, -1)$ $2v \times 2v$ CP matrices having growth greater than $2v$ is posed.

Notation 1. Write A for a matrix of order n whose initial pivots are derived from matrices with CP structure. Write $A(j)$ for the absolute value of the determinant of the $j \times j$ principal submatrix in the upper lefthand corner of the matrix A and $A[j]$ for the absolute value of the determinant of the $(n-j) \times (n-j)$ principal submatrix in the bottom righthand corner of the matrix A . Throughout this paper when we have used i pivots we then find all possible values of the $A(n-i)$ minors. Hence, if any minor is CP it must have one of these values. The magnitude of the pivots appearing after the application of GE operations on a CP matrix W is given by

$$p_j = A(j)/A(j-1), \quad j = 1, 2, \dots, n, \quad A(0) = 1. \quad (2)$$

In particular for a CP A , constructed from the \pm incidence matrices of 2-sds,

$$p_v = A(v)/A(v-1), \quad p_{v-1} = A(v-1)/A(v-2). \quad (3)$$

We use the notation M_j to denote the $j \times j$ minor of A .

2 Minors of Size $(2v - 1)$

We denote by $\Delta(h, i, j, k, m)$ the following matrix of order $2v$:

$$\Delta(h, i, j, k, m) = \begin{bmatrix} \overbrace{\begin{matrix} mI + (\Lambda - 1)(J - I) \\ (\Lambda + 1)J \\ -J \\ J \end{matrix}}^h & \overbrace{\begin{matrix} (\Lambda + 1)J \\ mI + (\Lambda - 1)(J - I) \\ J \\ -J \end{matrix}}^i & \overbrace{\begin{matrix} -J \\ J \\ mI + (\Lambda - 1)(J - I) \\ -J \end{matrix}}^j & \overbrace{\begin{matrix} J \\ -J \\ -J \\ mI + (\Lambda - 1)(J - I) \end{matrix}}^k \end{bmatrix}.$$

where $m = 2v = h + i + j + k$. Then by the Determinant Simplification Theorem [8]

$$\det \Delta(h, i, j, k, m) = (m-1)^{m-4} \begin{vmatrix} m + (h-1)(\Lambda-1) & (\Lambda+1)h & -h & h \\ (\Lambda+1)i & m + (i-1)(\Lambda-1) & i & -i \\ -j & j & m + (j-1)(\Lambda-1) & (\Lambda+1)j \\ k & -k & (\Lambda+1)k & m + (k-1)(\Lambda-1) \end{vmatrix}$$

To find the $(2v - 1) \times (2v - 1)$ minors we remove the first row and column of A to get D . The matrix DD^T is obtained from $\Delta(h, i, j, k, m)$ by removing a row and the corresponding column and performing on it appropriate permutations of rows and columns. Removing a row is equivalent by taking $m - 1$ rows and removing a column is equivalent by considering $h - 1$ or $i - 1$ or $j - 1$ or $k - 1$ columns. Thus $\det DD^T$ is $\det \Delta(h - 1, i, j, k, m - 1)$ or $\det \Delta(h, i - 1, j, k, m - 1)$ or $\det \Delta(h, i, j - 1, k, m - 1)$ or $\det \Delta(h, i, j, k - 1, m - 1)$.

In the sequel, we study the values of the $(2v - 1) \times (2v - 1)$ minors for the methods I-V mentioned in the Introduction.

Lemma 1 *The $(2v - 1) \times (2v - 1)$ minors of the matrix A given in (1), and constructed using Method II are:*

Proof. Here we use the $(1, -1)$ incidence matrices of the $2 - \{4t - 1; 2t - 1, 2t - 1; 2t - 1\}$ sds. PLEASE SPECIFY THESE VALUES

By the reasoning above, with $v = 4t - 1$, $h =$, $i =$, $j =$, $k =$, $m =$, substituted into $\det \Delta(h - 1, i, j, k, m - 1)$, $\det \Delta(h, i, j - 1, k, m - 1)$, $\det \Delta(h, i - 1, j, k, m - 1)$, and $\det \Delta(h, i, j, k - 1, m - 1)$ we obtain the result.

Specifically the $(2v - 1) \times (2v - 1)$ minor is the square root of the determinant and are given by one of

- 1) $\det \Delta(h - 1, i, j, k, m - 1) =$
- 2) $\det \Delta(h, i - 1, j, k, m - 1) =$
- 3) $\det \Delta(h, i, j - 1, k, m - 1) =$
- 4) $\det \Delta(h, i, j, k - 1, m - 1) =$

□

3 Minors of size $(2v - 2)$

We recall the partitioned matrix A of the design matrix, C , is composed from $2 - \{v; k_1, k_2; \lambda\}$ supplementary difference sets. Using the formula for the inner product of the rows of the $(1, -1)$ incidence matrix formed from these sds we see that the inner product is $2v - 4(k_1 + k_2 - \lambda) = \Lambda$.

We now return to A with two rows and columns removed to find the generic matrix. In expanded form this gives for C , simplified by the Determinant Simplification Theorem [8], the matrix D given by

$$\begin{bmatrix} N & \Lambda u_2 & \Lambda u_3 & (\Lambda + 2)u_4 & -2u_5 & 0 & 0 & 2u_8 \\ \Lambda u_1 & N & (\Lambda + 2)u_3 & \Lambda u_4 & 0 & -2u_6 & 2u_7 & 0 \\ \Lambda u_1 & (\Lambda + 2)u_2 & N & \Lambda u_4 & 0 & 2u_6 & -2u_7 & 0 \\ (\Lambda + 2)u_1 & \Lambda u_2 & \Lambda u_3 & N & 2u_5 & 0 & 0 & -2u_8 \\ -2u_1 & 0 & 0 & 2u_4 & N & 2u_6 & 2u_7 & (\Lambda + 2)u_8 \\ 0 & -2u_2 & 2u_3 & 0 & \Lambda u_5 & N & (\Lambda + 2)u_7 & \Lambda u_8 \\ 0 & 2u_2 & -2u_3 & 0 & \Lambda u_5 & (\Lambda + 2)u_6 & N & \Lambda u_8 \\ 2u_1 & 0 & 0 & -2u_4 & (\Lambda + 2)u_5 & \Lambda u_6 & \Lambda u_7 & N \end{bmatrix}.$$

where $N = 2v - 2u_i + (u_i - 1)\Lambda$ on the diagonal.

This gives the determinant of A with two rows and columns removed, as $(2v - \Lambda)^{v-5} \sqrt{\det D}$. Diagrammatically, we have used the matrix form

$$\begin{bmatrix} A_1 & A_2 \\ A_2^T & -A_1^T \end{bmatrix} = \begin{bmatrix} B_1 & B_2 \\ B_3 & B_4 \end{bmatrix}$$

For case I both rows and columns are removed from B_1 ; for case II one row is from B_1 and one from B_3 but both columns are from A_1 ; for case III one row is from B_1 and one from B_3 and one column is from B_1 and one column is from B_2 .

$\begin{array}{ccc} 1 & 1 & \\ \vdots & \vdots & \lambda_1 \\ 1 & 1 & \\ 1 & - & \\ \vdots & \vdots & k_1 - \lambda_1 \\ 1 & - & \\ - & 1 & \\ \vdots & \vdots & k_1 - \lambda_1 \\ - & 1 & \\ - & - & \\ \vdots & \vdots & v - 2k_1 + \lambda_1 \\ - & - & \end{array}$	$v - 2$ rows which have inner product Δ with rows one and two
$\begin{array}{ccc} 1 & 1 & \\ \vdots & \vdots & \lambda_2 \\ 1 & 1 & \\ 1 & - & \\ \vdots & \vdots & k_2 - \lambda_2 \\ 1 & - & \\ - & 1 & \\ \vdots & \vdots & k_2 - \lambda_2 \\ - & 1 & \\ - & - & \\ \vdots & \vdots & v - 2k_2 + \lambda_2 \\ - & - & \end{array}$	$\lambda = \lambda_1 + \lambda_2$ v rows which have inner product 0 with rows one and two

Number of Rows of Each Type I and II

u_1	u_2	u_3	u_4	u_5	u_6	u_7	u_8
$\lambda_1 - 2$	$k_1 - \lambda_1$	$k_1 - \lambda_1$	$v - 2k_1 + \lambda_1$	λ_2	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
$\lambda_1 - 1$	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1$	$v - 2k_1 + \lambda_1$	λ_2	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
$\lambda_1 - 1$	$k_1 - \lambda_1$	$k_1 - \lambda_1 - 2$	$v - 2k_1 + \lambda_1 - 1$	λ_2	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
$\lambda_1 - 1$	$k_1 - \lambda_1$	$k_1 - \lambda_1 - 2$	$v - 2k_1 + \lambda_1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
$\lambda_1 - 1$	$k_1 - \lambda_1$	$k_1 - \lambda_1 - 2$	$v - 2k_1 + \lambda_1 - 1$	λ_2	$k_2 - \lambda_2 - 1$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
$\lambda_1 - 1$	$k_1 - \lambda_1 - 2$	$k_1 - \lambda_1 - 2$	$v - 2k_1 + \lambda_1 - 1$	λ_2	$k_2 - \lambda_1$	$k_2 - \lambda_1$	$v - 2k_2 + \lambda_2$
λ_1	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1 - 1$	$v - 2k_1 + \lambda_1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
λ_1	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1$	$v - 2k_1 + \lambda_1 - 1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
λ_1	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1$	$v - 2k_1 + \lambda_1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
λ_1	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1 - 1$	$v - 2k_1 + \lambda_1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
λ_1	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1 - 1$	$v - 2k_1 + \lambda_1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
λ_1	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1 - 1$	$v - 2k_1 + \lambda_1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$
λ_1	$k_1 - \lambda_1 - 1$	$k_1 - \lambda_1 - 1$	$v - 2k_1 + \lambda_1$	$\lambda_2 - 1$	$k_2 - \lambda_2$	$k_2 - \lambda_2$	$v - 2k_2 + \lambda_2$

Table 1

Case III

To help understand Case III we recall that in this case one column removed comes from the columns with $k_1 + k_2$ ones per column and the other from the columns with $v - k_2 + k_1$ ones per column in the original design. This means the generic form of these two columns is

1		1	
1		\vdots	ρ
1		1	
1	k_1	—	
\vdots		\vdots	$k_1 - \rho$
1		—	
—		1	
\vdots		\vdots	$k_2 - \rho$
—	$v - k_1$	1	
—		—	
\vdots		\vdots	$v - k_1 - k_2 + \rho$
—		—	
1		1	
1		\vdots	$k_2 - \rho$
1		1	
1	k_2	—	
\vdots		\vdots	ρ
1		—	
—		1	
\vdots		\vdots	$v - k_1 - k_2 + \rho$
—	$v - k_2$	1	
—		—	
\vdots		\vdots	$k_1 - \rho$
—		—	

Note they have inner product zero.

Number of Rows of Each Type III							
u_1	u_2	u_3	u_4	u_5	u_6	u_7	u_8
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 1$	$k_1 - \rho - 1$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$
$\rho - 2$	$k_1 - \rho$	$k_2 - \rho$	$v - k_1 - k_2 + \rho$	$k_2 - \rho$	ρ	$v - k_1 - k_2 + \rho$	$k_1 - \rho$

Table 2

4 Pivot structure for family of design matrices

Conjecture

Let A be an $2v \times 2v$ CP D -optimal design of KKSSS family which is constructed from $2 - \{s^2 + s + 1; \frac{s(s-1)}{2}, \frac{s(s+1)}{2}; \frac{s(s-1)}{2}\}$ sds. Reduce A by GE. Set $\mathcal{P} = 2s^2 + 2s + 1$. Then we conjecture

- (i) $g(v, A) = \frac{s+1}{s}\mathcal{P}$, or $\frac{s}{s+1}\mathcal{P}$, or $\frac{s(s+1)}{s^2+s+1}\mathcal{P}$, or \mathcal{P} ;
- (ii) The last pivot is equal to $\frac{s+1}{s}\mathcal{P}$, or $\frac{s}{s+1}\mathcal{P}$, or $\frac{s(s+1)}{s^2+s+1}\mathcal{P}$, or \mathcal{P} ;
- (iii) The second last pivot can take the values given in Table 8.
- (iv) Every pivot before the last has magnitude at most $2v$;
- (v) The first four pivots are equal to $1, 2, 2, 4$;
- (vi) The fifth pivot may be 2 or 3.

We prove (ii) and (iii) in this paper. (v) and (vi) were proved for Brouwer's SBIBD($2s^2 + 2s + 1, s^2, \frac{1}{2}s(s-1)$) in [7] and we also show they hold for the KKSSS family.

v	growth	Pivot Pattern
7	8	(1, 2, 2, 4, 2, 4, 4, 2, 4, 4, 8, 4, 8, 8)
7	8	(1, 2, 2, 4, 3, $\frac{10}{3}$, 3.2, 4, 4, 4, 4, 4, 8, 8)
22	12	(1, 2, 2, 4, 3, $\frac{10}{3}$, $\frac{18}{5}$, 4, 3, 6, 6, 2, 4, 4, 8, 6, $\frac{20}{3}$, 7, 2, 8, 6, 12, 12)

Table 8: Growth Factors and Pivots Patterns for small CP KKSSS designs

References

- [1] L.D. Baumert, *Cyclic Difference Sets*, Lecture Notes in Mathematics, Vol. 182, Springer-Verlag, Berlin, Heidelberg, New York, 1971.
- [2] J. Day and B. Peterson, Growth in Gaussian elimination, *Amer. Math. Monthly*, 95 (1988), 489-513.
- [3] H. Ehlich, Determinantenabschätzungen für binäre matrizen, *Math. Z.*, 83 (1964), 123-132.
- [4] N. J. Higham, *Accuracy and Stability of Numerical Algorithms*, SIAM, Philadelphia, 1996.
- [5] C. Koukouvinos, S. Kounias, and J. Seberry, Supplementary difference sets and optimal designs, *Discrete Math.*, 88 (1991), 49-58.
- [6] C. Koukouvinos, M. Mitrouli and J. Seberry, Growth in Gaussian elimination for weighing matrices $W(n, n-1)$, *Linear Algebra and its Applications*, 306 (2000), 189-202.
- [7] C. Koukouvinos, M. Mitrouli and J. Seberry, Values of minors of $(1, -1)$ incidence matrices of SBIBDs and their application to the growth problem, *Designs, Codes and Cryptography*, 23 (2001), 267-281.

- [8] C. Koukouvinos, M. Mitrouli and J. Seberry, Values of minors of an infinite family of D -optimal designs and their application to the growth problem, *SIAM J. Matrix Anal. Appl.*, 23 (2001), 1-14.
- [9] C. Koukouvinos, M. Mitrouli and J. Seberry, Values of minors of an infinite family of D -optimal designs and their application to the growth problem: II, (submitted).
- [10] E. Lehmer, On residue difference sets, *Canad. J. Math.*, 5 (1953), 425-432.
- [11] R.E.A.C. Paley, On orthogonal matrices, *J. Math. and Phys.*, 12 (1933), 311-320.
- [12] J. Singer, A theorem in finite projective geometry and some applications to number theory, *Trans. Amer. Math. Soc.*, 43 (1938), 377-385.
- [13] E. Spence, Skew-Hadamard matrices of the Goethals-Seidel type, *Canad. J. Math.*, 27 (1975), 555-560.
- [14] J. Seberry Wallis, Some remarks on supplementary difference sets, *Colloquia Mathematica Societatis, Janos Bolyai*, No. 10, 1973, 1503-1526.
- [15] J. Seberry Wallis and A.L. Whiteman, Some classes of Hadamard matrices with constant diagonal, *Bull. Austral. Math. Soc.*, 7 (1972), 233-249.
- [16] A.L. Whiteman, A family of D -optimal designs, *Ars Combinatoria*, 30 (1990), 23-26.
- [17] J. H. Wilkinson, *The Algebraic Eigenvalue Problem*, Oxford University Press, London, 1988.
- [18] M. Yamada, Supplementary difference sets and Jacobi sums, *Discrete Math.*, 103 (1992), 75-90.