



UNIVERSITY  
OF WOLLONGONG  
AUSTRALIA

University of Wollongong  
Research Online

---

Faculty of Science - Papers (Archive)

Faculty of Science, Medicine and Health

---

2011

# Lutein from de-epoxidation of lutein epoxide replaces zeaxanthin to sustain an enhanced capacity for non-photochemical chlorophyll fluorescence quenching in avocado shade leaves in the dark

Britta Forster

*Australian National University*

Barry James Pogson

*Australian National University*

Charles Barry Osmond

*University of Wollongong, [cosmond@uow.edu.au](mailto:cosmond@uow.edu.au)*

---

## Publication Details

Forster, B., Pogson, B. & Osmond, C. B. (2011). Lutein from de-epoxidation of lutein epoxide replaces zeaxanthin to sustain an enhanced capacity for non-photochemical chlorophyll fluorescence quenching in avocado shade leaves in the dark. *Plant Physiology*, March 1-34.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: [research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

# Lutein from de-epoxidation of lutein epoxide replaces zeaxanthin to sustain an enhanced capacity for non-photochemical chlorophyll fluorescence quenching in avocado shade leaves in the dark

## Abstract

Leaves of avocado (*Persea americana* Mill.) that develop and persist in deep shade canopies have very low rates of photosynthesis but contain high concentrations of lutein epoxide (Lx) that are partially de-epoxidized to lutein (L) after one hour exposure to 120 to 350  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  increasing the total L pool by 5-10% ( $\Delta\text{L}$ ). De-epoxidation of Lx to L was near stoichiometric and similar in kinetics to de-epoxidation of violaxanthin (V) to antheraxanthin (A) and zeaxanthin (Z). Although the V pool was restored by epoxidation of A and Z overnight, the Lx pool was not. Depending on leaf age and pretreatment, the pool of  $\Delta\text{L}$  persisted for up to 72 h in the dark. Metabolism of  $\Delta\text{L}$  did not involve epoxidation to Lx. These contrasting kinetics enabled us to differentiate three states of the capacity for nonphotochemical chlorophyll fluorescence quenching (NPQ) in attached and detached leaves:  $\Delta\text{pH}$ -dependent (NPQ $\Delta\text{pH}$ ) before de-epoxidation; after de-epoxidation in the presence of  $\Delta\text{L}$ , A, and Z (NPQ $\Delta\text{LAZ}$ ); and after epoxidation of A+Z but with residual  $\Delta\text{L}$  (NPQ $\Delta\text{L}$ ). The capacity of both NPQ $\Delta\text{LAZ}$  and NPQ $\Delta\text{L}$  was similar and 45% larger than NPQ $\Delta\text{pH}$  but dark relaxation of NPQ $\Delta\text{LAZ}$  was slower. The enhanced capacity for NPQ was lost after metabolism of  $\Delta\text{L}$ . The near equivalence of NPQ $\Delta\text{LAZ}$  and NPQ $\Delta\text{L}$  provides compelling evidence that the small dynamic pool  $\Delta\text{L}$  replaces A+Z in avocado to “lock-in” enhanced NPQ. Results are discussed in relation to data obtained with other Lx-rich species and in mutants of *Arabidopsis* with increased L pools.

## Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

## Publication Details

Forster, B., Pogson, B. & Osmond, C. B. (2011). Lutein from de-epoxidation of lutein epoxide replaces zeaxanthin to sustain an enhanced capacity for non-photochemical chlorophyll fluorescence quenching in avocado shade leaves in the dark. *Plant Physiology*, March 1-34.

Lutein replaces zeaxanthin in photoprotection of avocado

Corresponding author:

Britta Förster

Plant Sciences Division, Research School of Biology, RN Robertson Building 46,

The Australian National University,

Canberra, ACT 0200, Australia.

Phone: +61-2-61254213

Email: [Britta.forster@anu.edu.au](mailto:Britta.forster@anu.edu.au)

Most appropriate journal research area:

Bioenergetics and photosynthesis

Lutein from de-epoxidation of lutein epoxide replaces zeaxanthin to sustain an enhanced capacity for non-photochemical chlorophyll fluorescence quenching in avocado shade leaves in the dark

Britta Förster<sup>\*1</sup>, Barry James Pogson<sup>2</sup>, Charles Barry Osmond<sup>1,3</sup>

<sup>1</sup> Plant Sciences Division, Research School of Biology, The Australian National University, Canberra, ACT 0200, Australia.

<sup>2</sup> ARC Centre of Excellence in Plant Energy Biology, Research School of Biology, The Australian National University Canberra, ACT 0200, Australia

<sup>3</sup> Present address: Institute of Conservation Biology and Environmental Management School of Biological Sciences, University of Wollongong, NSW 2522, Australia

Footnotes

This work was supported by the Australian Research Council (grant nos. DP0666289 to C.B.O. and CE056195 to B.J.P.)

\* Corresponding author

Britta Förster

Email: [britta.forster@anu.edu.au](mailto:britta.forster@anu.edu.au)

## ABSTRACT

Leaves of avocado (*Persea americana* Mill.) that develop and persist in deep shade canopies have very low rates of photosynthesis but contain high concentrations of lutein epoxide (Lx) that are partially de-epoxidized to lutein (L) after one hour exposure to 120 to 350  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$  increasing the total L pool by 5-10% ( $\Delta\text{L}$ ). De-epoxidation of Lx to L was near stoichiometric and similar in kinetics to de-epoxidation of violaxanthin (V) to antheraxanthin (A) and zeaxanthin (Z). Although the V pool was restored by epoxidation of A and Z overnight, the Lx pool was not. Depending on leaf age and pretreatment, the pool of  $\Delta\text{L}$  persisted for up to 72 h in the dark. Metabolism of  $\Delta\text{L}$  did not involve epoxidation to Lx. These contrasting kinetics enabled us to differentiate three states of the capacity for nonphotochemical chlorophyll fluorescence quenching (NPQ) in attached and detached leaves:  $\Delta\text{pH}$ -dependent ( $\text{NPQ}_{\Delta\text{pH}}$ ) before de-epoxidation; after de-epoxidation in the presence of  $\Delta\text{L}$ , A, and Z ( $\text{NPQ}_{\Delta\text{LAZ}}$ ); and after epoxidation of A+Z but with residual  $\Delta\text{L}$  ( $\text{NPQ}_{\Delta\text{L}}$ ). The capacity of both  $\text{NPQ}_{\Delta\text{LAZ}}$  and  $\text{NPQ}_{\Delta\text{L}}$  was similar and 45% larger than  $\text{NPQ}_{\Delta\text{pH}}$  but dark relaxation of  $\text{NPQ}_{\Delta\text{LAZ}}$  was slower. The enhanced capacity for NPQ was lost after metabolism of  $\Delta\text{L}$ . The near equivalence of  $\text{NPQ}_{\Delta\text{LAZ}}$  and  $\text{NPQ}_{\Delta\text{L}}$  provides compelling evidence that the small dynamic pool  $\Delta\text{L}$  replaces A+Z in avocado to “lock-in” enhanced NPQ. Results are discussed in relation to data obtained with other Lx-rich species and in mutants of *Arabidopsis* with increased L pools.

## INTRODUCTION

The violaxanthin cycle (V-cycle) based on reversible inter-conversion of zeaxanthin (Z), antheraxanthin (A) and violaxanthin (V) stabilizes non-photochemical chlorophyll fluorescence quenching (NPQ) and confers photoprotection in most green plants and algae (Demmig et al 1987; Demmig-Adams and Adams 1992; Niyogi 1999; Förster et al 2001). Since Bungard et al (1999) an additional xanthophyll cycle (the Lx-cycle), involving the  $\alpha$ -carotene ( $\alpha$ -C) pathway pigments lutein (L) and lutein epoxide (Lx), has been recognized in many diverse taxa (García-Plazaola et al 2003; Matsubara et al 2003, 2008).

The Lx-cycle is distinctive in that in some species it was reversible with kinetics similar to that of the V-cycle (Bungard et al 1999; Matsubara et al 2001; Esteban et al 2010), whereas in others it was only very slowly reversible (Matsubara et al 2005; García-Plazaola et al 2007; Förster et al 2009). Leaves that developed and persisted in deep shade canopies of avocado (*P. americana* Mill.) showed particularly high concentrations of Lx (Esteban et al 2008). Although de-epoxidation of Lx occurred during a few hours of exposure to sunlight and made a small addition to the much larger pool of L ( $\Delta$ L) little epoxidation of  $\Delta$ L to Lx was observed overnight and Lx concentrations were only restored after 3 to 5 weeks in the shade (Förster et al 2009). In contrast, the V-cycle was largely reversible on a diel basis.

Slow reversibility of the Lx-cycle in some species led to the hypothesis that augmentation of the L pool by de-epoxidation of Lx might “lock-in” photoprotection (García-Plazaola et al 2003; Matsubara et al 2005). Evidence for persistent, enhanced capacity for NPQ due to  $\Delta$ L, after A+Z had largely reverted to V, was reported during photosynthetic induction in-vivo with leaves of *Quercus rubra* and *Inga marginata* (García-Plazaola et al 2003; Matsubara et al 2008). These experiments provided the first direct evidence for potentially similar functional roles for  $\Delta$ L and Z in photoprotection (García-Plazaola et al 2007; Horton et al 2008) and presaged the recent demonstration of a Z-like radical cation that appeared when NPQ was engaged in a Z-free, L-enriched Arabidopsis mutant (Avenson et al 2007; Li et al 2009). However, as yet there have been no quantitative comparisons of the components of NPQ associated with  $\Delta$ L and A+Z.

Not surprisingly, shade leaves of avocado proved to be extremely responsive to light during treatment and assay. Photosynthetic induction curves at selected, often high, light intensities have been widely used to assess NPQ in relation to xanthophyll pigment composition in Arabidopsis and Chlamydomonas mutants (Niyogi et al 1998; Pogson et al 1998; Förster 2001), as well as in *Quercus* and *Inga*. These assays proved unsatisfactory for

quantitative comparisons of NPQ components in shade leaves of avocado, and were confounded by responses of stomata and CO<sub>2</sub> supply (Takayama et al 2008), by photosynthetic induction following treatment, and by xanthophyll de-epoxidation during the measurement. In contrast, the light response curve traditionally used to distinguish biochemical and physiological relationships in sun and shade plants (Björkman 1981), has found less application in these model systems (Russell et al 1996; Bailey et al 2001; Pérez-Bueno and Horton 2008). We tailored rapid light response curves (RLRC) using chlorophyll fluorescence (Schreiber et al 1994; White and Critchley 1999) to the requirements of avocado shade leaves. These minimized de-epoxidation during the assay and facilitated quantitative analysis of NPQ while monitoring opening of stomata and induction of photosynthetic metabolism following light treatments.

Rapid light response curves were used to differentiate and quantitatively compare three components of the capacity for NPQ in avocado leaves that are expressed following inter-conversions of xanthophyll pigments. Semantic confusion that has emerged from use of the terms qE and qN (Horton et al 1996), particularly in reference to different components of energy-dependent quenching, distinguished on different criteria, in different species and treatments led us to a simple designation the three types of NPQ in shade leaves of avocado with subscripts as follows:

(i) NPQ<sub>ΔpH</sub> in dark adapted, un-induced leaves with closed stomata that contain high concentrations of Lx and V and extremely low levels of A and Z. We presume this component is associated with development of ΔpH in chloroplast thylakoid membranes in the absence of external CO<sub>2</sub> (Krause et al 1982) during assay, prior to de-epoxidation of Lx and V.

(ii) NPQ<sub>ΔLAZ</sub> assayed in leaves after modest light exposures that open stomata, induce photosynthetic metabolism and lead to partial but stoichiometric de-epoxidation in the Lx- and V-cycles. We presume this component is associated with the stabilization of NPQ by both ΔL and A+Z.

(iii) NPQ<sub>ΔL</sub> in leaves treated as in (ii) but assayed after the 24-72 h in the dark following almost complete epoxidation of the V cycle (disappearance of A+Z) but not of Lx. We presume this component is associated with stabilization of NPQ by ΔL alone.

In this paper we provide the first direct evidence that the capacity of NPQ<sub>ΔL</sub> is quantitatively similar to NPQ<sub>ΔLAZ</sub> suggesting that a small addition (~5 - 10%) to the total L pool (ΔL) substitutes for A+Z and maintains the capacity of elevated NPQ for up to 72 h in the dark. The duration of elevated NPQ<sub>ΔL</sub> is determined by the rate of dark metabolism of ΔL by processes other than epoxidation to Lx. Importantly, dark relaxation of NPQ<sub>ΔL</sub> is faster



than that of  $\text{NPQ}_{\Delta\text{LAZ}}$  and similar to  $\text{NPQ}_{\Delta\text{PH}}$ . These observations are discussed in relation to a common mechanism of L and Z based NPQ recently proposed in other studies, and in relation to potential functions of L-enhanced photoprotection in shade leaves.

## RESULTS

### Optimization of light treatments to manipulate xanthophyll pigment composition in shade leaves of avocado and to avoid subsequent de-epoxidation during assays

Our previous studies of diel changes in Lx- and V-cycle pigment composition in avocado leaves suggested little de-epoxidation occurred at light intensities  $< 200 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  (Förster et al 2009). The optimal light intensities needed to bring about quantitatively similar de-epoxidation of Lx and V in mature leaves of shade-grown avocado plants without photoinhibition were explored with a single large shade-grown avocado leaf detached mid afternoon and arranged with the petiole in water in the open greenhouse. The top third of the leaf was covered by aluminum foil (Al-foil) as a dark control and neutral density filters were placed on the remainder of the leaf to give  $50 \times 50 \text{ mm}$  areas exposed to 80 and  $163 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , leaving other parts of the leaf exposed to the weak afternoon sunlight ( $300 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) for 90 min. As shown in Fig. 1, leaf discs from dark control areas contained very little A ( $1.6 \pm 0.2 \text{ mmol mol}^{-1} \text{ Chl}$ ), only traces of Z ( $0.5 \pm 0.5 \text{ mmol mol}^{-1} \text{ Chl}$ ). Note that the 3-5 fold larger pools of L are shown reduced by  $100 \text{ mmol mol}^{-1} \text{ Chl}$  (designated L\*) in all figures unless stated otherwise to facilitate comparison of stoichiometry with other xanthophylls. Whereas de-epoxidation of Lx to L was first evident after exposure to 163 and  $300 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  ( $\Delta\text{L} \sim 15 \text{ mmol mol}^{-1} \text{ Chl}$ ), quantitatively similar de-epoxidation of V to A+Z occurred already at  $80 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Therefore, all subsequent actinic light treatments to manipulate the pool sizes of both xanthophylls cycles were done at 120 and  $350 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  for 60 min.

Measurements of pigments in discs punched at random from control and treated areas of attached mature leaves in 9 experiments showed quantitatively similar, almost stoichiometric de-epoxidation of Lx and V (Table I). Initial levels of A were close to the limits of detection ( $0.8 \text{ mmol mol}^{-1} \text{ Chl}$ ) whereas Z was not detectable in most cases. Although levels of residual A+Z after 21 to 72 h recovery in the dark lab ( $0\text{-}5 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) were higher, residual A+Z and de-epoxidation ( $\text{DS} = \text{A} + \text{Z} / \text{V} + \text{A} + \text{Z}$ ) was about the same as that produced from de-epoxidation of V during the optimized RLRC assays

described below. Importantly, measurements of  $F_v/F_m$  showed these light exposures caused little photoinhibition (Table I).

Some de-epoxidation of V, but not of Lx, always occurred during chlorophyll fluorescence assays. For example, contrary to expectations of Takayama et al (2008), imaging photosynthetic induction for 20 min at  $100 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  caused substantial de-epoxidation of V and increased DS from 0.04 to 0.24 (data not shown). Assuming that each of the  $16 \times 1 \text{ s}$  saturating flashes used in this protocol delivered  $5,000 \mu\text{mol photons m}^{-2}$ , this assay delivered a total of  $200 \text{ mmol photons m}^{-2}$  which is equivalent to the photons delivered in a 6 min photosynthetic induction assay at  $300 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  with the MINI-PAM (Fig. 2 A). This problem had been mitigated previously by infiltration of leaf discs with DTT, an inhibitor of violaxanthin de-epoxidase (Adams et al 1990) but clearly this was impracticable in our measurements with large attached leaves. Moreover, infiltration with DTT had been reported to substantially depress  $F_v/F_m$  (Matsubara et al 2008). Seeking a compromise between the actinic light intensity needed to produce good resolution of kinetic responses and that needed to minimize de-epoxidation during assays (Fig. 2 B), we chose a rapid light response curve (RLRC) protocol delivering about  $60 \mu\text{mol photons m}^{-2}$  using  $8 \times 30 \text{ s}$  steps with actinic light from 0 to  $\sim 400 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ .

### **A persistent 5 to 10% addition ( $\Delta L$ ) to the L pool from de-epoxidation of Lx sustains enhanced capacity for NPQ after conversion of A+Z to V overnight**

The less intrusive RLRC assays enabled us to follow enhancement of NPQ in the presence of  $\Delta L$  under standardized conditions for up to 72 h in the dark. In addition, these curves routinely presented light response profiles of 1-qP (a measure of the redox state of the acceptor side of PSII) and photosynthetic electron transport (ETR). These data helped us ensure that other factors, such as stomatal opening and induction of photosynthesis during light treatments, did not confound the interpretation of NPQ data.

Light treatments caused an initial de-epoxidation of  $\sim 30\%$  of both Lx and V pools with practically no change in PSII efficiency indicated by steady  $F_v/F_m$  (Fig. 3 A). Pigment composition changed little during 1 h dark following treatment. V-cycle DS declined by one-third, mainly due to epoxidation of about half the Z pool, without change in the pools of A or L. Although 1-qP was depressed by 20% at low PFD immediately after exposure, the profiles of control and exposed areas were the same after 1 h in the dark (Fig. 3 B), consistent with the

initial increase in ETR that also returned to control levels after 1 h in the dark (Fig. 3 C). These responses of 1-qP to light treatment and their relaxation to controls within 1 h dark were similar in all experiments and are not presented in subsequent graphs. Light treatment led to a relatively larger increase in capacity for NPQ at lower light intensities during assay, as well as a higher level of NPQ at light saturation (Fig. 3 D). Notably, in the presence of A+Z NPQ relaxed more slowly in the dark during assay (Fig. 3 D). The small decline of the NPQ profile in exposed areas during 1 h dark may reflect the partial epoxidation of the Z pool (Fig. 3 A).

In general, L was not metabolized during 24 h in the dark, whereas epoxidation of A+Z was practically complete (Fig. 3 A). Photosynthetic parameters in the Al-foil controls after 21 h in the dark were almost identical to the dark controls at the beginning of the experiment. Profiles of 1-qP (cf. Fig. 3 B, E) and ETR (cf. Figs. 3 C, F) in control and recovered areas were the same after 1 h and 21 h dark. Enhanced capacity for NPQ persisted in exposed areas after 21 h dark (Fig. 3 G), in spite of the epoxidation of A+Z. Importantly the kinetics of dark NPQ relaxation now were much the same in control and exposed areas (Fig. 3 G). The slower decline of NPQ in the dark following de-epoxidation of Lx and V, and its fast relaxation after epoxidation of A+Z, was observed in all experiments with attached leaves. This showed that although  $\Delta L$  substituted for A+Z and sustained higher capacity for NPQ in the dark, it did not slow the dark relaxation of NPQ observed in the presence of A+Z.

### **Enhanced capacity for NPQ persists for up to 72 h in the dark without A+Z, while ever $\Delta L$ is detectable**

Younger, fully-expanded leaves with lower Lx content (Fig. 4 A) showed much the same de-epoxidation of Lx and V after light treatments, and commencement of epoxidation of A+Z during 1 h and 3 h in the dark. Photosynthetic induction in the light treatment increased ETR during assay which declined to near control levels after 3 h dark (Fig. 4 B). Non-photochemical quenching increased following light treatment then declined slightly as the level of Z began to decrease after 3 h in the dark, but the slower dark relaxation of NPQ persisted (Fig. 4 C). Epoxidation of A+Z was largely complete after 24 h dark but  $\Delta L$  was still detectable (Fig. 4 D). Although ETR of treated areas was somewhat lower than controls after 24 h dark (Fig. 4 E), persistent  $\Delta L$  sustained higher capacity for NPQ (Fig. 4 F) with relaxation kinetics similar to controls. Metabolism of  $\Delta L$  between 24 and 48 h re-established control levels of L (Fig. 4 C) and this was reflected in identical profiles of ETR and NPQ

(Fig. 4 G, H). Clearly, the capacity for enhanced NPQ persisted only as long as  $\Delta L$  was detectable, and interestingly, the metabolism of  $\Delta L$  between 24 and 48 h dark was not due to epoxidation to Lx.

The longest period of enhanced capacity for NPQ associated with sustained  $\Delta L$  in our experiments was found in isolated leaf discs 72 h after the start of the actinic light treatment (Fig. 5). Light treatment produced a persistent 30% decrease in Lx and a less than stoichiometric, but still substantial, increase in  $\Delta L$  after 48 and 72 h in the dark. The levels of V, A, and Z were the same as controls (Fig. 5 A) and A+Z formed during the light treatment was epoxidised to V within 24 h in the dark (data not shown). The ETR observed after 48 and 72 h in the dark was slightly greater than in controls (Fig. 5 B) showing that elevated NPQ capacity (Fig. 5 C) was not an artifact of reduced ETR. The relaxation of NPQ was the same as controls throughout.

We were able to eliminate sustained  $\Delta L$  accumulation and the associated enhanced NPQ <sub>$\Delta L$</sub>  by pretreatment of a plant in the dark lab for 11 days ( $0\text{-}5 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) prior to light treatment (1 h at  $200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). Prolonged dark pretreatment did not alter de-epoxidation of Lx and V during exposure, or epoxidation of A+Z overnight (Fig. 6 A). The response of ETR and NPQ to the light treatment was as found in all previous experiments (Fig. 6 B, C) as was the slower dark relaxation of NPQ in the presence of  $\Delta L$  and A+Z. However, prolonged dark pretreatment specifically accelerated metabolism of the pool of  $\Delta L$ . Both pools of  $\Delta L$  and A+Z were metabolized overnight and no enhancement of NPQ was detected during assays (Fig. 6 B, C) and as expected, relaxation kinetics of NPQ were the same as controls.

## DISCUSSION

Previous studies suggested that the presence of Lx- and V-cycles in shade canopy leaves of woody plants from Mediterranean (García-Plazaola et al 2003) and tropical American forests (Matsubara et al 2008) indicated that photoconversion of Lx to L may “lock-in” photoprotection overnight after restoration A+Z to V. The pronounced difference in the kinetics of epoxidation of L and A+Z in the Lx-rich shade leaves of avocado (Esteban et al 2008; Förster et al 2009) offered an ideal system in which to probe the role in photoprotection of the particular fraction of L ( $\Delta L$  from Lx) that persists after epoxidation of A+Z. This paper presents the first comprehensive and quantitative analysis using non-intrusive rapid light response curves to differentiate three states of the capacity for NPQ

associated with distinct xanthophyll pigment compositions in these plants and substantially extends our understanding of this enigmatic role of L in-vivo.

The three pigment composition states were established by exposing shade leaves of avocado to modest, non-photoinhibitory light treatments ( $120\text{-}350 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). The transitions between  $\text{NPQ}_{\Delta\text{pH}}$  (in the absence of de-epoxidation of xanthophylls), to  $\text{NPQ}_{\Delta\text{LAZ}}$  (after stoichiometric de-epoxidation of Lx and V) and to  $\text{NPQ}_{\Delta\text{L}}$  (persistence of  $\Delta\text{L}$  following overnight epoxidation of A+Z) were assayed using sensitive RLRC in which a range of 30 s exposures to increasing PFD was tailored to minimize de-epoxidation of xanthophyll pigments during assay. The non-intrusive assays were supported with simple tests, such as the “Vaseline patch”, that help monitor physiological (closure of stomata) and biochemical (induction of photosynthetic metabolism) processes with the potential to confound interpretations of NPQ capacity in relation to pigment composition in-vivo. We did not use routine photosynthetic induction curves to analyze changes in NPQ in shade-grown avocado leaves because they produced higher de-epoxidation of V (Fig. 2). As an aside, we strongly recommend that de-epoxidation should be routinely assessed during NPQ induction assays at high PFD.

Four functional aspects of photoprotection that have arisen from this investigation of avocado shade leaves will be discussed. First, we will explore comparative quantitative aspects of the substitution of A+Z by  $\Delta\text{L}$  to sustain enhanced capacity for NPQ for up to 72 h in the dark and expand earlier hypotheses that de-epoxidation of Lx to L “locks-in” photoprotection (García-Plazaola et al 2003, 2007; Matsubara et al 2005, 2007, 2008). Second and contrary to expectations, we briefly discuss the demonstration that the duration of enhanced NPQ capacity was determined by metabolism of  $\Delta\text{L}$  by processes other than its slow epoxidation to Lx. Third, we point out that in avocado leaves, de-epoxidation of V to A+Z slows the dark relaxation of NPQ during assay, whereas  $\Delta\text{L}$  from de-epoxidation of Lx does not affect NPQ relaxation kinetics. Fourth, we suggest some implications of sustained NPQ capacity with the above properties for photoprotection in shade canopies and light use efficiency during sun-flecks.

Past research has been focused on the extent to which Z-independent components of chlorophyll fluorescence quenching ( $\text{NPQ}_{\Delta\text{pH}}$ ) participate in NPQ even after de-epoxidation xanthophylls. There is a debate as to whether these components are additive or substitutive (Adams et al 1990; Bilger and Björkman 1994; Gilmore et al 1998; Li et al 2004), and whether common or separate mechanisms are involved (Pogson et al 1998; Finazzi et al

2004). Recent evidence suggests that both involve common antenna based mechanisms (Johnson et al 2009), and moreover, at least two distinct quenching mechanisms involving Z (type I and type II) seem to be involved. In the type I mechanism, Z in the L2 position of minor antenna complexes is a direct quencher of excited chlorophyll as demonstrated by a distinctive radical cation (Avenson et al 2008). In the type II mechanism, Z is an allosteric modulator of the  $\Delta$ pH-sensitivity of NPQ involving xanthophyll exchanges in the L1 position of Lhcs (Johnson et al 2009). It is beyond the scope of our data to speculate further on these mechanistic details.

We took a simple approach to quantitative estimation of the relative NPQ capacity using the total areas under RLRC of NPQ vs PFD to compare  $\text{NPQ}_{\Delta\text{pH}}$  and  $\text{NPQ}_{\Delta\text{LAZ}}$  in avocado leaves. Comparison of controls with those after 1 h exposure and after 1 h dark with similar amounts of  $\Delta$ L and A+Z, show an increase of  $45.1 \pm 1.5\%$  ( $n = 4$  experiments) in  $\text{NPQ}_{\Delta\text{LAZ}}$  compared to  $\text{NPQ}_{\Delta\text{pH}}$ . In other words, in these assays the capacity of Z (and  $\Delta$ L)-independent  $\text{NPQ}_{\Delta\text{pH}}$  in control leaves was about 69% of  $\text{NPQ}_{\Delta\text{LAZ}}$  after de-epoxidation of Lx and V. Compared to leaves with persistent  $\Delta$ L after 21-24 h dark recovery,  $\text{NPQ}_{\Delta\text{L}}$  was increased by  $36.6 \pm 2.5\%$  over  $\text{NPQ}_{\Delta\text{pH}}$ . Even after 48 h dark recovery  $\text{NPQ}_{\Delta\text{L}}$  was still 84% of  $\text{NPQ}_{\Delta\text{LAZ}}$ . Thus, our analyses provide evidence that  $\Delta$ L effectively and almost completely substitutes for A+Z in stabilizing capacity for NPQ in avocado.

The mechanistic aspects of substitution of A+Z by  $\Delta$ L are little known. Analyses of shade leaves of *Inga sapindoides* (Matsubara et al 2007) showed that Lx was the principal xanthophyll in LHCII complexes. In these experiments de-epoxidation of part of the Lx pool to L on exposure to  $200 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  for 30 min led to the replacement of Lx with L in the peripheral V1 site of LHCII trimers, as well as in the internal L2 site of both monomeric and LHCII trimers in both photosystems. It was concluded that the slowly reversible  $\text{Lx} \leftrightarrow \text{L}$  exchange took place at the same binding sites as reversible  $\text{V} \leftrightarrow \text{A+Z}$  exchange. We suggest the observed sustained high capacity for  $\text{NPQ}_{\Delta\text{L}}$  is based on substitution of A+Z with L in these sites, but further evidence for the association of  $\Delta$ L, A and Z with specific LHCII proteins in avocado is needed before this substitution can be integrated into general models of regulated excitation dissipation in antenna complexes (de Bianchi et al 2010).

Whereas the structural roles of L in the antennae of the photosynthetic apparatus have been the focus of much past research, the photoprotective function of L has remained enigmatic (Gilmore 2001). Analyses of Arabidopsis mutants for a decade or so had not

uncovered direct evidence of a role for L in NPQ. Although *lutein-deficient2* (*lut2*) mutants show less NPQ (Pogson et al 1998) and although other mutants with lower than wild type L content show impaired NPQ, mutants with slightly enhanced L and enhanced NPQ (Pogson and Rissler 2000; Niyogi et al 2001) have been difficult to evaluate, principally because none were markedly deficient in Z. A recently constructed suppressor of the *npq1* mutant *suppressor of zeaxanthinless1* (*slz1 npq1*) lacks Z, has low levels of V and A, accumulates more L and  $\alpha$ -carotene than wildtype, and retains about 50% wild type NPQ (Li et al 2009). The authors state “*Analysis of the carotenoid radical cation formation and leaf absorbance changes strongly suggest that the higher amount of lutein substitutes for zeaxanthin in qE, implying a direct role in qE,...*”. In the light of these observations in Arabidopsis it seems a compelling next step is to examine shade leaves of avocado in the NPQ $_{\Delta pH}$ , NPQ $_{\Delta LAZ}$  and NPQ $_{\Delta L}$  states to discover whether distinctive leaf absorbance changes, and changes in carotene radical cation formation (Avenson et al 2008; Li et al 2009), persist in-vivo following the substitution of A+Z by  $\Delta L$ .

A still unanswered key question is how the partitioning of the majority of the L pool to structural functions in Lhcs is regulated and how a small additional pool ( $\Delta L$ ) participates in excitation dissipation. This bears some analogy to the situation in the Arabidopsis mutants *lut2*, where V-cycle pigments substitute for the structural roles of L (Pogson et al 1996), and in *aba1-3* and *aba1-4* which are constitutively enriched in Z and depleted in V+A, but only a small fraction of the Z pool seems to be involved in NPQ photoprotection (Hurry et al 1997). Also, this may related to the still ill-defined fraction of the L and Z pools that mitigate photo-oxidation (Havaux and Niyogi 1999; Johnson et al 2007) and the contribution of these ROS scavenging activities to photoprotection (Förster et al 2005; Dall’Osto et al 2006). It remains to be seen if the recently described partitioning (Lepetit et al 2010) of xanthophylls in diatoms between protein-associated fractions (involved in NPQ) and lipid-dissolved fractions (involved in ROS scavenging) can be achieved with avocado thylakoids. Both L and Z serve similar roles in visual systems (Kim et al 2006), and photo-oxidation products of both xanthophylls have been identified in retinas of the human eye (Khachik et al 1997). We observed earlier that sudden exposure of shade-grown avocado leaves to sunlight was initially accompanied by a decline in the total L pool in spite of Lx de-epoxidation (Förster et al 2009), which could be explained by photo-oxidation of L. It would be of interest to investigate the occurrence of L (and Z) photo-oxidation products in avocado leaves under these conditions.

The transition from higher capacity  $\text{NPQ}_{\Delta\text{L}}$  to lower capacity  $\text{NPQ}_{\Delta\text{pH}}$  after prolonged darkness was associated with disappearance of  $\Delta\text{L}$ . Against expectation, the decline of  $\Delta\text{L}$  was not coupled to epoxidation of  $\Delta\text{L}$  to Lx and restoration of the Lx pool. Interestingly, metabolism of  $\Delta\text{L}$  occurred much faster when leaves were pretreated with 11 days in the dark lab prior to light exposure, when  $\Delta\text{L}$  was fully metabolized in 24 h darkness without epoxidation (Fig. 6). In previous experiments (Förster et al 2009) slow recovery of Lx pools in prolonged shade after de-epoxidation in sunlight was not simply related to epoxidation of L or accumulation of the biosynthetic precursor  $\alpha$ -carotene, as this pool remained constant and the total L pool declined more than twice as rapidly as the Lx pool recovered. Obviously, other pathways to metabolize  $\Delta\text{L}$  became dominant in prolonged dark. Little is known about degradation of xanthophyll pigments, although carotenoid cleavage dioxygenases and non-enzymatic degradation have been implicated in some tissues (Cazzonelli and Pogson 2010). At present we cannot explain the dark-induction of L metabolism.

Another intriguing aspect of the NPQ-pigment relationships was that relaxation kinetics of NPQ were delayed specifically by the presence of A+Z, whereas both  $\text{NPQ}_{\Delta\text{pH}}$  and  $\text{NPQ}_{\Delta\text{L}}$  relaxation rates were equally fast in the absence of A+Z. This seems consistent with previous evidence that Z slows the relaxation of qE in isolated chloroplasts (Nocter et al 1991). More recently, Johnson et al (2008) reported that in wildtype Arabidopsis and a Z-accumulating  $\beta$ -carotene hydroxylase overexpression line (*sChy B*), slower relaxation in the dark after assay was not related to the absolute Z pool itself, but to higher DS. Other contributing factors to the slow relaxation of  $\text{NPQ}_{\Delta\text{LAZ}}$  cannot be ruled out. For example, the slower decay of  $\text{NPQ}_{\Delta\text{LAZ}}$  might reflect post-illumination interactions with leaf respiratory metabolism and could reflect slowly declining energization of chloroplast thylakoids in darkness, when chloroplast and/or mitochondrial ATP pools sustain NPQ in the dark through a  $\Delta\text{pH}$  maintained by ATP hydrolysis rather than photosynthetic electron transport (Gilmore and Björkman 1995).

Persistent,  $\Delta\text{L}$ -enhanced capacity for ( $\text{NPQ}_{\Delta\text{L}}$ ) was first recognized in shade leaves of *Quercus* and *Inga* examined in the field (García-Plazaola et al 2003, Matsubara et al 2008) and has been confirmed in some *Inga* spp. in the Eden Project in Cornwall, UK (Nichol and Osmond, unpublished) as well as avocado shade leaves in orchards in tropical Eastern Australia (Osmond, Förster and Leonardi, unpublished). The deeply shaded canopies of the latter retain up to 30 leaves on shoots with photosynthetic capacities and pigment compositions that are comparable to those of shade-grown plants examined here. The modest



light treatments used to bring about xanthophyll pigment conversions in our greenhouse experiments correspond reasonably to changes in shade canopy light environment following upper canopy disturbance in storms, and to changes following routine orchard canopy pruning practices (Whiley et al 2002; Osmond, Förster and Leonardi, unpublished). The 30s steps in our RLRC curves reasonably correspond to sun light fleck experiences in tropical canopies (Percy 1990) and to sun fleck measurements in avocado orchards in California (Mickelbart, Stilwell, Arpaia and Heath, unpublished). The effect of  $\Delta L$  on “locking-in” higher capacity of  $NPQ_{\Delta L}$  in the shade for several days, without the penalty of slower relaxation associated with the presence of Z, may offer a lower penalty in light energy use efficiency during sequences of sun flecks than the slower relaxing  $NPQ_{\Delta LAZ}$  (Zhu et al 2004). These consequences of the slowly reversible Lx-cycle need to be evaluated in relation to the reversible engagement of V-cycle in these plants. For example, we found previously that sunlight exposure of avocado shade leaves results in sustained low  $F_v/F_m$  (Förster et al 2008) and persistently higher L and A+Z. Separation of the contributions of slowly reversible Z and/or L-related NPQ from xanthophyll-independent, low-efficiency photoinactivated PSII centers to these observations will be discussed in a subsequent paper.

In conclusion, this study established improved assay methods to analyze xanthophyll pigment-NPQ relationships in avocado shade leaves that resolved three different NPQ components based on the associated xanthophyll de-epoxidation state. Based on these distinctions we were able to provide novel, direct evidence for a special role of the fraction  $\Delta L$  of the L pool that had been generated by Lx de-epoxidation in conferring higher capacity for NPQ induction while allowing rapid dark relaxation, which implies a direct role of L in photoprotection and may improve the ability of avocado shade leaves to persist during frequent sun flecks. Evaluation of these processes during shade to sun acclimation in avocado shade leaves under laboratory and field conditions will be presented elsewhere.

## **MATERIALS AND METHODS**

### **Plant material and growth conditions**

Seedlings of avocado (*Persea americana* Mill. cv Edranol) were purchased from Vallance’s Nursery (Mullumbimby NSW 2482 Australia) and maintained in 20 L containers of potting soil with regular irrigation and additions of slow release nutrients. The seedlings were kept in a shaded section of a temperature controlled glasshouse for 24 months (18°C

night/ 29°C day), and pruned to the main stem 6-12 months prior to experiments. The light environment of upper and lower canopy leaves in the shade enclosure, measured with a LI-190 sensor ([www.licor.com](http://www.licor.com)), was 30-50 and 15-30  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  respectively, when horizontal full sun in the outer greenhouse was 1,100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  at noon.

Fully expanded leaves ranged from 20-35 cm in length and were thinner and deeper green in the shade than in the sun. Transverse sections confirmed that shade-grown avocado leaves were heterobaric to the level of 4<sup>th</sup> class veins with a single layer of palisade cells above 2-4 layers of spongy mesophyll cells and had numerous small stomata restricted to the underside. Imaging experiments showed this anatomy conferred markedly heterogeneous, small scale differences in NPQ during photosynthetic induction and light response curves. However this heterogeneity was effectively integrated over areas of 1-2  $\text{cm}^2$ , and similar results were obtained from imaging and spot measurement techniques (Takayama et al 2008). Photosynthetic parameters were routinely measured from the upper epidermis, but measurements from the lower epidermis were not significantly different.

### **Experimental protocols**

Small shade-grown avocado trees were moved to the dark lab overnight prior to light treatments to alter xanthophyll pigment composition, but one experiment was done with a plant maintained in the dark lab for 11 d. Sets of two or three adjacent attached leaves of similar size and age in the canopy were part covered with Al-foil and/or black cloth prior to illumination of the rest of the leaf at 120 to 350  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  for 60 min using wide-beam high-efficiency fluorescent spotlights (Philips PAR38 23W; 2700K warm white). A small leaf disc was punched from foil control and exposed areas of each leaf (or from the larger leaf discs) for pigment analysis before light treatment, and again at the end of the treatment, then after 1 to 3 h dark to relax photosynthetic induction following light treatment and after 21 to 72 h recovery in darkness. As found previously (Förster et al 2009), large avocado leaves were remarkably robust with respect to sampling for pigment analysis. Moreover, photosynthetic parameters measured adjacent to, or remote from, disc sampling areas were similar in both short-term (hours) and long-term (days) experiments.

Preliminary gas exchange measurements of photosynthesis in leaves of shade-grown plants during induction at 100  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  at 25 °C were very noisy, even at the slowest practicable flow rates. Stomata opened slowly to low conductance that was responsive to  $\text{CO}_2$  (20  $\text{mmol m}^{-2} \text{s}^{-1}$ , 7  $\text{mmol m}^{-2} \text{s}^{-1}$  and 3  $\text{mmol m}^{-2} \text{s}^{-1}$  in 100, 400 and 700ppm  $\text{CO}_2$

respectively after 20 min). Photosynthetic CO<sub>2</sub> assimilation saturated at 1.5 to 2.0 μmol m<sup>-2</sup> s<sup>-1</sup> after 5 min in 700 ppm CO<sub>2</sub>, reached 1.0 μmol m<sup>-2</sup> s<sup>-1</sup> after 20 min in 400 ppm CO<sub>2</sub>, and remained negative in 100 ppm CO<sub>2</sub>. Chlorophyll fluorescence imaging during the above experiments confirmed that NPQ was sensitive to [CO<sub>2</sub>] and increased most rapidly to highest values in 100 ppm (Takayama et al 2008). Mature leaves taken from the shade enclosure during the light period showed higher NPQ when a “Vaseline patch” was applied to the lower epidermis. Indeed, we used this simple technique to confirm stomata were closed in leaves dark adapted overnight (no response of ETR or NPQ to a “Vaseline patch”) and that stomata were open and photosynthesis was induced following the above actinic light treatments (decreased ETR and increased NPQ in response to a “Vaseline patch”).

Chlorophyll fluorescence assays of photosynthetic parameters in mature leaves of shade-grown avocado were made with the Photosynthesis Yield Analyzer MINI-PAM fitted with leaf clip holder 2030-B ([www.Walz.com](http://www.Walz.com)). Photosynthetic induction curves with *Arabidopsis* grown under similar light environments have been routinely run at 1,000 to 1,500 μmol photons m<sup>-2</sup> s<sup>-1</sup> (Pogson et al 1998). Previous light response curve assays using chlorophyll fluorescence have ranged from dwell times of 10 s at PFD from 0 to 2,500 μmol photons m<sup>-2</sup> s<sup>-1</sup> (White and Critchley 1999; peas grown at 200 μmol photons m<sup>-2</sup> s<sup>-1</sup>) to 180 s at PFD from 0 to 2,000 μmol photons m<sup>-2</sup> s<sup>-1</sup> (Pérez-Bueno and Horton 2008; *Arabidopsis* grown at 120 μmol photons m<sup>-2</sup> s<sup>-1</sup>). Leaves of shade-grown avocado were extremely responsive to actinic light intensity and duration during measurement and these protocols led to photoinhibition, manifest as reductions in ETR during photosynthetic induction curves with >300 μmol photons m<sup>-2</sup> s<sup>-1</sup> and in rapid light response curves with dwell times of 60 to 120 s at each PFD. More important was our observation that measureable de-epoxidation of V, but not of Lx, occurred during both assays and was always greater during induction assays. A compromise was reached between the actinic light intensity needed to produce acceptable kinetic responses in ETR and NPQ during RLRC assays and that causing the least de-epoxidation of V. We settled for eight steps of 30 s duration over the range from 0 to 400 μmol photons m<sup>-2</sup> s<sup>-1</sup>, followed by measurements at four intervals during 240 s in the dark.

### **Calculation of photosynthetic parameters**

The automated PIC and RLRC protocols in the MINI-PAM fluorimeter measured intrinsic chlorophyll fluorescence (F) and maximum fluorescence yield during a saturating flash (F<sub>m</sub>) in dark adapted leaves to calculate the maximum quantum yield of photochemical

energy conversion  $F_v/F_m = (F_m - F)/F_m$ . Photosynthetic electron transport was calculated from fluorescence yield in saturating flashes under actinic light ( $F_m'$ ) from the quantum yield of photochemical energy conversion  $\Delta F/F_m' = F_m' - F/F_m'$ , and the PFD measured at that spot using the quantum sensor of the leaf clip (adjusted for absorptance of 0.85 and assuming equal light absorption in PSII and PSI). After light exposures to alter the xanthophyll pigment composition of leaves all NPQ data were recalculated using  $F_m$  measured on the leaf kept in the dark overnight and/or  $F_m$  from Al-foil shaded areas of the leaf during and after treatment. Likewise,  $1 - qP = 1 - ((F_m' - F)/(F_m' - F_o'))$  was recalculated using the minimum value of  $F$  obtained within 100 s after actinic light was switched off as  $F_o'$ .

### **Pigment analyses**

Leaf discs (1 cm diameter) were punched from treated and control areas of leaves at the times specified in each experiment. Discs were wrapped in foil and immediately frozen in liquid nitrogen. Pigments were extracted from individual discs in a microfuge tube with 0.6 mL of ethyl acetate:acetone (60:40 v/v) and shaken at 30 Hz for 2 min with a stainless steel ball (2 mm diameter) before addition of 0.5 mL of water prior to 5 min centrifugation at 13,000 rpm. The pigment containing upper layer was transferred to a fresh microfuge tube, centrifuged as before, and 0.1 mL of the pigment solution placed in vials for HPLC analysis on an Agilent 1100 fitted with Waters Spherosorb ODS2 column, using a linear gradient from 100-33% acetonitrile:water (90:10 v/v with 0.1% triethanolamine) into ethyl acetate over 31 min. Pigments were identified by retention times and spectra, and carotenoid concentrations were calculated using conversion factors for absorbance at 440 nm obtained with pure pigments, determined by Dr Shizue Matsubara, ICG-III: Phytosphäre, Forschungszentrum Jülich, 52425 Jülich, Germany. The L pool in shade leaves of avocado was three to five-fold greater than that of the other xanthophylls, so values for L have been reduced by 100  $\mu\text{mol mol}^{-1}$  Chl (marked L\* in graphical presentations) to facilitate more sensitive comparisons of stoichiometric relationships. As found previously (Förster et al 2009) there were scarcely detectable changes in neoxanthin,  $\alpha$ - or  $\beta$ -carotene pool or Chl *a/b* ratio in the course of any of the 24 to 72 h experiments. In most cases only Lx- and V-cycle pigment concentrations are reported.

## **ACKNOWLEDGEMENTS**

The authors are grateful to Kotaro Takayama for his preliminary investigations of the photosynthetic properties of shade-grown avocado leaves using chlorophyll fluorescence imaging, to Steve Dempsey for maintenance of the plant material, and to Fred Chow for the use of his dark laboratory facilities.

## LITERATURE CITED

- Adams WW III, Demmig-Adams B, Winter K** (1990) Relative contributions of zeaxanthin-related and zeaxanthin-unrelated types of 'high-energy-state' quenching of chlorophyll fluorescence in spinach leaves exposed to various environmental conditions. *Plant Physiol* **92**: 302-309
- Avenson TJ, Ahn TK, Zigmantis D, Niyogi KK, Li Z, Ballottari M, Bassi R, Flemming GR** (2008) Zeaxanthin radical cation formation in minor light-harvesting complexes of higher plant antenna. *J Biol Chem* **283**: 3550-3558
- Bailey S, Walters RG, Jansson S, Horton P** (2001) Adaptation of *Arabidopsis thaliana* to the light environment: existence of low light and high light responses. *Planta* **213**:794-801
- Bilger W, Björkman O** (1994) Relationships among violaxanthin deepoxidation, thylakoid membrane conformation, and nonphotochemical chlorophyll fluorescence quenching in leaves of cotton (*Gossypium hirsutum* L.). *Planta* **193**: 238-246
- Björkman O** (1981) Responses to different quantum flux densities. *In* OL Lange, PS Nobel, CB Osmond, H Ziegler, eds, *Encyclopedia of Plant Physiology. Physiological Plant Ecology: I. Responses to the Physical Environment*, Vol 12A. Springer, Berlin, pp 57-107
- Bungard RA, Ruban AV, Hibberd JM, Press MC, Horton P, Scholes JD** (1999) Unusual carotenoid composition and a new type of xanthophyll cycle in plants. *Proc Natl Acad Sci USA* **96**: 1135-1139
- Cazzonelli CI, Pogson BJ** (2010) Source to sink: regulation of carotenoid biosynthesis in plants. *Trends Plant Sci* **15**: 266-274
- Dall'Osto L, Lico C, Alric J, Giuliano G, Havaux M, Bassi R** (2006) Lutein is needed for efficient chlorophyll fluorescence quenching in the major LHCII antenna complex of higher plants and effective photoprotection *in vivo* under strong light. *BMC Plant Biol* **6**: 32
- de Bianchi S, Ballottari M, Dall'Osto L, Bassi R** (2010) Regulation of plant light harvesting by thermal dissipation of excess energy. *Biochem Soc Trans* **38**: 651-660
- Demmig B, Winter K, Krüger A, Czygan F-C** (1987) Photoinhibition and zeaxanthin formation in intact leaves; a possible role of the xanthophyll cycle in the dissipation of excess light energy. *Plant Physiol* **84**: 218-224
- Demmig-Adams B, and Adams WW III** (1992) Photoprotection and other responses of plants to high light stress. *Annu Rev Plant Physiol Plant Molec Biol* **43**: 599-626

- Esteban R, Soledad-Jiménez M, Morales D, Jiménez E, Hormaetxe K, Becerril J-M, Osmond B, García-Plazaola J-I** (2008) Short and long-term modulation of the lutein-epoxide and violaxanthin cycles in two species of the Lauraceae: sweet bay laurel (*Laurus nobilis* L.) and avocado (*Persea americana* Mill). *Plant Biol* **10**: 288-297
- Esteban R, Matsubara S, Soledad-Jiménez M, Morales D, Brito P, Lorenzo R, Fernández-Marin B, Becerril J-M, García-Plazaola, JI** (2010) Operation and regulation of the lutein epoxide cycle in seedlings of *Ocotea foetans*. *Func Plant Biol* **37**: 859-869
- Finazzi G, Johnson GN, Dall'Osto L** (2004) A zeaxanthin-independent nonphotochemical quenching mechanism localized in the photosystem II core complex. *Proc Natl Acad Sci USA* **101**: 12375-12380
- Förster B, Osmond CB, Boynton JE** (2001) Very high light resistant mutants of *Chlamydomonas reinhardtii*: responses of photosystem II, nonphotochemical quenching and xanthophyll pigments to light and CO<sub>2</sub>. *Photosynth Res* **67**: 5-15
- Förster B, Osmond CB, Pogson BJ** (2005) Improved survival of very high light and oxidative stress is conferred by spontaneous gain-of-function mutations in *Chlamydomonas*. *Biochim Biophys Acta* **1709**: 45-57
- Förster B, Osmond CB, Pogson BJ** (2009) De-novo synthesis and degradation of Lx- and V-cycle pigments during shade and sun acclimation in avocado leaves (*P. americana*). *Plant Physiol* **149**: 1179-1195
- García-Plazaola J-I, Olano JM, Hernández A, Becerril JM** (2003) The operation of the lutein epoxide cycle correlates with energy dissipation. *Func Plant Biol* **30**: 319-324
- García-Plazaola J-I, Matsubara S, Osmond CB** (2007) The lutein epoxide cycle in higher plants: its relationship to other xanthophyll cycles and possible functions. *Func Plant Biol* **34**: 754-779
- Gilmore AM** (2001) Xanthophyll cycle-dependent nonphotochemical quenching in Photosystem II: mechanistic insights gained from *Arabidopsis thaliana* L. mutants that lack violaxanthin deepoxidase activity and/or lutein. *Photosynth Res* **67**: 89-101
- Gilmore AM, Björkman O** (1995) Temperature sensitive coupling and uncoupling of ATPase-mediated nonradiative energy dissipation: similarities between chloroplasts and leaves. *Planta* **197**: 646-654

- Gilmore AM, Shinkarev VP, Hazlett TL, Govindjee** (1998) Quantitative analysis of intrathylakoid pH and xanthophyll cycle pigments on chlorophyll *a* fluorescence lifetime distributions and intensity in thylakoids. *Biochemistry* **37**: 13582-13593
- Havaux M, Niyogi KK** (1999) The violaxanthin cycle protects plants from photooxidative damage by more than one mechanism. *Proc Natl Acad Sci USA* **96**: 8762-8767
- Horton P, Ruban AV, Walters RG** (1996) Regulation of light harvesting in green plants. *Annu Rev Plant Physiol Plant Molec Biol* **47**: 655-684
- Horton P, Johnson MP, Perez-Bueno ML, Kiss, AZ, Ruban AV** (2008) Photosynthetic acclimation: does the dynamic structure and macro-organisation of photosystem II in higher plant grana membranes regulate light harvesting states? *FEBS J* **275**: 1069-1079
- Hurry V, Anderson JM, Chow WS, Osmond CB** (1997) Replacement of epoxy-xanthophylls by zeaxanthin in the ABA-deficient mutants of *Arabidopsis thaliana* L (Heynh.) does not affect chloroplast fluorescence quenching, or sensitivity to photoinhibition in vivo. *Plant Physiol* **113**: 639-648
- Johnson MP, Havaux M, Triantaphylidès C, Ksas B, Pascal AA, Robert B, Davidson PA, Ruban AV, Horton P** (2007) Elevated zeaxanthin bound to oligomeric LHCII enhances resistance of *Arabidopsis* to photo-oxidative stress by a lipid protective, anti-oxidant mechanism. *J Biol Chem* **282**: 22605-22618
- Johnson MP, Davidson PA, Ruban AV, Horton P** (2008) The xanthophyll pool size controls the kinetics of nonphotochemical chlorophyll quenching in *Arabidopsis thaliana*. *FEBS Lett* **582**: 262-266
- Johnson MP, Pérez-Bueno ML, Zia A, Horton P, Ruban A** (2009) The zeaxanthin-dependent and zeaxanthin-independent qE components of nonphotochemical quenching involve common conformational changes within the photosystem II antenna in *Arabidopsis*. *Plant Physiol* **149**: 1061-1075
- Khachik F, Bernstein PS, Garland DL** (1997) Identification of lutein and zeaxanthin oxidation products in human and mouse retinas. *Investig Ophthalmol Visual Sci* **38**: 1802-1811
- Kim SR, Nakanishi K, Itagaki Y, Sparrow JR** (2006) Photooxidation of A2-PE, a photoreceptor outer segment fluorophore, and protection by lutein and zeaxanthin. *Exp Eye Res* **82**: 828-839



- Krause GH Vernotte C, Briantais** (1982) Photoinduced quenching of chlorophyll fluorescence in intact chloroplasts and algae. Resolution of two components. *Biochim Biophys Acta* **679**: 116-124
- Lepetit B, Volke D, Gilbert M, Wilhelm C, Goss R** (2010) Evidence for the existence of one antenna-associated, lipid-dissolved and two protein-bound pools of diadinoxanthin pigments in diatoms. *Plant Physiol* **154**: 1905-1920
- Li X-P, Gilmore AM, Caffarri S, Bassi R, Golan T, Kramer D, Niyogi KK** (2004) Regulation of light harvesting involves lumen pH sensing by the PsbS protein. *J Biol Chem* **279**: 22866-22874
- Li Z, Ahn TK, Avenson TJ, Ballottari M, Cruz JA, Kramer DM, Bassi R, Fleming GR, Keasling JD, Niyogi KK** (2009) Lutein accumulation in the absence of zeaxanthin restores nonphotochemical chlorophyll quenching in the *Arabidopsis thaliana npq1* mutant. *Plant Cell* **21**: 1798-181
- Matsubara S, Gilmore AM, Osmond CB** (2001) Diurnal and acclimatory responses of violaxanthin and lutein epoxide in the Australian mistletoe *Amyema miquelii*. *Aust J Plant Physiol* **28**: 793-800
- Matsubara S, Morosinotto T, Bassi R, Christian AL, Fischer-Schliebs E, Lüttge U, Orthen B, Franco AC, Scarano FR, Förster B, Pogson BJ, Osmond CB** (2003) Occurrence of the lutein-epoxide cycle in mistletoes of the Loranthaceae and Viscaceae. *Planta* **217**: 868-879
- Matsubara S, Naumann M, Martin R, Nichol C, Rascher U, Morosinotto T, Bassi R, Osmond B** (2005) Slowly reversible de-epoxidation of lutein-epoxide in deep shade leaves of a tropical tree legume may “lock in” lutein-based photoprotection during acclimation to strong light. *J Exp Bot* **56**: 461-468
- Matsubara S, Morosinotto T, Osmond B, Bassi R** (2007) Short- and long-term operation of the lutein-epoxide cycle in light-harvesting antenna complexes. *Plant Physiol* **144**: 926–941
- Matsubara S, Krause GH, Seltmann M, Virgo A, Kursar TA, Jahns P, Winter K** (2008) Lutein epoxide cycle, light harvesting and photoprotection in species of the tropical tree genus *Inga*. *Plant Cell Environ* **31**: 548-561

- Matsubara S, Krause GH, Aranda J, M, Virgo A, Beisel KG, Jahns P, Winter K** (2009) Shade-sun patterns of leaf carotenoid composition in 86 species of neotropical forest plants. *Func Plant Biol* **36**: 20-36
- Niyogi KK, Grossman AR, Björkman O** (1998) Arabidopsis mutants define a central role for the xanthophyll cycle in the regulation of photosynthetic energy conversion. *Plant Cell* **10**: 1121-1134
- Niyogi KK** (1999) Photoprotection revisited: Genetic and molecular approaches. *Annu Rev Plant Physiol Plant Molec Biol* **50**: 333-359
- Niyogi KK, Shih C, Chow WS, Pogson BJ, DellaPenna D, Björkman O** (2001) Photoprotection in a zeaxanthin and lutein-deficient double mutant of Arabidopsis. *Photosynth Res* **67**: 139-145
- Nocter G, Rees D, Young AJ, Horton P** (1991) The relationship between zeaxanthin, energy-dependent quenching of chlorophyll fluorescence, and trans-thylakoid pH gradient in isolated chloroplasts. *Biochim Biophys Acta* **1057**: 320-330
- Pearcy RW** (1990) Sunflecks and photosynthesis in plant canopies. *Annu Rev Plant Physiol Plant Molec Biol* **41**: 421-453
- Pérez-Bueno ML, Horton P** (2008) The role of lutein in the acclimation of higher plant chloroplast membranes to suboptimal conditions. *Physiol Plant* **134**:227-236
- Pogson BJ, McDonald K, Truong M, Britton G, DellaPenna D** (1996) Arabidopsis carotenoid mutants demonstrate lutein is not essential for photosynthesis in higher plants. *Plant Cell* **8**: 1627-1639
- Pogson BJ, Niyogi KK, Björkman O, DellaPenna D** (1998) Altered xanthophyll compositions adversely affect chlorophyll accumulation and nonphotochemical quenching in Arabidopsis mutants. *Proc Natl Acad Sci USA* **95**: 13324-13329
- Pogson BJ, Rissler HM** (2000) Genetic manipulation of carotenoid biosynthesis and photoprotection. *Phil Trans R Soc B* **355**: 1395-1403
- Ruban AV, Horton P** (1995) Regulation of non-photochemical quenching of chlorophyll fluorescence in plants. *Aust J Plant Physiol* **22**: 221-230
- Russell AW, Critchley C, Robinson SA, Franklin LA, Seaton GGR, Chow WS, Anderson JM, Osmond CB** (1995) Photosystem II regulation and dynamics of the

chloroplast D1 protein in Arabidopsis leaves during photosynthesis and photoinhibition. *Plant Physiol* **107**: 943-952

**Schreiber U, Bilger W, Neubauer C** (1994) Chlorophyll fluorescence as a non intrusive indicator for rapid assessment of in-vivo photosynthesis. *In* E-D Schulze, MM Caldwell, *Ecophysiology of Photosynthesis*. Springer, Berlin, pp.49-70

**Takayama K, Osmond B, Omasa K** (2008) Imaging heterogeneity of xanthophyll independent non-photochemical quenching during photosynthetic induction in shade-grown leaves of avocado (*Persea americana* L.). *In* JF Allen, E Gant, JH Golbeck, B Osmond B, eds, *Photosynthesis: Energy from the Sun*. 14<sup>th</sup> International Congress on Photosynthesis. Springer, Dordrecht. pp. 681-685

**Whiley AW, Schaffer B, Wolstenholme BN** (2002) Eds. *The Avocado: Botany, Production and Uses*. CABI Publishing, Oxford, UK.

**White AJ, Critchley C** (1999) Rapid light curves: a new fluorescence method to assess the state of the photosynthetic apparatus. *Photosynth Res* **59**: 63-72

**Zhu XG, Ort DR, Whitmarsh J, Long SP** (2004) The slow reversibility of photosystem II thermal energy dissipation on transfer from high to low light may cause large losses in carbon gain by crop canopies. *J Exp Bot* **55**: 1167-1175

## FIGURE LEGENDS

**Figure 1:** Light intensity dependence of Lx- and V-de-epoxidation in a detached avocado leaf on water. Note that values for L\* have been reduced by 100 mmol mol<sup>-1</sup> Chl to facilitate stoichiometric presentation of data (i.e., the scale for L\* corresponds to 100-160 mmol mol<sup>-1</sup> Chl). Parts of the leaf were shaded with Al-foil as control (dark), neutral density filters were used to vary light intensity (80 and 163 μmol photons m<sup>-2</sup> s<sup>-1</sup>), and the remainder of the leaf was un-shaded during 90 min exposure to weak afternoon sunlight (300 μmol photons m<sup>-2</sup> s<sup>-1</sup>) in a glasshouse at 28°C. (Mean ± SE; n = 3).

**Figure 2:** De-epoxidation of V in detached avocado leaves during chlorophyll fluorescence assay of photosynthetic parameters. Pigment composition of mature shade leaves before and after (A) photosynthetic induction assay at 300 μmol photons m<sup>-2</sup> s<sup>-1</sup> for 6 min and (B) rapid light response curves (RLRC) assay from 0 to 410 μmol photons m<sup>-2</sup> s<sup>-1</sup>. Note that values for L\* have been reduced by 100 mmol mol<sup>-1</sup> Chl to facilitate stoichiometric presentation of data as in Fig. 1 (i.e., the scale for L\* corresponds to 100-160 mmol mol<sup>-1</sup> Chl). (Mean ± SE; n = 4).

**Figure 3:** Pigment composition in foil covered (cont) and light exposed areas of attached avocado leaves (A) measured immediately following exposure to 150 μmol photons m<sup>-2</sup> s<sup>-1</sup> for 1 h (exp 1h), after recovery for 1 h dark (exp +1h dk), and in foil covered (cont 21h) and exposed areas (rec 21h) again after 21 h in the dark. Note the scale for L\* corresponds to 100 to 170 mmol mol<sup>-1</sup> Chl. Changes in photosynthetic parameters measured in RLRC at each of these pigment sampling times are shown in (B, E) 1-qP, (C, F) ETR and (D, G) NPQ, with the time course of NPQ relaxation in the dark shown in panels on the right of (D) and (G). (Mean ± SE, n = 3; error bars appear whenever SE exceeds symbol size).

**Figure 4:** Enhancement of NPQ is lost when ΔL is metabolized after 48 h dark. (A) Pigment composition measured in Al-foil control (cont) after exposure to 120 μmol photons m<sup>-2</sup> s<sup>-1</sup> (exp) and after 1 h and 3 h in the dark. Note the scale for L\* corresponds to 100 to 160 mmol mol<sup>-1</sup> Chl. (B) ETR and (C) NPQ assayed in RLRC in control and treatment at times as in (A). Pigment compositions (D) and ETR (E) and NPQ (F) measured again after 24 h and (G) and (H) 48 h recovery in the dark. The time course of NPQ relaxation in the dark following the RLRC is shown in columns on the right of (C), (F) and (H) (Mean ± SE; n = 4; error bars appear whenever SE exceeds symbol size).


**Figure 5:** Enhanced NPQ and retention of  $\Delta L$  from Lx persists for up to 72 h in the dark in leaf discs in the absence of A+Z. Discs from four leaves on a plant dark adapted overnight were floated lower epidermis uppermost and exposed to white LED from below (1 h at  $220 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) then allowed to recover on water in the dark. (A) Pigment composition of controls and of samples after 48 and 72 h; samples were collected after RLRC. Note that  $L^*$  has been adjusted by  $50 \text{ mmol mol}^{-1} \text{ Chl}$ , so the scale for L corresponds to 0 to  $120 \text{ mmol mol}^{-1} \text{ Chl}$ . (B) Photosynthetic parameters assayed in RLRC on controls and after 48 and 72 h recovery in the dark. (Mean  $\pm$  SE;  $n = 4$ ; error bars appear whenever SE exceeds symbol size).

**Figure 6:** Prolonged dark pretreatment promotes metabolism of  $\Delta L$  overnight and prevents sustained NPQ. Pigment compositions (A) show  $\Delta L$  is metabolized within 24 h. Note that the scale for  $L^*$  corresponds to 100 to  $150 \text{ mmol mol}^{-1} \text{ Chl}$ . Measurements of photosynthetic parameters in RLRC confirm normal induction of ETR (B) and enhanced NPQ (C) following de-epoxidation. However, following overnight metabolism of  $\Delta L$  without de-epoxidation and epoxidation of A+Z, ETR and NPQ return to control levels (D, E). The time course of NPQ relaxation in the dark is shown in panels on the right of C and E. (Mean  $\pm$  SE;  $n = 3$ ; error bars appear whenever SE exceeds symbol size).


**Table I:** Changes in Lx and V-cycle pigments in attached mature leaves of shade-grown avocado following light treatments (1 h at  $223 \pm 27 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) in nine experiments. (Means  $\pm$  SE; n = 9-13; 2-3 replicate leaves in each experiment).

Parameter	Pigments, $\text{mmol mol}^{-1}$ Chl			
	$\Delta\text{Lx}$	$\Delta\text{L}$	$\Delta\text{V}$	$\Delta\text{A+Z}$
De-epoxidation (1 h)	$-11.8 \pm 2.1$	$+10.2 \pm 0.6$	$-12.5 \pm 1.1$	$+11.7 \pm 1.1$
Epoxidation (>24 h)	A+Z $\text{mmol mol}^{-1}$ Chl	$\frac{\text{A+Z}}{\text{V+A+Z}}$	$F_v/F_m$	
Initial dark control	$0.8 \pm 0.3$	$0.03 \pm 0.01$	$0.808 \pm 0.002$	
After treatment	$12.5 \pm 0.6$	$0.37 \pm 0.03$	$0.787 \pm 0.002$	
After recovery	$3.4 \pm 0.6$	$0.11 \pm 0.03$	$0.789 \pm 0.003$	

**Figure 1**




**Figure 2**







**Figure 3**



**Figure 4**



**Figure 5**



**Figure 6**

