



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science - Papers (Archive)

Faculty of Science, Medicine and Health

2010

Spatial variability of cadmium, copper, manganese, nickel and zinc in the Port Curtis Estuary, Queensland, Australia

Brad M. Angel

University of Wollongong, bma06@uow.edu.au

Leigh T. Hales

CSIRO Land and Water

Stuart L. Simpson

CSIRO Land and Water, stuart.simpson@csiro.au

Simon C. Apte

CSIRO Land and Water

Anthony A. Chariton

CSIRO Land and Water

See next page for additional authors

Publication Details

Angel, B., Hales, L. T., Simpson, S. L., Apte, S. S., Chariton, A. A., Shearer, D. & Jolley, D. F. (2010). Spatial variability of cadmium, copper, manganese, nickel and zinc in the Port Curtis Estuary, Queensland, Australia. *Marine and Freshwater Research*, 61 (2), 170-183.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Spatial variability of cadmium, copper, manganese, nickel and zinc in the Port Curtis Estuary, Queensland, Australia

Abstract

Port Curtis is a rapidly growing industrialised and urbanised harbour in Central Queensland, Australia. Water sampling surveys were undertaken in late 2003 and 2004, accompanied by additional sediment sampling to investigate the sources and behaviour of trace metals, and the effects of pH on metal partitioning between dissolved and particulate forms. Sampling and analyses of trace metals in waters and suspended particulates were undertaken along axial transects extending away from possible point-sources within the harbour. Additional sampling was undertaken in selected inlets and major freshwater sources to Port Curtis, including the Fitzroy River. Most dissolved metal concentrations were significantly elevated in Port Curtis compared to the concentrations measured in the adjacent coastal waters. Dissolved cadmium did not exhibit any discernible spatial trend, with concentrations measured in Port Curtis in the range

Keywords

spatial, queensland, estuary, curtis, port, australia, copper, variability, zinc, cadmium, nickel, manganese, CMMB

Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

Angel, B., Hales, L. T., Simpson, S. L., Apte, S. S., Chariton, A. A., Shearer, D. & Jolley, D. F. (2010). Spatial variability of cadmium, copper, manganese, nickel and zinc in the Port Curtis Estuary, Queensland, Australia. *Marine and Freshwater Research*, 61 (2), 170-183.

Authors

Brad M. Angel, Leigh T. Hales, Stuart L. Simpson, Simon C. Apte, Anthony A. Chariton, Damon A. Shearer, and Dianne F. Jolley

1 Spatial variability of cadmium, copper, manganese, nickel,
2 and zinc in the Port Curtis Estuary, Queensland, Australia

3
4
5 **Brad M. Angel^{A,B}, Leigh T. Hales^A, Stuart L. Simpson^A, Simon C. Apte^A,**
6 **Anthony A. Chariton^A, Damon A. Shearer^C, Dianne F. Jolley^B**
7

8
9
10 ^A Centre for Environmental Contaminants Research, CSIRO Land and Water, Private Mailbag
11 7, Bangor, NSW 2234, Australia

12 ^B GeoQuest, Department of Chemistry, University of Wollongong, NSW 2522, Australia

13 ^C CRC for Coastal Zone, Estuary and Waterway Management, Centre for Environmental
14 Management, Central Queensland University, Gladstone, QLD 4680, Australia

15
16 Corresponding author: Brad Angel

17
18 Word count: 8400 Number of Figures: 7 Number of tables: 3
19
20

21 **Abstract** Port Curtis is a rapidly growing industrialised and urbanised harbour in Central
22 Queensland, Australia. Water sampling surveys were undertaken in late 2003 and 2004,
23 accompanied by additional sediment sampling to investigate the sources and behaviour of
24 trace metals, and the effects of pH on metal partitioning between dissolved and particulate
25 forms. Sampling and analyses of trace metals in waters and suspended particulates were
26 undertaken along axial transects extending away from possible point-sources within the
27 harbour. Additional sampling was undertaken in selected inlets and major freshwater
28 sources to Port Curtis, including the Fitzroy River. Most dissolved metal concentrations
29 were significantly elevated in Port Curtis compared to the concentrations measured in the
30 adjacent coastal waters. Dissolved cadmium did not exhibit any discernible spatial trend,
31 with concentrations measured in Port Curtis in the range <1-38 ng/L. Dissolved copper and
32 zinc concentrations ranged from <19 to 800 and <31 to 580 ng/L respectively, and maxima
33 were observed in inner harbour waters adjacent to the southern entrance to the Narrows
34 and in close proximity to anthropogenic sources. Dissolved nickel concentrations were
35 measured in the range of 110 to 900 ng/L, and exhibited a maximum concentration in the
36 central to northern Narrows, in an area that was not adjacent to anthropogenic sources.
37 Dissolved manganese concentration maxima were measured in close proximity to the
38 dissolved nickel maxima. Measurements of marginally lower pH and higher salinities in
39 the inner harbour and Narrows may partially explain the trends in metal concentrations. By
40 contrast, dissolved metal concentrations in oceanic waters adjacent to Port Curtis were low
41 and comparable to those measured in the open Pacific Ocean and uncontaminated coastal
42 sites around Australia. It appears likely that the elevated dissolved metal concentrations
43 measured in Port Curtis and the Narrows were not caused by a single point source, and are
44 the result of several factors, including extensive industrial activity around the foreshore,
45 fluxes from sediment-porewater, low flushing durations, lower water pH, and possibly
46 episodic inputs from the flooding of adjacent rivers.

47

48

49

50

51 **Additional keywords:** baseline study, spatial trends, copper, manganese, nickel, and zinc

52

53

54 **Introduction**

55 Port Curtis estuary is situated on the eastern coast of central Queensland, Australia,
56 adjacent to the city of Gladstone and the World Heritage-listed Great Barrier Reef Marine
57 Park (GBRMP) (Figure 1). Substantial industrialisation has occurred in the region over the
58 last century, which has resulted in increased urbanisation, industrialisation, and shipping
59 activities in the area. Port Curtis is one of Australia's largest ports for the export of raw
60 materials such as coal. Its foreshore is lined with a number of heavy industries, including
61 the world's largest alumina refinery, aluminium smelters, a 1680-megawatt coal-fired
62 power station, and several chemical plants (Jones *et al.*, 2005).

63 A recent screening-level risk assessment of metals in the Port Curtis estuary indicated
64 that dissolved concentrations of metals in the waters were in the low- or sub- $\mu\text{g/L}$ range
65 and generally below levels of regulatory concern in Australia (ANZECC/ARMCANZ,
66 2000; Jones *et al.*, 2005). However, the concentrations of many metals were elevated
67 above background levels, and local seagrass, oyster, and mud whelk species were enriched
68 in one or more of the metals cadmium, copper, nickel, and zinc (Andersen and Norton,
69 2001; Andersen, 2004; Jones *et al.*, 2005). A shell disease in local mud crabs has also been
70 shown to be associated with copper enrichment (Andersen and Norton, 2001). The
71 pathways for metal enrichment were unclear, and further investigation of metal sources
72 was recommended to improve the information available for environmental planning and
73 decision making processes. Surprisingly, there are few data on trace metal concentrations
74 in the adjacent Great Barrier Reef Marine Park (GBRMP), despite the acknowledged
75 ecological significance of the location and the increased risk of contamination from
76 anthropogenic activities in the region.

77 The aim of this study was to investigate trace metal behaviour and sources in Port
78 Curtis and surrounding waters by characterising the spatial distributions of the cadmium,
79 copper, nickel, and zinc. Two sampling surveys were undertaken, first along axial transects
80 extending away from possible point sources within the harbour and through the Narrows,
81 and then, following analysis of results, targeted sampling at selected sites (Figure 1). This
82 included a repeat axial transect through Port Curtis, and temporal sampling at two selected
83 locations having elevated metal concentrations. Laboratory experiments using sediment
84 from the Narrows were conducted to investigate the effects of pH on metal partitioning
85 between dissolved and particulate phases in sediment re-suspension and digestion tests.

86

87 **Materials and methods**

88 *Study area*

89 In Port Curtis, the major component estuaries, the Calliope and Boyne Rivers, the
90 Narrows and Auckland Creek merge into a naturally sheltered 30 km long deepwater
91 harbour protected by Curtis Island and Facing Island (Figure 1). The main entrances that
92 the estuary has with the ocean are located either side of Facing Island, although the estuary
93 is also connected to the ocean at Keppel Bay in the north-west via the Narrows. Most
94 samples taken during this study were divided into one of either five zones, in order to make
95 a statistical comparison of metal concentrations. These included Port Curtis (PC) and the
96 Narrows (TN), and the Keppel Bay (KB), Facing Island (FI), and Rodd's Bay (RB)
97 offshore sampling zones. The Keppel Bay, Facing Island, and Rodd's Bay offshore zones
98 included sites from each respective transect, which were greater than 10 km from the
99 mainland.

100 *Water sample collection and treatment*

101 Two water sampling surveys were conducted during December 2003 (axial transect
102 survey) and 2004 (targeted and temporal sampling survey). The axial transect sampling
103 was conducted in waters through the Port Curtis estuary and extending to surrounding
104 coastal waters to identify trends in metal concentrations that may indicate possible metal
105 sources and also characterise background trace metal concentrations in the adjacent coastal
106 waters (Figure 1). Water samples were collected from 49 sites approximately 4 km apart,
107 along 4 transects: (i) Keppel Bay through the Narrows and Port Curtis towards Rodd's Bay
108 (samples A1.1 to A1.21), with most sites along this transect included in either the Port
109 Curtis (PC) or the Narrows (TN) sample zones, (ii) continuing transect in point (i) east into
110 Rodd's Bay, and then north-east into offshore waters (A2.1 to A2.8), with all sites included
111 in the Rodd's Bay offshore area (RB), (iii) north-east from the seaward side of Facing
112 Island into offshore waters (A3.1 to A3.10), with all but one site included in the Facing
113 Island offshore area (FI), (iv) north east from the Keppel Bay into waters beyond Great
114 Keppel Island (A2.1 to A2.10), with all but one site included in the Keppel Bay offshore
115 area (KB). The offshore transects extended to approximately 45 km from the shore.

116 Targeted, transect, and temporal sampling was conducted during December 2004 in
117 waters of Port Curtis, the Narrows, and the Fitzroy Estuary to aid the identification of
118 metal sources. Water samples were collected from 51 sites and included: (i) an

119 approximate repeat of the Survey 1 transect (A1 series) through Port Curtis (PC) and the
120 Narrows (TN) (B1.1 to B1.15), (ii) targeted sampling in selected inlets and creeks of Port
121 Curtis and the Narrows (B2.1 to B2.9), (iii) targeted sampling of larger inlets in Keppel
122 Bay between the entrance of the Fitzroy estuary and the northern entrance of the Narrows
123 (B3.1 to B3.5), and the Fitzroy estuary and River (B4.1 to B4.8). Temporal sampling was
124 also undertaken at a location approximately 3 km south of Ramsay's Crossing (December
125 7-8, 2004, site B1.6) and near Fisherman's Landing (December 8-9, 2004, site B2.4 - 5) at
126 approximately hourly intervals for 7 h, and after 24 h.

127 To minimise sample contamination, all water samples were taken from the front of the
128 boat while the boat was gently cruising forward into the current. On the vessel, care was
129 taken to ensure that contamination was minimised at all times by ensuring that staff
130 handling the samplers and sample bottles wore powder-free disposable vinyl gloves.
131 Samples were taken from surface water at a single depth of 0.5 m because the waters of
132 Port Curtis are considered well mixed (Herzfeld *et al.*, 2003).
133 Low-density polyethylene (Nalgene) bottles were used for water sampling in the field and
134 for holding filtrates in the laboratory. The bottles were cleaned rigorously before use using
135 a three-stage sequential washing procedure, involving sequential soaking in 2% (v/v)
136 Extran 300 detergent, 10% (v/v) high purity nitric acid (Merck Tracepur), and 1% high
137 purity nitric acid (Merck Tracepur) and five rinses with deionised water (18 M Ω ·cm, Milli-
138 Q, Millipore) between different cleaning agents (Apte and Day, 1998). The bottles were
139 placed in two zip-lock bags during transportation to the field. At each site the samples were
140 collected by placing bottles into a holder installed on a purpose-built Perspex pole-sampler
141 and using ultratrace sampling techniques (Ahlers *et al.* 1990; Apte and Day, 1998). During
142 water sampling, for purposes of quality assurance and quality control (QA/QC), at least
143 10% site duplicates and 10% field blanks (air blanks from site filled with deionised water
144 in the laboratory) were collected (USEPA 1996; Apte *et al.*, 1998).

145 Following collection, samples were stored in ice-filled cooler boxes until return to the
146 laboratory and were generally filtered within 8 h, and always filtered within 24 h of
147 collection. Filtrations were undertaken using polycarbonate filtration units (Sartorius)
148 which were acid-washed (10% HNO₃, Tracepur, 24 h), and fitted with pre-weighed 0.45
149 μ m membrane filters (50 mm, type HA, Millipore) that were rinsed with 50 mL of 10%
150 (v/v) HNO₃ (Merck, Tracepur) followed by two 100 mL volumes of deionised water before
151 use (Hatje *et al.*, 2001; Hatje *et al.*, 2003; Teasdale *et al.*, 2003).

152 The initial 50 mL of filtrate was used to rinse the 0.5 L acid-washed bottles (low-
153 density polyethylene, Nalgene) and discarded, and the remaining filtrate was then
154 transferred into these bottles. The filtrates were preserved by addition of concentrated
155 HNO₃ (Merck Suprapur, 2 mL/L final concentration) in a class-100 clean laboratory. The
156 filtration rig and filter were rinsed with deionised water and the suspended particulate
157 matter (SPM) concentration was determined by drying and re-weighing the material
158 retained on the filters. The total particulate metal concentrations were determined by
159 digesting the dry material and filter in 120 mL polycarbonate vials according to the aqua-
160 regia digestion method described by Simpson *et al.* (2004), and diluting with deionised
161 water (18 MΩ·cm, Milli-Q, Millipore) to a 10% HNO₃ matrix.

162 The salinity of waters collected in Survey 1 was determined using the micro-scale
163 chlorinity titration method of Grasshoff *et al.* (1983). A water quality analyser multi-probe
164 (YSI 6600) was used in Survey 2 to determine water temperature, salinity, pH, and
165 dissolved oxygen concentrations, which was calibrated immediately before use following
166 the manufacturer's instructions. Both methods used to measure salinity were cross checked
167 using a laboratory seawater sample of known salinity.

168 *Sediment manipulations used to investigate metal partitioning*

169 At low tide, sediment samples (~ 1 kg) were collected from the upper 10 cm of
170 sediment at intertidal sites in the mid- and southern-Narrows to investigate metal
171 partitioning in the laboratory. The sediment was collected with an acid-washed Teflon
172 spatula and placed into double Zip-lock bags with the air squeezed out, and transferred to
173 the laboratory in ice boxes. In the laboratory, the sediment was stored in a refrigerator (2-5
174 °C) for no more than two weeks before use to minimise physico-chemical changes. An
175 aqua-regia digestion of these sediments was performed to measure particulate-bound
176 metals at the different locations (Simpson *et al.*, 2004).

177 The relationship between water pH and metal mobilisation from sediment-bound
178 phases was investigated using sediments collected from the mid-Narrows and waters of
179 pH 4 to 9. The sediment was homogenised with a clean plastic spatula in a nitrogen-filled
180 glove box. Volumes of 0.5 L of filtered seawater (Stanwell Park beach, NSW) were
181 adjusted to pre-determined pHs, by spiking with 1-10% NaOH or 1-10% HNO₃, so that the
182 overlying water was at the desired pH 24 h after mixing the sediment with the water. The
183 seawater was added to wet sediments 10 g/L (dry weight), shaken vigorously for 30 sec,

184 and allowed to stand at room temperature for 24 h. After 24 h a sub-sample of water was
185 taken to measure pH, and the remaining water was filtered and acidified to 2 mL HNO₃/L
186 (Merck, Tracepur), and retained for measurement of dissolved metals. A large water pH
187 range and high sediment-to-water ratio were chosen to investigate metal mobilisation to
188 reflect the conditions that may occur in areas such as sediment porewaters, organism
189 burrows, salt marshes, and areas with hotspots of acid-sulfate soils.

190 *Chemical analyses*

191 The metal concentrations in filtered samples and particulate digests were measured
192 using inductively coupled plasma-atomic emission spectrometry (ICP-AES) (Spectroflame
193 EOP, Spectro Analytical Instruments) calibrated with matrix-matched standards (QCD
194 Analysts, Eaglewood, FL, USA). As concentrations of dissolved cadmium, copper, nickel,
195 and zinc were below the detection limit of the ICP-AES in the filtered seawater samples, a
196 dithiocarbamate complexation/solvent extraction method (Hatje *et al.*, 2003) was used to
197 concentrate metals and remove salt prior to analysis. The extraction solution (modified
198 from Magnusson and Westerlund, 1981) combined sodium bicarbonate buffer, ammonium
199 pyrrolidine dithiocarbamate (APDC) and diethyldithiocarbamic acid (DDC) complexing
200 reagents, which were extracted into 1,1,1-trichloroethane in place of Freon (1,1,2-
201 trichlorotrifluoroethane) (Apte and Gunn 1987; Hatje *et al.*, 2003). The metals were back-
202 extracted into 1 mL of concentrated nitric acid (Tracepur), diluted to a final volume of 10
203 mL with deionised water (18 MΩ·cm, Milli-Q, Millipore), and analysed by graphite
204 furnace-atomic absorption spectrometry (GF-AAS) (Perkin Elmer, 4100ZL) using Zeeman
205 effect background correction and operating conditions recommended by the manufacturer.

206 For purposes of quality assurance/quality control (QA/QC), method blanks, method
207 duplicates and spike recoveries were performed on at least 10% of the filtered samples.
208 The certified reference material (CRM) seawater (CASS-4) and particulate (PACS-2)
209 (National Research Council of Canada) were processed as part of the routine quality
210 control procedures.

211 *Data analysis*

212 Differences in the mean concentrations of dissolved and suspended solid-bound
213 metals, pH, salinity, and suspended solids measured in the Port Curtis, the Narrows, and
214 the Keppel Bay, Facing Island, and Rodd's Bay offshore sampling zones were examined
215 using a one-way analysis of variance (ANOVA). For calculation purposes, samples that

216 had metal concentrations below the detection limit were assigned a value of half the
217 detection limit. An analysis of variance in the mean concentrations of dissolved and
218 particulate metals in Port Curtis and the Narrows between Survey 1 (December, 2003) and
219 Survey 2 (December, 2004) was also performed. Student-Newman-Keuls test (SNK) was
220 used as a post-hoc procedure to test for multiple comparisons upon means.

221 **Results**

222 *General water quality parameters*

223 The Port Curtis system was dominated by highly saline waters (Figure 2). The
224 salinities were significantly ($p < 0.05$) higher in the Narrows (mean \pm SD = 38 ± 1 PSU) and
225 Port Curtis (37 ± 1 PSU) sampling zones than in the Keppel Bay (36 ± 1 PSU), Rodd's Bay
226 (35 ± 1 PSU), and Facing Island (35 ± 1 PSU) offshore sampling zones in Survey 1. The only
227 salinities that were lower than typical oceanic levels were measured in the Calliope River
228 (22.8 PSU) and the Fitzroy River adjacent to the city of Rockhampton (< 0.2 - 11.5 PSU).

229 During the targeted sampling (Survey 2), the pH of waters in Port Curtis, the Narrows,
230 and adjacent inlets and Rivers ranged from 7.73-8.24 (Figure 3). The pH exhibited a broad
231 minima in the range pH 7.95-8.05 between the Yarwun trade waste outlet at Fisherman's
232 Landing in the inner harbour and Ramsay's Crossing in the Narrows. The lowest water
233 pHs measured in this survey occurred in the Upper Fitzroy (pH 7.73-8.12) and Calliope
234 Rivers (pH 7.95), that were the sites with the lowest salinities.

235 The concentrations of suspended particulate matter (SPM) in this study were in the
236 range < 0.5 - 89 mg/L (data not shown). They were highest in the Fitzroy River near the
237 entrance to Keppel Bay, and decreased in the order: Keppel Bay $>$ the Narrows $>$ Port
238 Curtis Harbour. The higher SPM measured in waters of the lower Fitzroy estuary and
239 Keppel Bay (15 - 89 mg/L) were the result of the outflows of the Fitzroy River, which is
240 known to have high and variable SPM concentrations (20 - 1000 mg/L) (Ford *et al.*, 2005).

241 *Dissolved metal concentrations*

242 The analytical detection limits of dissolved cadmium, copper, nickel, and zinc were 1, 19,
243 48, and 31 ng/L (parts per trillion), respectively, and were similar to other ultratrace metal
244 studies (Apte *et al.*, 1998; Mackey *et al.*, 2002). Concentrations of metals measured in the
245 reference material CASS-4 were within 86 to 99% of the certified value for each metal,
246 and spike recoveries were in the range 88-102%. The mean percentage relative standard
247 deviations (RSDs) of site and method duplicates were typically 2-15%.

248 The concentrations of dissolved metals measured in the two surveys are shown in Tables 1
249 and 2, and also in Figures 4 and 5. The metal concentrations were generally lowest at
250 offshore sites, particularly north-east of Facing Island, where the concentrations ranged
251 from <1-1.5, <19-68, 120-160, and <31-130 ng/L, for Cd, Cu, Ni, and Zn, respectively.
252 There were distinct gradients in trace metal concentrations with the highest concentrations
253 being measured closest to Port Curtis and the Narrows.

254 The dissolved cadmium concentrations measured in the waters of Port Curtis, the
255 Narrows, and the multiple inlets between the Fitzroy River and the northern mouth of the
256 Narrows were in the range <1-38 ng/L, and did not exhibit an observable spatial trend
257 (Table 1). In Survey 1, the mean dissolved cadmium concentrations measured in Port
258 Curtis (7 ± 5 ng/L) and the Narrows (4 ± 1 ng/L) were significantly ($p<0.05$) higher than
259 those measured in the offshore sampling zones (Figure 1) of Keppel Bay (2 ± 1 ng/L),
260 Rodd's Bay (3 ± 3 ng/L), and Facing Island (1 ± 0 ng/L) (Table 2).

261 Dissolved copper and nickel concentrations ranged between 270-740 and 120-910
262 ng/L, respectively (Table 1 and 2, and Figures 5 and 6). In Survey 2, the concentrations of
263 dissolved copper and nickel measured in inlets adjacent to the Narrows were similar to
264 those in the Narrows, which may indicate there was not a single point-source of copper and
265 nickel in these inlets at the time of this survey. Similar trends in dissolved copper and
266 nickel were measured in Port Curtis Harbour and the Narrows in Surveys 1 and 2. The
267 mean dissolved copper concentrations measured in Port Curtis in surveys 1 and 2 (530 ± 80
268 ng/L and 500 ± 200 ng/L, respectively) and the Narrows (560 ± 40 ng/L and 500 ± 100 ng/L,
269 respectively) were not significantly ($p<0.05$) different. However, the mean dissolved nickel
270 concentrations measured in the Narrows in surveys 1 and 2 (730 ± 140 ng/L and 600 ± 200
271 ng/L) were nearly twice the concentrations found in Port Curtis (350 ± 60 ng/L and
272 300 ± 100 ng/L).

273 In Survey 1, the dissolved copper concentrations measured in the waters of Port Curtis
274 and the Narrows were above 500 ng/L over a broad area, and exhibited a maximum of 640
275 ng/L in the southern Narrows. In Survey 2, the dissolved copper concentration measured
276 along this transect exhibited a more pronounced trend, with a maximum concentration of
277 650 ng/L, in the area of the inner-harbour between Fisherman's Landing and the southern
278 entrance to the Narrows (Figures 4 and 5).

279 Dissolved nickel and manganese measured along the transect through Port Curtis and
280 the Narrows, exhibited similar pronounced concentration maxima in both surveys (Figures
281 4 and 5). The peak dissolved nickel concentrations measured along this transect were 910
282 ng/L (situated approximately 6 km north of Ramsay's Crossing) and 800 ng/L (situated
283 approximately 2 km south of Ramsay's Crossing) during Surveys 1 and 2, respectively.
284 Correspondingly, the peak dissolved manganese concentrations in Surveys 1 and 2 were
285 7000 (situated approximately 2 km north of Ramsay's Crossing) and 6400 ng/L (situated
286 approximately 2 km south of Ramsay's Crossing), respectively. The dissolved manganese
287 concentration measured at one location in the Targinie inlet in the southern Narrows was
288 19000 ng/L, which was nearly three times higher than the maximum concentration
289 measured in the Narrows (6400 ng/L).

290 Dissolved zinc concentrations in Port Curtis Harbour and the Narrows ranged from 60-
291 310 ng/L (Tables 1 and 2, and Figures 4 and 5). The dissolved zinc concentration was
292 broadly elevated in waters between the outer harbour of Port Curtis and the southern
293 entrance to the Narrows in Surveys 1 and 2 (Figures 4 and 5). There was no significant
294 difference in the mean concentrations of dissolved zinc measured in Port Curtis (170 ± 40
295 ng/L) and the Narrows (110 ± 30 ng/L) during Survey 1. However, the mean dissolved zinc
296 concentration measured in Port Curtis (170 ± 90 ng/L) was significantly ($p < 0.05$) higher
297 than the Narrows (100 ± 40 ng/L) during the Survey 2. In Survey 2, the concentrations of
298 dissolved zinc measured in the Narrows and in adjacent inlets were similar, likely
299 indicating there was not a single point-source of zinc in these inlets.

300 The dissolved copper and nickel measured in Survey 1 exhibited similar spatial trends
301 between offshore sampling zones, with the Keppel Bay zone having significantly higher
302 ($p < 0.05$) dissolved concentrations than the Rodd's Bay zone, and both of these zones had
303 significantly higher ($p < 0.05$) dissolved concentrations than the Facing Island offshore
304 zone. The higher mean dissolved copper and nickel concentrations measured in the Keppel
305 Bay offshore sampling zone is probably due to water inputs from the Fitzroy River. The
306 higher mean dissolved copper, nickel, and zinc measured in the Rodd's Bay offshore
307 sampling zone than the Facing Island offshore sampling zone is likely due to tidal
308 movement of waters in Port Curtis towards Rodd's Bay.

309 The site sampled in Boat Creek was in close proximity to the Clinton Coal Wharf and
310 Fisherman's Landing in the inner harbour, and had marginally higher dissolved copper
311 (770 ng/L) and the Calliope River had marginally higher dissolved copper (670-730 ng/L)

312 and zinc (340-500 ng/L) than the maximum concentration of these metals measured in the
313 transect through waters of Port Curtis and the Narrows (Table 2). The highest
314 concentrations of dissolved copper, nickel, and zinc were measured in the Fitzroy River
315 adjacent to the city of Rockhampton during Survey 2, and ranged from 1200-1410, 1430-
316 1760, and 140-580 ng/L, respectively (Table 2 and Figure 5), possibly due to
317 anthropogenic sources from the city and surrounding agriculture and geological sources
318 (such as derelict mines in the region) (Jones 2000; Jones *et al.*, 2002; Moss and Costanzo,
319 1998; Rolfe *et al.*, 2004). The concentrations of dissolved copper, nickel, and zinc in the
320 lower estuary near the entrance to Keppel Bay were lower than upstream and ranged from
321 650-690, 980-1010, and 100-140 ng/L, respectively, probably due to dilution from tidal
322 mixing (Baeyens *et al.*, 2005), and scavenging of dissolved metals by SPM in the lower
323 estuary (Ford *et al.*, 2005; Sanudo-Wilhelmy *et al.*, 1996; Teasdale *et al.*, 2003).

324 *Metals associated with suspended particulate matter*

325 The concentrations of Al, Fe, and Mn bound to suspended particulate matter (SPM)
326 measured in waters along the transect through Port Curtis and the Narrows ranged from
327 4400-9600 µg/g, 7050-16200 µg/g, and 190-950 µg/g, respectively (Table 1 and 2). The
328 concentrations of suspended aluminium and iron (SPM-Al and SPM-Fe) did not exhibit a
329 strong trend in Port Curtis and the Narrows, although SPM-Mn exhibited a broad
330 concentration maximum in the inner harbour and the southern Narrows (Figure 4). The
331 SPM-Cu and SPM-Zn concentrations were in the range of <3-50 µg/g and <3-230 µg/g,
332 and did not exhibit strong trends in either survey. The SPM-Cd and SPM-Ni were below
333 the analytical detection limit. The concentrations of SPM-bound metals were generally
334 lower in the offshore sampling zones and often below detection limits.

335 *The release of sediment-bound metals with changes in seawater pH*

336 Results from aqua-regia metal digests of benthic sediments collected in Port Curtis and
337 the Narrows indicated the central Narrows had significantly ($p < 0.05$) higher concentrations
338 of particulate copper and nickel than the mid-harbour. Concentrations in the central
339 Narrows were approximately double those in the mid-harbour (data not shown). The
340 concentrations of dissolved copper, nickel, and zinc released from the sediments re-
341 suspended in seawater of varying pH are shown in Figure 6. As expected, the highest
342 concentrations of dissolved copper, nickel, and zinc were measured following resuspension
343 at pH 4 (Figure 6 A). The release of metals in resuspension tests in the pH 7.75-8.25 range
344 (Figure 6 B) were more representative of waters in Port Curtis and the Narrows (pH 7.95-

345 8.20), although lower pH's may occur in the mud flats adjacent to the Narrows. Dissolved
346 nickel and zinc increased by 58% and 34%, respectively, as the pH decreased from pH
347 8.25 to pH 7.75. In contrast, dissolved copper decreased as pH decreased from pH 8.25 to
348 pH 7.75, and the copper concentration only increased significantly ($p < 0.05$) above that
349 measured for pH 8.25 waters in re-suspension treatments that employed waters with a pH
350 equal to or less than pH 5.

351 **Discussion**

352 Port Curtis is different to many estuaries, as it is connected to the ocean at multiple
353 locations (the Narrows and openings between Curtis and Facing Islands), and has highly
354 variable, spasmodic inputs of freshwater due to the wet season/dry season climate
355 dynamics of the catchment area (Ford *et al.*, 2005; Jones *et al.*, 2005). The hydrodynamics
356 of water movement and the general water quality parameters (pH, salinity, SPM) in Port
357 Curtis, the Narrows, and the Fitzroy River may have a significant influence on the
358 concentrations of metals. A recent hydrodynamic model predicted that water in Port Curtis
359 undergoes tidal movement over large distances, but little net displacement occurs, as the
360 water returns close to its original location (Herzfeld *et al.*, 2003). The model proposed a
361 flushing duration in Port Curtis of 22-26 days, although poor tidal mixing in some
362 protected inlets and bays of Port Curtis and the Narrows suggests that these waters have
363 longer flushing durations. The model did not include all of the Narrows and it was not
364 entirely clear whether hydrodynamic forces predominantly direct waters from Port Curtis
365 north through the Narrows into Keppel Bay or from Keppel Bay south through the
366 Narrows into Port Curtis. The direction of water flow in the Narrows fluctuates (north-west
367 or south-east) through the tidal cycle making it difficult to predict metal dispersion from
368 potential sources, such as the point sources from the Yarwun trade waste outlet and the
369 Calliope Power Station, or diffuse sources from urban runoff and leaching from sediments.

370 The broad pH minimum measured between the inner harbour and Ramsay's Crossing
371 in the Narrows was in agreement with past pH measurements (Apte *et al.*, 2006). The
372 position of the pH minimum was in close proximity to the entrance of the Calliope River
373 and the Yarwun trade waste outlet, and both these areas may potentially contribute water
374 of lower pH into Port Curtis. The lack of a decrease in salinity indicates that freshwater
375 inputs were minimal during the sampling period. The discharges from the Yarwun Trade
376 Waste outlet were not believed to be acidic (Leonie Anderson, Central Queensland
377 University, personal communication), so the relatively small outflows (110 kg Cu per year,

378 410 kg Zn per year) of these waters were unlikely to have a significant influence on the
379 water pH in Port Curtis Harbour (National Pollution Inventory, 2008).

380 The foreshore of the Narrows is predominantly lined with mangrove mud flats and salt
381 marshes. The sediment surface area that comes into contact with water is relatively high
382 because of the large tidal range (4-5 m, WBM Oceanics 1992) and flatness of the adjacent
383 sediments. The water depth is also quite shallow in the Narrows and there is a relatively high ratio
384 between the surface area of sediment that comes into contact with water and the volume of water.
385 Several studies have reported lower pHs in waters lined with mangrove foreshores due to
386 factors that result in the release of H^+ into waters including, mangrove systems having
387 faster rates of diagenesis and organism-facilitated aerobic oxidation of organic matter
388 because of the larger volumes of organic carbon (Clark *et al.*, 1998; Kristensen, 2000;
389 Otero and Macias, 2002; Van Cappellen and Wang, 1996). The lower pH of waters in the
390 Narrows may also be generated by adjacent acid-sulfate soils (ASS), which are common
391 along the east coast of Queensland and northern NSW, with hotspots of ASS activity
392 identified in the vicinity of the Narrows (Powell and Martens, 2005; Ross, 2002). Mud
393 flats and salt marshes often have higher water temperatures than the adjacent water column
394 that may allow greater rates of chemically driven oxidation (e.g. air oxidation of sulfides),
395 resulting in lower water pH (Otero and Macias, 2002).

396 The salinity of the waters of Port Curtis and the Narrows was in the range 33 to 39
397 PSU, and did not decrease at sites in the inner harbour and the Narrows where several
398 creeks and rivers may potentially input fresh water into the system. On the contrary, the
399 salinity was higher in the estuary near the outflows of these inlets than the offshore waters.
400 Typical estuarine waters have salinities that increase as the estuary approaches the sea and
401 are approximately 35 PSU at the seawater end-member (Hatje *et al.*, 2003; Sanudo-
402 Wilhelmy *et al.*, 1996). The chemical analysis of major ions that may contribute to salinity
403 indicated that the spatial trends in dissolved Ca, Mg, K, and Na were the same as salinity
404 (data not shown), indicating no particular species were responsible for the increased
405 salinity.

406 The mangrove mud flats and salt marshes along the foreshore of the Narrows and inner
407 harbour are subject to evaporation (Saenger, 1995). It is likely that the hypersaline
408 conditions measured in the Narrows and the inner harbour were the result of evaporative
409 water losses. Average daily rainfall measurements in the catchment of Port Curtis
410 (Calliope, Mount Larcon, and Gladstone Airport stations) in the thirty days before and

411 during the two surveys were <5 mm, and most days had <1 mm, indicating low freshwater
412 inputs from rainfall into the estuary (Australian Bureau of Meteorology, 2008).
413 Hypersaline conditions may be typical of many tropical Australian estuaries and may be
414 assisted by evapotranspiration occurring in saline tolerant vegetation (water removed by
415 plants leading to concentration of salt in estuarine system) (Largier *et al.*, 1997; Ridd and
416 Stieglitz, 2002; Wolanski, 1986).

417 Considering the ecological importance of the World Heritage listed GBRMP, there
418 was surprisingly few data on the concentrations of trace metals in these waters. The
419 increased industrial and shipping activity in this region of Australia also means that it is
420 important to have knowledge of natural (baseline) concentrations for future comparison. A
421 study by Denton and Burdon-Jones, (1986), that used Chelex-100 extraction and analysis
422 of cadmium, copper, nickel, and zinc in waters of the GBRMP reported concentrations of
423 <10-60, 110-240, 60-160 and 30-350 ng/L, respectively, but the results had high replicate
424 variability and the method had poor extraction efficiencies.

425 The lowest concentrations of trace metals were measured in the Facing Island offshore
426 sampling zone. This zone was considered the most representative of regional coastal water
427 concentrations because Facing Island restricts water flow and thus, metal transfer from the
428 harbour into these waters. The mean offshore concentrations of trace metals measured in
429 this study were similar to those measured in offshore sites in Northern and Eastern
430 Australia and the Western Pacific, and may be considered metal depleted relative to waters
431 in the northern Atlantic which receive significantly greater inputs from atmospheric and
432 riverine sources (Apte *et al.*, 1998; Bruland *et al.*, 1994; Kremling and Pohl, 1989; Jickells,
433 1995; Mackey *et al.*, 2002; Williams *et al.*, 1998) (Table 3). The similar concentrations of
434 trace metal concentrations measured in offshore waters around Northern and Eastern
435 Australia, indicate that the offshore metal concentrations measured in the current study are
436 likely to be representative of those found in the GBRMP. The current study achieved
437 relatively low detection limits and excellent quality control results, and to our knowledge,
438 is the most recent and accurate data set of trace metal concentrations representative of
439 those found in waters of the GBRMP.

440 The dissolved metal concentrations in Port Curtis did not exceed the current Australia
441 and New Zealand water quality guideline trigger values for ecosystem protection
442 (ANZECC/ARMCANZ 2000) (Table 3). The highest concentrations of dissolved Cd, Cu,
443 Ni, and Zn measured in Port Curtis and the Narrows were 40, 740, 910, and 310 ng/L,

444 respectively. These concentrations are at the lower end of the concentration range of these
445 metals measured in industrialised harbours around the world, indicating that Port Curtis
446 was not contaminated to the same degree (Table 3). The level of metal enrichment in Port
447 Curtis relative to regional concentrations was investigated by comparing mean metal
448 concentrations from Port Curtis to those from the Facing Island offshore sampling zone.
449 The ratios of Cd, Cu, Ni, and Zn measured in Port Curtis compared to the Facing Island
450 offshore sampling zone were 2.4, 6.9, 3.0, and 1.7, respectively. The ratio of dissolved
451 copper in Port Curtis compared to offshore sampling zones was larger than the other
452 metals, indicating greater enrichment in the harbour. The copper concentrations were also
453 higher than those measured in previous studies of pristine estuaries, such as Bathurst
454 Harbour, Tasmania or Sea lochs in Scotland (Hall *et al.*, 1996; Mackey *et al.*, 1996).

455 The presence of a strong spatial separation of concentration maxima for dissolved
456 copper, manganese, nickel, and zinc in this study is likely to indicate that there were
457 multiple metal sources or parameters influencing metal concentrations. The dissolved
458 copper and zinc concentration maxima were located in the inner harbour and southern
459 Narrows. The foreshore of these waters was lined with urban, commercial, and industrial
460 centres, which means there were likely to be multiple sources of metals (Comber *et al.*,
461 1995; Hatje *et al.*, 2003; Laslett and Balls, 1995; Owens and Balls, 1997). For instance,
462 the National Pollution Inventory website of the Commonwealth of Australia reports that
463 copper and zinc are discharged into waters of the Calliope River and the Yarwun trade
464 waste outlet at Fisherman's Landing, by industries such as the Gladstone power station and
465 sewage works (National Pollution Inventory, 2008). The leachate from reclaimed land
466 containing fly ash from the nearby power station and antifouling paints from boats may
467 also contribute to the higher concentrations of dissolved copper and zinc in the inner
468 harbour (Jones *et al.*, 2005; Warnken *et al.*, 2004). The dissolved concentrations of Cu, Ni,
469 and Zn exhibited low temporal variation at the Ramsay's Crossing and Fisherman's
470 Landing sites over the 24-h duration of the sampling period (data not shown).

471 In contrast, the dissolved nickel concentration maximum was located in the central to
472 northern Narrows and is not in close proximity to anthropogenic sources. The dissolved
473 nickel concentration maximum of each survey was located within close proximity (2-4 km)
474 to the dissolved manganese maxima, and strong correlations of 0.67 ($p < 0.001$) and 0.92
475 ($p < 0.0001$) were measured in Surveys 1 and 2, respectively (Figure 7). The close
476 association of nickel and manganese has been shown in several previous studies, indicating

477 that the same process may be responsible for the release of dissolved manganese and nickel
478 from soils and sediments (Hatje *et al.*, 2003; Laslett and Balls, 1995; Sanudo-Wilhelmy
479 and Gill, 1999).

480 Many trace metals are associated with manganese and iron (hydr)oxides in sediments.
481 The chemically or bacterially driven reduction of manganese and iron (hydr)oxides may
482 result in the release of manganese, iron, and trace metals into the dissolved phase, although
483 iron would rapidly re-oxidise and be removed from the dissolved phase upon entering the
484 overlying water (Laslett and Balls, 1995; Sanudo-Wilhelmy and Gill, 1999; Simpson and
485 Batley, 2003; Zhang *et al.*, 2002). The reduction of manganese and iron hydr(oxides) is a
486 natural process that has been reported to be enhanced in warmer temperatures, when
487 biological activity is increased and oxygen concentrations are generally lower (Laslett and
488 Balls, 1995; Otero and Macias, 2002). Natural geochemical weathering and leaching
489 processes may also release trace metals including nickel from particulate-bound forms into
490 the dissolved phase, and these processes are likely to be facilitated in the Narrows because
491 of ASS hotspots. Processes that cause the release of nickel from sediments into the
492 dissolved phase are likely to have a greater influence on sources of nickel in the Narrows
493 than Port Curtis, because of the higher particulate nickel in the sediments of the Narrows
494 than the mid-harbour measured previously and in the current study (Vincente-Beckett and
495 Shearer, 2005).

496 The water pH may also influence the behaviour of nickel in Port Curtis and the
497 Narrows, as shown by the increase in dissolved nickel with decreasing water pH in re-
498 suspension experiments (Figure 6 B). However, the results from these experiments were
499 indicative of nickel release from SPM in highly turbid conditions or in the benthic water
500 layer, as the SPM concentrations were approximately 3 orders of magnitude higher than
501 the SPM measured in the Narrows. Therefore, the results are expected to greatly
502 overestimate the nickel released from SPM due to the decreased pH of water in the
503 overlying water of the Narrows, but may be representative of metal release in sediment
504 porewater and the benthic water layer of mudflats in this area.

505 The Narrows was unlikely to act as a significant source of metals to Port Curtis
506 Harbour because of the difference in concentrations was not particularly great, and the
507 relatively small water volume in the Narrows compared with the harbour. The Calliope
508 River may also contribute natural and/or anthropogenic metals to the waters of this area of
509 the harbour, although rainfall was low before and during both surveys and it is unlikely

510 that outflows from this river had a significant influence on the concentrations during this
511 study. The sites sampled in rivers, creeks, and inlets adjacent to Port Curtis and the
512 Narrows had similar metal concentrations to those measured along adjacent transects,
513 indicating that these waters were probably not acting as substantial metal sources at the
514 time of the survey. It is unlikely that atmospheric deposition contributes significant inputs
515 of metals to the waters sampled, as this would be expected to cause higher concentrations
516 over a broad area (Jickells, 1995; Kremling and Pohl, 1989).

517 **Conclusions**

518 Overall, the waters of Port Curtis and the Narrows had a marginally lower water pH and
519 higher salinity, and had elevated metal concentrations compared to the surrounding coastal
520 waters. The concentrations measured in the GBRMP and adjacent waters were similar to
521 concentrations measured in regional waters on the north east coast of Australia. Strong
522 spatial separations of maxima for dissolved copper, manganese, nickel, and zinc were
523 measured, indicating that there was not one particular source responsible for the elevated
524 metal concentrations. Copper and zinc are likely to originate from anthropogenic sources,
525 whereas, nickel and manganese are likely to be released by natural processes, such as
526 reduction of manganese (hydr)oxides and leaching by the lower sediment and water pH.
527 All metal concentrations measured in waters of Port Curtis were below regulatory
528 guideline values. However, dissolved copper was significantly ($p < 0.05$) elevated above the
529 concentrations in the adjacent coastal waters and pristine estuarine systems, and may be of
530 potential concern if the concentrations in Port Curtis and the Narrows continue to increase.

531

532 **Acknowledgements**

533 The authors acknowledge the financial support from the University of Wollongong,
534 CSIRO and the Cooperative Research Centre for Coastal Zone Estuary and Waterway
535 Management. Leonie Anderson and Andrew Davis (Central Queensland University) are
536 thanked for logistical support and coordination of field work, and David Spadaro (CSIRO)
537 for assistance in the laboratory

538

539


540 **References**

- 541 Andersen, L. E., Norton, J. H. (2001). Port Curtis mud crab shell disease. Final report. Central QLD
542 University, Department of Primary Industry, Fisheries Research and Development Corporation.
- 543 Andersen, L.E. (2004). Imposex: a biological effect of TBT contamination in Port Curtis, Queensland. *Aust.*
544 *J. Ecotoxicol.* 10, 105-113.
- 545 ANZECC/ARMCANZ. (2000). Australian and New Zealand guidelines for fresh and marine water quality,
546 Australia and New Zealand Environment and Conservation Council/Agricultural and Resource
547 Management Council of Australia and New Zealand., Canberra, Australia.
- 548 Apte, S. C., and Day, G. M. (1998). Dissolved metal concentrations in the Torres Strait and Gulf of Papua.
549 *Marine Pollution Bulletin* **30**, 298-304.
- 550 Apte, S. C., and Gunn, A. M. (1987). Rapid determination of copper, nickel, lead and cadmium in small
551 samples of estuarine waters by liquid/liquid extraction and electrothermal atomic absorption
552 spectrometry. *Analytica Chimica Acta* **193**, 147-156.
- 553 Apte, S., Anderson, L., Andrewartha, J., Angel, B, Shearer, D., Simpson, S., Stauber, J., and Vicente-Beckett,
554 V. (2006). Contaminant pathways in Port Curtis. Cooperative Research Centre for Coastal Zone,
555 Estuary and Waterway Management. Technical Report No. 75.
- 556 Ashlers, W. W., Reid, M. R., Kim, J. P., and Hunter, K. A. (1990). Contamination-free sample collection and
557 handling protocols for trace elements in natural waters. *Marine and Freshwater Research* **41**, 713-
558 720.
- 559 Baeyens, W., Goeyens, M., Monteny, F., and Elskens, M. (1998). Effect of organic complexation on the
560 behaviour of dissolved cadmium, copper, and zinc in the Scheldt estuary. *Hydrobiologia* **366**, 15-43.
- 561 Baeyens, W., Leermakers, M., De Gieter, M., Nguyen, H. L., Parmentier, K., Panutrakul, S., and Elskens, M.
562 (2005). Overview of trace metal contamination in the Scheldt estuary and effect of regulatory
563 measures. *Hydrobiologia* **540**, 141-154.
- 564 Balzer, W. (1982). On the distribution of iron and manganese at the sediment/water interface:
565 thermodynamic versus kinetic control. *Geochimica et Cosmochimica. Acta* **46**, 1153-1161.
- 566 Benoit, G., Oktay-Marshall, S. D., Cantu, H. A., Hood, E. M., Coleman, C. H., Corapcioglu, M. G., and
567 Santschi, P. H. (1994). Partitioning of Cu, Pb, Ag, Zn, Fe, Al, and Mn between filter-retained
568 particles, colloids and solution in six Texas estuaries. *Marine Chemistry* **45**, 307-336.
- 569 BOM (2008). Australian bureau of meteorology. Weather station directory.
570 <http://www.bom.gov.au/climate/cdo/about/sitedata.shtml> (accessed 14/02/2008).
- 571 Bruland, K. W., Orians, K. J., and Cowen, J. P. (1994). Reactive trace metals in the stratified central North
572 Pacific. *Geochimica et Cosmochimica. Acta* **58**, 3171-3182.

- 573 Clark, M. W., McConchie, D., Lewis, D. W., and Saenger, P. (1998). Redox stratification and heavy metal
574 partitioning in *Avicennia*-dominated mangrove sediments: a geochemical model. *Chemical Geology*
575 **149**, 147-171.
- 576 Comber, S. D. W., Gunn, A. M., and Whalley, C. (1995). Comparison of the partitioning of trace metals in
577 the Humber and Mersey Estuaries. *Marine Pollution Bulletin* **30**, 851-860.
- 578 Denton, G. R. W., and Burdon-Jones, C. (1986). Trace metals in surface waters from the Great Barrier
579 Reef. *Marine Pollution Bulletin* **19**, 96-98.
- 580 Fabris, G. L., and Monahan, C. A. (1995). Characterisation of toxicants in waters from Port Phillip Bay:
581 metals. CSIRO. INRE. Port Phillip Bay Environmental Study Technical Report No. 18., Melbourne.
- 582 Ford, P., Tillman, P., Robson, B., and Webster, I. T. (2005). Organic carbon deliveries and their flow related
583 dynamics in the Fitzroy estuary. *Marine Pollution Bulletin* **51**, 119-127.
- 584 Grasshoff, K., Ehrhardt, M., and Kremling, K. (1983). Methods of seawater analysis. Second Edition, Verlag
585 Chemie, Weinheim.
- 586 Hall, I. R., Hydes, D. J., Statham, P. J., and Overnell, J. (1996). Dissolved and particulate trace metals in a
587 Scottish Sea Loch: an example of a pristine environment. *Marine Pollution Bulletin* **32**, 846-854.
- 588 Hatje, V., Apte, S.C., Hales, L.T., and Birch, G.F. (2003). Dissolved trace metal distributions in Port Jackson
589 estuary (Sydney Harbour), Australia. *Marine Pollution Bulletin* **46**, 719-730.
- 590 Hatje, V., Birch, G.F., and Hill, D.M. (2001). Spatial and temporal variability of particulate trace metals in
591 Port Jackson Estuary, Australia. *Estuarine, Coastal and Shelf Science* **53**, 63-77.
- 592 Herzfeld, M., Parslow, J., Andrewartha, J., Sakov, P., and Webster, I.T. (2004). Hydrodynamic modeling of
593 the Port Curtis region. CRC for Coastal Zone, Estuary and Waterway Management, Technical
594 Report no. 7 Brisbane.
- 595 Jickells, T. (1995). Atmospheric inputs of metals and nutrients to the oceans: their magnitude and effects.
596 *Marine Chemistry* **48**, 199-214.
- 597 Jones, M.-A., Stauber, J.L., Apte, S.C., Simpson, S.L. Vincente-Beckett, V., Johnson, R. and
598 Duivenvoorden, L. (2005). A risk assessment approach to contaminants in Port Curtis, Queensland,
599 Australia. *Marine Pollution Bulletin* **51**, 448-458.
- 600 Jones, M., Duivenvoorden, L., Choy, S., and Moss, A. (2000). National land and water resources report 3,
601 theme 7 – catchment health, Fitzroy implementation project, Queensland Department of Natural
602 Resources, Brisbane.
- 603 Kremling, K., and Pohl, C. (1989). Studies on the Spatial and Seasonal Variability of Dissolved Cadmium,
604 Copper, and Nickel in North-East Atlantic Surface Waters. *Marine Chemistry* **27**, 43-60.
- 605 Kristensen, E. (2000). Organic matter diagenesis at the oxic/anoxic interface in coastal marine sediments
606 with emphasis on the role of burrowing animals. *Hydrobiologia* **426**, 1-24.

- 607 Largier, J. L., Hollibaugh, J. T., and Smith, S. V. (1997). Seasonally hypersaline estuaries in Mediterranean-
608 climate regions. *Estuarine, Coastal and Shelf Science* **45**, 789-797.
- 609 Laslett, R. E., and Balls, P. W. (1995). The behaviour of dissolved Mn, Ni and Zn in the Forth an
610 industrialized partially mixed estuary. *Marine Chemistry* **48**, 311-328.
- 611 Laslett, R. E., and Balls, P. W. (1995). The behaviour of dissolved Mn, Ni and Zn in the Forth an
612 industrialized partially mixed estuary. *Marine Chemistry* **48**, 311-328.
- 613 Mackey, D. J., O'Sullivan, R. J., Watson, G., and Pont, D. (2002). Trace metals in the Western Pacific:
614 temporal and spatial variability in the concentrations of Cd, Cu, Mn, and Ni. *Deep-Sea Research*
615 *Part 1* **49**, 2241-2259.
- 616 Mackey, D. J., Butler, C. V., Carpenter, P. D., Higgins, H. W., O'Sullivan, R. J., and Paschke, R. B.
617 (1996). Trace elements and organic matter in a pristine environment: Bathurst Harbour, South
618 western Tasmania. *Science of the Total Environment* **191**, 137-151.
- 619 Magnusson, B., and Westerlund, S. (1981). Solvent extraction procedures combined with back-extraction for
620 trace metal determinations by atomic absorption spectrometry, *Analytica Chimica Acta* **131**, 63-72.
- 621 Moss, A., and Costanzo, S. (1998). Levels of heavy metals in sediments of Queensland rivers, estuaries, and
622 coastal waters, Environmental technical report no 20. Queensland Environment Protection
623 Authority, Brisbane.
- 624 Munksgaard, N. C., and Parry, D. L. (2001). Trace metals, arsenic and lead isotopes in dissolved and
625 particulate phases of North Australian coastal and estuarine seawater. *Marine Chemistry* **75**, 165-
626 184.
- 627 National Pollution Inventory (2009). Australian Government Department of the environment, water, heritage
628 and the arts. Tracking emissions across Australia. (<http://www.npi.gov.au>, accessed 22 March 2008)
- 629 Otero, X. L., and Macias, F. (2002). Spatial and seasonal variation in heavy metals in interstitial water of salt
630 marsh soils. *Environmental Pollution* **120**, 183-190.
- 631 Owens, R. E., and Balls, P. W. (1997). Dissolved trace metals in the Tay Estuary. *Estuarine, Coastal and*
632 *Shelf Science* **44**, 421-434.
- 633 Powell, B., and Martens, M. (2005). A review of acid sulfate soil impacts, actions and policies that impact on
634 water quality in Great Barrier Reef catchments, including a case study on remediation at East
635 Trinity. *Marine Pollution Bulletin* **51**, 149-164.
- 636 Ridd, P. V., and Stieglitz, T. (2002). Dry season salinity changes in arid estuaries fringed by mangroves and
637 saltflats. *Estuarine, Coastal and Shelf Science* **54**, 1039-1049.
- 638 Rolfe, J., Alam, K., and Windle, J. (2004). Overview of the Fitzroy basin and opportunities for offset
639 trading: Establishing the Potential for offset trading in the lower Fitzroy Rivers research reports.
640 Research Report No. 1, Central Queensland University.


- 641 Ross, D. J. (2002). Acid sulfate soils, Tannum Sands to St. Lawrence, central Queensland coast. Department
642 of National Resources and mines, Rockhampton.
- 643 Saenger, P. (1995). Ecology of Mangroves of Port Curtis: Regional Biogeography, Productivity and
644 Demography. Mangroves - a resource under threat: An issue for the central Queensland coast,
645 Central Queensland University, Gladstone.
- 646 Sanudo-Wilhelmy, S. A., Rivera-Duarte, I., and Flegal, A. R. (1996). Distribution of colloidal trace metals in
647 the San Francisco Bay estuary. *Geochimica et Cosmochimica. Acta* **60**, 4933-4944.
- 648 Sanudo-Wilhelmy, S. A., and Gill, G. A. (1999). Impact of the Clean Water Act on the levels of toxic metals
649 in urban estuaries: the Hudson River estuary revised. *Environmental Science and Technology* **33**,
650 3477-3481.
- 651 Simpson, S.L., Angel, B.M., and Jolley, D.F. (2004). Metal equilibration in laboratory-contaminated (spiked)
652 sediments used for the development whole-sediment toxicity tests. *Chemosphere* **54**, 597-609.
- 653 Simpson, S. L., and Batley, G. E. (2003). Disturbances to metal partitioning during toxicity testing of iron
654 (II)-rich estuarine pore waters and whole sediments. *Environmental Toxicology and Chemistry* **22**,
655 424-432.
- 656 Teasdale, P. R., Apte, S. C., Ford, P. W., Batley, G. E., and Khoenken, L. (2003). Geochemical cycling and
657 speciation of copper in waters and sediments of Macquarie Harbour, Western Tasmania. *Estuarine,
658 Coastal and Shelf Science* **57**, 475-487.
- 659 Van Cappellen, P., and Wang, Y. (1996). Cycling of iron and manganese in surface sediments: A general
660 theory for the coupled transport and reaction of carbon, oxygen, nitrogen, sulfur, iron, and
661 manganese. *American Journal of Science* **296**, 197-243.
- 662 Vincente-Beckett, V., Shearer, D. (2005). Contaminant sources in benthic sediments of Port Curtis.
663 Milestone report IC10.1. CRC for Coastal Zone Estuary and Waterway Management, Brisbane.
- 664 USEPA, (1996). Sampling ambient water for trace metals at EPA water quality criteria levels. Method 1669.
665 US Environmental Protection Agency, Office of Water Engineering and Analysis Division,
666 Washington, DC.
- 667 Warnken, J., Dunn, R. J. K., and Teasdale, P. R. (2004). Investigation of recreational boats as a source of
668 copper at anchorage sites using time-integrated diffusive gradients in thin films and sediment
669 measurements. *Marine Pollution Bulletin* **49**, 833-843.
- 670 Williams, M. R., Millward, G. E., Nimmo, M., and Fones, G. (1998). Fluxes of Cu, lead and Mn to the
671 North-Eastern Irish Sea: the importance of sedimental and atmospheric inputs. *Marine Pollution
672 Bulletin* **36**, 366-375.
- 673 Wolanski, E. (1986). An evaporation-driven salinity maximum zone in Australian tropical estuaries.
674 *Estuarine, Coastal and Shelf Science* **22**, 415-424.
- 675 Zhang, H., Davison, W., Mortimer, R. J. G., Krom, M. D., Hayes, P. J., and Davies, I. M. (2002). Localised
676 remobilization of metals in a marine sediment. *Science of the Total Environment* **296**, 175-187.



677


678 **Fig. 1.** Map of Port Curtis estuary and surrounding waters showing sampling sites for Survey 1
 679 (○), Survey 2 (◇), and both surveys (▲). The different areas that were compared statistically were
 680 PC = Port Curtis, TN = The Narrows, KB = Keppel Bay offshore area, FI = Facing Island offshore
 681 area, and RB = Rodd's Bay offshore area.

682



683


684 **Fig 2.** Measured salinity of Port Curtis and surrounding waters.
 685



686

687 **Fig 3.** Measured pH of Port Curtis and surrounding waters during Survey 2, where the symbols
 688 FL, SE, RC, and NE refer to the locations of Fisherman's Landing, the southern entrance to the
 689 Narrows, Ramsay's Crossing, and the northern entrance to the Narrows, respectively. The entrance to
 690 the harbour was defined as site B1.15 (Survey 2).
 691

691




692

693


694 **Fig 4.** The concentrations of dissolved copper ($\blacktriangle \triangle$), manganese ($\bullet \circ$), nickel ($\diamond \blacklozenge$), and zinc
 695 ($\blacksquare \square$) measured along the transect direct from the outer harbour of Port Curtis into the Narrows
 696 during Survey 1 ($\blacktriangle \bullet \blacklozenge \blacksquare$) and Survey 2 ($\triangle \circ \diamond \square$). The entrance to the harbour was defined as
 697 site A1.17 (Survey 1) and site B1.15 (Survey 2). The symbols FL, SEN, RC, and NEN refer to the
 698 locations of Fisherman's Landing, the southern entrance to the Narrows, Ramsay's Crossing, and
 699 the northern entrance to the Narrows, respectively.
 700

700



701


702 **Fig 5.** Dissolved copper, nickel and zinc concentrations (ng/L) in Port Curtis estuary and
 703 surrounding waters.



704

705 **Fig 6.** Dissolved (<0.45 μm) metals measured following re-suspension and 24-h storage of central
 706 Narrows (sampling site B1.5) sediment (10000 mg/L) for water pH in the range (A) pH 3.0-10.0
 707 and (B) pH 6.5-9.5.

708



709

710 **Fig 7.** The relationship between dissolved manganese and nickel measured in Port Curtis and the
 711 Narrows in Surveys 1 and 2.

712

713
714

Table 1. Dissolved (<0.45 µm) and particulate metal concentrations in Port Curtis Estuary and surrounding waters in Survey 1

Site	Area	Dissolved metals (ng/L)					Suspended particulate metals (µg/g)					
		Cd	Cu	Mn	Ni	Zn	Al	Mn	Fe	Cu	Zn	
The Narrows and Port Curtis (Keppel Bay → Rodd's Bay)												
A1.1		3.0	380	<200	460	61	7000	420	12000	13	26	
A1.2		3.5	430	<200	500	57	7000	400	12000	11	22	
A1.3	TN	Keppel Bay	6.5	520	<200	610	150	4300	240	7000	7.7	19
A1.4	TN	northern Narrows	3.5	540	<200	700	55	6800	360	9500	8.8	26
A1.5	TN	↓	5.8	580	1400	880	100	7800	460	9700	9.2	32
A1.6	TN	↓	5.8	560	3400	910	76	8400	560	11000	5.8	44
A1.7	TN	Ramsay's Crossing	4.0	550	7000	840	130	8500	640	14000	9.4	43
A1.8	TN	↓	3.5	510	5200	780	92	9600	670	16000	14	55
A1.9	TN	↓	3.0	610	2900	620	91	8400	670	13000	17	57
A1.10	TN	southern Narrows	3.0	640	1600	520	140	9300	940	15000	19	55
A1.11	PC	↓	5.0	620	1500	470	210	8400	950	14000	14	61
A1.12	PC	Fisherman's Landing	2.0	550	740	330	130	7800	750	12000	16	30
A1.13	PC	mid-harbour	15	610	1100	360	200	8700	880	15000	22	56
A1.14	PC	↓	4.5	520	620	330	130	9200	920	15000	24	46
A1.15	PC	↓	4.5	500	<200	350	160	6300	600	11000	15	29
A1.16	PC	outer-harbour	15	460	<200	310	230	4500	480	8200	10	26
A1.17	PC	↓	6.3	410	<200	280	130	8800	770	14000	20	61
A1.18		↓	15	300	<200	230	190	2200	190	3600	46	27
A1.19		↓	2.0	250	<200	200	70	6300	470	8600	14	88
A1.20		↓	3.8	190	360	200	64	2400	170	2900	3.0	34
A1.21		Rodd's Bay	3.0	220	490	190	57	n/a	n/a	n/a	n/a	n/a
Rodd's Bay → offshore												
A2.1	RB	Rodd's Bay	10	66	-	170	140	n/a	n/a	n/a	n/a	n/a
A2.2	RB	↓	1.8	84	-	140	61	6400	350	7500	<3	71
A2.3	RB	↓	3.0	70	-	120	92	2600	130	2500	16	19
A2.4	RB	↓	<1.5	69	-	150	42	500	90	600	4.2	53
A2.5	RB	↓	3.0	60	-	110	95	600	90	730	<3	44
A2.6	RB	↓	<1.5	41	-	120	68	370	50	510	<3	51
A2.7	RB	Offshore	2.0	39	-	160	140	n/a	n/a	n/a	n/a	n/a
A2.8	RB	↓	<1.5	63	-	160	41	n/a	n/a	n/a	n/a	n/a
Facing Island → offshore												
A3.1		Facing Island	<1.5	120	-	170	61	n/a	n/a	n/a	n/a	n/a
A3.2	FI	↓	1.5	68	-	150	<31	n/a	n/a	n/a	n/a	n/a
A3.3	FI	↓	1.5	38	-	150	41	n/a	n/a	n/a	n/a	n/a
A3.4	FI	↓	<1.5	51	-	120	130	230	160	400	<3	<3
A3.5	FI	↓	<1.5	19	-	140	<31	420	280	640	<3	<3
A3.6	FI	↓	<1.5	<19	-	130	37	n/a	n/a	n/a	n/a	n/a
A3.7	FI	↓	<1.5	30	-	140	67	n/a	n/a	n/a	n/a	n/a
A3.8	FI	↓	1.5	41	-	140	43	n/a	n/a	n/a	n/a	n/a
A3.9	FI	Offshore	<1.5	22	-	140	<31	n/a	n/a	n/a	n/a	n/a
A3.10	FI	↓	<1.5	35	-	160	<31	n/a	n/a	n/a	n/a	n/a
Keppel Bay → offshore												
A4.1		Keppel Bay	3.0	430	-	540	48	9300	410	14000	14	31
A4.2	KB	↓	3.0	370	-	440	64	4600	230	7100	7.5	26
A4.3	KB	↓	2.0	270	-	340	46	8000	510	12000	10	68
A4.4	KB	↓	<1.5	190	-	250	82	8300	460	11000	10	82
A4.5	KB	↓	<1.5	170	-	260	<31	7300	400	7600	7.9	80
A4.6	KB	↓	<1.5	160	-	260	<31	4200	460	4500	6.6	110
A4.7	KB	↓	<1.5	120	-	190	<31	1700	360	2300	<3	100
A4.8	KB	Offshore beyond	4.0	90	-	170	100	420	80	510	<3	18
A4.9	KB	Great Keppel	<1.5	60	-	160	<31	1700	260	2000	<3	69
A4.10	KB	Island	<1.5	70	-	180	33	n/a	n/a	n/a	n/a	n/a

715
716
717
718

Where, PC refers to Port Curtis, TN refers to The Narrows, RB refers to Rodd's Bay, FI refers to Facing Island, KB refers to Keppel Bay, "-" refers to data not measured, and "n/a" refers to SPM-bound metal concentrations that are not available because the amount of SPM recovered for digestion was too low.

719
720
721
722

723
724

Table 2. Dissolved (<0.45 µm) and particulate metal concentrations in Port Curtis estuary and surrounding waters in Survey 2

Site	Area	Dissolved metals (ng/L)					Suspended particulate metals (µg/g)					
		Cd	Cu	Mn	Ni	Zn	Al	Mn	Fe	Cu	Zn	
The Narrows and Port Curtis (Keppel Bay → the mid-harbour)												
B1.1	TN	Keppel Bay	9.6	330	490	380	76	5200	250	7800	8.8	69
B1.2	TN	northern Narrows	3.1	350	<200	390	79	-	-	-	-	-
B1.3	TN	↓	4.1	400	1700	530	61	4400	280	6200	7.3	75
B1.4	TN	↓	38	450	3100	640	89	-	-	-	-	-
B1.5	TN	Ramsay's Crossing	4.8	470	3900	700	87	4700	290	6900	8.1	75
B1.6	TN	↓	15	560	6400	800	83	-	-	-	-	-
B1.7	TN	↓	11	610	4600	800	180	5800	410	9700	11	68
B1.8	TN	southern Narrows	12	600	2800	700	100	-	-	-	-	-
B1.9	PC	↓	5.4	640	2100	530	170	4200	330	6800	11	77
B1.10	PC	Fisherman's Landing	13	640	1400	480	150	-	-	-	-	-
B1.11	PC	mid-harbour	6.4	650	560	370	260	4900	390	8600	13	78
B1.12	PC	↓	12	580	400	300	310	4600	300	7600	11	74
B1.13	PC	↓	4.2	520	<200	280	150	3900	280	6300	12	78
B1.14	PC	outer-harbour	2.3	270	<200	200	94	-	-	-	-	-
B1.15	PC	↓	2.3	180	<200	150	62	-	-	-	-	-
Creeks and inlets in direction: Port Curtis → the Narrows												
B2.1		Boat Creek	5.6	770	-	480	140	-	-	-	-	-
B2.2		Caliope 1	5.6	670	-	330	340	5300	370	10000	17	78
B2.3		Caliope 2	9.2	730	6900	430	500	-	-	-	-	-
B2.4		Fishermans Landing 1	4.6	740	-	410	220	-	-	-	-	-
B2.5		Fishermans Landing 2	4.1	710	-	450	170	4800	470	8800	14	72
B2.6		Graham's Creek 1	3.9	630	-	510	94	-	-	-	-	-
B2.7		Graham's Creek 2	4.2	630	-	600	140	-	-	-	-	-
B2.8		Targinie Creek	5.5	610	19000	640	120	-	-	-	-	-
B2.9		Blackswan Creek	4.1	560	2500	790	84	-	-	-	-	-
Delta of inlets in direction: northern end of the Narrows → mouth of Fitzroy estuary												
B3.1		Division Point	5.2	430	-	670	120	4500	300	6600	8.1	74
B3.2		Connors Creek	16	440	-	660	93	-	-	-	-	-
B3.3		Port Alma	6.8	530	-	760	90	5600	260	8100	12	67
B3.4		Casuarina 1	7.0	540	-	710	260	7400	430	13000	15	55
B3.5		Casuarina 2	20	570	-	770	120	-	-	-	-	-
B3.5		Cattle Point	8.8	410	-	470	120	6100	260	9000	10	71
Fitzroy River / estuary and Dee River samples: in direction river → estuary												
B4.1		Fitzroy River 1	4.1	1200	-	1600	360	8500	2500	15000	22	160
B4.2		Fitzroy River 2	4.3	1300	-	1800	580	-	-	-	-	-
B4.3		Fitzroy River 3	2.0	1200	-	1400	140	10000	2700	18000	29	230
B4.4		Fitzroy estuary 1	7.2	690	-	1100	96	6700	440	11000	13	55
B4.5		Fitzroy estuary 2	8.9	650	-	980	140	-	-	-	-	-
B4.6		Dee River	22000	6E+06	-	65000	3E+06	-	-	-	-	-

725
726

Where, PC refers to Port Curtis, TN refers to The Narrows, and “-“ refers to data not measured.

727 **Table 3. Comparison of dissolved trace metal concentrations in Port Curtis with published data for other estuarine and marine waters**

Location	Metal concentration, ng/L				Reference
	Cadmium	Copper	Nickel	Zinc	
Port Curtis, offshore	<1.5 (<1.5-4.0)	40 (<19-85)	150 (11-190)	40 (<30-140)	This study
Port Curtis Harbour	7.0 (2.0-15.0)	510 (410-620)	340 (280-470)	170 (130-230)	This study
The Narrows	8.0 (3.0-6.5)	530 (520-640)	650 (470-910)	110 (60-210)	This study
NSW coast	2.5	30	180	<22	Apte <i>et al.</i> 1998
Western Pacific Ocean	1-101	40-280	120-522	-	Mackey <i>et al.</i> 2002
North Pacific Ocean	0.3-112	-	-	15-520	Bruland <i>et al.</i> 1994
Great Barrier Reef	<10-60	110-240	60-160	30-350	Denton and Burdon-Jones. 1986
Bathurst Harbour, Tasmania	2.2	140	140	392	Mackey <i>et al.</i> 1996
North Atlantic	0.7	68	-	-	Kremling and Pohl. 1989
Scottish sea loch	<10	180	20	-	Hall <i>et al.</i> 1996
Port Jackson, Australia	6-104	932-2550	175-1610	3270-9660	Hatje <i>et al.</i> 2003
Torres Straight & Gulf of Papua	<1-29	36-986	940-4600	-	Apte and Day, 1998
Port Phillip Bay, Australia	<5-70	400-630	540-1100	250-1050	Fabris and Monahan, 1995
Nine estuaries, northern Australia	1.4-72	150-5500	120-4250	<10-11300	Munksguard and Parry, 2001
Tweed estuary, UK	7-33	490-4700	-	430-1900	Laslett 1995
Humber estuary, UK	80-450	180-10100	2500-12000	3000-20500	Comber <i>et al.</i> 1995
Mersey estuary, UK	10-110	800-4950	2000-10500	6500-28000	Comber <i>et al.</i> 1995
Tay estuary, Scotland	<3-85	250-1550	250-900	<100-3200	Owens and Balls, 1997
Forth estuary, Scotland	-	-	290-1470	460-10130	Laslett and Balls, 1995
Scheldt estuary, Netherlands	15-100	750-1800	1000-6800	1000-10000	Baeyens <i>et al.</i> 2005
San Francisco Bay estuary, USA	22-123	315-2230	140-2410	160-1960	Sanudo-Wilhelmy <i>et al.</i> 1996
Six estuaries, Texas, USA	-	100-3200	-	300-18000	Benoit <i>et al.</i> 1994
Australian Guideline values (95% protection)	700	1300	7000	15000	ANZECC/ARMCANZ 2000

728