



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Commerce - Papers (Archive)

Faculty of Business

2008

'Primary care' presentations at emergency departments - rates and reasons by age and sex

Peter M. Siminski

University of Wollongong, siminski@uow.edu.au

Andrew J. Bezzina

bezzina@uow.edu.au

L. P. Lago

University of Wollongong, lago@uow.edu.au

Kathy Eagar

University of Wollongong, keagar@uow.edu.au

Publication Details

This article was originally published as Siminski P, Bezzina A, Lago L, Eagar K, 'Primary care' presentations at emergency departments - rates and reasons by age and sex', *Australian Health Review* 32 (4), 2008, 700-709.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

'Primary care' presentations at emergency departments - rates and reasons by age and sex

Abstract

'Primary care' presentations at Emergency Departments (EDs) have been the subject of much attention in recent years. This paper is a demographic analysis of such presentations in New South Wales EDs and of self-reported reasons for presentation.

Disciplines

Business | Health Policy | Social and Behavioral Sciences

Publication Details

This article was originally published as Siminski P, Bezzina A, Lago L, Eagar K, 'Primary care' presentations at emergency departments - rates and reasons by age and sex', Australian Health Review 32 (4), 2008, 700-709.

Editorial Manager(tm) for Australian Health Review
Manuscript Draft

Manuscript Number: AHR581R1

Title: 'Primary Care' Presentations at Emergency Departments - Rates and Reasons by Age and Sex

Article Type: Research Note

Corresponding Author: Mr Peter Siminski, BMathEcon(Hons), BA

Corresponding Author's Institution: University of Wollongong

First Author: Peter Siminski, BMathEcon(Hons), BA

Order of Authors: Peter Siminski, BMathEcon(Hons), BA; Andrew J Bezzina, MB BS Dip RACOG FACEM;
Luise P Lago, BMath (Hons); Kathy Eagar, PhD

Manuscript Region of Origin: New South Wales

Abstract: Objectives

'Primary care' presentations at Emergency Departments (EDs) have been the subject of much attention in recent years. This paper is a demographic analysis of such presentations in New South Wales EDs and of self-reported reasons for presentation.

Methods

The demographic characteristics of 'potential primary care' attendances are summarised using administrative data from the Emergency Department Information System (EDIS) for 2005, which covers 76 per cent of emergency attendances in New South Wales. Age and sex differences in the reasons given by patients for such presentations are analysed using data from a survey of patients conducted in a subset of EDs in 2004.

Results

The rate of 'potential primary care' presentations varies greatly with age and to a lesser extent with sex. Almost half (47%) of these presentations are made by people under 25 years of age. Children aged 0-4 years, account for 14% of the total. The pattern is distinctly different to the corresponding rate of ED presentations that do not fit the 'potential primary care' definition. Reasons given for 'potential primary care' presentations are strikingly consistent across all age groups. These reflected self assessed urgency, access to diagnostics and self assessed complexity. Older 'primary care' patients are particularly unlikely to give reasons associated with GP affordability or availability for their presentations. Young adults' responses are consistent with the overall population and children under the age of 5 seem most susceptible to availability issues.

Conclusion

Despite differences in the presentation rates, patients in all demographic groups were most likely to identify self-assessed urgency; being able to see the doctor and having diagnostics done in the same place; and self-assessed seriousness or complexity as the reasons for presentation.

Abstract

Objectives

‘Primary care’ presentations at Emergency Departments (EDs) have been the subject of much attention in recent years. This paper is a demographic analysis of such presentations in New South Wales EDs and of self-reported reasons for presentation.

Methods

The demographic characteristics of ‘potential primary care’ attendances are summarised using administrative data from the Emergency Department Information System (EDIS) for 2005, which covers 76 per cent of emergency attendances in New South Wales. Age and sex differences in the reasons given by patients for such presentations are analysed using data from a survey of patients conducted in a subset of EDs in 2004.

Results

The rate of ‘potential primary care’ presentations varies greatly with age and to a lesser extent with sex. Almost half (47%) of these presentations are made by people under 25 years of age. Children aged 0-4 years, account for 14% of the total. The pattern is distinctly different to the corresponding rate of ED presentations that do not fit the ‘potential primary care’ definition. Reasons given for ‘potential primary care’ presentations are strikingly consistent across all age groups. These reflected self assessed urgency, access to diagnostics and self assessed complexity. Older ‘primary care’ patients are particularly unlikely to give reasons associated with GP affordability or availability for their presentations. Young adults’ responses are consistent with the overall population and children under the age of 5 seem most susceptible to availability issues.

Conclusion

Despite differences in the presentation rates, patients in all demographic groups were most likely to identify self-assessed urgency; being able to see the doctor and having diagnostics done in the same place; and self-assessed seriousness or complexity as the reasons for presentation.

Objectives

Background

Less urgent presentations at Emergency Departments (EDs) have been the subject of much attention in recent years. As a result of well publicised problems of access to care in emergency departments^{1,2} they have been perceived as an issue for concern in ED management not only in Australia but in places as disparate as Canada³, Spain⁴, Britain^{5,6,7,8}, France⁹, Holland¹⁰, New Zealand¹¹ and many others^{12,13}. A recent study focussed on the reasons that 'potential primary care' (PPC) patients give for presenting to EDs rather than to general practitioners¹⁴. The main finding was that patients identified "very appropriate and sensible reasons for coming to the ED – urgency, complexity and being able to have the diagnostic tests they had anticipated would be required". It was argued that improvements to GP affordability and availability would hence be unlikely to affect the numbers of such attendances in a large way.

Importance

Recent publications in the Australian context have focussed on illustrating the small proportion of overall presentations for which this patient group accounts^{15, 16, 17, 18}. They further emphasise that urgency Category 4 and 5 patients do not equate to primary care patients. If, despite definitional issues, strategies are to be developed to influence patients in this group into altering their pattern of accessing health care then a broad set of factors must be considered. This includes any discrepancy between self-assessed and clinician assessed urgency. Further, there needs to be an understanding of variation both between PPC presentations and other presentations (NonPPC), and within the group of PPC cases.

Goals of this investigation

This paper explores presentation patterns specific to PPC cases, how they compare to Non-PPC presentations and whether differences exist in reasons for presentation between age and sex subgroups of potential primary care cases.

Methods

The paper draws on two data sources. One to provide the data on patterns of presentation and the other to provide insight into the possible drivers for any differences between PPC presentations and non PPC presentations.

The first is an administrative data set - EDIS (Emergency Department Information System), which at December 2005 covered 61 EDs in New South Wales, representing 76 per cent of New South Wales ED attendances¹⁹. The EDs covered by this system include all major departments in the state of New South Wales. The departments not included are a selection of small, rural, GP run services and some very small metropolitan units. The second data source is a survey of patients conducted in 2004, described by [reference to own work suppressed]¹⁴. Patients completed the survey in the Emergency Departments of the Illawarra region of New South Wales at the time of presentation. The survey included 5 EDs representing all levels of facility within the state from rural, GP run service through to major regional referral. Patients were offered 20 possible responses as to reasons for their choice to attend the ED and the option of further comment. Any number of responses could be selected.

In both sources, the analysis focussed on a group of patients that would represent PPC attendances. Based on a review of the literature²⁰ attendances were classified as “potential primary care” in the survey when they met all of the criteria below:

- Low urgency and/or acuity, indicated by being classified as Triage Category 4

or 5 on the Australasian Triage Scale

- did not arrive by ambulance
- were self-referred. By definition, patients referred by GP/ community primary medical services are not primary care cases because a primary care service has referred them on
- were presenting for a new episode of care and
- were not expected to be admitted (according to the assessment of staff in the ED).

The same definition was used in EDIS, with two exceptions. 'Not admitted' was used as a criterion instead of 'not expected to be admitted' since this was a retrospective analysis. Source of referral was not available in EDIS. Irrespective of the definition used they reflect a group that is only potentially appropriate to manage in a primary care setting rather than an Emergency Department. The breadth of the definition in either instance will mean that there is a significant overestimate of cases. De-identified EDIS data were selected for the 2005 calendar year, tabulated by potential primary care status, sex and age in 5 year bands. The number of presentations and presentation rates were calculated from EDIS data and the estimated resident population for NSW at June 2005 ²¹.

The survey involved a convenience sample of 400 PPC patients invited to participate between 14/1/04 and 19/4/04. Of these only three refused, a response rate of over 99%. Approximately half the participants (those not accompanied by friends or family) were assisted in responding to the questionnaire by the nurse researcher.

The reasons given by patients in the survey were analysed by age and sex. The selection of age categories for the survey analysis was informed by the presentation rate results. The age groups analysed were 0-4 years, 5-14 years, 15-29 years, 30-64

years and 65+ years. Particular attention is given to the 0-4, 15-29 and 65+ year categories, for reasons discussed below. Only the first 18 reasons for presentation were considered, because the sample of after-hours presentations was too small to analyse by age. The proportions of people selecting each reason as very important or moderately important were tabulated. The mean number of very important or moderately important reasons were also analysed.

Results

Presentations by age and sex

The number of ED presentations in 2005 by age and potential primary care status (PPC or non-PPC) is shown in Figure 1. Some 0.5% of records were not classifiable as PPC or non-PPC due to missing data on one or more of the criteria. These records were excluded from the analysis. Almost half (47%) of PPC presentations were by people aged under 25 years. By far the largest number of PPC presentations was by children aged 0-4 years, accounting for 14% of the total. This pattern contrasts with the profile of non-PPC presentations. The non-PPC profile is characterised by a relatively even age distribution, though it includes a high number of presentations by 0-4 year olds. Thus in raw numbers PPC presentations are dominated by younger age groups, while non-PPC presentations have an even age distribution.

Figure 1 is partly a function of the age distribution of the population. It is thus informative to consider the rates of ED presentations, equal to the number of presentations in each sex-age group divided by its population. PPC presentation rates are shown in Figure 2. Overall, the male rates are 18% higher than the female rates on an age-standardised basis. The PPC presentation rate is clearly highest among the youngest age group (0-4 years) (244 and 210 per 1000 people for males and females,

respectively). This is more than twice as high as the overall rate (105). For both sexes, the rate is also relatively high amongst persons aged 15-29. It decreases for subsequent age groups to 55-59. The rate levels off for older females, but increases slightly for older males. This shows that the dominance of younger age groups persists even after the age distribution of the population is accounted for.

People aged 65 and over account for only 8.9% of PPC presentations in 2005. However, whilst the overall PPC presentation rate fell between 1999 and 2006, it increased amongst people aged 65 and over.²² Thus despite relatively low PPC presentation rates, older age groups are of particular interest when developing access strategies because of further projected population ageing and their relatively large apparent increase in PPC presentation rates.

For comparative purposes, non-PPC presentation rates are shown in Figure 3. Unlike the PPC rate, the non-PPC rate increases greatly with age from about 60 years. The non-PPC rate for persons aged 85 and over is 531, more than 5 times higher than for 45-49 year olds, and 8 times higher than the PPC rate for people aged 85 and over. Thus older ages are utilising EDs much more frequently overall and predominantly for non-PPC issues. As in the PPC rates, males also have a higher non-PPC presentation rate than females (16% higher on an age-standardised basis), though the difference is close to zero in most child-bearing age groups. The 15-29 year age groups have slightly higher non-PPC presentation rates than immediately younger and immediately older age groups, but this spike is not as large as it is for PPC rates.

Reasons for Presenting

The reasons given by PPC patients for presentation at an ED, as identified in the survey, are analysed primarily by age. The difference between males and females was not statistically significant for any of the eighteen reasons even at the 10% level,

regardless of whether 'very important' reasons are considered in isolation, or 'very important' and 'moderately important' reasons are considered together. This is an important finding in that it suggests that there is no evidence of a gender based element to this care choice.

Results are shown for all age groups but attention is focussed on:

- Infants (0-4 years) because they have the highest presentation rates;
- Young adults (15-29 years) because they have the next highest presentation rates, which have a sharper spike than corresponding non-PPC rates. They were also hypothesised to be susceptible to issues of GP availability and affordability.
- Older people (65 years and over) because they appear to have the fastest growth in presentations.

In interpreting the results, it is important to note that the average number of reasons selected by patients differs with age. Younger patients (or their proxies) selected more reasons than older patients (or their proxies) (Table 1). The sample size of each group is also shown. All subsequent results should be interpreted in this context. The sample size is particularly small for children aged less than 5 years. The complete set of results by age is shown in Table 2. The most striking finding is the consistency of the most prevalently selected reasons across all age groups. Regardless of age, Q1, Q7 and Q2 (in that order) were selected as very important by the greatest proportion of people. These reflect self assessed urgency, access to diagnostics and self assessed complexity. When very important and moderately important reasons were analysed together, the same finding is observed, with the exception that for 15-29 year olds, Q1 and Q7 ranked equal first. For all age groups, these three reasons stood out from the other reasons.

Attention is now turned to the subset of reasons that relate to primary care availability or affordability and to the age groups selected for particular attention. Questions 8, 9, 10 and 11 all related to availability. Questions 12 and 13 are related to affordability. The results (which include both very important and moderately important reasons) are shown in Figure 4.

Older patients were very unlikely to select affordability or availability reasons. In fact, of 74 respondents aged 65 or over, not a single person selected an affordability reason as being important. This is perhaps unsurprising, as older people are more likely to be bulk-billed (no direct fee paid by the patient) than others²³. Older people were also unlikely to select issues of availability. Of all age groups, they were the least likely to select questions 8, 9 or 10. The proportion of older people to select question 11 was similar to the rest of the population. This may reflect a high reliance on public or community transport. Availability can also be analysed as a single factor, calculated as the sum of Q8, Q9, Q10 and Q11. The average of this variable is significantly smaller ($p < 0.001$) amongst older people (0.41) than for the other age groups combined (0.78). These results are not a function of older people's apparent tendency to select fewer reasons (Table 1). In fact, these six questions, combined, account for most of the difference in average number of reasons selected between older people and the full sample.

A second observation relates to the unremarkable responses of those aged 15-29. It was hypothesised that this group may be particularly susceptible to issues of availability and affordability. This does not appear to be the case. Their responses to the availability questions were quite similar to those of the full sample. Their responses to the affordability questions were very similar to those made on behalf of children aged under 15 (who were slightly more likely to select these reasons than

those aged 30 and over). The unexceptional responses of young adults are not confined to questions of affordability and availability, as can be seen from a closer inspection of Table 2.

Finally, children aged under 5 years appear to be slightly more vulnerable to availability issues, particularly in relation to waiting time for a GP appointment (Q9). The percentage of persons selecting this as a reason on behalf of children under 5 (44% = 8/18) was almost twice as high as the rest of the sample (23%). Considered in isolation, this difference is statistically significant ($p < 0.05$). However, the general tendency of such respondents to select more reasons than other respondents weakens the strength of the conclusion. When availability is considered as a factor, the difference between the responses for children under 5 and the rest of the sample is not statistically significant. At best, this finding constitutes a weak suggestion that GP waiting time may be a particular issue for children aged under 5. If true this may be because parents are likely to escalate their judgement of urgency for infants and they are hence more likely to seek immediate medical attention. This is supported by the observation that urgency was selected as a reason by almost all (95%) people responding on behalf of children aged under 5 years.

Limitations

As discussed above, this study used data from two data sources. This creates two possible issues. The first is that the survey participants are not a random sample of NSW EDIS cases. However, these results for Illawarra patients may be regarded as a useful indicator for broader inference because of geographic and other characteristics, as discussed elsewhere and repeated in summary form here¹⁴. EDs in the Illawarra span all types, from a major referral hospital to small community hospitals. On a number of ED, General Practice and socioeconomic indicators, Illawarra is very

similar to NSW overall. These include ED and General Practice utilisation rates, average ED waiting times by triage and bulk-billing rates. A slightly higher proportion of Illawarra's patients were affected by access block than for NSW overall. Of the seventeen NSW health areas at the time of the survey, Illawarra ranked near the middle in remoteness and in socio-economic status.

The second issue is that the definitions of PPC differed between the sources in two ways. Source of referral was used as a criterion for the survey, but not for the EDIS analysis, because the data item is incomplete. For a subset of hospitals where this variable is complete (Wollongong, Shellharbour and Shoalhaven over 2002-03 and 2003-04), we find that 6% of cases otherwise identified as 'PPC' were actually referred, and thus excluded from our preferred definition. This percentage is higher for older age groups (as high as 14% for 70-89 year olds) than younger age groups (as low as 4% for 0-9 year olds), but there was almost no difference by sex. Thus the EDIS results presented here are likely to overestimate PPC presentations relative to non-PPC presentations, especially among older people.

In addition, only those patients who were not admitted were in scope for the EDIS analysis, while patients who were *not expected* to be admitted were in scope for the survey. The EDIS data will thus inevitably include some patients who would not have been selected for the survey. This could occur for a number of clinical reasons such as a complex presentation that makes the initial assessment difficult or a change in the clinical picture following presentation (such as an abnormal but uncomplicated cardiac rhythm that settles and thus does not require admission).

The same holds true in reverse where patients who are not expected to be admitted ultimately end up admitted for unexpected reasons such as rapidly progressive illness.

If the premise is taken that expectations over admission are a more appropriate criterion than admission itself, then the definition used in the EDIS data may introduce some error. Given the purposes of the paper, however, such definitional issues are only a problem to the extent that they affect age and sex groups differently. This is expected to be more of an issue for particularly complex older patients and injury cases, though this is difficult to quantify. Nevertheless, this is unlikely to fully explain the pattern of the findings. The apparent primary care attendance rates of the elderly are already low, even if they are overestimated. The age spikes for 15 – 29 year olds may again, to some extent, be explained due to the overestimation of potential primary care cases related to trauma. But this issue is unlikely to explain it entirely and the results are consistent with epidemiological data in reviews of minor injury presentations to any ED^{24, 25, 26}.

The EDIS database is representative of the population of emergency presentations within NSW as a whole, except for small rural hospitals with less than 5000 admissions per year.

Conclusion

EDIS data reveal that the age profile of potential primary care attendances at EDs is considerably different to that of other attendances. The rates of both potential primary care and non- potential primary care attendances are higher amongst men than amongst women, and both are relatively high amongst infants. Amongst older people, however, potential primary care rates are much lower than non- potential primary care rates. Despite this, the rate of potential primary care attendances amongst older people appears to have grown the fastest of all age groups in recent years, and coupled with the structural ageing of the population, this age group is of particular interest despite its relatively low presentation rate. In this context, it is interesting that older people

are reportedly unresponsive to the characteristics of GP services (availability and affordability) in the decision to attend EDs for less urgent cases since it would suggest strategies that focus on changing these aspects are unlikely to succeed.

There is a spike in the rate of PPC presentations amongst young adults (aged 15-29), both male and female. This spike is more distinct than in the corresponding pattern of non- PPC presentations. The patient survey data were utilised to examine whether the reasons for presentation might explain this spike. In fact there was very little difference between the responses of young adults and the rest of the sample. In particular, a similar proportion of young adults identified availability and affordability issues as important reasons as people of other ages. It is possible their higher rate of potential primary care attendances reflects a higher rate of minor accidents. Another possible explanation is that young adults are perhaps less likely to have established trusting relationships with general practitioners. This issue was not directly investigated in the survey, and is worthy of further investigation.

Similarly, there are no significant differences between the reasons given by males and females. Thus the higher rate of PPC attendances by males also appears unrelated to GP characteristics or other reasons for presentation. Again it may reflect a higher rate of minor accidents amongst males.

Overall, however, the main conclusion is clear. While there are differences by age and sex, patients in all age groups were most likely to identify self-assessed urgency; being able to see the doctor and having tests or X-rays done in the same place; and self-assessed seriousness or complexity as the reasons for presentation to ED. These reasons stand out from all other reasons, regardless of age or sex. The implication here is that utilisation of EDs by these patients is, irrespective of age or sex, premised on reasonable decision making processes and as such may not be

amenable to commonly promoted education programs focussed on clarifying service roles.

References


1. Iemma's after hours promise for patients in a hurry. *Sydney Morning Herald* March 5th, 07.
2. GP clinics will take the strain off casualty teams. *Sydney Morning Herald* June 7th, 2006.
3. Boushy D, Dubinsky I. Primary Care Physician and Patient Factors that result in patients seeking emergency care in a hospital setting: The patient's perspective. *J Emerg. Med.* 1999; 17 (3): 405 – 12.
4. Vazquez Quiroga B, Pardo Moreno G, Fernandez Cantalejo G. et al Why do our patients go to hospital emergency departments? *Atencion Primaria.* 2000; 25(3):172-5,.
5. Coleman P, Irons R, Nicholl J. Will alternative immediate care services reduce demands for a non urgent treatment at accident and emergency? *Emerg Med J* 2001; 18 (6): 482 – 7.
6. Davies T. Accident Department or General Practice? *Br Med J (Clin Res Ed)* 1986; 292 (6915): 241 – 3.
7. Rajpar SF, Smith MA, Cooke MW. Study of Choice between accident and emergency departments and general practice centres for out of hours primary care problems. *J Accid Emerg Med* 2000; 17 (1):18 – 21.
8. Lowy A, Kohler B, Nicholl J. Attendance at Accident and Emergency departments: unnecessary or inappropriate? *J Public Health Med* 1994; 16 (2): 134 –
9. Lang T, Davido A, Diakite B et al Using the hospital emergency department as a regular source of care. *Eur J Epidemiol.* 1997; 13 (2):223-8.

10. Rieffe C, Oosterveld P, Wijkkel D et al Reasons why patients bypass their GP to visit a hospital emergency department. *Accid Emerg Nurs*. 1999; 7(4):217-25.
11. Richardson S. Emergency Departments and the inappropriate attender – is it time for a reconceptualisation of the role of primary care in emergency facilities? *Nurs Pract NZ* 1999; 14 (2): 13 – 20.
12. Diesburg- Stanwood A, Scott J, Oman K et al Non-emergent ED patients referred to Community Resources after Medical Screening Examination: Characteristics, Medical Condition After 72 hours, and Use of Follow-up Services. *J Emerg Nurs*. 2004; 30 (4): 312-7.
13. Krakau I, Hassler E. Provision for clinic patients in the ED produces more non-emergency visits. *Am J Emerg Med* 1999; 17 (1): 18 -20.
14. [reference to own work suppressed]
15. Dent A.W., Phillips G., Chenhall, and McGregor L.R. The heaviest repeat users of an inner city Emergency Department are not general practice patients. *Emerg Med* 2003; **15** (4), 322 – 29.
16. Nagree Y. Ercleve TN. Sprivulis PC. After-hours general practice clinics are unlikely to reduce low acuity patient attendances to metropolitan Perth emergency departments. *Australian Health Review*. 28(3):285-91, 2004 Dec 13.
17. Australasian College for Emergency Medicine, *Fact Sheet: Re Urban Emergency Services – ATS 4 and 5 patients*. 2001.
www.acem.org.au/media/ats_4_5_factsheet.pdf
18. Australasian College for Emergency Medicine, *The Relationship Between Emergency Department Overcrowding and Alternative After Hours GP Services*. 2004. www.medeserv.com.au/acem/open/documents/after_hours_gp.pdf

19. NSW Health (2006) <http://www.health.nsw.gov.au/hospitalinfo/perfnon.html>
Accessed 2 March, 2006.
20. [reference to own work suppressed]
21. Australian Bureau of Statistics *Population by Age and Sex, Australian States and Territories, Jun 2005*, ABS Cat. No. 3201.0.
22. [reference to own work suppressed]
23. Abbott, T. (2006) *New GP bulk-billing records set for young and rural patients*
Media Release, Minister for Health and Ageing, May 12 2006. [cited 8 June, 2006]. Available from URL:
[http://www.health.gov.au/internet/ministers/publishing.nsf/Content/7D78299BBB F63ACECA25716C0008E68C/\\$File/abb066.pdf](http://www.health.gov.au/internet/ministers/publishing.nsf/Content/7D78299BBB F63ACECA25716C0008E68C/$File/abb066.pdf)
24. Li L, Ozanne-Smith J. Injury hospitalisation rates in Victoria, 1987-97: trends, age and gender patterns. *Aust NZ J Public Health*. 2000;24(2):158-65.
25. Watson WL, Ozanne-Smith J. Injury surveillance in Victoria, Australia: developing comprehensive injury incidence estimates. *Accid Anal Prev*. 2000; 32(2):277-86.
26. Schmertmann M, Williamson A. A brief overview of injury in New South Wales. *N S W Public Health Bull* 2002; 13(4): 66-70

Figure 1


Figure 1 ED Presentations ('000s) by Potential Primary Care (PPC) status and Age group – NSW, 2005 (EDIS)^a



a) Only includes presentations at EDs with the EDIS system

Figure 2


Figure 2 ED Potential Primary Care (PPC) Presentation Rates by Sex and Age (per 1000 people)
– NSW, 2005 (EDIS) ^a



a) Only includes presentations at EDs with the EDIS system.

Figure 3


Figure 3 ED Non-PPC Presentation Rates by Sex and Age (per 1000 people) – NSW, 2005
(EDIS)^a



a) Only includes presentations at EDs with the EDIS system.

Figure 4

Figure 4 Reasons associated with GP availability of affordability by selected age groups: per cent of all valid responses[†]



† Includes very important and moderately important reasons

Table 1

Table 1 Sample size and average number of reasons selected by age of patient

Age Group	Sample size	Average number of reasons ¹		
		Very important	Moderately important	Very important or moderately important
less than 5	19	2.8	2.1	4.8
5-14	36	2.3	1.3	3.6
15-29	105	2.1	2.0	4.1
30-64	154	2.2	1.3	3.5
65+	74	2.2	0.9	3.0
All ages ²	388	2.4	1.4	3.7

(1): Questions 19 and 20 are excluded

(2) 'All ages' excludes those records with missing age to conform to its components

Table 2

Table 2 Very important and moderately important reasons why patients presented to an ED by age: per cent of valid responses¹

Summary reason	Age group (years)					
	less than 5	5-14	15-29	30-64	65 and over	All ages ²
Q1: Problem too urgent	95	86	83	75	81	80
Q2: Problem too serious/complex	68	49	50	50	60	53
Q3: Medical treatment better at ED	42	31	35	34	33	34
Q4: Second opinion	21	8	16	13	13	14
Q5: Did not want GP to know	11	0	2	3	1	2
Q6: Prefer doctor I don't know	11	0	11	4	3	6
Q7: See doctor and have tests/X-rays done in same place	83	69	83	71	70	74
Q8: Not able to see GP as books are closed	17	14	19	19	5	16
Q9: Not happy with GP waiting time	44	22	20	31	11	24
Q10: Do not like making appointments	28	6	19	12	1	12
Q11: Easier to get to the ED	28	28	24	17	23	21
Q12: No charge to see a doctor	17	14	15	8	0	9
Q13: No charge for X-rays or medicine	17	14	18	9	0	10
Q14: Female doctor	0	0	5	2	0	2
Q15: Doctor or interpreter who speaks my language	0	0	5	2	0	2
Q16: Aboriginal health staff	0	0	5	2	0	2
Q17: Prefer ED environment	6	11	8	4	1	5
Q18: Traditional use by family	11	8	14	7	4	9

Notes:

- (1) The reasons are in summary format. A copy of the survey can be found in [reference to own work suppressed]
- (2): 'All ages' excludes those records with missing age to conform to its components

Acknowledgments

Acknowledgments

We acknowledge the comments made by an anonymous reviewer. The State Commonwealth Research Issues Forum (SCRIF) through the NHMRC funded this study. New South Wales Health provided EDIS data. The joint University of Wollongong, Illawarra Area Health Service Health Research Ethics Committee approved the patient survey.