

University of Wollongong

Research Online

Faculty of Engineering and Information
Sciences - Papers: Part A

Faculty of Engineering and Information
Sciences

1-1-2011

Regularity of Monge-Ampere equations in optimal transportation

Jiakun Liu

Princeton University, jiakunl@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/eispapers>



Part of the [Engineering Commons](#), and the [Science and Technology Studies Commons](#)

Recommended Citation

Liu, Jiakun, "Regularity of Monge-Ampere equations in optimal transportation" (2011). *Faculty of Engineering and Information Sciences - Papers: Part A*. 618.
<https://ro.uow.edu.au/eispapers/618>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Regularity of Monge-Ampere equations in optimal transportation

Abstract

We consider some recent regularity results for the Monge-Ampère equation arising in the optimal transportation problem. The Monge-Ampère equation under consideration has the following type

Keywords

regularity, transportation, optimal, equations, ampere, monge

Disciplines

Engineering | Science and Technology Studies

Publication Details

Liu, J. (2011). Regularity of Monge-Ampere equations in optimal transportation. *Bulletin of the Australian Mathematical Society*, 83 173-176.

REGULARITY OF MONGE–AMPÈRE EQUATIONS IN OPTIMAL TRANSPORTATION

JIAKUN LIU

(Received 17 November 2010)

2010 *Mathematics subject classification*: primary 35J60; secondary 35B45.

Keywords and phrases: Monge–Ampère equation, optimal transportation.

We consider some recent regularity results for the Monge–Ampère equation arising in the optimal transportation problem. The Monge–Ampère equation under consideration has the following type

$$\det\{D^2u(x) - A(x, Du)\} = f(x) \quad \text{in } \Omega, \quad (1)$$

where $\Omega \subset \mathbb{R}^n$ is a bounded domain and $A = \{A_{ij}\}$ is an $n \times n$ symmetric matrix defined in $\Omega \times \mathbb{R}^n$.

In optimal transportation, u is the potential function, and the matrix A and the right-hand side f are given by

$$A(x, Du) = D_x^2c(x, T_u(x)), \quad (2)$$

$$f = |\det\{D_{xy}^2c\}| \frac{\rho}{\rho^* \circ T_u}, \quad (3)$$

where $c(\cdot, \cdot)$ is the cost function, $T_u : x \rightarrow y$ is the optimal mapping determined by $Du(x) = D_xc(x, y)$, and ρ and ρ^* are mass distributions in the initial domain Ω and the target domain Ω^* , respectively.

We assume that the cost function $c \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n)$ and satisfies the following conditions.

(A1) For any $x, p \in \mathbb{R}^n$, there is a unique $y \in \mathbb{R}^n$ such that $D_xc(x, y) = p$; and for any $y, q \in \mathbb{R}^n$, there is a unique $x \in \mathbb{R}^n$ such that $D_yc(x, y) = q$.

(A2) For any $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$, $\det\{D_{xy}^2c(x, y)\} \neq 0$.

(A3) For any $x, p \in \mathbb{R}^n$, and any $\xi, \eta \in \mathbb{R}^n$ with $\xi \perp \eta$,

$$A_{ij,kl}(x, p)\xi_i\xi_j\eta_k\eta_l \geq c_0|\xi|^2|\eta|^2, \quad (4)$$

for some constant $c_0 > 0$, where $A_{ij,kl} = D_{p_k p_l}^2 A_{ij}$ and A is given by (2).

Thesis submitted to The Australian National University, April 2010. Degree approved, July 2010.
Supervisors: Professor Xu-Jia Wang and Professor Neil S. Trudinger.

© 2011 Australian Mathematical Publishing Association Inc. 0004-9727/2011 \$16.00

Under the above conditions on the cost function, the optimal mapping T_u is uniquely determined by the corresponding potential function u . Therefore, in order to study the regularity of optimal mapping it suffices to study the regularity of potential functions, that is, regularity of elliptic solutions of (1).

In the special case when the cost function is the Euclidean distance squared, equation (1) becomes the standard Monge–Ampère equation, and the various regularity results have been obtained by Caffarelli [1, 2], Delanoë [5], Urbas [12] and many other mathematicians. In particular, the interior $C^{1,\alpha}$, $C^{2,\alpha}$, $W^{2,p}$ estimates are due to Caffarelli [3]. The regularity of optimal transportation with general costs is an important open problem in the area, as pointed out by Caffarelli [4] and Villani [13]. Our goal is to establish the corresponding regularity results for general cost functions satisfying conditions (A1)–(A3), assuming the mass distributions are merely measurable or Hölder continuous, [7–9]. When the mass distributions are smooth, the C^3 smooth regularity has been obtained in [11].

The first result is the $C^{1,\alpha}$ regularity for potentials, [7]. A similar estimate was previously obtained by Loeper [10]. We give a completely different proof and our exponent is optimal when the inhomogeneous term $f \in L^\infty$.

THEOREM 1. *Let u be a potential function in the optimal transportation problem. Assume that the cost function c satisfies conditions (A1)–(A3), Ω^* is c -convex with respect to Ω , and $f \geq 0$, $f \in L^p(\Omega)$ for some $p \in (\frac{1}{2}(n+1), +\infty]$. Then $u \in C^{1,\alpha}(\bar{\Omega})$, where $\alpha = \beta(n+1)/(2n^2 + \beta(n-1))$ and $\beta = 1 - (n+1)/2p$.*

In particular, when $p = \infty$, our Hölder exponent $\alpha = 1/(2n-1)$ is optimal.

Here and below, we say that the target domain Ω^* is c -convex with respect to Ω if $D_x c(x, \Omega^*)$ is convex for any $x \in \Omega$.

The second result gives the Hölder and more general continuity estimates for second derivatives of solutions u , when the inhomogeneous term f is Hölder or Dini continuous, together with corresponding regularity results for potentials, [9].

THEOREM 2. *Assume that the cost function c satisfies (A1)–(A3) and f satisfies $C_1 \leq f \leq C_2$ for some positive constants $C_1, C_2 > 0$. Let $u \in C^2(\Omega)$ be the potential function satisfying (1). Then for all $x, y \in \Omega_\delta$, we have the estimate*

$$|D^2u(x) - D^2u(y)| \leq C \left[d + \int_0^d \frac{\omega_f(r)}{r} + d \int_d^1 \frac{\omega_f(r)}{r^2} \right], \quad (5)$$

where $d = |x - y|$, $\Omega_\delta = \{x \in \Omega : \text{dist}(x, \partial\Omega) > \delta\}$, $\omega_f(r) = \sup\{|f(x) - f(y)| : |x - y| < r\}$ is the oscillation of f , and the constant $C > 0$ depends only on $n, \delta, C_1, C_2, A, \sup |Du|$, and the modulus of continuity of Du .

It follows that:

- (i) if f is Dini continuous, that is, $\int_0^1 (\omega_f(r)/r) dr < \infty$, then the modulus of continuity of D^2u can be estimated by (5) above;

(ii) if $f \in C^\alpha(\Omega)$ for some $\alpha \in (0, 1)$, then

$$\|u\|_{C^{2,\alpha}(\Omega_\delta)} \leq C \left[1 + \frac{\|f\|_{C^\alpha(\Omega)}}{\alpha(1-\alpha)} \right]; \quad (6)$$

(iii) if $f \in C^{0,1}(\Omega)$, then

$$|D^2u(x) - D^2u(y)| \leq Cd[1 + \|f\|_{C^{0,1}}|\log d|] \quad \forall x, y \in \Omega_\delta. \quad (7)$$

We remark that for standard Monge–Ampère equations, these estimates were obtained by Caffarelli for Hölder continuous f [1] and by Wang for Dini continuous f [14]. In particular, the estimate (5) for standard Monge–Ampère equations was obtained by Jian and Wang [6], and previously by Wang for uniform elliptic equations [15].

The third result is the following $W^{2,p}$ estimate, which is obtained by making a more detailed study of local geometry of potential and cost functions [8].

THEOREM 3. *Assume that the cost function c satisfies (A1)–(A3), f is continuous, $0 < C_1 \leq f \leq C_2$, and Ω^* is c -convex with respect to Ω . Let u be the potential function satisfying (1). Then $D^2u \in L^p(\Omega')$ for any $p \geq 1$, $\Omega' \Subset \Omega$, and we have the estimate*

$$\|u\|_{W^{2,p}(\Omega')} \leq C,$$

where C depends on $n, p, C_1, C_2, \Omega, \Omega', \Omega^*$, and the modulus of continuity of f .

In the proof of Theorem 3, we show that in a proper normalization process, the cost function is uniformly smooth and converges locally smoothly to the linear function $x \cdot y$, and the potential function converges to a quadratic function. As an application, we also obtain the following sharp $C^{1,\alpha}$ estimate for the potentials.

COROLLARY 4. *Let u be the potential function satisfying (1). Suppose that the cost function c satisfies (A1)–(A3) and Ω^* is c -convex with respect to Ω . Then if*

$$|f - 1| \leq \varepsilon, \quad (8)$$

we have $u \in C_{\text{loc}}^{1+\alpha}(\Omega)$ for some $\alpha \in (1 - C_1\varepsilon, 1]$, and for all $\Omega' \Subset \Omega$,

$$\|u\|_{C^{1+\alpha}(\Omega')} \leq C,$$

where the constants $C, C_1 > 0$ depend only on ε, n , and C depends also on $\text{dist}(\Omega', \partial\Omega)$.

Note that in Corollary 4 we do not assume the continuity of f and we have the linear relation

$$\alpha \geq 1 - C_1\varepsilon.$$

Moreover, our proof of Theorem 3 and Corollary 4 also implies a related result. That is, for all $p < \infty$, there exists $\varepsilon = \varepsilon(p)$ such that if f satisfies (8), then $u \in W_{\text{loc}}^{2,p}(\Omega)$. See [8] for more details.

References

- [1] L. A. Caffarelli, ‘Interior $W^{2,p}$ estimates for solutions of the Monge–Ampère equation’, *Ann. of Math.* (2) **131** (1990), 135–150.
- [2] L. A. Caffarelli, ‘Some regularity properties of solutions of Monge–Ampère equation’, *Comm. Pure Appl. Math.* **44** (1991), 965–969.
- [3] L. A. Caffarelli, ‘The regularity of mappings with a convex potential’, *J. Amer. Math. Soc.* **5** (1992), 99–104.
- [4] L. A. Caffarelli, ‘Non linear elliptic theory and the Monge–Ampère equation’, *Proceedings of the ICM*, Vol. 1 (Beijing, 2002) (Higher Ed. Press, Beijing, 2002), pp. 179–187.
- [5] Ph. Delanoë, ‘Classical solvability in dimension two of the second boundary value problem associated with the Monge–Ampère operator’, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **8** (1991), 443–457.
- [6] H. Y. Jian and X.-J. Wang, ‘Continuity estimates for the Monge–Ampère equation’, *SIAM J. Math. Anal.* **39** (2007), 608–626.
- [7] J. Liu, ‘Hölder regularity of optimal mappings in optimal transportation’, *Calc. Var. Partial Differential Equations* **34** (2009), 435–451.
- [8] J. Liu, N. S. Trudinger and X.-J. Wang, ‘On asymptotic behaviour and $W^{2,p}$ regularity of potentials in optimal transportation’, submitted.
- [9] J. Liu, N. S. Trudinger and X.-J. Wang, ‘Interior $C^{2,\alpha}$ regularity for potential functions in optimal transportation’, *Comm. Partial Differential Equations* **35** (2010), 165–184.
- [10] G. Loeper, ‘On the regularity of maps solutions of optimal transportation problems’, *Acta Math.* **202** (2009), 241–283.
- [11] X. N. Ma, N. S. Trudinger and X.-J. Wang, ‘Regularity of potential functions of the optimal transportation problem’, *Arch. Ration. Mech. Anal.* **177** (2005), 151–183.
- [12] J. Urbas, ‘On the second boundary value problem of Monge–Ampère type’, *J. reine angew. Math.* **487** (1997), 115–124.
- [13] C. Villani, *Topics in Optimal Transportation*, Graduate Studies in Mathematics, 58 (American Mathematical Society, Providence, RI, 2003).
- [14] X.-J. Wang, ‘Remarks on the regularity of Monge–Ampère’, in: *Proc. Inter. Conf. Nonlinear PDE*, (eds. G. C. Dong and F. H. Lin) (Academic Press, Beijing, 1992), pp. 257–263.
- [15] X.-J. Wang, ‘Schauder estimate for elliptic and parabolic equations’, *Chin. Ann. Math. Ser. B* **27** (2006), 637–642.

JIAKUN LIU, Department of Mathematics, Princeton University, Fine Hall,
Washington Road, Princeton, NJ 08544-1000, USA
e-mail: jiakunl@math.princeton.edu