



UNIVERSITY  
OF WOLLONGONG  
AUSTRALIA

University of Wollongong  
Research Online

---

Faculty of Health and Behavioural Sciences - Papers  
(Archive)

Faculty of Science, Medicine and Health

---

2012

# Practice guidelines need to address the 'how' and the 'what' of implementation

Kenneth D. Walsh

*University of Wollongong, kenw@uow.edu.au*

Jackie Crisp

*University Of Technology Sydney*

Ann McKillop

---

## Publication Details

McKillop, A., Crisp, J. & Walsh, K. (2012). Practice guidelines need to address the 'how' and the 'what' of implementation. *Primary Health Care Research & Development*, 13 (1), 48-59.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:  
[research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

# Practice guidelines need to address the 'how' and the 'what' of implementation

## **Abstract**

**Aim:** The aim of this study was to explore the realities of everyday nursing practice associated with the implementation of a guideline for the assessment and management of cardiovascular risk. **Background:** The use of clinical practice guidelines is pivotal to improving health outcomes. However, the implementation of guidelines into practice is complex, unpredictable and, in spite of much investigation, remains resistant to explanation of what works and why. Exploration of the nature of guideline implementation has the potential to illuminate the complexities of guideline implementation by focussing on the nature of practice. Nurses are well placed at the front line of primary health care to contribute to an understanding of how guideline implementation plays out in their everyday practice. **Methods:** Qualitative description was used, involving focus groups and interviews with 32 participants (20 nurses, four doctors, five managers and three funder/planners), to explore the use of a guideline in everyday primary health-care practice. Thematic analysis of data was managed through an inductive process of familiarisation, coding, categorising and generation of themes. **Findings:** Four themes were generated from the data portraying the realities of guideline implementation for primary health-care nurses: self-managing patient, everyday nursing practice, developing new relationships in the health team and impact on health-care delivery. The findings reveal that, even with the best of intentions to implement the guideline, health professionals were frustrated and at a loss as to how to achieve that in practice. Consequently, cardiovascular risk assessment and management was uneven and fragmented. Primary health-care practice environments vary so much that solutions to the difficulties of implementing evidence into practice requires context-specific solution-finding through collaborative teamwork. Furthermore, the attention of guideline developers, health-care policymakers, funders and researchers requires direct focus on the 'how' and the 'what' of evidence implementation.

## **Keywords**

Practice, guidelines, need, address, implementation

## **Disciplines**

Arts and Humanities | Life Sciences | Medicine and Health Sciences | Social and Behavioral Sciences

## **Publication Details**

McKillop, A., Crisp, J. & Walsh, K. (2012). Practice guidelines need to address the 'how' and the 'what' of implementation. *Primary Health Care Research & Development*, 13 (1), 48-59.

# Practice guidelines need to address the ‘how’ and the ‘what’ of implementation

Ann McKillop<sup>1</sup>, Jackie Crisp<sup>2,3</sup> and Kenneth Walsh<sup>4</sup>

<sup>1</sup>Senior Lecturer, Faculty of Medical and Health Sciences, School of Nursing, The University of Auckland, New Zealand

<sup>2</sup>Professor of Child and Adolescent Nursing, Sydney Children’s Hospital and Faculty of Nursing, Midwifery and Health, University of Technology, Sydney

<sup>3</sup>Conjoint Professor, University of New South Wales, Australia

<sup>4</sup>Professor of Nursing, Practice Development, Nursing Development and Research Unit, South East Sydney and Illawarra Area Health Service and School of Nursing, Midwifery and Indigenous Health, University of Wollongong, Australia

**Aim:** The aim of this study was to explore the realities of everyday nursing practice associated with the implementation of a guideline for the assessment and management of cardiovascular risk. **Background:** The use of clinical practice guidelines is pivotal to improving health outcomes. However, the implementation of guidelines into practice is complex, unpredictable and, in spite of much investigation, remains resistant to explanation of what works and why. Exploration of the nature of guideline implementation has the potential to illuminate the complexities of guideline implementation by focussing on the nature of practice. Nurses are well placed at the front line of primary health care to contribute to an understanding of how guideline implementation plays out in their everyday practice. **Methods:** Qualitative description was used, involving focus groups and interviews with 32 participants (20 nurses, four doctors, five managers and three funder/planners), to explore the use of a guideline in everyday primary health-care practice. Thematic analysis of data was managed through an inductive process of familiarisation, coding, categorising and generation of themes. **Findings:** Four themes were generated from the data portraying the realities of guideline implementation for primary health-care nurses: self-managing patient, everyday nursing practice, developing new relationships in the health team and impact on health-care delivery. The findings reveal that, even with the best of intentions to implement the guideline, health professionals were frustrated and at a loss as to how to achieve that in practice. Consequently, cardiovascular risk assessment and management was uneven and fragmented. Primary health-care practice environments vary so much that solutions to the difficulties of implementing evidence into practice requires context-specific solution-finding through collaborative teamwork. Furthermore, the attention of guideline developers, health-care policymakers, funders and researchers requires direct focus on the ‘how’ and the ‘what’ of evidence implementation.

**Key words:** clinical practice guidelines; implementation of evidence; primary health care; qualitative description

*Received 13 December 2010; accepted 7 August 2011; first published online 13 October 2011*

---

Correspondence to: Dr Ann McKillop, Senior Lecturer, School of Nursing, Faculty of Medical and Health Sciences, The University of Auckland, Private Bag 92019, Auckland Mail Centre, Auckland 1142, New Zealand. Email: a.mckillop@auckland.ac.nz

## Introduction

The implementation of evidence-based guidelines in primary health care is expected to bring about

improved health outcomes, and early detection and reduction of the burden of disease. However, the precise nature of getting evidence into practice is so resistant to explanation that it has been described as occurring in a 'black box' (Sullivan *et al.*, 2008; Factor-Litvak and Sher, 2009). Consequently, and in spite of rigorously developed evidence-based guidelines being available and widely disseminated for a vast range of health problems, there is still a gap between evidence and practice (Miller and Kearney, 2004; Kitson, 2009). Because no 'magic bullets' for guideline implementation have been discovered (Shojania and Grimshaw, 2004; Brouwers *et al.*, 2009; Kitson, 2009), the uptake of evidence into practice remains poorly understood.

Approaches to the implementation of evidence have evolved over time from a primary focus on the nature of evidence itself ('science push') to the implementation behaviours of individual clinicians ('demand pull'), and then a combination of 'push-pull'. 'Science push' is a unidirectional process in which the benefit of using research evidence pushed out for clinicians to receive passively is expected to be self-evident (McWilliam *et al.*, 2009; Randell *et al.*, 2009). Passive distribution methods are still being used, perhaps for want of effective alternatives (Graham *et al.*, 2006). The consequences of the 'science push' approach to implementation has resulted in a lack of attention to implementation as a process and has contributed to the persisting gap between evidence and practice (Rycroft-Malone, 2006).

'Science push' has been combined with 'demand pull' in which sophisticated efforts were made to educate clinicians to pull in evidence that has been presented to them in increasingly palatable forms (Dopson *et al.*, 2002; Kuronen *et al.*, 2010). In 235 studies comparing 309 interventions, some improvements in care were associated with reminders (14.1%), some with educational materials (8.1%), audit and feedback (7%) and a mere 6% showed improvement using multifaceted educational outreach (Grimshaw *et al.*, 2004). However, with no definitive behavioural strategies found to improve the uptake of evidence, other approaches are required to inform the implementation of guidelines into practice.

Push and/or pull approaches to the implementation of evidence have failed to consistently influence practice decisions because they fail to

accommodate the messy world of health-care practice. The facilitation of evidence into practice requires methods that acknowledge the chaos and unpredictability of health-care settings and multiple layers of interaction and change cycles (Rycroft-Malone, 2008). Barriers to the adoption of evidence into practice have been attributed to a lack readiness of health professionals to change their practice, rather than a lack of willingness to change (McCormack *et al.*, 2008). Readiness to change practice is impaired by a lack of awareness, familiarity, agreement, self efficacy, motivation and expectations of success (Grol and Wensing, 2004; Brouwers *et al.*, 2009).

Evidence-based guidelines that address major health problems have high priority for implementation. Therefore, the implementation of a guideline for reducing cardiovascular risk was chosen as the focus of this study for the likelihood that practitioners would be familiar with it and be actively using it. Cardiovascular disease is the biggest killer worldwide (World Health Organization, 2008b) and is associated globally with marked health inequity linked to ethnicity and socioeconomic status (Thurston *et al.*, 2005; Kim *et al.*, 2008; Beauchamp *et al.*, 2010). New Zealand's indigenous Māori people, and indigenous peoples elsewhere (Baum, 2007; Marmot, 2007), suffer worse health overall than non-Māori and are disproportionately affected by cardiovascular disease, the main cause of the increasing difference in life expectancy between Māori and non-Māori (Tobias *et al.*, 2006). Because of the imperative and urgency to improve cardiovascular health, implementation of the Assessment and Management of Cardiovascular Risk (AMCVR) guideline (New Zealand Guidelines Group, 2003) is of high priority. Implementation of that guideline was chosen for this study because of its high profile. This study aimed to explore the realities of everyday nursing practice associated with the implementation of a guideline.

## Method

### Design

A qualitative research approach was used in this study in order to access rich descriptions of what are complex primary health-care nursing practices. Focus groups were held with primary health-care

nurses, doctors and managers associated with the implementation of the AMCVR guideline. Interviews were held with district health board primary health-care funder/planners. A general inductive approach was used to guide systematic condensing and summarising of textual data, and then linking of the summary of findings to the research objective. This method provides for interpretations that can be traced back to the words of the participants (Thomas, 2006).

### **Setting**

The setting of this study is a region of New Zealand with one small city, several towns and remote rural areas. Each of these locations has characteristics that impact differently on primary health-care delivery, and therefore participants were chosen from a variety of practice settings to gain maximum variation of perspectives. The population found in the study area was believed to be at higher risk of cardiovascular disease for two reasons: a higher median age (38.9 versus 35.9) and higher proportion of Māori (31.7% versus 14.6%) than the general New Zealand population (Statistics New Zealand, 2006).

### **Participants**

There were 33 participants, including 20 primary health-care nurses, four GPs, five primary health-care managers and three funder/planners. Primary health-care nurses were eligible for inclusion if their work was associated with the implementation of the AMCVR guideline. Doctors, managers and funder/planners were included for their perspectives of nurses' work as they implement the guideline. All participants were working in a region of high health need with significant health inequity for Māori, particularly cardiovascular health, so that their work with the guideline was expected to have a high profile. Potential participants who were nominated by their colleagues indicated their willingness to be included and were selected to gain maximum variation of perspectives of those working with the guideline.

### **Data collection**

Ethics approval was gained from the relevant regional New Zealand Ministry of Health Ethics Committee, the University of Technology, Sydney, Human Research Ethics Committee. Ethical

standards were maintained throughout the study. Seven focus groups of three to five participants (five groups of nurses, one of doctors and one of managers), and three individual interviews with funder/planners were held to discuss the topics identified in Table 1 below. The timing and venues for data collection were at the discretion of participants; all data collection took place in meeting rooms convenient to their work places and schedules. One researcher (AM) conducted all focus groups of an average of 65 minutes for nurses' groups, and 30 minutes for the GPs' and managers' groups, and three individual interviews with funder/planners of 30 minutes each. The groups and interviews were recorded and transcribed verbatim.

### **Data analysis**

A general inductive approach to data analysis was used to access and distil the informational content of the transcribed recordings of focus groups and interviews. The purposes of such an approach are to condense raw text into a summarised format, to establish clear and transparent links between the research aim and the findings and to produce a valid, overall representation of participants' perceptions (Thomas, 2006).

Data analysis involved five steps of induction informed by Thomas (2006) and Pope *et al.* (2000). Initially, the transcriptions were read through several times to get an overall sense of what was said. The second step began with systematic annotation of the transcripts identifying the text associated with topics mentioned and naming them provisionally for their informational content. Indexing followed, with the construction of a table for each transcript including assigning code names to segments of text and then aggregating codes into categories according to similarity of content. Constant comparison of codes to verbatim data and with each other ensured the best fit for aggregating codes firstly into 29 categories that were reduced to 27 categories on further review. Categories were then aggregated for similarity of content into four provisional themes. Criteria were developed for the informational content of each theme and the goodness of fit was reviewed by tracking back from categories to codes to the verbatim data. Allocation of codes to categories and categories to themes was reviewed once more

**Table 1** Interview topics

---

Sources of clinical information used currently in primary health care for assessment and management of cardiovascular risk
Ease of access to information (adapted for funder/planners and managers to contribution to guideline dissemination and utilisation)
Usefulness of the AMCVR guideline in everyday practice
Planning for and adaptation of guidelines for implementation
Primary health-care nurses' contribution to implementation of the AMCVR guideline
Supports and challenges of guideline implementation

---

AMCVR = Assessment and Management of Cardiovascular Risk.

(AM) for fit against the criteria for each theme and reviewed by another author for appropriate fit (JC). Through further review and discussion of the nature of the four themes, the three authors agreed on a further level of abstraction as an overarching concept that encompassed each of the four themes. Participants were not asked to check whether the results coincided with their personal views because the overall synthesis and abstraction involved in data interpretation and analysis produced findings in which individual participants may not recognise themselves specifically or their particular narrative (Morse *et al.*, 2002).

## Results

Data analysis generated four themes: self-managing patient, everyday nursing practice, developing new relationships in the health team and impact on health-care delivery. Each theme represents a facet of nurses' work with the cardiovascular guideline, and collectively they reveal the realities of everyday nursing practice associated with the implementation of the guideline. The concept of collaborative relationships drew together the essence of each theme and is presented as the overarching theoretical construct (Table 2).

### Self-managing patient

Nurses were adamant that implementing the guideline requires health care that empowers patient self-management. The 'self' in self-management could involve a range of family/whanau (Māori for family, usually wider than nuclear family) and/or

self/other partnerships. Two characteristics of patient self-management were revealed: empowerment and satisfaction. Nurse focus groups stressed the importance of 'letting the patient determine priorities' to 'set goals' and 'take control of their own health'. Conversation about patient self-management included acknowledgement of the complexity of the journey, both for patients and health professionals, and that readiness for change was an important factor in empowerment:

'You can't just fix it. You've got to allow people to decide what they want ... One of my kaumatua [Māori elder] says to me, "Why do you want to keep fixing me?"'  
(Practice Nurse)

Empowerment was valued highly by all participants as foundational to self-management:

'For the patient to become self-managing, that's the goal. It's knowing how to get there. A journey in itself...'  
(Practice Nurse)

Even though a patient's choices may not be optimally cardio-protective, nurse participants voiced their respect for a patient's right to decide the timing and pace of change: 'You can't just fix it. You've got to allow people to decide what they want ...' (Practice Nurse). Patient satisfaction was apparent from nurses' observations that patients appreciate intensive one-on-one education and planning associated with implementing the guideline.

'The feedback from the patients is just amazing and that something they mention a lot is that they really value their wellness plan because they own it. They're setting their own achievable small goals ... Instead of just going to their GP to get their three monthly prescriptions for meds'  
(Practice Nurse)

This theme underscored nurses' patient-centred approach to implementing the AMCVR guideline and their awareness of the importance of building relationships to optimise patients' self-care.

### Everyday nursing practice

Everyday nursing practice represented how nurses worked towards the integration of guideline recommendations into usual health care overall

**Table 2** Overarching theoretical construct with associated themes and categories

Collaborative relationships			
Self-managing client	Everyday nursing work	Developing new relationships in health team	Impact on health-care delivery
Client satisfaction	Nurses relating to nurses	Focusing on and accessing population health needs	Māori health providers work differently
Client empowerment	Taking a whole person approach Partnering for client buy-in Practicing in tune with community Brokering knowledge and resources Making the most of experience	Using a systematized approach Working together differently Communicating with the team New understandings Using the guideline tools target population	Funding issues Difficulties, challenges Changes in service delivery Regional cohesion Challenges to workforce capacity

so that it was embedded in normal practice. Nurses spoke about the need to use knowledge gained from experience and to tap into a wide variety of community-based resources to help patients make lifestyle changes. Cardiovascular risk assessment was seen by nurses as just one aspect of overall health assessment.

Nurses were aware that the level of deprivation that a person experienced was closely associated with their level of cardiovascular risk. The whole person view required an understanding of overall life circumstances, support needs and pace of change. One nurse group spoke about offering comprehensive care:

‘I don’t care about cv risk [whispering] I care about the relationship I have with them [patients]. Because for us we’re a community development team. We look at all of the family ... so if there’s any older people in the house, “Do you need anything? We have a home-based nursing service. Do you need some home-care for your elderly?”’

Goal setting had to be realistic, and for some patients multi-generational poverty had created a sense of hopelessness:

‘There are a lot of social factors you know. People who are unemployed. They never really get a good job. They never have that steady income. There are a lot of things that are out of their control’

(Practice Nurse)

Cardiovascular health as a life-long goal required ongoing support for patients. Participants agreed that sustained support following risk assessment requires enduring relationships. Nurse participants appreciated the importance of being available and accessible to their patients:

Nurse 1 – ‘And they come here because they know that we care about them’.

Nurse 2 – ‘We’ll listen to them’.

Nurse 3 – ‘We’ll take the time’.

However, they were also concerned that mass screening would create heavy demands not only on health-care services, but would also expose the need for coordination of community-based help for people:

‘... you have to have services to refer people to. No point in identifying a risk if you can’t do anything about it. And it has been about making sure that everyone has access to the range of services ... I mean it’s not just a matter of saying “You’ve got a risk here. Take a pill”; “You’ve got transport? Na”- those sort of things ...’

(Manager)

In spite of their concerns about implementing the guideline:

‘... nurses have just jumped at this whole new approach.... It’s as if the nurses are doing what they became a nurse for ... They’ve actually become, like “I’m home”’

(Practice Nurse)

Nurse focus groups spoke about how they valued their relationships with each other, but were concerned about a lack of communication about how to implement the guideline. The guideline recommended what to do but not how to do it and they were not sharing the know-how of their practice.

### **Developing new relationships in the health team**

Implementation of the guideline in a changing health-care environment has resulted in clinicians working together differently. The sheer volume of work involved in cardiovascular screening and follow-up has been one factor that has contributed to a realisation of the largely untapped potential of nurses in general practice. Changes in practice associated with implementing the guideline were more positive for nurses than for doctors. As nurses were taking on a more satisfying role in direct patient care, doctors also needed to change their practice in a number of ways that required new understandings in the health-care team.

As primary health-care nurses adopted a more active role in patient consultations, there was a noticeable change in how doctors and nurses were relating:

‘I think for me it’s meant that you work with the GP as a team and I think over the last few years, with a lot of the things that we’ve been doing it’s become a team effort... gives you a feeling of trust and respect’

(Practice Nurse)

Working together differently resulted in better teamwork and made a difference to patients’ knowledge and confidence. One of the nurse groups found that:

‘People with chronic diseases don’t come in asking about their pills. With better education they’re more at ease about how things are going because they are more in control. So the doctors are rapt because it saves them time. So it’s good teamwork’

(Practice Nurse)

In addition, nurse-led clinics were working well for cardiovascular risk assessment and ongoing self-management:

‘We’ve run a nurse-led clinic for many years so it allows some independence for a lot of

the nurses and I’d say a lot of the CVD risk is initiated by the practice nurse’

(Practice Nurse)

Even though GPs were aware of the health promotion needs of their enrolled populations, they were overwhelmed with treating those with illness, especially in areas of high health need where there are:

‘... just over 70% Māori and a fairly high level of deprivation and high morbidity which puts a heavy workload to cover that. And lots of comorbidities. ... our doctor-nurse ratio is pretty high and our patient-doctor ratio is pretty high ... we’re stretched’

(GP)

The GP group indicated they were at a loss as to how to encourage healthy people to come forward for cardiovascular screening. They were more at ease with ‘explaining to them (patients who present for treatment) what they should do’ (GP). Māori health-care providers had more active relationships with the entire population they served (mainly Māori) than general practice providers who tend to see only those patients who came through the door of their clinic with a problem or needing a prescription repeat. As one funder/planner put it:

‘Māori providers themselves know how to access and link [with the Māori population]. [But for others] there are cross-cultural issues getting the right service to the right people at the right time’

(Funder/planner)

### **Impact on health-care delivery**

The last theme concerned the impact on guideline implementation of difficulties in communication, integration and coordination of care across the region and the lack of resources available to services. All the clinician groups conveyed a sense of despair that they did not have the capacity for integrated and coordinated cardiovascular risk profiling, mass cardiovascular screening and the follow-up that was required after assessment.

A common concern and frustration for all participant groups was the lack of funding to implement the guideline. The GP group was

particularly despondent about the lack of financial assistance:

Doctor 1 – ‘That’s what the problem is. You know the guideline is being pushed to save money at the other end. It makes sense doesn’t it, economically?’

Doctor 2 – ‘But we really need to be resourced at this end. I’m amazed. ... No resourcing at this end! You must be mad. It’s just a crazy situation! It’s not going to work! It’s all gonna fall flat!’

A practice nurse group was equally at loss as to how they were going to implement the guideline within existing resources:

Nurse 2 – ‘There’s no point screening if you can’t follow up with what’s required ...’

Nurse 3 – ‘... there was nothing actually about what you gonna do when you’ve found them! Who’s gonna pay for all this extra work that is going to be done?’

A lack of coordination and data sharing among practices interfered with community-based mass screening of the target population. Screening was a waste of time if the clinical data and risk score were not accessible to those who would follow-up with the patient. At least when screening was held at a GP clinic, the patients coming through the door were likely to be enrolled at that practice. As one nurse participant explained:

‘... going into work places, they may not be our patients ... If you identify a risk, how are you gonna manage non-registered patients?’

All focus groups were concerned about the capacity of the primary health-care workforce to implement all of the guideline recommendations and were at a loss to know how to cope especially with screening and early detection. Nurses, doctors and managers spoke about the challenges of clinical staff shortages and the need for upskilling health professionals. The GP group acknowledged that although most of the work involved in implementing the guideline could be carried out by practice nurses (eg, risk assessment, patient education and follow-up), there were not enough of them to do that.

The four themes depict motivated, person-centric, health professional teams who agree with

the intentions of the guideline, but are struggling to find ways to implement the recommendations for prevention and screening, given their existing workload. The notion of collaborative relationships was central to each of the themes and formed an overarching connection among them. In the first theme, a mutually trusting relationship between clinicians and patients in supporting self-management was central to all other relationships involved in implementing the guideline. The second theme, everyday nursing practice, focussed on the need to build professional know-how through sharing insights gained from weaving the guideline recommendations into usual care. The third theme, developing new relationships in the health team, revealed that interdisciplinary collaboration enhanced patient–clinician collaboration and engendered better doctor–nurse teamwork. The fourth theme captured the frustration of clinicians struggling to gain the resources required for the communication, integration and coordination of care to support guideline implementation.

## Discussion

Even though the AMCVR guideline provided high-quality evidence for what to do, it failed to satisfy health-care professionals’ need to know how to implement the recommendations. In spite of their clear willingness to work with the guideline, a lack of knowing how to do this successfully has caused confusion and frustration and led to fragmentation of services. The importance of collaborative relationships between all players involved in implementation of the guideline in this study has been well represented in the literature (D’Amour *et al.*, 2005; Reeves *et al.*, 2009; Shultz and Kitson, 2010; Gaboury *et al.*, 2011; Jackson and Bluteau, 2011; Schmidt, 2011). This study has underscored not only the importance of collaborative relationships between health professionals, but also the crucial role of relationships with patients, their families and agencies outside a general practice in the implementation of this guideline.

Person-centric values and beliefs about concordance, empowerment and working together with patients were found to be pivotal in this study to implementing the AMCVR guideline because, ultimately, enacting the recommendations is at the patient’s discretion. Concordance refers to

partnership in decision-making between health professionals and patients in which patients' preferences are elicited, valued and respected in the context of co-interpretation and co-conceptualisation (Parker, 2005). An essential element of concordant relationships is cultural competence (Betancourt *et al.*, 2005). Cultural competence is vitally important in health care and has special meaning in New Zealand in light of the imperative to reduce health inequity for Māori (Cram *et al.*, 2003; Hefford *et al.*, 2005). A holistic and relational Māori view of health is an essential element in health and healing, including the importance of wairua (spirit) and whanau (family). From a Māori patient's point of view, health care that has been designed to be culturally appropriate, affordable and accessible gains a high level of patient satisfaction (Maniapoto and Gribben, 2003). Further research is required to develop knowledge about culturally competent care, which addresses cardiovascular health inequity through implementing the AMCVR guideline.

The role of the patient as principal implementer of evidence in primary health care has been underscored in this study, a factor scarcely recognised in the guideline implementation literature that has so far been more focussed on the behaviours and attitudes of health professionals than on relationships with patients (Grol and Grimshaw, 2003; DiCenso *et al.*, 2005; Kuronen *et al.*, 2010), even though reliable predictors of change in professional practice have not been found (Hulscher *et al.*, 2005). In particular, there is a paucity of research about primary health-care practice that leads to health promoting actions of patients (Hrisos *et al.*, 2009).

Ideally, primary health care for individuals has a population-based approach that addresses inequity at the level of the social determinants of health (Keller *et al.*, 2004; Neuwelt *et al.*, 2009; Edgcombe and Stephens, 2010). Primary health care is ineffective if it fails to address socio-economic disparities as the most direct means of improving population health (Gervas *et al.*, 2008; World Health Organization, 2008a; Goldberg, 2009). Participants in this study were aware that treating the biophysical effects of disease characterises a 'short-term, problem specific, individual-based "downstream" approach' (Cypress, 2004: 249). Upstream actions are required that strengthen the determinants of health – education, employment, housing, nutrition, income and working conditions

(Marmot, 2003; Keller *et al.*, 2004; Sharpe, 2006; World Health Organization, 2008a). However, in spite of many guidelines for primary health care, professionals are frustrated and powerless to know how to implement evidence in a way that influences the social determinants of health in vulnerable populations.

A traditional model of clinic-based, illness-focussed primary health care is not conducive to the provision of care that includes health promotion and disease prevention (Workforce Taskforce, 2008; Kuronen *et al.*, 2010). Even with the best of intentions, such a care environment is bound to cater best for patients attending for treatment rather than delivering on population health outcomes (Daniels *et al.*, 2000; Marmot, 2003; Goldberg, 2009). The people, processes and structures that impact on care co-create the culture of a clinical environment and are equally implicated in cultural change (Smith, 2009; McCormack and McCance, 2010). Cultural change is difficult to achieve in any clinical setting; however, geographic distances between numerous small general practices in rural settings present challenges to communication and collaboration.

Solutions to difficult clinical practice challenges require active processes of facilitated support (Bandura, 2000; Bandura, 2006; Manley, 2008; McCormack *et al.*, 2008; Walsh *et al.*, 2008). Health professionals' awareness of the interactions between their work activities and the social context of their practice positions them well to envision change (Manley *et al.*, 2009; Nairn, 2009). Clinicians in this study recognised the benefits of working more collegially and extending their everyday relationships to include other health and social services professionals. The positive association that has been found between team-level job satisfaction and quality of care (Mohr *et al.*, 2011; Roblin *et al.*, 2011), as well as the role of interagency collaboration in problem solving (Daley, 2009) underlines the importance of collaborative relationships within a team and across organisations.

How things are done and what counts as important reflects the inherent values and beliefs that shape a clinical context (Walsh *et al.*, 2006; McCormack and McCance, 2010), and the work involved in co-creating new ways of working results in improved collegial relationships (Manley, 2004). Active collaboration across several general practices at some distance apart may be achieved

through the development of communities of practice in which clinicians use video conferencing, web-based social networking and file sharing to support case-review for the purpose of making sense of their experiences, share concerns, reflect on approaches to care and develop new care strategies (Soubhi *et al.*, 2010). Further research is required to explore the process and outcome/s of social participation, collective sense-making and bridging the gap between knowing and doing in communities of practice (Soubhi *et al.*, 2010).

Not all barriers to the successful implementation of a guideline can be addressed by clinicians in their workplace. Some problems require infra-structural, multi-sectorial, governance, policy and funding interventions (Workforce Taskforce, 2008). For example, health inequity in indigenous peoples is a worldwide issue that must be addressed at all levels in health-care systems (Bacal *et al.*, 2006). This study has emphasised the problems of capacity and capability of the primary health-care workforce to address health inequity for their Māori patients. Others have called for workforce development specifically focussed at recruiting, retaining and strengthening a culturally appropriate health and disability workforce (Ramsden, 1997; Durie, 2003; Ratima *et al.*, 2007). Furthermore, the structural and socio-economic causes of health inequity must be addressed at the highest level of governance, funding and policy (Marmot, 2010).

The findings have revealed the realities of the implementation of the AMCVR guideline in primary health care. Nurses were committed to improving their practice in spite of environmental constraints on their work. However, there was a pervading sense of despair and frustration about implementing the AMCVR guideline because of a lack of knowing how to do that. The purpose of this study was to provide detailed understanding of guideline implementation for these participants rather than to produce generalisable findings. However, as suggested by Ritchie and Lewis (2003), when there is homogeneity of the clinical work of several participants, a degree of theoretical transferability is possible.

## Conclusions and implications

The findings of this study have provided insight into the essential nature of collaborative relationships

*Primary Health Care Research & Development* 2012; 13: 48–59

in knowing not only what is recommended in a guideline, but also how to weave the recommendations into everyday practice. Health professionals in this study showed that they value the guideline, have a person-centred orientation to care and are willing to implement the guideline, but that skilled facilitation is required for the discovery of new ways of working that are appropriate to the context of their everyday work. The most obvious implication for clinicians is the absolute need for support, processes and resources to develop evidence that informs how to implement guideline recommendations, in addition to recommendations for health-care interventions. Health-care funders and policymakers need to find long-term solutions to the problems of the current funding and service model of primary health care, which addresses the failure to adequately compensate and incentivise health-care providers for the implementation of clinical practice guidelines, especially for preventative health-care interventions. A further implication of this study is that further investment is required in development of clinical leadership at all levels of primary health-care governance and provision.

## Acknowledgement

The Ministry of Health, New Zealand, provided funding support for data collection for this study.

## References

- Bacal, K., Jansen, P. and Smith, K.** 2006: Developing cultural competency in accordance with the Health Practitioners Competence Assurance Act. *New Zealand Family Practice* 33, 305–09.
- Bandura, A.** 2000: Exercise of human agency through collective efficacy. *Current Directions in Psychological Science* 9, 75–78.
- Bandura, A.** 2006: Toward a psychology of human agency. *Perspectives on Psychological Science* 1, 164–80.
- Baum, F.** 2007: Cracking the nut of health equity: top down and bottom up pressure for action on the social determinants of health. *Global Health Promotion* 14, 90–95.
- Beauchamp, A., Peeters, A., Wolfe, R., Turrell, G., Harriss, L.R., Giles, G.G., English, D.R., McNeil, J., Magliano, D., Harrap, S., Liew, D., Hunt, D. and Tonkin, A.** 2010: Inequalities in cardiovascular disease mortality: the role of behavioural, physiological and social risk factors. *Journal of Epidemiology and Community Health* 64, 542–48.

- Betancourt, J.R., Green, A.R., Carrillo, J.E. and Park, E.R.** 2005: Cultural competence and health care disparities: key perspectives and trends. *Health Affairs* 24, 499–508.
- Brouwers, M., Hanna, S., Abdel-Motagally, M. and Yee, J.** 2009: Clinicians' evaluations of, endorsements of, and intentions to use practice guidelines change over time: a retrospective analysis from an organized guideline program. *Implementation Science* 4, 34. DOI:10.1186/1748-5908-4-34.
- Cram, F., Smith, L. and Johnstone, W.** 2003: Mapping the themes of Maori talk about health. *New Zealand Medical Journal* 116. Retrieved 10 October 2009 from <http://www.nzma.org.nz/journal/116-1170/357/>
- Cypress, M.** 2004: Looking upstream. *Diabetes Spectrum* 17, 249–53.
- D'Amour, D., Ferrada-Videla, M., San Martin Rodriguez, L. and Beaulieu, M.-D.** 2005: The conceptual basis for interprofessional collaboration: core concepts and theoretical frameworks. *Journal of Interprofessional Care* 19, 116–31.
- Daley, D.M.** 2009: Interdisciplinary problems and agency boundaries: exploring effective cross-agency collaboration. *Journal of Public Administration Research and Theory* 19, 477–93.
- Daniels, N., Kennedy, B. and Kawachi, I.** 2000: *Is inequality bad for our health?*. Boston, MA: Beacon Press.
- Dicenso, A., Guyatt, G. and Ciliska, D.** 2005: *Evidence-based nursing: a guide to clinical practice*. St. Louis, MO: Elsevier Mosby.
- Dopson, S., Fitzgerald, L., Ferlie, E., Gabbay, J. and Locock, L.** 2002: No magic targets! Changing clinical practice to become more evidence-based. *Health Care Management Review* 27, 35–47.
- Durie, M.H.** 2003: The health of indigenous people. *British Medical Journal* 326, 510–11.
- Edgecombe, G. and Stephens, R.** 2010: Healthy communities: the evolving roles of nursing. In Daly, J., Speedy, S. and Jackson, D., editors, *Contexts of Nursing*, third edition. Sydney, Australia: Churchill Livingstone, 274–86.
- Factor-Litvak, P. and Sher, A.** 2009: Invited commentary: coming out of the box. *American Journal of Epidemiology* 169, 1179–81.
- Gaboury, I., M. Lapierre, L., Boon, H. and Moher, D.** 2011: Interprofessional collaboration within integrative healthcare clinics through the lens of the relationship-centered care model. *Journal of Interprofessional Care* 25, 124–30.
- Gervas, J., Starfield, B. and Heath, I.** 2008: Is clinical prevention better than cure? *The Lancet* 372, 1997–99.
- Goldberg, D.S.** 2009: In support of a broad model of public health: disparities, social epidemiology and public health causation. *Public Health Ethics* 2, 70–83.
- Graham, I.D., Logan, J., Harrison, M.B., Straus, S., Tetroe, J., Casswell, W. and Robinsons, N.** 2006: Lost in knowledge translation: time for a map? *Journal of Continuing Education in the Health Professions* 26, 13–24.
- Grimshaw, J.M., Thomas, R., MacLennan, G., Fraser, C., Ramsay, C., Vale, L., Whitty, P., Eccles, M., Matowe, L., Shirran, L., Wensing, M., Dijkstra, R. and Donaldson, C.** 2004: Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health Technology Assessment* 8, iii–72.
- Grol, R. and Grimshaw, J.** 2003: From best evidence to best practice: effective implementation of change in patients' care. *The Lancet* 362, 1225–29.
- Grol, R. and Wensing, M.** 2004: What drives change? Barriers to and incentives for achieving evidence-based practice. *Medical Journal of Australia* 180, S57–60.
- Hefford, M., Crampton, P. and Foley, J.** 2005: Reducing health disparities through primary care reform: the New Zealand experiment. *Health Policy* 72, 9–23.
- Hrisos, S., Eccles, M.P., Francis, J.J., Dickinson, H.O., Kaner, E.F., Beyer, F. and Johnston, M.** 2009: Are there valid proxy measures of clinical behaviour? A systematic review. *Implementation Science* 4, 37. DOI:10.1186/1748-5908-4-37.
- Hulscher, M.E.J., Wensing, M., Weijden, T.V.D. and Grol, R.** 2005: Interventions to implement prevention in primary care. *The Cochrane Database of Systematic Reviews* 4. CD000362.
- Jackson, A. and Bluteau, P.** 2011: Interprofessional education, collaborative practice and primary care. *InnovAiT* 4, 230–35.
- Keller, L., Strohschein, S., Lia-Hoagberg, B. and Schaffer, M.** 2004: Population-based public health interventions: practice-based and evidence-supported. Part 1. *Public Health Nursing* 21, 453–68.
- Kim, D., Kawachi, I., Vanderhoorn, S. and Ezzati, M.** 2008: Is inequality at the heart of it? Cross-country associations of income inequality with cardiovascular diseases and risk factors. *Social Science and Medicine* 66, 1719–732.
- Kitson, A.** 2009: Knowledge translation and guidelines: a transfer, translation or transformation process? *International Journal of Evidence-Based Healthcare* 7, 124–39.
- Kuronen, R., Jallinoja, P., Airola, R. and Patja, K.** 2010: Educational implementation programme of guidelines on cardiovascular risk factors: an analysis of changes in familiarity, use and attitudes. *Primary Health Care Research & Development* 11, 241–49.
- Maniapoto, T. and Gribben, B.** 2003: Establishing a Maori case management clinic. *New Zealand Medical Journal* 116, 328, available at <http://www.nzma.org.nz/journal/116-1169/328/>
- Manley, K.** 2004: Transformational culture: a culture of effectiveness. In McCormack, B., Manley, K. & Garbett, R., editors, *Practice development in nursing*. Oxford, England: Blackwell, 51–82.
- Manley, K.** 2008: 'The way things are done around here' – developing a culture of effectiveness: a pre-requisite to individual and team effectiveness in critical care. *Australian Critical Care* 21, 83–85.
- Manley, K., Titchen, A. and Hardy, S.** 2009: Work-based learning in the context of contemporary health care education and practice: a concept analysis. *Practice Development in Health Care* 8, 87–127.
- Marmot, M.** 2003: Understanding social inequalities in health. *Perspectives in Biology and Medicine* 46, S9–23.

- Marmot, M.** 2007: Achieving health equity: from root causes to fair outcomes. *The Lancet* 370, 1153–63.
- Marmot, M.** 2010: *Fair society, healthy lives: the Marmot review*. London: Department of Health.
- McCormack, B.** and **McCance, T.** 2010: *Person-centred nursing*. Oxford: Wiley-Blackwell.
- McCormack, B., Manley, K.** and **Walsh, K.** 2008: Person-centred systems and processes. In Manley, K., McCormack, B. and Wilson, V., editors, *International practice development in nursing and healthcare*. Oxford, England: Blackwell.
- McWilliam, C.L., Kothari, A., Ward-Griffin, C., Forbes, D.** and **Leipert, B.** 2009: Evolving the theory and praxis of knowledge translation through social interaction: a social phenomenological study. *Implementation Science* 4, 26. DOI: 10.1186/1748-5908-4-26.
- Miller, M.** and **Kearney, N.** 2004: Guidelines for clinical practice: development, dissemination and implementation. *International Journal of Nursing Studies* 41, 813–21.
- Mohr, D.C., Young, G.J., Meterko, M., Stolzmann, K.L.** and **White, B.** 2011: Job satisfaction of primary care team members and quality of care. *American Journal of Medical Quality* 26, 18–25.
- Morse, J., Barrett, M., Mayan, M., Olson, K.** and **Spiers, J.** 2002: Verification strategies for establishing reliability and validity in qualitative research. *International Journal of Qualitative Methods* 1, 13–22.
- Nairn, S.** 2009: Social structure and nursing research. *Nursing Philosophy* 10, 191–202.
- Neuwelt, P., Matheson, D., Arroll, B., Dowell, A., Winnard, D., Crampton, P., Sheridan, N.** and **Cumming, J.** 2009: Putting population health into practice through primary health care. *New Zealand Medical Journal* 122. Retrieved 6 July 2010 from <http://www.nzma.org.nz/journal/122-1290/3484/>
- New Zealand Guidelines Group.** 2003: *The assessment and management of cardiovascular risk*. Wellington, New Zealand.
- Parker, M.** 2005: False dichotomies: EBM, clinical freedom, and the art of medicine. *Journal of Medical Ethics; Medical Humanities* 31, 23–30.
- Pope, C., Ziebland, S.** and **Mays, N.** 2000: Qualitative research in health care: Analysing qualitative data. *BMJ* 320, 114–16.
- Ramsden, I.** 1997: Cultural safety: implementing the concept. The social force of nursing and midwifery. In Te Whaiti, P., McCarthy, M. and Durie A., editors, *Mai i Rangiatea: Mapri well being and development*. Auckland: Auckland University Press and Bridget Williams Books, 113–25.
- Randell, R., Mitchell, N., Thompson, C., McCaughan, D.** and **Dowding, D.** 2009: From pull to push: understanding nurses' information needs. *Health Informatics Journal* 15, 75–85.
- Ratima, M., Brown, R., Garrett, N., Wikaire, E., Ngawati, R., Aspin, C.** and **Potaka, U.** 2007: *Rauringa Raupa recruitment and retention of Māori in the health and disability workforce*. Auckland, New Zealand, Taupua Waiora: Division of Public Health and Psychosocial Studies.
- Reeves, S., Zwarenstein, M., Goldman, J., Barr, H., Freeth, D., Hammick, M.** and **Koppel, I.** 2009: Interprofessional education: effects on professional practice and health care outcomes (Review). *The Cochrane Library*, Issue 4, available at <http://www.thecochranelibrary.com>
- Ritchie, J.** and **Lewis, J.** 2003: *Qualitative research practice: a guide for social science students and researchers*. London/Thousand Oaks/New Delhi: Sage.
- Roblin, D.W., Howard, D.H., Junling Ren,** and **Becker, E.R.** 2011: An evaluation of the influence of primary care team functioning on the health of medicare beneficiaries. *Medical Care Research and Review* 68, 177–201.
- Rycroft-Malone, J.** 2006: The politics of the evidence based practice movements: legacies and current challenges. *Journal of Research in Nursing* 11, 95–108.
- Rycroft-Malone, J.** 2008: Evidence-informed practice: from individual to context. *Journal of Nursing Management* 16, 404–08.
- Schmidt, K.** 2011: *The organization of cooperative work*. London: Springer eBook, Springer.
- Sharpe, N.** 2006: Heart health has an adverse future forecast in New Zealand: an alarm call to action across the continuum. *New Zealand Medical Journal* 119. Retrieved 12 October 2010 from <http://www.nzma.org.nz/journal/119-1232/1936/>
- Shojania, K.G.** and **Grimshaw, J.M.** 2004: Still no magic bullets: pursuing more rigorous research in quality improvement. *American Journal of Medicine* 116, 778–80.
- Shultz, T.J.** and **Kitson, A.L.** 2010: Measuring the context of care in an Australian acute care hospital: a nurse survey. *Implementation Science* 5. available at <http://www.biomedcentral.com/content/pdf/1748-5908-5-60.pdf>
- Smith, J.** 2009: *Critical analysis of the implementation of the primary health care strategy and framing of issues for the next phase*. Wellington, New Zealand: Ministry of Health.
- Soubhi, H., Bayliss, E.A., Fortin, M., Hudon, C., Van den Akker, M., Thivierge, R., Posel, N.** and **Fleischer, D.** 2010: Learning and caring in communities of practice: using relationships and collective learning to improve primary care for patients with multimorbidity. *Annals of Family Medicine* 8, 170–77.
- Statistics New Zealand.** 2006: QuickStats About Northland Region.
- Sullivan, G., Blevins, D.** and **Kauth, M.** 2008: Translating clinical training into practice in complex mental health systems: toward opening the 'Black Box' of implementation. *Implementation Science* 3, 33. DOI:10.1186/1748-5908-3-33.
- Te Ara.** 2009: *Northland region population*. New Zealand Government. Retrieved 21 May 2004 from <http://www.teara.govt.nz/en/northland-region/4>
- Thomas, D.R.** 2006: A general inductive approach for analyzing qualitative evaluation data. *American Journal of Evaluation* 27, 237–46.
- Thurston, R.C., Kubzansky, L.D., Kawachi, I.** and **Berkman, L.F.** 2005: Is the association between socio-economic position and

coronary heart disease stronger in women than in men? *American Journal of Epidemiology* 162, 57–65.

**Tobias, M., Sexton, K., Mann, S. and Sharpe, N.** 2006: How low can it go? Projecting ischaemic heart disease mortality in New Zealand to 2015. *New Zealand Medical Journal* 119. Retrieved 8 September 2009 from <http://www.nzma.org.nz/journal/119-1232/1932/>

**Walsh, K., Moss, C. and Fitzgerald, M.** 2006: Solution-focused approaches and their relevance to practice development. *Practice Development in Health Care* 5, 145–55.

**Walsh, K., Moss, C., Lawless, J., McKelvie, R. and Duncan, L.** 2008: Puzzling practice: a strategy for working with clinical

practice issues. *International Journal of Nursing Practice* 14, 94–100.

**Workforce Taskforce.** 2008: *Working together for better primary health care: Overcoming barriers to workforce change and innovation.* Wellington, New Zealand: Ministry of Health.

**World Health Organization.** 2008a: *Primary health care – now more than ever. The World Health Report.* Geneva: World Health Organization.

**World Health Organization.** 2008b: *The top ten causes of death* [Online]. Retrieved 10 June 2010 from [http://www.who.int/mediacentre/factsheets/fs310\\_2008.pdf](http://www.who.int/mediacentre/factsheets/fs310_2008.pdf)