

2012

Pesticide removal by a mixed culture of bacteria and white-rot fungi

Faisal I. Hai

University of Wollongong, faisal@uow.edu.au

Oska Modin

University of Tokyo

Kazuo Yamamoto

University Of Tokyo

Kensuke Fukushi

University Of Tokyo

Fumiyuki Nakajiam

*University Of Tokyo**See next page for additional authors*

<http://ro.uow.edu.au/engpapers/4417>

Publication Details

Hai, F. I., Modin, O., Yamamoto, K., Fukushi, K., Nakajiam, F. & Nghiem, L. D. (2012). Pesticide removal by a mixed culture of bacteria and white-rot fungi. *Journal of the Taiwan Institute of Chemical Engineers*, 43 (3), 459-462.

Authors

Faisal I. Hai, Oska Modin, Kazuo Yamamoto, Kensuke Fukushi, Fumiyuki Nakajiam, and Long Nghiem

Pesticide Removal by a Mixed Culture of Bacteria and White Rot Fungi

Faisal Ibney Hai^{1,4,* α , ζ} , Oskar Modin^{2, β , ζ} , Kazuo Yamamoto¹, Kensuke Fukushi³, Fumiyuki Nakajima¹ and Long Duc Nghiem⁴

¹Environmental Science Center, The University of Tokyo,
7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan.

²Department of Urban Engineering, The University of Tokyo,
7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-8656, Japan.

³Integrated Research System for Sustainability Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku,
Tokyo, 113-8654, Japan.

⁴Strategic Water Infrastructure Laboratory, School of Civil, Mining and Environmental Engineering, University
of Wollongong, NSW 2522, Australia

* Corresponding author: Faisal Ibney Hai

^{α} Current address: Faisal Ibney Hai

School of Civil, Mining and Environmental Engineering

The University of Wollongong, NSW 2522, Australia

Email: faisal@uow.edu.au; Ph +61 2 4221 31054

^{β} Current address: Oskar Modin

Department of Civil and Environmental Engineering, Water Environment Technology section

Chalmers University of Technology, Göteborg, Sweden

Email: oskar.modin@chalmers.se; Ph +46 31 772 2138

^{ζ} Equal contribution

Abstract

Combining activated sludge cultures with microbes harboring specific degradation pathways could constitute a relevant process for the removal of toxic and recalcitrant organic substances from wastewater. Enhanced removal of three widely used recalcitrant pesticides from their liquid mixture was demonstrated by implementing a non-acclimated mixed culture of bacteria and white rot fungus. During an incubation period of 14 days, the mixed fungus-bacteria culture achieved 47, 98, and 62% removal of aldicarb, atrazine and alachlor from the liquid phase, respectively. This compared favorably to batches containing only non-acclimated fungus or previously published removal rates with non-acclimated bacterial cultures. Biosorption along with biodegradation was responsible for the removal of the pesticides from the liquid phase. Potential application modes of the studied biodegradation process were also discussed.

Keywords: bioaugmentation, biodegradation, biosorption, pesticide, white rot fungi

Hai et al. (2012) Journal of the Taiwan Institute of Chemical Engineers, (43), 459-462

1 **1. Introduction**

2 The use of pesticides is ubiquitous in modern agriculture and is important to increase crop yield and
3 reduce post-harvest losses. However, indiscriminate and excessive use of agricultural pesticides can
4 lead to contamination of land and water. Emissions of pesticides come from both diffuse and point
5 sources. The latter include mixing and loading facilities on the farm where spillages and leakages
6 from the filling operation and spray equipment, and water from rinsing and cleaning of the
7 equipment may contribute to pesticide contamination [1]. Wastewater generated in vegetable
8 washing facilities and pesticide manufacturing plants are also important point sources of pollution.

9 Many pesticides are recalcitrant compounds and persist for long periods of time in the
10 environment. Pesticides have been detected in ground- and surface waters used for potable water
11 supply and have been linked to adverse human health effects [2]. The adverse effects of agricultural
12 pesticide contamination extend to birds, wild animals and plants in the aquatic habitat. Because of
13 the widespread use of pesticides in modern agricultural practice and their potential threat to the
14 environment, the formulation of treatment methods for the removal of these pesticides from
15 contaminated water is essential.

16 Biological treatment technologies are usually inexpensive and effective for a range of
17 organic contaminants. However, owing to the recalcitrant structures of the pesticides, only specific
18 bacterial and fungal species have been reported to be capable of degrading them [3]. Interestingly,
19 in the soil environment, bacteria and fungi seem to adopt different but complementary metabolic
20 pathways for the degradation of these recalcitrant pesticides [4]. For instance, the contributions of
21 both bacteria and fungi to the degradation of a chloroaromatic fungicide - chlorothalonil - in soil
22 were confirmed by a selective inhibition method [5].

23

24 To date, the number of studies investigating novel treatment techniques for the removal of
25 pesticides from contaminated agricultural wastewater remains limited. The bacteria-dominated
26 conventional activated sludge process has been proved to be ineffective for pesticide removal.

27 While the importance of a mixed microbial community to initiate and complete pesticide removal
28 in the soil environment has been convincingly demonstrated by several researchers, studies
29 concerning the removal of pesticides from water/wastewater have been predominantly focused on
30 selected bacterial or fungal species separately [6]. For instance, white rot fungi have been
31 demonstrated to possess excellent capacity to initiate degradation of a wide variety of pesticides by
32 means of their nonspecific extracellular enzyme systems. A few studies have explored the
33 bioaugmentation synergy of enriched bacterial cultures with the conventional activated sludge [7-
34 8]. However, there appears to be no report on the combination of fungi and bacteria to remove
35 pesticide from the aqueous phase.

36 With the aim to address the important knowledge gap illustrated above, in this study the
37 removal of three pesticides from their mixture using a mixed culture of bacteria and white rot fungi
38 was investigated. The removal efficiency of the dissolved parent compounds were compared to a
39 culture with only white-rot fungi and to previously published values with bacterial cultures. The
40 importance, limitations, and potential application modes of the studied biodegradation process were
41 also discussed.

42 **2. Materials and methods**

43 *2.1 Selected pesticides*

44 Three pesticides, namely, aldicarb, atrazine, and alachlor were selected in view of their
45 widespread use and reported linkage to water pollution. Aldicarb is an N-methyl carbamate; it is an
46 insecticide/ nematocide used on a wide range of crops including cotton, potatoes, sugarcane, and is
47 applied as granules to the soil [3]. Atrazine is a triazine and is used as a herbicide on, for example,
48 forests, grass, corn, and sorghum. Alachlor is a chloroacetanilide and also used as a herbicide on,
49 for example, corn and beans. Stock solutions (500 mg L⁻¹) of each of the three pesticides were
50 prepared in pure methanol.

51 *2.2 Preparation of bioaugmented culture and nutrient medium*

52 The white-rot fungus *Coriolus versicolor*, NBRC 9791 obtained from the NITE Biological
53 Resource Center (NBRC), Japan was used for this study. Full-grown and enzyme secreting large
54 flocs of fungi [9] were disintegrated into fine grains by sonication (Branson sonifier 450, CT, USA)
55 for 5 minutes under sterile condition. The resulting biomass was then incubated for a week into 500
56 mL beakers each containing 100 mL growth medium at the optimum growth temperature of 28 °C
57 on a shaker (BR-300LF, Taitec rotary bio-shaker, Japan) at 80 rpm. At the end of the incubation
58 period approximately 0.5 cm granules exhibiting an enzymatic activity of approximately 30 μM
59 substrate $\text{min}^{-1}\text{g}^{-1}$ were obtained. On the other hand, activated sludge sample collected from
60 Shibaura wastewater treatment plant, Tokyo, Japan was utilized for obtaining the mixed culture.
61 The sludge was centrifuged under 2150 x g and reconstructed with Milli-Q water. The
62 bioaugmented culture was obtained by adding equal amounts (0.05 g, dry wt.) of pure fungus
63 granules and reconstituted activated sludge. Neither the mixed bacterial culture nor the fungus was
64 acclimatized before incubation with pesticide.

65 A nutrient-sufficient growth medium previously optimized by Kapdan *et al.* [10] for *C.*
66 *versicolor* was modified by adding a lesser amount of glucose (1 g L^{-1}) and using ammonium
67 nitrate (0.13 g L^{-1}) as the nitrogen source instead of urea. The initial pH of the media was 4.5.

68 2.3 Batch test description

69 The test solution was prepared by adding specified amounts from each of the pesticide stock
70 solutions to autoclaved ($121 \text{ }^\circ\text{C}$, 0.2 MPa, 15 min) empty beakers (8.5 cm diameter, 500 mL
71 capacity), allowing the methanol to evaporate, and then adding 100 mL of the autoclaved nutrient
72 medium. The final concentration of each of the pesticides in the mixture was 10 mg L^{-1} . The
73 bioaugmented culture (0.1 g (dry weight); reconstructed activated sludge: fungus = 1:1) was
74 aseptically transferred to the test solution. The beakers were loosely covered with aluminium foil
75 and then placed under $28 \text{ }^\circ\text{C}$ on a shaker at 80 rpm. Beakers containing only fungus (0.1 g) and
76 only activated sludge (0.1 g) were also incubated in order to assess the atrazine removal
77 performance by single cultures. Same amount of autoclaved fungus and bacteria were incubated

78 separately to assess the extent of atrazine removal owing to biosorption only. Atrazine was added
79 into the nutrient media at a concentration of $1200 \pm 200 \mu\text{g L}^{-1}$ during the experiments with single
80 cultures. The batch tests were carried out in duplicate and each batch test lasted 14 days.

81 *2.4 Analytical methods*

82 The pesticide concentrations were measured using HPLC coupled to a diode array detector
83 (Hewlett Packard, 1100 series). The compounds were separated on a Novapak C18, 60 Å, 4 μm , 3.9
84 x 150 mm column with an acetonitrile / water eluent at the wavelength of 220 nm (atrazine) and
85 205 nm, respectively. Pesticide concentrations above 0.05 mg L^{-1} could be quantified. Total organic
86 carbon (TOC) and total nitrogen (TN) concentrations were measured with a TOC-TN analyzer
87 (TOC-V, Shimadzu, Japan). The activity of fungal enzyme (Laccase) was measured by monitoring
88 the change in absorbance at 468 nm due to the oxidation of the substrate (2,6-dimethoxyphenol) by
89 enzyme at room temperature over a 2 min period as described in a previous study [9].

90 **3. Results and discussion**

91 *3.1 Pesticide removal*

92 Figure 1a depicts the reduction of aqueous phase concentration of each pesticide from the
93 mixture containing the unacclimated fungus-bacteria and fungus-only cultures. Within 7 days, the
94 concentrations of aldicarb, atrazine, and alachlor dropped by 82, 77, and 67%, respectively, in the
95 mixed cultures, and by 79, 81, and 59% in the fungus-only cultures. In comparison, in a previous
96 study, we reported the removal efficiencies of these pesticides by an unacclimated bacterial culture
97 under similar experimental conditions to be only 17, ~0, and 35% for aldicarb, atrazine, and
98 alachlor, respectively (Modin et al., 2008).

99 **[Figure 1]**

100 The concentration of all three pesticides in the fungus-only batches increased after day 7 of the
101 incubation period. Results reported in Figure 1a suggests that the pesticides adsorbed temporarily
102 onto the fungus biomass and later got released back to the aqueous phase. It is noteworthy that the
103 selected pesticides are moderately hydrophobic. Hydrophobic interaction along with other
Hai et al. (2012) Journal of the Taiwan Institute of Chemical Engineers, (43), 459-462

104 mechanism such as ion exchange, surface complexation and hydrogen bonding can play a
105 significant role in sorption of these compounds to solid substrate. Therefore, in good agreement
106 with several previous studies [11], biosorption of the selected pesticides to the fungus biomass is
107 not entirely unexpected. However, the observed increase in liquid phase concentration at the later
108 stage may be attributed to desorption of the non-biodegraded portion of the initially biosorbed
109 pesticides.

110 A different trend was observed in the fungus-bacteria mixtures. The aldicarb concentration
111 increased slightly suggesting release by the biomass after day 7. Atrazine, on the other hand, was
112 almost completely removed from the aqueous phase. The alachlor concentration remained steady
113 between day 7 and day 14. The high removal of atrazine by the fungus-bacteria mixture is
114 especially interesting since among the three tested pesticides, atrazine is generally considered the
115 most persistent with a half-life typically over 100 days [12]. In contrast, in this study the removal of
116 atrazine by the bioaugmented culture reached over 98% in two weeks. In additional experiments,
117 where the extent of biosorption and biodegradation was assessed separately, it was confirmed that
118 biodegradation of atrazine did not occur in fungus-only and activated sludge-only cultures (Figure
119 1b).

120 The success of the fungus-bacterial mixture for removal of atrazine and alachlor can be
121 explained in analogy to a relevant observation made in the soil environment by Levanon [13]. It
122 was reported that the mineralization of alkyl-side chains of alachlor and alkyl-amino-side chains of
123 atrazine was mainly due to fungal activity. However, neither heterocyclic ring-labelled atrazine nor
124 aromatic ring-labelled alachlor were degraded when fungi or bacteria were separately inhibited.
125 The importance of combining fungi and bacteria can be further recognized by comparing the
126 removal rates observed in this study with that by individual bacterial and fungal species or enriched
127 mixed bacterial cultures reported in the literature. For instance, the fungus *C. versicolor* was
128 reported to take 42 days to reach the maximum 86% degradation of atrazine in liquid culture [14].
129 Knapp et al. [15] reported a half life ranging 16-122 days of alachlor in surface water environment.

130 While the culture conditions can be significantly different in individual studies, thereby restricting
131 direct comparison of removal efficiencies, it can generally be stated that the mixed fungus-bacteria
132 culture implemented in this study exhibited high pesticide removal capacity, despite not being
133 acclimated to the pesticides. In practice, the possibility to use non-acclimated microbial cultures for
134 pesticide degradation may be beneficial since the use of pesticides, and thus the pesticide load in
135 wastewater, vary significantly during a year.

136 *3.2 Fungi morphology and extent of biosorption*

137 For the fungus-bacteria mixture, the initial distinct fungus granules appeared disintegrated
138 at the end of the incubation period. Disintegration, but to a lesser extent, was also observed when
139 the fungus was incubated alone (Figure 2). These results show that the presence of bacteria had a
140 clear effect on the morphology of the fungus.

141 **[Figure 2]**

142 The mechanisms of pesticide removal from the liquid phase in this study are not entirely
143 clear. Physical adsorption or uptake by the biomass without biochemical degradation of the
144 pesticides appears to be a major removal mechanism by the fungus-only cultures. The pesticides
145 were initially taken up by the fungus but later released, perhaps as a result of the change in
146 morphology (Figure 2) and also the depletion of nutrients in the growth medium as demonstrated
147 by the TOC and TN concentration over the incubation period (Figure 3).

148 **[Figure 3]**

149 For the fungus-bacteria mixed cultures, biochemical degradation and physical
150 adsorption/uptake both appears to have played roles for removing pesticides from the liquid phase.
151 In the case of aldicarb, there is a slight increase in concentration between day 7 and day 12.
152 However, for atrazine there is a clear decrease to nearly non-detectable levels, and for alachlor the
153 concentration remains stable between day 7 and day 14 (Figure 1a). This suggests that it was not
154 biosorption alone that was responsible for the pesticide removal, but biochemical degradation also
155 played a role, at least for atrazine. No biodegradation or biosorption of atrazine was noticed in

156 bacteria-only culture. On the other hand, temporary adsorption of atrazine onto fungus was
157 observed, but again no biodegradation was detected in fungus-only culture (Figure 1b). Therefore
158 synergistic biodegradation of atrazine by the fungus-bacteria mixed culture following its initial
159 biosorption appears to be very likely. Thus despite comparatively more pronounced disintegration
160 of the biomass as well as the reduction of residual nutrient levels in test solution (Figures 2 and 3)
161 the bioaugmented culture demonstrated better removal.

162 As described in section 3.1, the pesticide removal rate by the fungi-bacteria culture was
163 high compared to the previously published values by our group [12] for bacterial cultures under
164 similar conditions. The results from the current study suggest that the presence of bacteria has a
165 clear effect on the fungus morphology (Figure 2) and nutrient depletion from the test solution
166 (Figure 3). Further research is necessary to establish how bacteria affect fungus morphology and
167 fungal activity (enzyme secretion). For further insight into the reported results, the kinetics of
168 pesticide removal and its toxic effect on bioaugmented culture need to be studied under a range of
169 pesticide dosage and varying ratios of fungus and activated sludge masses. Nevertheless, the results
170 presented in the current study do suggest that combining fungus and bacteria for removing
171 recalcitrant compounds such as pesticides from wastewater is beneficial.

172 **4. Conclusions**

173 This study showed that mixed cultures of bacteria and fungus can remove the pesticides
174 aldicarb, atrazine, and alachlor from their mixture. Particularly atrazine was very effectively
175 removed. The reported data demonstrated the complementary effect of combining bacteria and a
176 fungal culture for improving the removal rates of the three selected recalcitrant pesticides. The
177 results have important implications towards the development of a scaled-up treatment system for
178 pesticide contaminated water.

179 **5. Acknowledgements**

180 A part of this study was carried out during the Japan Society for promotion of Science
181 fellowship tenure of Dr. Faisal I. Hai at the University of Tokyo, Japan. Oskar Modin was

182 supported by the Monbukagakusho scholarship from MEXT, Japan. Shufan Yang, Shaza Raini and
183 Luong N. Nguyen of University of Wollongong, Australia are thanked for their assistance in some
184 of the analyses.

185 **6. References**

- 186 [1] Carter A. How pesticides get into water - and proposed reduction measures. *Pestic Outlook* 2000;
187 11(4):149-56.
- 188 [2] Gilliom RJ. Pesticides in the hydrologic system—what do we know and what's next? *Hydrol Process*
189 2001; 15(16):3197-201.
- 190 [3] Navaratna D, Shu L, Jegatheesan V. Existence, impacts, transport and treatments of herbicides in great
191 barrier reef catchments in Australia. In: Virkutyte JV, Rajender S.; Jegatheesan, Veeriah, editor. *Treatment*
192 *of micropollutants in water and wastewater*. London, UK: IWA publishing; 2010. p. 425-57.
- 193 [4] Ronhede S, Jensen B, Rosendahl S, Kragelund BB, Juhler RK, Aamand J. Hydroxylation of the Herbicide
194 Isoproturon by Fungi Isolated from Agricultural Soil. *Appl Environ Microbiol* 2005; 71(12):7927-32.
- 195 [5] Mori T, Fujie K, Katayama A. Bacterial and fungal contributions to chlorothalonil degradation in soil. *Soil*
196 *Sci Plant Nutr* 1998; 44(3):297-304.
- 197 [6] Osman KA, Ibrahim GH, Askar AI, Aba Alkhail ARA. Biodegradation kinetics of dicofol by selected
198 microorganisms. *Pestic Biochem Physiol* 2008; 91(3):180-5.
- 199 [7] Kontchou CY, Gschwind N. Biodegradation of s-triazine compounds by a stable mixed bacterial
200 community. *Ecotoxicol Environ Saf* 1999; 43(1):47-56.
- 201 [8] Liu C, Huang X, Wang H. Start-up of a membrane bioreactor bioaugmented with genetically engineered
202 microorganism for enhanced treatment of atrazine containing wastewater. *Desalination*. 2008; 231(1-
203 3):12-9.
- 204 [9] Hai FI, Yamamoto K, Nakajima F, Fukushi K. Factors governing performance of continuous fungal reactor
205 during non-sterile operation - The case of a membrane bioreactor treating textile wastewater.
206 *Chemosphere*. 2009; 74(6):810-7.
- 207 [10] Kapdan IK, Kargi F. Simultaneous biodegradation and adsorption of textile dyestuff in an activated
208 sludge unit. *Process Biochem* 2002; 37:973-81.
- 209 [11] Bouju H, Buttiglieri G, Malpei F. Perspectives of persistent organic pollutants (POPs) removal in an
210 MBR pilot plant. *Desalination*. 2008; 224(1-3):1-6.
- 211 [12] Modin O, Fukushi K, Yamamoto K. Simultaneous removal of nitrate and pesticides from groundwater
212 using a methane-fed membrane biofilm reactor. *Water Sci Technol* 2008; 58(6):1273-9.
- 213 [13] Levanon D. Roles of fungi and bacteria in the mineralization of the pesticides atrazine, alachlor,
214 malathion and carbofuran in soil. *Soil Bio Biochem* 1993; 25(8):1097-105.
- 215 [14] Bending GD, Friloux M, Walker A. Degradation of contrasting pesticides by white rot fungi and its
216 relationship with ligninolytic potential. *FEMS Microbiol Lett* 2002; 212(1):59-63.
- 217 [15] Knapp CW, Graham DW, Berardesco G, deNoyelles F, Cutak BJ, Larive CK. Nutrient level, microbial
218 activity, and alachlor transformation in aerobic aquatic systems. *Water Res* 2003; 37(19):4761-9.

219
220
221
222
223
224
225
226
227
228
229
230

231

232


FIGURES

233


(a)

Fungus □ Fungus-bacteria ●


234



238




242



247

(b)



253

254

255

256

257

258

Figure 1. Change in liquid phase concentration of the pesticide(s) over the incubation period. Error bars indicate the deviations between two replicates. (a) The batch test solution comprised of three pesticides (each at 10 mg L^{-1}) and a nutrient medium. (b) The test solution comprised of atrazine ($1200 \pm 200 \text{ } \mu\text{g L}^{-1}$) and a nutrient medium. Controls inoculated with heat-killed (autoclaved) biomass were added to assess biosorption separately.

259
260
261
262
263
264




Figure 2. Difference between change of morphology of the fungus-only and the fungi-bacteria culture over the incubation period. (a) Firm pure fungus granules at the start of the experiment (b) Fungus granules appearing black due to being covered with activated sludge (day 3 of incubation of mixed culture batches). (c) Disintegrated fungus granules (smaller in size and deformed) still covered with activated sludge at the end of the mixed culture incubation. Presence of significant amount of fibrous structures along with granules noticeable. (d) Fungus granules (relatively more intact as compared to 'c') along with some fungus in fibrous morphology at the end of the fungus-only incubation.




Figure 3. Concentration of TOC and TN in the batch solutions at the start and the end of the experiment showing the effect of presence of bacteria on consumption of TOC and TN. The batch test solution comprised of three pesticides and a nutrient medium.