



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science - Papers (Archive)

Faculty of Science, Medicine and Health

2012

Isolation of tuberospironine A, a novel croomine derivative from *Stemona tuberosa* Lour.

Pratiwi Pudjiastuti

Airlangga University

Stephen G. Pyne

University of Wollongong, spyne@uow.edu.au

S Sugiyanto

Bureau of Cultivation and Plantation Protections, Ambon, Molucas

Wilford Lie

University of Wollongong, wilford@uow.edu.au

Publication Details

Pudjiastuti, P., Pyne, S. G., Sugiyanto, S. & Lie, W. (2012). Isolation of tuberospironine A, a novel croomine derivative from *Stemona tuberosa* Lour.. *Phytochemistry Letters*, 5 (2), 358-360.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Isolation of tuberospironine A, a novel croomine derivative from *Stemona tuberosa* Lour.

Abstract

A novel croomine derivative, tuberospironine A (3-*epi*-tuberospironine) was isolated from the root extracts of *Stemona tuberosa* Lour. found growing on Seram Island, Moluccas Province, Indonesia. The structure of this novel alkaloid, with unprecedented configuration at C-3 for a croomine derivative, was determined from interpretation of its NMR spectroscopic data.

Keywords

tuberosa, lour, croomine, novel, stemona, tuberospironine, derivative, isolation, CMMB

Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

Pudjiastuti, P., Pyne, S. G., Sugiyanto, S. & Lie, W. (2012). Isolation of tuberospironine A, a novel croomine derivative from *Stemona tuberosa* Lour.. *Phytochemistry Letters*, 5 (2), 358-360.

Isolation of tuberospironine A, a novel croomine derivative from *Stemona tuberosa* Lour.

Pratiwi Pudjiastuti^{a,*}, Stephen G. Pyne^{b,*}, Sugiyanto^c, Wilford Lie^b

^aDepartment of Chemistry, Airlangga University, Surabaya, Indonesia

^bSchool of Chemistry, University of Wollongong, Wollongong, NSW, 2522, Australia

^cInstitute of Cultivation and Plantation Protections, Ambon, Moluccas, Indonesia

*Corresponding author: Tel: +61-02-4221 3511; fax: +61-02-4221 4287

E-mail address: spyne@uow.edu.au

Keywords:

Stemona alkaloid

Croomine

Stemona tuberosa

ABSTRACT

A novel croomine derivative, tuberospironine A (3-*epi*-tuberospironine) was isolated from the root extracts of *Stemona tuberosa* Lour. found growing on Seram Island, Moluccas Province, Indonesia. The structure of this novel alkaloid, with unprecedented configuration at C-3 for a croomine derivative, was determined from interpretation of its NMR spectroscopic data.

1. Introduction

The *Stemona* family of alkaloids includes more than one hundred different natural products (Pilli et al., 2005; Greger, 2006) which have been structurally classified by Pilli into eight different groups. The pyrrolo[1,2-*a*]azepine nucleus is common to six of these groups,

while a pyrido[1,2-*a*]azepine ring system is found in the more recently discovered Stemocurtisine group of *Stemona* alkaloids (Pyne et al., 2007). A miscellaneous group comprising five *Stemona* alkaloids has also been identified (Pilli et al. 2005). Greger has recently classified the *Stemona* alkaloids into three skeletal types based on their proposed biosynthetic origins (Greger, 2006). The root extract of *Stemona tuberosa* Lour. has been extensively studied and has been shown to contain croomine and/or stichoneurine-type alkaloids (Greger, 2006) including, croomine **1** (Lin et al. 2008; Schinnerl et al., 2005, 2007), 6-hydroxycroomine **2** (Jiang et al., 2006a, Lin et al., 2008; Schinnerl et al., 2005, 2007), 10-hydroxycroomine **3** (Lin et al., 2008) and tuberospironine **4** (Jiang et al., 2006b) and tuberostemonine **5** (Götz et al. 1961; Hu, et al., 2009; Jiang et al., 2006; Lin et al., 2008; Schinnerl et al., 2005, 2007; Xu; Y. T. et al., 2010; Xu; R-S. et al. 1982) and tuberostemonine A **6** (Schinnerl et al., 2005, 2007), respectively (Figure 1). We report here the isolation of a novel croomine-like alkaloid, tuberospironine A **7** (3-*epi*-tuberospironine), from the root extracts of *Stemona tuberosa* Lour., which has the opposite configuration at C-3 and C-9 to that of croomine **1** itself (Figure 1).




Figure 1. The structures of representative croomine and tuberostemonine alkaloids.

2. Results and Discussion

The roots of *Stemona tuberosa* Lour. were collected from Seram Island, Moluccas Province, Indonesia. The ethanol extract of the dried and powdered roots was concentrated *in vacuo* and the resulting residue was acidified to pH 1-2 and then partitioned between water and CH₂Cl₂. The aqueous layer was made basic with ammonia and then extracted with CH₂Cl₂. The crude alkaloid mixture was subjected to successive separations by column chromatography and finally preparative TLC to give a pure sample of tuberospironine A **7**.

Compound **7** was obtained as a pale brown, amorphous solid. Its molecular formula was determined to be C₁₈H₂₇NO₅ from its HRESIMS, (*m/z* 338.1953 [M+H]⁺), the same molecular formula as 10-hydroxycroomine **3** and tuberospironine **4** (Figure 1). The IR spectrum of **7** showed bands for the presence of a γ -lactone (1730 cm⁻¹) and a hydroxyl group (3420 cm⁻¹). The ¹H NMR spectrum of **7** (Table 1) showed a downfield resonance at δ 4.37 (m, 1H, H-14) and two methyl doublet signals (δ 1.27 (d, 3H, *J* = 7.0 Hz, H-18 and δ 1.29 (d, 3H, *J* = 7.5 Hz, H-13) typical resonances for the spirocyclic (at C-9) and the appended (at C-3) γ -lactone moieties of croomine-derived alkaloids. The ¹H NMR spectrum of **7** (Table 1) showed a doublet resonance at δ 3.95 (d, *J* = 10.5 Hz, H-10) diagnostic of a C-10 hydroxyl group. The chemical shift and coupling constant of H-10 were different to that of 10-hydroxycroomine **3** (δ 4.32 (d, *J* = 7.8 Hz, H-10) (Lin et al.) and tuberospironine **4** (δ 4.09 (d, *J* = 10.3 Hz, H-10) (Jiang et al.) but were closer in magnitude to those of tuberospironine **4** which has the opposite configuration at C-9 to the alkaloids **1-3** and at C-10 to 10-hydroxycroomine **3** (Figure 1). The ¹H, COSY, ¹³C NMR and HSQC and HMBC spectra of **7** (Table 1) indicated that this compound was a diastereomer of **3** and **4**.

The relative configuration of **7** was determined from NOESY NMR experiments (Figure 2), with the absolute configuration at C-9a assumed to be the same as that of croomine **1**. The NOESY correlations between H-10 and H-9a and H-1 α indicated that the C-9 configuration

of **7** was opposite to that of the alkaloids **1-3** and was the same as that of tuberospirone **4** (Figure 1). The NOESY correlation between H-10 and the C-11 methyl group (H-13) indicated their relative *syn*-relationship, as observed in tuberospirone **4**, which was further supported by the aforementioned similarity in magnitude of $J_{10,11}$ in **7** (10.5 Hz) and **4** (10.3 Hz). The NOESY correlations between H-3 and H-15 α and H-14 and the two H-5 protons, and H-15 β (Figure 2, see Figure 1 for the definition of 15 α and 15 β) indicated the β -configuration of H-3, opposite that found in all previous naturally occurring croonine-like alkaloids (Pilli et al., 2005; Greger, 2006). This C-3 configuration is rare and has been only found in two tuberostemonines, tuberostemonine A **6** (Schinnerl et al., 2005, 2007) and tuberostemonine M (Sastraraji, 2006).

In conclusion, we have isolated a novel 10-hydroxycroonine derivative from the root extracts of *Stemona tuberosa* Lour. which we have named tuberospirone A **7** based on its same C-9/C-9a relative configuration to tuberospirone **4**. The structure of this novel alkaloid, with unprecedented configuration at C-3 for a croonine derivative, was determined from interpretation of its NMR spectroscopic data.

Table 1 ^{13}C NMR (125 Hz) and ^1H NMR (500 MHz) spectroscopic data for compound **7** in CDCl_3 .

Position	δ_{C} (DEPT)	δ_{H} (mult., J (Hz))	HMBC
1 α	26.5 (CH ₂)	1.97 (m)	H-2, H-3, H-9a
1 β		1.73 (m)	
2 α	26.6 (CH ₂)	1.89 (m)	H-1, H-9a, H-14
2 β		1.54 (m)	
3	66.2 (CH)	3.42 (dd, 7.5, 14.5)	H-5 (weak), H-9a (weak), H-14
5 α	47.8 (CH ₂)	3.09 (m)	H-3, H-6, H-9a
5 β		3.30 (br d, 13.5)	
6a	26.9(CH ₂)	1.72 (m)	H-5
6b		1.64 (m)	
7a	22.1 (CH ₂)	1.77 (m)	H-8a
7b		1.69 (m)	
8a	30.3 (CH ₂)	2.06 (m)	
8b		1.80 (m)-	
9	87.8 (C)	-	H-8, H-9a, H-10
9a	67.8 (CH)	3.58 (dd, 7.5, 8.0)	H-1, H-2, H-5, H-6, H-9, H-10
10	78.0 (CH)	3.95 (d, 10.5)	H-8, H-9, H-9a, H-11
11	40.2 (CH)	2.65 (m)	
12	175.5 (C)	-	H-13
13	12.7 (CH ₃)	1.31 (d, 7.5)	H-12
14	79.4 (CH)	4.37 (m)	H-3
15 α	35.0 (CH ₂)	1.52 (m)	H-17
15 β		2.39 (m)	
16	35.5 (CH)	2.61 (m)	
17	179.3 (C)	-	15 β
18	14.9 (CH ₃)	1.27 (d, 7.0)	,
OH	-	1.93 (br s)	




Figure 2. Selected NOESY correlations (shown as double headed arrows) for compound **7** based on a SPARTAN (AM3) generated structure.

3. Experimental

3.1 General Experimental Procedure

The IR spectra were recorded on a MIRacle 10 Shimadzu Spectrometer and optical rotations on a Jasco P-2000 polarimeter. The ESIMS and HRESIMS were recorded on Micromass Platform LCZ and factory modified Waters QToF *Ultima* Mass Spectrometer (Wynteshawe, UK). NMR spectra were recorded on Variant-500 MHz NMR spectrometer. Silica gel was used for column chromatography and TLC was carried out on silica gel 60 GF254 plates Merck HX1 15287. The TLC spots were viewed at 254 nm and visualized by Dragendorff's reagent.

3.2 Plant material

The roots of *S. tuberosa* Lour were collected from Seram Island, Moluccas Province, Indonesia in July 2011. The material was identified at the Conservation Institute of Purwodadi Botanical Garden, Pasuruan, East of Java, Indonesia, where a voucher specimen (No. IV.D.IV.14) was deposited.

3.3 Extraction and isolation

The roots of *S. tuberosa* (3.0 kg) were dried and ground into a powder (840 g) and extracted with 95% ethanol three times for 12 days. The filtrate was evaporated under vacuum at 50° C temperature, to give 148.0 g of crude extract. The crude extract was acidified with 5% of HCl until pH 1-2 and partitioned between H₂O and CH₂Cl₂. The aqueous fraction was basified with NH₄OH until pH 10-11 and extracted using CH₂Cl₂ to give 4.13 g of crude alkaloid extract. The crude alkaloid extract (4.0 g) was subjected to column chromatography over silica gel with gradient elution from CH₂Cl₂/MeOH (100:0) to CH₂Cl₂/MeOH (70:30) to give 27 fractions. Fractions 6-8 (1.864 g) were combined and separated by column chromatography and then preparative TLC to afford tuberospironine A **7** (13.8 mg).

3.4 Tuberospironine A **7**

Pale brown, amorphous, $[\alpha]_D^{24}$ -6.32 (*c* 0.005, CHCl₃); IR ν_{\max} (cm⁻¹); 3420, 2924, 2861, 1730, 1453, and 1156; ¹H and ¹³C NMR data are shown in Table 1; HRESIMS *m/z* 338.1953 [M+H]⁺, calcd for C₁₈H₂₈NO₅ 338.1962.

Acknowledgements

The authors are grateful to the Directorate General of Higher Education, Indonesia for supporting this project. We would like to thank University of Wollongong as the counterpart university. Thanks also to Karin Maxwell, Phurpha Wangkuk, Ari Satya Nugraha and Dr. Sudhir Sangule.

References

- Goetz, M.; Boegri, T.; Gray, A. H. 1961. Structural studies of the major alkaloid of *Stemona tuberosa*, tuberostemonine (I). *Tetrahedron Lett.* , 707-715.
- Greger, H. 2006. Structural Relationships, Distribution and Biological Activities of *Stemona* Alkaloids. *Planta Med.* 72, 99-113.
- Hu, J.-P.; Yang, D.-H.; Lin, W.-H.; Cai, S.-Q. 2009. Alkaloids from the Roots of *Stemona tuberosa*. *Helv. Chim. Acta* 92, 2125-2133.
- Jiang, R.-W.; Hon, P.-M.; Xu, Y.-T.; Chan, Y.-M.; Xu, H.-X.; Shaw, P.-C.; But, P. P.-H. 2006a. Isolation and chemotaxonomic significance of tuberostemospironine-type alkaloids from *Stemona tuberosa*. *Phytochemistry*, 67, 52-57
- Jiang, R.-W.; Hon, P.-M.; Zhou, Y.; Chan, Y.-M.; Xu, Y.-T.; Xu, H.-X.; Greger, H.; Shaw, P.-C.; But, P. P.-H. 2006b. Alkaloids and Chemical Diversity of *Stemona tuberosa*. *J. Nat. Prod.* 69, 749-754.
- Lin, L.-G.; Leung, H. P.-H.; Zhu, J.-Y.; Tang, C.-P.; Ke, C.-Q.; Rudd, J. A.; Lin, G.; Ye, Y. 2008. Croomine- and tuberostemonine-type alkaloids from roots of *Stemona tuberosa* and their antitussive activity. *Tetrahedron* 64, 10155-10161.
- Pilli, R. A.; Rosso, G. B. de Oliveira, M. C. F. 2005. *The Alkaloids*, vol. 62; Cordell, G. A. Ed; Elsevier; San Diego, **2005**; Chapter 2, pp 77-173.
- Pyne S. G.; Ung A. T.; Jatisatienr A.; Mungkornasawakul P. 2007. The Pyrido[1,2-*a*]azepine *Stemona* alkaloids. *Maejo Intern. J. of Sci. Technol.* 1, 157-165.
- Sastraruji, T.; Jatisatienr, A.; Issakul, K.; Pyne, S. G.; Ung, A. T.; Lie, W.; Williams, M. C. 2006. Phytochemical studies on *Stemona* plants: isolation of new tuberostemonine and stemofoline alkaloids. *Nat. Prod. Commun.* 1, 813-818.
- Schinnerl, J.; Kaltenecker, E.; Pacher, T.; Vajrodaya, S.; Hofer, O.; Greger, H. 2005. New Pyrrolo[1,2-*a*]azepine Type Alkaloids from *Stemona* and *Stichoneuron* (Stemonaceae) *Monatsh. Chemie* 136, 1671-1680.

- Schinnerl, J.; Brem, B.; But, P. P.-H.; Vajrodaya, S.; Hofer, O.; Greger, H. 2007. Pyrrolo- and pyridoazepine alkaloids as chemical markers in *Stemona* species. *Phytochemistry* 68, 1417-1427.
- Xu, R. S.; Lu, Y. J.; Chu, J. H.; Iwashita, T.; Naoki, H.; Naya, Y.; Nakanishi, K. 1982. Studies on some new *Stemona* alkaloids. A diagnostically useful proton NMR line-broadening effect. *Tetrahedron*. 38, 2667-2670.
- Xu Y.-T.; Shaw P.-C.; Jiang R.-W.; Hon P.-M.; Chan Y.-M.; But P. P.-H. 2010. Antitussive and central respiratory depressant effects of *Stemona tuberosa*. *J. Ethnopharm.* 128, 679-684.