

1-1-2011

What do older patients want? Understanding older patients' attitudes towards general practice trainees

Andrew D. Bonney
University of Wollongong, abonney@uow.edu.au

Sandra C. Jones
University of Wollongong, sandraj@uow.edu.au

Donald Iverson
University of Wollongong, iverson@uow.edu.au

Christopher Magee
University of Wollongong, cmagee@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/medpapers>



Part of the [Medicine and Health Sciences Commons](#)

Citation

Bonney, Andrew D.; Jones, Sandra C.; Iverson, Donald; and Magee, Christopher, 2011, What do older patients want? Understanding older patients' attitudes towards general practice trainees.
<https://ro.uow.edu.au/medpapers/64>

What do older patients want? Understanding older patients' attitudes towards general practice trainees

Abstract

Context: Older patients constitute an increasing proportion of the caseloads in general practice (GP) training practices, but are relatively reluctant to consult trainees. Understanding their attitudes is a first step in improving older patient-trainee interaction. Objectives: Characterise the attitudes of older patients to GP trainees. Design: Cross-sectional survey; exploratory factor analysis; logistic regression. Setting: Randomised, stratified sample of 38 training practices across five Australian states. Participants: Patients aged 60 years and over (N=911; response rate 47.9%). Instrument: Questionnaire for self-completion: previously piloted and subjected to factor analysis. Main and secondary outcome measures: Identification of attitude factors; patient characteristics predicting high factor scale scores; patient responses predicted by factor scale scores. Results: Cronbach's $\alpha = .791$. Three factors identified: interpersonal trust (IPT); system trust (ST); and interpersonal continuity of care (IPC). Female gender and age ≥ 75 predicted high IPT scores [OR 1.95, 95% CI 1.39-2.72, $p < .001$; OR 1.41, 95% CI 1.01-1.96, $p = .044$]. Female gender and chronic illness predicted high IPC scores [OR 2.88, 95% CI 1.66-5.00, $p < .001$; OR 2.17, 95% CI 1.93-3.94, $p = .011$]. Self-rated health score $\geq 4/5$ predicted high ST scores [OR 1.91, 95% CI 1.38-2.64, $p < .001$]. High IPT reduced odds of satisfaction with trainees [OR .51, 95% CI .32-.81, $p = .004$], and both high IPT and IPC reduced odds of comfort with trainee chronic/complex care [OR .61, 95% CI .40-.92, $p = .02$; OR .31, 95% CI .18-.56, $p < .001$]. High IPC predicted reduced odds of seeing a trainee more than once or twice [OR .42, 95% CI .24-.74, $p = .002$]. High ST predicted increased satisfaction with trainees [OR 6.96, 95% CI 4.47-10.83, $p < .001$]. Conclusions: Three underlying constructs (IPT, ST and IPC) appear important in influencing older patients' interactions with trainees. Patients with high IPT and/or IPC scores demonstrate significantly reduced acceptance of trainees. Strategies aimed to improve trust and continuity, e.g. shared continuity of care with their regular GP, may improve acceptance for these patients.

Keywords

older, do, patients, practice, want, trainees, understanding, attitudes, towards, general

Disciplines

Medicine and Health Sciences

Publication Details

Bonney, AD, Jones, SC, Iverson, D & Magee, C, What do older patients want? Understanding older patients' attitudes towards general practice trainees. In The 2010 NAPCRG Annual Meeting, Nov 2010, Seattle, Washington. Family Medicine, 43, Suppl1, 2011.



Society of Teachers of Family Medicine

- About
- Conferences & Education
- Groups & Initiatives
- Advocacy
- Member Resources
- Publications
- Foundation

Family Medicine

Vol 43 (Suppl 1)
January 2011

Presentations Given At The 2010 NAPCRG Annual Meeting

Search For A Specific Presentation

Enter a search criteria (*the presentations displayed below will automatically be filtered to match your criteria as you type*):

Author's Name:

Title:

Or Browse All Presentations

Page 1 of 1 - Presentations 1 - 2 of 2

Different Needs, Same Solution: The Effect of Visible Supervision On Older Patients' Attitudes to General Practice Trainee Chronic/Complex Care

Andrew D Bonney, MBBS, MFPM; Sandra C Jones, PhD; Don Iverson, PhD; Christopher Magee, PhD

[Hide Abstract](#)

Context: The ageing population requires training practices to provide appropriate training for future general practitioners (GPs) in chronic disease management, whilst recognising the particular needs of older patients. There has been little research to assist practices meet these requirements. Objectives: To assess older patients' attitudes to GP trainee management of a chronic condition with varying levels of visible supervision. Design: Cross-sectional survey; exploratory factor analysis; logistic regression. Setting: Randomised, stratified sample of 38 training practices across five Australian states. Participants: Patients aged 60 years and over (N=911; response rate 47.9%). Instrument: Questionnaire for self-completion: previously piloted and subjected to factor analysis; clinical vignette assessing responses to trainee chronic/complex care with varying supervision. Main and secondary outcome measures: Identification of attitude factors; patient responses to vignette according to factor scale scores. Results: Cronbach's $\alpha = .791$ for main attitude scale, $\alpha = .780$ for vignette. Three factors identified: interpersonal trust (IPT); system trust (ST); and interpersonal continuity of care (IPC). High ST factor scale score predicted not feeling the need for a regular GP [OR 2.50, 95% CI 1.43-4.37, $p < .001$]. There was noticeable variation between the IPT, ST and IPC high score groups in their attitudes to unsupervised GP trainee chronic/complex care [Not at all comfortable: IPT 42.9% vs. ST 15.1% vs. IPC 23.7%]. The responses significantly improved for all groups if their regular GP checked on management [Not at all comfortable: IPT 7.2%, $p < .001$; ST 2.2%, $p < .001$; IPC 3.9%, $p < .001$]. Conclusions: Older patients' attitudes to trainees are heterogeneous and have varying influences upon their acceptance of trainee chronic/complex care. Beyond the educational and safety issues, all three groups characterised by their factor scores would benefit from supervision for chronic disease management: the IPT and IPC groups to address their trust and continuity concerns; and the ST group as it appears they are significantly less likely to arrange for continuity in their care of their own accord.

What Do Older Patients Want? Understanding Older Patients' Attitudes Towards General Practice Trainees

Andrew D Bonney, MBBS, MFPM; Sandra C Jones, PhD; Don Iverson, PhD; Christopher Magee, PhD

[Hide Abstract](#)

Context: Older patients constitute an increasing proportion of the caseloads in general practice (GP) training practices, but are relatively reluctant to consult trainees. Understanding their attitudes is a first step in improving older patient-trainee interaction. Objectives: Characterise the attitudes of older patients to GP trainees. Design: Cross-sectional survey; exploratory factor analysis; logistic regression. Setting: Randomised, stratified sample of 38 training practices across five Australian states. Participants: Patients

aged 60 years and over (N=911; response rate 47.9%). Instrument: Questionnaire for self-completion: previously piloted and subjected to factor analysis. Main and secondary outcome measures: Identification of attitude factors; patient characteristics predicting high factor scale scores; patient responses predicted by factor scale scores. Results: Cronbach's α =.791. Three factors identified: interpersonal trust (IPT); system trust (ST); and interpersonal continuity of care (IPC). Female gender and age \geq 75 predicted high IPT scores [OR 1.95, 95% CI 1.39-2.72, $p < .001$; OR 1.41, 95% CI 1.01-1.96, $p = .044$]. Female gender and chronic illness predicted high IPC scores [OR 2.88, 95% CI 1.66-5.00, $p < .001$; OR 2.17, 95% CI 1.93-3.94, $p = .011$]. Self-rated health score \geq 4/5 predicted high ST scores [OR 1.91, 95% CI 1.38-2.64, $p < .001$]. High IPT reduced odds of satisfaction with trainees [OR .51, 95% CI .32-.81, $p = .004$], and both high IPT and IPC reduced odds of comfort with trainee chronic/complex care [OR .61, 95% CI .40-.92, $p = .02$; OR .31, 95% CI .18-.56, $p < .001$]. High IPC predicted reduced odds of seeing a trainee more than once or twice [OR .42, 95% CI .24-.74, $p = .002$]. High ST predicted increased satisfaction with trainees [OR 6.96, 95% CI 4.47-10.83, $p < .001$]. Conclusions: Three underlying constructs (IPT, ST and IPC) appear important in influencing older patients' interactions with trainees. Patients with high IPT and/or IPC scores demonstrate significantly reduced acceptance of trainees. Strategies aimed to improve trust and continuity, e.g. shared continuity of care with their regular GP, may improve acceptance for these patients.

Page 1 of 1 - Presentations 1 - 2 of 2

[Top](#) | [Home](#) | [Contact Us](#) | [Jobs](#) | [Join](#) | [Renew](#) | [Member Log In](#) | [Search](#)
[About Us](#) | [Conferences](#) | [Groups](#) | [Initiatives](#) | [Advocacy](#) | [Member Resources](#) | [Publications](#)

© 2008 Society of Teachers of Family Medicine | [Privacy Policy](#) | [Refund Policy](#)

Administrative Office:

11400 Tomahawk Creek Parkway, Ste 540, Leawood, KS 66211
 Phone: 800-274-2237, ext. 5410 Fax: 913-906-6096

Government Affairs Office:

2021 Massachusetts Ave, NW, Washington, DC 20036
 Phone: (202) 986-3309 Fax (202) 232-9044