University of Wollongong

Research Online

Faculty of Science, Medicine and Health -
Papers: part A

Faculty of Science, Medicine and Health

1-1-2013

Sediment disturbance associated with trampling by humans alters species assemblages on a rocky intertidal seashore

Todd E. Minchinton

University of Wollongong, tminch@uow.edu.au

Katie J. Fels

University of Adelaide

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>

Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Minchinton, Todd E. and Fels, Katie J., "Sediment disturbance associated with trampling by humans alters species assemblages on a rocky intertidal seashore" (2013). *Faculty of Science, Medicine and Health - Papers: part A*. 77.

<https://ro.uow.edu.au/smhpapers/77>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Sediment disturbance associated with trampling by humans alters species assemblages on a rocky intertidal seashore

Abstract

Our observations reveal that species living on rocky intertidal reefs can be positively and negatively associated with increasing sediment load. We therefore tested the hypothesis that sediment disturbance, including increased sediment loads associated with trampling by humans, alters the abundance of macroinvertebrates on a sheltered rocky intertidal platform in southern Australia. First we trampled the reef in shallow water at several disturbance intensities, simulating different numbers of people walking on the platform on a rising tide, and determined that sediment load was directly related to trampling intensity. Trampling displaced sediment, and up to an order of magnitude more sediment accumulated on the reef near intensively trampled areas compared to those with natural sedimentation. We then manipulated sediment load on the reef to mimic increased sedimentation due to trampling and other potential human (e.g. terrestrial run-off) and natural (e.g. storms) disturbances, and monitored changes to dominant species. Sediment addition increased the cover and depth of sediment on the reef. Increased sediment load negatively impacted barnacles, but not mussels, which occur naturally bound in a sediment matrix in small depressions on the platform. The dominant grazing gastropods *Nerita atramentosa* and *Bembicium nanum* were negatively influenced by increased sediments, although not at levels associated with trampling, whereas other gastropods were not affected or responded positively to increased sediment load. Changing sediment loads, including the cumulative effects of small-scale disturbances such as trampling, can alter macroinvertebrate species assemblages on rocky reefs and favour species that tolerate a range of environmental conditions and habitat types.

Keywords

sediment, disturbance, associated, trampling, humans, alters, species, assemblages, rocky, intertidal, seashore

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Minchinton, T. E. & Fels, K. J. (2013). Sediment disturbance associated with trampling by humans alters species assemblages on a rocky intertidal seashore. *Marine Ecology: Progress Series*, 472 (N/A), 129-140.

SEDIMENT DISTURBANCE AND TRAMPLING BY HUMANS ALTERS SPECIES
ASSEMBLAGES ON A ROCKY INTERTIDAL SEASHORE

(running head: Sediment disturbance on rocky reefs)

Todd E. Minchinton^{1,*} and Katie J. Fels^{2,3}

¹Institute for Conservation Biology and School of Biological Sciences

University of Wollongong, New South Wales 2522, Australia

²School of Earth and Environmental Sciences

The University of Adelaide, South Australia 5005, Australia

*Corresponding author: Tel: +61 (2) 4221 5188; Fax +61 (2) 4221 4135; tminch@uow.edu.au

³Current address: Kellogg Brown & Root, 186 Greenhill Road, Parkside, South Australia
5067, Australia

ABSTRACT: Our observations reveal that species living on rocky intertidal reefs can be positively and negatively associated with increasing sediment load. We therefore tested the hypothesis that sediment disturbance, including increased sediment loads associated with trampling by humans, alters the abundance of macroinvertebrates on a sheltered rocky intertidal platform in southern Australia. First we trampled the reef in shallow water at several disturbance intensities simulating different numbers of people walking on the platform on a rising tide, and determined that sediment load was directly related to trampling intensity. Trampling displaced sediment and up to an order of magnitude more sediment accumulated on the reef near intensively trampled areas compared to those with natural sedimentation. We then manipulated sediment load on the reef to mimic increased sedimentation due to trampling and other potential human (e.g. terrestrial run-off) and natural (e.g. storms) disturbances, and monitored changes to dominant species. Sediment addition increased the cover and depth of sediment on the reef. Increased sediment load negatively impacted barnacles, but not mussels that occur naturally bound in a sediment matrix in small depressions on the platform. The dominant grazing gastropods *Nerita atramentosa* and *Bembicium nanum* were negatively influenced by increased sediments, although not at levels associated with trampling, whereas other gastropods were not affected or responded positively to increased sediment load. Changing sediment loads, including the cumulative effects of small-scale disturbances such as trampling, can alter macroinvertebrate species assemblages on rocky reefs and favour species that tolerate a range of environmental conditions and habitat types.

KEY WORDS: Australia; Grazing gastropods; *Hormosira*; Human impacts; Intertidal macroalgae; Marine invertebrates; Mussel beds; Resilience

INTRODUCTION

Increased sedimentation on coastal reefs due to human activity is an emerging environmental problem (see review by Airolidi 2003). Changing sediment loads resulting from natural disturbances, such as storm waves and flooding, have been observed to alter dramatically species assemblages on rocky intertidal shores (e.g. Daly & Mathieson 1977, Robles 1982, Littler et al. 1983, Branch et al. 1990). Indeed, studies manipulating sediment loads have demonstrated changes to ecological processes and species assemblages on intertidal and subtidal rocky reefs (e.g. Airolidi 1998, Irving & Connell 2002, Connell 2005, Airolidi & Hawkins 2007, Balata et al. 2007, Huff & Jarett 2007, Vaselli et al. 2008, Atalah & Crowe 2010). A significant source of increased sediment deposition on coastal reefs is run-off from terrestrial habitats, particularly in urban and agricultural landscapes (Rogers 1990, Airolidi 2003, Thompson et al. 2002). Less recognised but perhaps equally important is the cumulative impacts of numerous small-scale disturbances from human recreational activities on intertidal and shallow subtidal reefs (e.g. Zedler 1978, Kingsford et al. 1991, Addressi 1994, Fletcher & Frid 1996, Keough & Quinn 1998, Thompson et al. 2002, Hardiman & Burgin 2010, Luna-Pérez et al. 2011).

Trampling by humans during the collection of food and bait and recreational activities is an important small-scale agent of disturbance that has been shown to alter species assemblages in a diversity of marine habitats (rocky intertidal reefs: Beauchamp & Gowing 1982, Ghazanshahi et al. 1983, Bally & Griffiths 1989, Povey & Keough 1991, Brosnan & Crumrine 1994, Quinn & Keough 1998, Araújo et al. 2009; coral reefs: Liddle & Kay 1987, Neil 1990, Liddle 1991; seagrass: Eckrich & Homquist 2000; mud and sand flats: Wynberg & Branch 1997, Rossi et al. 1997; mangroves: Ross 2006; salt marshes: Headley & Sale 1999). Studies of trampling on intertidal reefs have largely focused on direct impacts, particularly the crushing and dislodgement of intertidal organisms when the reef is exposed at low tide (e.g.

Povey & Keough 1991, Keough & Quinn 1998). Trampling in coastal habitats is not only done when intertidal and shallow subtidal reefs are exposed at low tide, they are also trampled when people wade or swim through shallow water over the reef (e.g. walkers, anglers, swimmers, snorkelers, SCUBA-divers) (e.g. Addison et al 1998, Thompson et al. 2002). Walking when the reef is submerged and swimming can indirectly impact reef organisms through increased sedimentation stress as sediment is resuspended with movement through shallow water.

Studies examining how human movement affects sedimentation are uncommon. Neil (1990) showed that walking by humans on coral reefs mobilised sediment in the water column and increased sediment deposition on the reef at loads that had the potential to cause moderate stress in corals. Zedler (1978) showed that trampling by humans reduced the abundance of coralline algal turf, resulting in a loss of sand associated with these mats of algae and in turn the organisms that rely on these sediments for habitat (see also Brown & Taylor 1999). On rocky intertidal reefs, increased sedimentation is predicted to have dramatic impacts on filter-feeders and grazers because sediments can also smother and obstruct the feeding apparatus of sessile barnacles and mussels, and impede movement and reduce the amount of emergent rock substratum necessary for feeding by snails and limpets (e.g. Airoidi & Virgilio 1998, Pulfrich et al. 2003, Schiel et al. 2006). For example, Airoidi and Hawkins (2007) demonstrated that a 1 mm thick layer of sediment could substantially reduce the grazing rates of intertidal limpets. In contrast, other intertidal species may respond positively to the presence of sediments (see Thompson et al. 2002). For example, Huff & Jarett (2007) found that psammophilic snails increased in abundance following the addition of sand to intertidal coralline turf.

In many coastal landscapes, rocky intertidal seashores are adjacent to sandy beaches and soft-sediment subtidal habitats, resulting in the influx and deposition of sediment on these rock platforms, which can be naturally dynamic in space and time (e.g. Littler et al. 1983, Airoidi 2003). In sheltered regions, these rock platforms often accumulate sand, particularly in shallow depressions and pools common in the rock surface and within and around turfing and foliose macroalgae and invertebrates that bind sediment (Airoidi 2003), and the platform can shift from exposed rock to being covered by sand. Human activity in these regions, and particularly trampling, can potentially influence the distribution and load of sediment on the rock platform. Importantly, sediment is often integrally associated with species, such as mussel-sediment or algal turf-sediment matrixes, that form an integral part of the reef habitat, which may in turn influence the abundance and composition of infauna as well as mobile and sessile species present (Airoidi 2003). Here we investigated how sediment deposition affects species assemblages on a sheltered, rocky intertidal platform in southern Australia dominated by mats of mussels, barnacles and a suite of gastropods. We tested the hypotheses that trampling by humans indirectly influences the abundance of macroinvertebrates on the platform through increased sediment load. We did this by simulating various intensities of trampling on a reef when it was covered by shallow water and then measuring sedimentation on the reef. Based on these estimates and a range of potential sediment loads that might occur on the reef due to human (e.g. terrestrial run-off from land clearing) and natural disturbance (e.g. storms), we then manipulated sediment load on the reef and monitored the response of the dominant macroinvertebrate species.

MATERIALS AND METHODS

Study location and intertidal communities

The study was done over the austral summer on a rocky intertidal platform at Coobowie on the Yorke Peninsula, which is a rural area approximately 70 km southwest of Adelaide, South

Australia (35°03'16" S, 137°43'45" E). The platform is on the southern side of a sheltered inlet (Salt Creek Bay) and experiences minimal wave action. Typical coastal vegetation borders the landward edge of the rock platform and the seaward edge gradually leads into soft-bottom, subtidal habitat, which is the likely source of sediments (Airoidi 2003). The rock platform is almost flat and extends 2 km alongshore and 0.8 km from landward to seaward. Maximal tidal range during the study was 2.41 m.

Sediments (primarily sand) transported onto the platform are typically associated with two types of intertidal communities that are prevalent in the region: one dominated by the habitat-forming macroalga *Hormosira banksii* (hereafter referred to as macroalgal habitat), and the other, which lacks canopy-forming macroalgae, has less sediment, and occurs at slightly higher tidal elevations, dominated by patches of the mussel *Xenostrobus inconstans* with byssus bound in a sediment matrix (hereafter referred to as mussel habitat). The entire rock platform is pitted with small shallow depressions, allowing patches of sediment to accumulate on the platform at low tide. These habitats comprise species that are common and abundant in sheltered to moderately exposed bays throughout the region and more generally in southern Australia, including mussels, barnacles, macroalgae, and a suite of grazing, predatory and scavenging gastropods (e.g. Keough & Quinn, 1998, Underwood 1998).

Relationships between sediment cover and macroinvertebrate abundance

The relationship between the cover of sediment and abundance of dominant macroinvertebrate species was quantified in the mussel habitat by sampling quadrats (1 m × 1 m) with varying levels of sediment cover (0%, 20%, 40%, 60%, 80%; $n = 3$ per cover level). The primary substratum of the mussel habitat is dominated by the mussel *X. inconstans* and the barnacle *E. modestus*, and there are small patches of sediment in a thin layer covering the bare rock. Common species of gastropods include the grazers *Nerita atramentosa*, *Bembicium*

nanum, *Austrocochlea porcata* and *A. constricta*, the predator *Haustrum vinosum* (formerly *Lepsiella vinosa*) and *Cacozeliana granarium* (formerly *Bittium granarium*), which feeds on microalgae and particulate matter.

Abundances of bare rock, sediment overlying the rock, and the mussel *X. inconstans* were estimated by their percent covers using the point-intercept method, with 100 uniformly spaced points over the 1 m × 1 m quadrat. The depth of sediment was also estimated at three randomly selected places in each quadrat. Densities for the majority of gastropods were also quantified in the 1 m × 1 m quadrat. Due to their great abundances and patchy distribution, however, densities of *Bembicium nanum* and *Elminius modestus* were randomly sampled in 16 smaller (9 cm × 9 cm) quadrats (an area of 1296 cm²) within the larger 1 m² quadrat. Linear regression with untransformed data ($n = 15$ observations: $n = 3$ replicates for each of 5 sediment cover levels) was used to examine the relationship between percent cover of sediment and dependent variables (sediment depth, cover of bare rock, macroinvertebrate abundance).

Sediment load due to trampling

Changes to the loading (i.e. net effect of removal and deposition) of sediment on the rock platform due to trampling by humans were determined in both the macroalgal and mussel habitats. A 50 m × 100 m area was selected in each of the macroalgal and mussel habitats. Twenty quadrats (0.3 m × 1.0 m), separated by a least 2 m, were located in each of these habitats and permanently marked. To minimise initial variation in the abundances of organisms and sediment dynamics, quadrats in each habitat were randomly selected where there was a relatively similar cover of the dominant, habitat-forming species: $53 \pm 2.1\%$ (mean $\pm$ se; $n = 20$) cover of *H. banksii* in the macroalgal habitat; and $41 \pm 1.7\%$ (mean $\pm$ se; $n = 20$) cover of *X. inconstans* in the mussel habitat. Four intensities of trampling were

randomly allocated to the 20 quadrats ($n = 5$ quadrats per trampling intensity): 0, 15, 30 or 60 trampling passes. A trampling pass consisted of one person (weight 75 kg, height 175 cm), wearing rubber-soled shoes (the most commonly worn footwear in intertidal areas; Bally & Griffiths 1989) walking at a normal pace across the length of the quadrat (i.e. one or two steps in the 1 m long quadrat per trampling pass). Trampling was done when water covered the platform to a depth of 15 to 30 cm (ankle to calf depth) and within two hours of the platform being covered by water during a rising tide. Trampling at this time was most likely to alter sediment deposition and to simulate accurately visitation to the reef for recreational activities (e.g. walking, fishing, swimming).

There were six trampling days from mid November to mid December, with trampling on each of two consecutive days over three successive low tide periods (i.e. each two-day trampling event was separated by approximately 14 days). Intensities, frequency and timing of trampling were chosen to represent weekend visitation to metropolitan rocky intertidal reefs with the onset of summer (e.g. Kingsford et al. 1991). The effect of trampling on loading of sediment onto the rock platform in each habitat was quantified at two times: immediately after the third and fifth days of trampling. Therefore, sediment loads reflected the effects of trampling of a single trampling day. Sedimentation was quantified by placing two, circular trays (diameter = 18 cm, area = 254.5 cm², depth = 2.5 cm) 0.5 m away from and at opposite ends of each of the 20 quadrats, trampling on a rising tide, and then collecting the sediment in the trays. Trays were collected in the same order as they were trampled, allowing each tray to collect sediment for approximately 45 min. Due to logistical constraints associated with having the same person trample each quadrat while ensuring constant water depth while trampling, trays in the mussel habitat had to be left overnight (approximately 16 h) and collected the following morning. Trays were collected by placing them in plastic bags, and then sediment was rinsed and filtered to remove large pieces of organic matter, dried to a

constant weight in an oven at 60°C, and weighed. The total value from the two sediment trays was used to estimate sediment load per quadrat (i.e. g sediment per 509 cm² area of sediment tray). Differences in sediment load among trampling intensities and habitats for each of the two trampling times were analysed with two-factor analysis of variance (ANOVA) using untransformed data.

The cover of sediment in all quadrats in the macroalgal habitat was also measured before and after the six days of trampling to determine if trampling intensity influenced the removal or accumulation of sediment on the rock platform. Differences in sediment cover in the macroalgal habitat before and after six days of trampling was analysed with one-factor ANOVA using untransformed data. For both the cover and loading of sediment, student-Newman-Keuls (SNK) multiple comparisons tests were used to determine location of differences among means after significant ANOVA.

Effect of sediment load on macroinvertebrate abundance

To determine the effect of sediment load on species living in the mussel habitat, various amounts of sediment were added to the rock platform from early January to mid February and the response of dominant macrofaunal species was assessed before the first addition of sediment until early March. Forty-two quadrats (1 m × 1 m), each separated by at least 2 m, were randomly located in areas where there was 20% to 30% cover of the mussel *X. inconstans*. Six intensities of sediment disturbance were randomly allocated to the 42 quadrats: 0 g, 125 g, 250 g, 500 g, 1000 g, and 2000 g of dry sediment ($n = 7$ replicates per intensity). These sediment loads were chosen to mimic disturbances, ranging from those recorded here due to trampling by humans (up to 313 g / m² after a trampling day; see Sediment load due to trampling subsection in Results) to natural and anthropogenic processes such as those resulting from storms and urban run-off (see Branch et al. 1990, Airolidi et al.

1996, Airoidi 2003, Connell 2005, Airoidi & Hawkins 2007). Natural sediment loading into the habitat (quantified as described in the previous subsection) for two periods spanning the duration of the experiment ranged from 13.2 to 17.6 g / m² per day. Sediment was added three times at approximately fortnightly intervals, which simulated three pulse disturbances resulting from trampling by humans during weekend visits to the rock platform during summer (or other larger sediment disturbances). Sediment was collected from the intertidal platform, dried, and applied at low tide. Sediments were distributed evenly throughout the quadrat using a coarse grade sieve (i.e. the sieve did not retain any of the sediment) held about 0.5 m above the rock platform.

The response of the dominant macroinvertebrate species was quantified at three times: (1) before the first addition of sediment (and there were no significant differences among sediment addition treatments for any of the dependent variables; unpublished data); (2) approximately two weeks after the first sediment addition (and before the second sediment addition); and (3) approximately two weeks after the third and final sediment addition (about six weeks after the first sediment addition). The effects of sediment addition were quantified in each quadrat by sampling the percent cover of sessile species occupying the substratum (except for *E. modestus* where density was measured) and the densities of all dominant macroinvertebrate species. Abundances of bare rock, sediment overlying the rock, and the mussel *X. inconstans* were estimated by their percent covers using the point-intercept method, with 100 uniformly spaced points over the 1 m × 1 m quadrat. The depth of sediment was also estimated at three randomly selected places within each quadrat. Densities for the majority of gastropods were also quantified in the 1 m × 1 m quadrat. Due to their small numbers, densities of *Austrocochlea porcata*, *A. constricta*, *Haustrum vinosum* and *Cacozeliana granarium* (all of which showed positive associations with sediment cover; see Results) were combined and collectively referred to as “other gastropods”. Again, densities of *B. nanum* and

E. modestus were randomly sampled in 16 smaller (9 cm × 9 cm) quadrats (an area of 1296 cm²) within the larger 1 m² quadrat. Differences in percent cover (sediment, bare rock, the mussel *X. inconstans*; all arcsine-transformed), sediment depth (log-transformed), and macroinvertebrate densities (*E. modestus*, *N. atramentosa*, *B. nanum*, other gastropods; all log-transformed) among sediment disturbance intensities were analysed six weeks after the first addition of sediment (i.e. two weeks after the third addition of sediment) with one-factor ANOVA, and SNK multiple comparisons tests were used to determine location of differences among means after significant ANOVA.

RESULTS

Relationships between sediment cover and macroinvertebrate abundance

The cover and depth of sediments on the rock platform were strongly correlated (Fig. 1), with some areas with 80% cover having sediments 1 cm deep. Not surprisingly, areas of the platform with increasing sediment cover had declining availability of bare rock (Fig. 1). There were both positive and negative relationships between the cover of sediment on the rock platform and the abundance of macroinvertebrates, and these were dependent on functional group and species. The abundance of filter feeders was negatively associated with sediment cover (Fig. 1), including cover of the mussel *Xenostrobus inconstans*, and the barnacle *Elminius modestus* was not present in areas with greater than about 20% sediment cover. The direction and strength of associations of gastropods with sediment cover were highly variable. The grazers *Nerita atramentosa* and *Bembicium nanum* were negatively related to increasing sediment cover, whereas the grazer *Austrocochlea porcata*, the grazer *A. constricta*, the predator *Haustorium vinosum*, and *Cacozeliana granarium* were positively related to sediment cover (Fig. 1).

Sediment load due to trampling

In both the macroalgal and mussel habitats, all intensities of trampling increased the deposition of sediment onto the rock platform above natural levels of sedimentation, with the most intensive trampling resulting in more than an order of magnitude more sediment deposited on the platform (Fig. 2). After the third trampling day, sediment load was directly related to trampling intensity in both habitats, and was consistently higher in the macroalgal than mussel habitat (Time 1 in Fig. 2; trampling intensity: $F_{3,16} = 136.7$, $P < 0.001$; SNK: $60 > 30 > 15 > 0$ trampling passes; habitat: $F_{1,16} = 7.35$, $P = 0.015$). Indeed, maximum sediment loads were 313 g m^{-2} and 283 g m^{-2} in the macroalgal and mussel habitats, respectively.

In contrast, after five trampling days, substantial components of the species assemblage were removed due to direct trampling impacts, and sediment load in trampled areas was reduced (and more variable) than after three trampling days (compare Time 1 with Time 2 in Fig. 2). Nevertheless, in both habitats sediment load in trampled areas remained greater than in areas with natural levels of sedimentation (Time 2 in Fig. 2; trampling intensity: $F_{3,16} = 5.8$, $P = 0.007$; SNK: $60 = 30 = 15 > 0$ trampling passes). In the macroalgal habitat, there was a trend for increased sediment load with trampling intensity, but sediment load in the mussel habitat was extremely variable and reduced at the highest compared to the more moderate levels of trampling intensity. Regardless, there was no difference in sediment load between habitats (Time 2 in Fig. 2; habitat: $F_{1,16} \ll 0.1$, $P = 0.993$).

Trampling in the macroalgal habitat dramatically reduced sediment cover, with close to a 90% reduction in the most intensively trampled areas (0 passes = $21.4 \pm 5.5\%$; 15 passes = $5.9 \pm 1.9\%$; 30 passes = $7.7 \pm 3.8\%$; 60 passes = $2.3 \pm 2.3\%$; mean $\pm$ se, $n = 5$). There was no significant difference in sediment cover among trampling intensities before sampling ($F_{3,16} = 1.7$, $P = 0.213$), and sediment cover in control quadrats without trampling remained

remarkably consistent before ($20.0 \pm 3.1\%$; mean $\pm$ se, $n = 5$) and after trampling, whereas six trampling days significantly reduced sediment cover in trampled areas ($F_{3,16} = 5.2$, $P = 0.011$; SNK: $0 > 15 = 30 = 60$ trampling passes).

Effect of sediment load on macroinvertebrate abundance

There was a natural influx of sediment onto the rock platform due to a storm in the first two weeks of the experiment: sediment cover (but not depth) tripled in control areas where sediment was not added but then gradually returned to levels at the start of the experiment after six weeks, demonstrating the dynamic nature of sediment disturbance on this reef (Fig. 3). Interestingly, trends in abundance in relation to sediment addition, if not their magnitudes, were generally the same after two and six weeks (Fig. 3). After six weeks, there was a strong, positive relationship between the cover and depth of sediment on the rock platform with the addition of sediments to the mussel habitat (Fig. 3; sediment cover: $F_{5,36} = 32.8$, $P < 0.001$; sediment depth: $F_{5,36} = 85.9$, $P < 0.001$), indicating that sediment accumulated on the platform over time as more sediment was added. Interestingly, although the cover of sediment on the platform declined from two to six weeks after application, the depth of sediment increased, indicating that sediments initially formed a relatively thin layer but then were swept into and accumulated in particular areas on the platform over time. Not surprisingly, as sediment accumulated on the platform with each application, the cover of bare space decreased, with about 50% less available space at the greatest level of sediment load compared to areas where no sediment was added (Fig. 3; $F_{5,36} = 22.0$, $P < 0.001$).

Cover of the rock by patches of the mussel *X. inconstans* was remarkably consistent over the entire study period and, despite the addition of sediment, even increased marginally from that at the start of the experiment (Fig. 3; $F_{5,36} = 2.4$, $P = 0.053$). There was also no influence of sediment addition on the abundance of the only other dominant sessile species in the mussel

habitat, the barnacle *E. modestus* (Fig. 3; $F_{5,36} = 1.3$, $P = 0.274$). Nevertheless, there was a trend for decreased barnacle abundance with sediment addition, but abundances were highly variable and the patterns were strongly influenced by densities in quadrats where 125 g of sediment was added (Fig. 3).

The mobile gastropods responded in generally the same manner to sediment addition as observed when examining their patterns of abundance in association with sediment cover (see Fig. 1). There was an influx in the abundance of the grazing gastropods *N. atramentosa* and *B. nanum* to all quadrats after the start of the experiment, except those receiving the largest loads of sediment (Fig. 3). After six weeks densities of these grazers generally declined with increasing sediment load, but this reduction was most pronounced only at the highest level of sediment addition (Fig. 3; *N. atramentosa*: $F_{5,36} = 8.1$, $P < 0.001$; SNK: $0 = 125 = 250 > 1000 = 2000$ g sediment added; *B. nanum*: $F_{5,36} = 11.1$, $P < 0.001$; SNK: $0 = 125 = 250 = 500 = 1000 > 2000$ g sediment added). In contrast to *N. atramentosa* and *B. nanum*, the small densities of other gastropods (*A. porcata*, *A. constricta*, *H. vinosum* and *B. granarium*) were positively influenced by sediment load, with numbers increasing at the two highest levels of sediment addition (Fig. 3; $F_{5,36} = 3.0$, $P = 0.219$; SNK: cannot be resolved).

DISCUSSION

Our study has shown that sediment disturbance impacts the abundance of species living on this sheltered rocky reef, and while increasing sediment load negatively influenced most species, some displayed positive responses. Trampling by humans resulted in the redistribution of sediment on the rock platform, but loading of sediment at levels associated with trampling had only minor effects on species' abundances. In contrast, at higher levels of sediment addition there were dramatic effects on the abundance of the majority of species, suggesting potential shifts in species composition on the rock platform from species sensitive

to higher sediment environments to those that tolerate or favour increased sediment loads (Thompson et al. 2002, Airoidi 2003).

Trampling when shallow water covered the rock platform increased sedimentation by up to an order of magnitude beyond natural levels. It was clear that trampling disturbed sediment on the rock platform, suspended it in the water column, and then it was deposited back onto the rock platform (Neil 1990). Sediment load increased outside of trampled areas and sediment cover was reduced and the availability of bare rock increased within trampled areas, indicating that trampling redistributes (and possibly results in a net loss of) sediment on the reef. Brown and Taylor (1999) also found that trampling by humans on a rock platform in New Zealand reduced the amount of sand on the platform, and this was primarily due to losses of coralline algal turfs with which sand is associated. In our study we also noticed losses of sediment associated with reductions in cover of the macroalgal *H. banksii* and the mussels *X. inconstans*, species that trap and bind sediment (Povey & Keough 1991, Airoidi 2003).

The effect of trampling on sedimentation is complex because trampling not only resuspends sediment directly but also alters sediment dynamics by removing species that bind sediment. For example, after the first trampling event, sedimentation (i.e. suspension of sediment due to trampling and deposition) on the rock platform in both habitats was directly related to trampling intensity. Loading of sediment onto the rock platform following the second trampling event was, however, less than following the first trampling event, and the trend for increased sedimentation with trampling intensity had changed, with the most severely trampled areas having reduced loading of sediment compared to areas that had been less intensive trampled. These differences in the relationship between trampling intensity and sediment load occurred because the absolute amount of sediment on the platform that could

be resuspended was reduced with increasing trampling and the loss of sediment-binding species. Therefore, the influence of trampling intensity on redistribution of sediment load across the rock platform is likely non-linear, with sedimentation in the local area increasing with increasing trampling intensity, and then declining when trampling is so intense that unbound sediment and sediment bound within the biotic community is removed from the platform. Heavily trampled platforms would therefore be expected to be largely free of sediment and species that depend on it, such as epi- and infauna associated with sediment-bound biological communities (e.g. Brown & Taylor 1999).

Abundances of sessile species, particularly the mussel *X. inconstans* and the barnacle *E. modestus*, were not influenced by sediment addition. We initially predicted that sessile species would be most impacted by sediment addition because they are not able to move away from impacted areas. Indeed, previous studies have shown that burial by sediment on rocky reefs can cause mortality of mussels and barnacles (Seapy & Littler 1982, Branch et al. 1990, Menge et al. 1994), although burial was not common in our study. The amount of sediment added in our study was less than that associated with extreme storm events that bury and kill sessile intertidal invertebrates (Daly & Mathieson 1977, Littler et al. 1983), and this might account for differences here (although the trend for reduced barnacle abundances with sediment addition should be noted). It is possible that, at the highest sediment loads applied in our experiment, sessile species experienced minimal and only transient burial given the dynamic sediment environment on this rocky reef. Indeed, over time the cover of sediment diminished, but sediment became deeper and accumulated in particular areas on the platform, such as depressions in the rock or was bound in the byssus of mussels. Consequently, although cover of sediment increased with sediment addition, much of the rock platform was free of substantial amounts of sediment or merely covered by a thin sediment layer, even at the highest levels of sediment addition. We would nevertheless predict that the barnacle *E.*

modestus, which was killed by burial in some places at highest sediment loads, would be more greatly affected by increased sedimentation than the mussel *X. inconstans*. The mussel *X. inconstans* not only inhabits rocky intertidal shores, but also lives in sheltered sedimentary habitats (Buschbaum et al. 2009), such as salt marshes where it is fed upon by *H. vinosum* (Morton 2004). Moreover, *X. inconstans* binds sediment and lives within or at the surface of the sediment-mussel matrix, providing habitat for infauna such as polychaete worms (authors' personal observations, Buschbaum et al. 2009), and thus may come to dominate the rock platform should increased sediment loading shift the rocky shore to a soft-sediment habitat.

Changes to the load of sediment and availability of bare rock on the platform strongly contributed to changes in the abundance of gastropods. Both in mensurative and experimental studies the grazing gastropods *Nerita atramentosa* and *Bembicium nanum* were negatively influenced by increasing sediment load (Daly & Mathieson 1977, Seapy & Littler 1982, Littler et al. 1983). Both of these species graze on microalgae that is typically refreshed on the rock substratum with each successive tide (Underwood 1975, 1984), and individuals were thus likely to reduce feeding activity or move out of areas that received sediment and reduced food resources (Airolidi & Hawkins 2007, Huff & Jarett 2007, Atalah & Crowe 2010). Similarly, Airolidi & Hawkins (2007) experimentally manipulated sediment load in the laboratory and found that 1 mm thick layers of sediment equivalent to 500 g / m² were sufficient to reduce substantially the grazing activity of the limpet *Patella vulgata*. Effects on *N. atramentosa* and *B. nanum* in our study were most pronounced at the highest levels of sediment addition, indicating that smaller magnitudes of sediment deposition and thin layers of sediment on the rock, such as those recorded due to trampling here, are likely tolerated by these species. Indeed, *N. atramentosa* appeared to bulldoze the thin layers of sediment (Bertness 1984). Importantly, sediment was mostly comprised of sand, although precise grain size structure was not determined, and fine fractions have been found to have greater impacts

on grazers than coarse sediments (Airoldi & Hawkins 2007). Higher sediment loads change the physical character of the substratum and create microhabitats well known to influence the movement of such gastropods.

Some gastropods displayed positive associations and responses to increasing cover and addition of sediment on the rock platform, including the grazers *Austrocochlea porcata* and *Austrocochlea constricta*, the predator *Haustrum vinosum*, and *Cacozeliana granarium* (Huff & Jarett 2007). These associations were confirmed by experimental addition of sediment, particularly for the predator *H. vinosum* and *B. granarium*, but numbers were small. Some of these rocky shore gastropods, such as *H. vinosum*, also inhabitant soft-sediment habitats with hard substrata, such as mangrove forests and salt marshes where they prey on barnacles and the mussel *X. inconstans*, respectively (e.g. Bayliss 1982, Morton 2004), and maybe thus able to cope or respond opportunistically to the presence of sediment and perhaps the paucity of other gastropods. Results here therefore suggest that on this sheltered rock platform increased sediment loads might lead to changes in the composition of gastropods and other species, favouring those that tolerate or are facilitated by sediment (Daly & Mathieson 1977, Brown & Taylor 1999, Thompson et al. 2002, Huff & Jarett 2007). More generally, species that are flexible in their use of habitat (e.g. mussels such as *X. inconstans* and the predator *H. vinosum*) are more likely to be represented across habitats that are subjected to environmental change, such as rocky coasts subjected to increased sedimentation or variable sediment regimes.

A key element of this study was that we applied the trampling disturbance when shallow water covered the rock platform. The vast majority of studies examining the impacts of trampling by humans on reefs have been done on emergent rock at low tide when there is no water covering the platform, but it is common for people to walk, fish and swim on the rock

platform when it is covered by water, particularly in sheltered areas where sediment loads might be expected to be greatest (e.g. Addison et al. 1998, Thompson 2002). Neil (1990), for example, showed that sediment resuspension and deposition is an important component of the overall impact of trampling on coral reefs, with people trampling in 30 cm of water substantially increasing sedimentation on the reef. Our findings across two different habitat types confirm the influence of trampling on sediment dynamics, but reveal that, at least for levels of trampling used in this study, sediment deposition associated with trampling has only small impacts on the benthic species assemblage in the mussel habitat (and less than those of the direct impacts of trampling). Nevertheless, we have demonstrated that trampling has the potential to impact rocky reef species at higher sediment loads, and trampling could create such impacts if sediment load was in naturally greater abundances on the rock platform, trampling intensities were increased, or trampling was done under different conditions (water depth, wave exposure, etc.). Therefore, the importance of tidal state, which produced these indirect effects on the species assemblage through changes in sediment load, need to be incorporated into the suite of human visitation and trampling impacts, particularly across rocky shores of varying wave exposure (sheltered versus exposed platforms have different sediment dynamics) and habitats with varying sediment environments (e.g. rocky reefs, coral reefs, mangrove forests). For example, the loss of species that trap and bind sediment might be particularly important to maintaining species composition on the rock platform.

This study shows the dramatic effects that sediment disturbance can have on species assemblages living on rocky shores. Patterns of sedimentation in response to trampling were generally similar across both the macroalgal and mussel habitats, suggesting that such disturbances might be generally important across a range of habitats (Neil 1990). Although experiments were only done at one site in the mussel habitat, thus limiting their generality, results of experimental manipulations were consistent with patterns of species abundance in

the habitat in relation to sediment cover, providing support for the impacts of sedimentation on species assemblages.

The cumulative effect of small-scale disturbances such as trampling might have similar impacts to larger-scale disturbances due to sediment run-off from the terrestrial environment or storm events. The persistence of the current species assemblage in our study is likely due to the dynamic nature of sedimentation across the shore and features of the platform, such as depressions in the rock and the presence of species that retain sediment, that influence sediment accumulation. With increasing sediment loading of coastal waterways from urban and agricultural development, however, we might predict that species assemblages on rocky reefs may shift to be dominated by species that can resist or thrive in multiple habitats, particularly those with fluctuating sediment regimes (e.g. Taylor & Littler 1982, Branch et al. 1990, Irving & Connell 2002). To predict the influence on sedimentation on coastal habitats, ecologists will need to quantify the magnitude and character of sediment loaded into coastal communities in space and time so that mensurative and manipulative experiments can be done to assess impacts on and understand mechanisms driving changes in species assemblages (Airolidi 2003).

Acknowledgements. We thank S. Barker, C. Wilson and P. Goodsell for help in the field. G. Ganf and K. Walker made comments on early drafts of the manuscript. R. McWilliam assisted with the literature search. We particularly thank J. Tanner for helpful suggestions on all aspects of the study. Funding was provided by the University of Adelaide and the Institute for Conservation Biology at the University of Wollongong.

LITERATURE CITED

- Addessi L (1994) Human disturbance and long-term changes on a rocky intertidal community. *Ecol Appl* 4:786–797
- Addison PFE, Koss RS, O'Hara TD (1998) Recreational use of a rocky intertidal reef in Victoria: implications for ecological research and management. *Australas J Environ Manage* 15:169–179
- Airolidi L (1998) Roles of disturbance, sediment stress, and substratum retention on spatial dominance in algal turf. *Ecology* 79:2759–2770
- Airolidi L (2003) The effects of sedimentation on rocky coast assemblages. *Oceanogr Mar Biol Annu Rev* 41:161–263
- Airolidi L, Hawkins SJ (2007) Negative effects of sediment deposition on grazing activity and survival of the limpet *Patella vulgata*. *Mar Ecol Prog Ser* 332:235–240
- Airolidi L, Virgilio M (1998) Responses of turf-forming algae to spatial variations in the deposition of sediments. *Mar Ecol Prog Ser* 165:271–282
- Airolidi L, Fabiano M, Cinelli F (1996) Sediment deposition and movement over a turf assemblage in a shallow rocky coastal area on the Ligurian Sea. *Mar Ecol Prog Ser* 133:241–251
- Araújo R, Vaselli S, Almeida M, Serrão E, Sousa-Pinto I (2009) Effects of disturbance on marginal populations: human trampling on *Ascophyllum nodosum* assemblages at its south distribution limit. *Mar Ecol Prog Ser* 378:81–92
- Atalah J, Crowe TP (2010) Combined effects of nutrient enrichment, sedimentation and grazer loss on rock pool assemblages. *J Exp Mar Biol Ecol* 388:51–57
- Balata D, Piazzzi L, Cinelli F (2007) Increase of sedimentation in a subtidal system: Effects on the structure and diversity of macroalgal assemblages. *J Exp Mar Biol Ecol* 351:73–82

- Bally R, Griffiths CL (1989) Effects of human trampling on an exposed rocky shore. *Int J Environ Stud* 34:115–125
- Bayliss DE (1982) Switching by *Lepsiella vinosa* (Gastropoda) in South Australian mangroves. *Oecologia* 54:212–226
- Beauchamp KA, Gowing MM (1982) A quantitative assessment of human trampling effects on a rocky intertidal community. *Mar Environ Res* 7:279–293
- Bertness MD (1984) Habitat and community modification by an introduced herbivorous snail. *Ecology* 65:370–381
- Branch GM, Eekhout S, Bosman AL (1990) Short-term effects of the 1988 Orange River floods on the intertidal rocky shore communities of the open coast. *Trans R Soc Arr* 47:331–354
- Brosnan DM, Crumrine LL (1994) Effects of human trampling on marine rocky shore communities. *J Exp Mar Biol Ecol* 177:79–97
- Brown PJ, Taylor RB (1999) Effects of trampling by humans on animals inhabiting coralline algal turf in the rocky intertidal. *J Exp Mar Biol Ecol* 235:45–53
- Buschbaum C, Dittman S, Hong J-S, Hwand I-S, Strasser M, Thiel M, Valdivia N, Yoon S-P, Reise K (2009) Mytilid mussels: global habitat engineers in coastal sediments. *Helgol Mar Res* 63:47–58
- Connell SD (2005) Assembly and maintenance of subtidal habitat heterogeneity: synergistic effects of light penetration and sedimentation. *Mar Ecol Prog Ser* 289:53–61
- Daly MA, Mathieson AC (1977) The effects of sand movement on intertidal seaweeds and selected invertebrates at Bound Rock, New Hampshire, USA. *Mar Biol* 43:45–55
- Eckrich CE, Holmquist JG (2000) Trampling in a seagrass assemblage: direct effects, response of associated fauna, and the role of substrate characteristics. *Mar Ecol Prog Ser* 201:199–209

- Fletcher H, Frid CLJ (1996) Impact and management of visitor pressure on rocky intertidal algal communities. *Aquat Conserv: Mar Freshw Ecosys* 6:287–297
- Ghazanshahi J, Huchel TD, Devanny JS (1983) Alteration of southern California rocky shore ecosystems by public recreational use. *J Environ Manage* 16:379–394
- Hardiman N, Burgin S (2010) Recreational impacts on the fauna of Australian coastal marine ecosystems. *J Environ Manage* 91:2096–2108
- Headley AD, Sale P (1999) The impact of trampling by student groups on saltmarsh vegetation. *Field Stud* 9:513–530
- Huff TM, Jarett JK (2007) Sand addition alters the invertebrate community of intertidal coralline turf. *Mar Ecol Prog Ser* 345:75–82
- Irving AD, Connell SD (2002) Sedimentation and light penetration interact to maintain heterogeneity of subtidal habitats: algal versus invertebrate dominated assemblages. *Mar Ecol Prog Ser* 245:83–91
- Keough MJ, Quinn GP (1998) Effects of periodic disturbances from trampling on rocky intertidal algal beds. *Ecol Appl* 8:141–161
- Kingsford MJ, Underwood AJ, Kennelly SJ (1991) Humans as predators on rocky reefs in New South Wales, Australia. *Mar Ecol Prog Ser* 72:1–14
- Liddle MJ (1991) Recreation ecology: effects of trampling on plants and corals. *Trends Ecol Evol* 6:13–17
- Liddle MJ, Kay AM (1987) Resistance, survival and recovery of trampled corals on the Great Barrier Reef. *Biol Conserv* 42:1–18
- Littler MM, Martz DR, Littler DS (1983) Effects of recurrent sand deposition on rocky intertidal organisms: importance of substrate heterogeneity in a fluctuating environment. *Mar Ecol Prog Ser* 11:129–139

- Luna-Pérez B, Valle-Pérez C, Sánchez-Lizaso JL (2011) *Halocynthia papillosa* as SCUBA diving impact indicator: An in situ experiment. *J Exp Mar Biol Ecol* 398:33–39
- Morton B (2004) Predator-prey interactions between *Lepsiella vinosa* (Gastropoda: Muricidae) and *Xenostrobus inconstans* (Bivalvia: Mytilidae) in a southwest Australian marsh. *J Moll Stud* 70:237–245
- Neil D (1990) Potential for coral stress due to sediment resuspension and deposition by reef walkers. *Biol Conserv* 52:221–227
- Povey A, Keough MJ (1991) Effects of trampling on plant and animal populations on rocky shores. *Oikos* 61:355–368
- Pulfrich A, Parkins CA, Branch GM, Bustamante RH, Velásquez CR (2003) The effects of sediment deposits from Namibian diamond mines on intertidal and subtidal reefs and rock-lobster populations. *Aquat Conserv Mar Freshw Ecosyst* 13:257–278
- Purcell SW (2000) Association of epilithic algae with sediment distribution on a windward reef in the northern Great Barrier Reef, Australia. *B Mar Sci* 66:199–214
- Robles C (1982) Disturbance and predation in an assemblage of herbivorous *Diptera* and algae on rocky shores. *Oecologia* 54:23–31
- Rogers CS (1990) Responses of coral reefs and reef organisms to sedimentation. *Mar Ecol Prog Ser* 62:185–202.
- Ross PM (2006) Macrofaunal loss and microhabitat destruction: The impact of trampling in a temperate mangrove forest, NSW Australia. *Wetlands Ecol Manage* 14:167–184
- Rossi F, Forster RM, Montserrat F, Ponti M, Terlizzi A, Ysebaert T, Middelburg JJ 2007. Human trampling as short-term disturbance on intertidal mudflats: effects on macrofaunal biodiversity and population dynamics of bivalves. *Mar Biol* 151:2077–2090

- Schiel DR, Wood SA, Dunmore RA, Taylor DI (2006) Sediment on rocky intertidal reefs: effects on early post-settlement stages of habitat-forming seaweeds. *J Exp Mar Biol Ecol* 331:158–172
- Taylor PR, Littler MM (1982) The roles of compensatory mortality, physical disturbance, and substrate retention in the development and organization of a sand-influenced, rocky-intertidal community. *Ecology* 63:135–146.
- Thompson RC, Crowe TP, Hawkins SJ (2002) Rocky intertidal communities: past environmental changes, present status and predictions for the next 25 years. *Environ Conserv* 29:168–191
- Underwood AJ (1975) Comparative studies on the biology of *Nerita atramentosa* Reeve, *Bembicium nanum* (Lamarck), and *Cellana tramoserica* (Sowerby) (Gastropoda: Prosobranchia) in S.E. Australia. *J Exp Mar Biol Ecol* 18:153–172
- Underwood AJ (1984) Microalgal food and the growth of the intertidal gastropods *Nerita atramentosa* and *Bembicium nanum* at four heights on a shore. *J Exp Mar Biol Ecol* 79:277–291
- Underwood AJ (1998) Grazing and disturbance: an experimental analysis of patchiness in recovery from a severe storm by the intertidal alga *Hormosira banksii* on rocky shores in New South Wales. *J Exp Mar Biol Ecol* 231:291–306
- Vaselli S, Bertocci I, Maggi E, Benedetti-Cecchi L (2008) Effects of mean intensity and temporal variance of sediment scouring events on assemblages of rocky shores. *Mar Ecol Prog Ser* 364:57–66
- Wynberg RP, Branch GM (1997) Trampling associated with bait-collection for sandprawns *Callinassan kraussi* Stebbing: effects on biota of an intertidal sandflat. *Environ Conserv* 24:139–148

Zedler JB (1978) Public use effects in the Cabrillo National Monument intertidal zone;
project report. San Diego State University, Biology Department, San Diego, California,
USA

Figure captions

Fig. 1. Relationships between sediment cover (%) on the rock platform and sediment depth (mm), bare rock cover (%), *Xenostrobus inconstans* cover (%), *Elminius modestus* density (no. ind. 1296 cm⁻²), *Nerita atramentosa* density (no. ind. 1 m⁻²), *Bembicium nanum* density (no. ind. 1296 cm⁻²), *Austrocochlea porcata* (no. ind. 1 m⁻²), *Austrocochlea constricta* (no. ind. 1 m⁻²), *Haustrum vinosum* (no. ind. 1 m⁻²), *Cacozeliana granarium* (no. ind. 1 m⁻²) in the mussel habitat. Points represent means (± 1 se) of three replicate quadrats for each level of sediment cover. Lines depict significant linear relationships (r^2 and P values given in panels, $n=15$ quadrats for each).

Fig. 2. Mean (± 1 se) sediment load (1 g 509 cm⁻² $\approx$ 19.6 g m⁻²) in macroalgal and mussel habitats following various trampling intensities (0, 15, 30 or 60 passes) for each of two trampling times: immediately after the third trampling day (Time 1) and immediately after the fifth trampling day (Time 2). Note differences in scale on y-axes.

Fig. 3. Mean (± 1 se) sediment cover (%), sediment depth (mm), bare rock cover (%), *Xenostrobus inconstans* cover (%), *Elminius modestus* density (no. ind. 1296 cm⁻²), *Nerita atramentosa* density (no. ind. m⁻²), *Bembicium nanum* density (no. ind. 1296 cm⁻²) and other gastropods (*Austrocochlea porcata*, *Austrocochlea constricta*, *Haustrum vinosum* and *Cacozeliana granarium*; no. ind. m⁻²) in the mussel habitat after approximately two weeks (i.e. two weeks after the first sediment addition) and six weeks (i.e. two weeks after the third sediment addition) for various sediment loads (0, 125, 250, 500, 1000 or 2000 g m⁻²). Horizontal line in each graph is the mean value for all 42 quadrats at the beginning of the experiment.

Fig. 1.

Fig. 2.

Fig. 3.