

2012

Physical activity during school recess: a systematic review

Nicola D. Ridgers
Deakin University

Jo Salmon
Deakin University

Anne-Maree Parrish
University of Wollongong, aparrish@uow.edu.au

Rebecca M. Stanley
University Of South Australia

Anthony D. Okely
University of Wollongong, tokely@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/sspapers>



Part of the [Education Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Ridgers, Nicola D.; Salmon, Jo; Parrish, Anne-Maree; Stanley, Rebecca M.; and Okely, Anthony D., "Physical activity during school recess: a systematic review" (2012). *Faculty of Social Sciences - Papers*. 106.
<https://ro.uow.edu.au/sspapers/106>

Physical activity during school recess: a systematic review

Abstract

Context: Interest has increased in examining the physical activity levels of young people during school recess. Identifying correlates of their recess physical activity behaviors is timely, and would inform school-based physical activity programming and intervention development. The review examined the correlates of children's and adolescent's physical activity during school recess periods. **Evidence acquisition:** A systematic search of six electronic databases, reference lists, and personal archives identified 53 studies (47 focused on children) published between January 1990 and April 2011 that met the inclusion criteria. Data were analyzed in 2011. Correlates were categorized using the social-ecological framework. **Evidence synthesis:** Forty-four variables were identified across the four levels of the social-ecological framework, although few correlates were studied repeatedly at each level. Positive associations were found of overall facility provision, unfixed equipment, and perceived encouragement with recess physical activity. Results revealed that boys were more active than girls. **Conclusions:** Providing access to school facilities, providing unfixed equipment, and identifying ways to promote encouragement for physical activity have the potential to inform strategies to increase physical activity levels during recess periods.

Keywords

activity, during, physical, school, recess, review, systematic

Disciplines

Education | Social and Behavioral Sciences

Publication Details

Ridgers, N. D., Salmon, J., Parrish, A., Stanley, R. M. & Okely, A. D. (2012). Physical activity during school recess: a systematic review. *American Journal of Preventive Medicine*, 43 (3), 320-328.

Physical Activity during School Recess: A Systematic Review

Nicola D. Ridgers, PhD, Jo Salmon, PhD, Anne-Maree Parrish, PhD, Rebecca M. Stanley, BAppSc (Hons), Anthony D. Okely, EdD

From the Centre for Physical Activity and Nutrition Research (Ridgers, Salmon), Deakin University, Melbourne, the Faculty of Health and Behavioural Sciences (Parrish), Interdisciplinary Educational Research Institute (Okely), University of Wollongong, Wollongong, Health and Use of Time Group (Stanley), School of Health Sciences, University of South Australia, Adelaide, Australia

Address correspondence to: Nicola Ridgers, PhD, Centre for Physical Activity and Nutrition Research, Deakin University, 221 Burwood Highway, Burwood, 3125, Australia. E-mail: nicky.ridgers@deakin.edu.au.

Context: There has been increased interest in examining the physical activity levels of young people during school recess periods. Identifying correlates of their physical activity behaviors is timely, and would inform school-based physical activity programming and intervention development. The aim of this narrative review was to examine the correlates of children's and adolescent's physical activity during school recess periods.

Evidence acquisition: A systematic search of six electronic databases, reference lists and personal archives identified 53 studies (47 of which focused on children) published between January 1990 and April 2011 that met the inclusion criteria. Data were analyzed in 2011. Correlates were categorized using the social-ecological framework.

Evidence synthesis: Forty-four variables were identified across the 4 levels of the social-ecological framework, though few correlates were repeatedly studied at each level of influence. Positive associations were found between overall facility provision, unfixed equipment, and perceived encouragement and recess physical activity. Results also revealed that boys were more active than girls.

Conclusions: Providing access to school facilities, providing unfixed equipment, and identifying ways to promote encouragement for physical activity have the potential to inform strategies to increase physical activity levels during recess periods.

Introduction

Physical activity is positively associated with psychological well-being, bone health and motor skill development, and negatively associated with waist circumference and clustering of cardiovascular disease risk factors.¹⁻⁴ However, as children and adolescents may not be engaging in sufficient physical activity to benefit their health,¹ the promotion of physical activity is a public health priority.⁵

Children and adolescents spend a significant proportion of their waking at school. Non-curricular time, such as school recess periods (recess and lunchtime) and after-school programs, provide opportunities for children to be physically active within the school environment.⁶ Of these contexts, recess periods may provide the single greatest opportunity during the school year to impact on youth physical activity levels.⁷ However, in recent years there has been a trend to reduce the frequency and duration of school recess, or remove it from the school day altogether, often due to academic pressures. Consequently, it is important that school recess is included in school-based physical activity programming and policy, and that the recess environment is conducive for youth to make physically active choices.⁸ Whilst the scheduling and duration of recess periods vary between countries, schools should provide social and physical environments that facilitate enjoyable and safe physical activity engagement in this context.⁹ A number of reviews have examined correlates of preschool, children's and adolescents' physical activity,¹⁰⁻¹² yet these have typically focused on factors associated with whole-day activity. Since physical activity is a multidimensional behavior influenced by numerous factors across several domains,¹⁰ it is logical to also consider specific contexts in which children and adolescents are active. Few reviews have focused on the correlates of specific parts of the school day, such as school recess periods, yet synthesizing

such information could help to inform both school-based physical activity programming and intervention development aimed at increasing children's physical activity levels in this context. Furthermore, this information has the potential to inform future policy concerning recess periods and their role within the school day.

The aim of this review was to examine correlates of children's and adolescents' physical activity during school recess periods. From here on in, physical activity during school recess will be referred to as recess physical activity. For the purpose of this review, school recess was defined as the non-curriculum time allocated by schools between lessons for youth to engage in leisure activities.¹³

Methods

A systematic literature search of papers was conducted in 6 electronic databases (Pubmed, SportsDiscus, Web of Science, Proquest, Cochrane, and Scopus). Search strategies for the different databases included the following key words in three main areas: population (child, infant, youth, adolescent), school (school, primary, elementary, middle school, high school, secondary school), and recess (breaktime, break time, school recess, recess, playtime, lunchtime, free play). Only papers that had been published in peer-reviewed journals were considered for inclusion. Abstracts, expert opinion, case studies, conference proceedings and dissertations were not included. In addition to electronic searches, bibliographies of retrieved studies and authors' personal collections were also searched.

For inclusion in the review, studies were required to: a) include participants aged 5-18 years; b) have a measure of physical activity as the outcome variable; c) specifically examine physical activity during recess and/or lunchtime; d) have examined associations between

physical activity and other variables; e) be published between January 1990 and April 2011; and f) be published in peer-reviewed journals in English. School recess periods included morning recess, lunchtime, and afternoon recess. A previous review of recess physical activity levels reported that only two studies were published before 1990,¹³ therefore this was used as the start date for the current review. Data were collected and analyzed in 2011. Studies that reported total daily recess (e.g. sum of morning recess and lunchtime) or investigated specific recess periods were included. Before school time and after-school programs conducted on school sites were not considered as recess periods in this review. Studies that reported findings before/after school time and during school time were included where the results were presented separately for the recess periods. Data from control groups, baseline intervention studies, and intervention studies where the correlate was reported separately as a confounding variable were included. Longitudinal studies were included if at least one measurement period was conducted during elementary school and results at this time point reported. Studies that only reported the effectiveness of recess-based interventions on physical activity levels were excluded. Studies that investigated physical activity levels during recess periods in special populations (e.g. autistic spectrum disorders, attention deficit disorders) were also included in the review, as these studies collected data from schools where children with and without particular disorders had access to the same school environment and had recess and/or lunchtime together. When comparisons were made between children with and without particular disorders and no differences were found, all data were included for review. In circumstances where differences occurred, data from the children without the disorders were considered to enable comparisons with other studies. Lastly, studies that solely investigated physical activity levels of special populations during recess periods were included but noted accordingly in the results.

Results obtained from the initial literature search were independently assessed. Potentially relevant papers were selected by screening a) the title; b) the abstract; and c) the full paper if insufficient detail was provided in the abstract. Data extraction was then undertaken using standardized forms and included: country of study, study design, sample characteristics (e.g. sample size, age, gender), the physical activity measure used and the outcome variable, recess period(s) studied, recess characteristics (e.g. duration, supervision, environment), and results. Appendix A provides an overview of the characteristics of the studies that met the inclusion criteria. Data are presented separately for children (5-12 years) and adolescents (13-18 years). Due to some variability across studies in the age that children transition from primary (elementary) and/or middle schools to secondary (high) schools, if children were aged 11-12 but the study was conducted in secondary schools, they were classed as adolescents.

Categorization of variables

For consistency and comparability with previous reviews of correlates of youth physical activity,¹⁰⁻¹² a social-ecological framework was used. The social-ecological model posits that multiple levels of influence interact to promote or constrain participation in physical activity.¹⁴ Identified variables were categorized into four groups according to the social-ecological framework; individual, social, physical environment, and organizational/policy. Conceptually similar variables were combined (e.g. different facilities available for use during recess periods) to enable consistency with interpretation.

A range of measures were used to examine recess physical activity (see Appendix A). Children's physical activity was largely measured using objective measures, with accelerometry the most commonly used method (36% of all studies). Adolescents' physical activity was typically measured using subjective measures (e.g. self-report questionnaires;

71% of adolescent studies). Multiple physical activity measures were used in six children's studies and one adolescent study, respectively. When moderate (MPA) and vigorous (VPA) activity were reported separately and the associations obtained were in different directions, they were treated as separate results and noted accordingly. If the MPA and VPA results were reported in the same direction, the results were combined to form one overall result (i.e. MVPA). For studies that used two or more objective measures of physical activity, a combined result was reported when results from both of these measures were in the same direction.¹⁰ Where differences in the direction of the association occurred between measures, the results were presented separately and noted accordingly. The same approach was adopted when studies examined school recess periods separately (e.g. morning recess and lunchtime) and reported different findings. A variety of statistical techniques were used to evaluate the associations (e.g. correlations, multilevel modeling, ANOVA). A number of studies reported univariate analyses, though some also reported multivariate analyses with adjustments for confounding variables. Where possible, the findings reported are from the fully adjusted models.

The results were coded using the procedure outlined in previous reviews where the association between an identified variable and recess physical activity was determined by the number of findings that support the direction of the association.^{2,10,11} The summary column identifies the number of studies finding positive (+), negative (-) or no (0) associations between recess physical activity and the identified variable. The summary coding column indicates the percentage of findings that demonstrated a significant association between recess physical activity and the identified variable. When 0%-33% of the findings supported an association, the result was classified as no association (0). An indeterminate/inconclusive (?) classification was determined if 34%-59% of the findings supported an association, whilst

a positive (+) or negative (-) association was determined if 60%-100% of the findings supported the direction of the association (Table 1). If a variable was investigated four or more times and supported an association as positive, the result was coded ++, while negative associations were coded as -- and no associations coded as 00.

[INSERT TABLE 1 ABOUT HERE]

Results

Of the 2718 studies identified, 53 papers were included in the review (Figure 1). The majority of studies were cross-sectional (n=42), focused on children (n = 47), and reported MVPA as the outcome variable (n=26). Only one study included children and adolescents and reported the results separately.¹⁵ Eight studies used self-report measures to assess physical activity, with seven and five of these reporting the validity and/or reliability of the measures used, respectively. Due to the limited number of adolescent studies, data from children and adolescents are reported together to obtain the summary coding variables reported below. The majority of studies were conducted in the USA (n=16)¹⁶⁻³¹ or the United Kingdom and Northern Ireland (n=12).³²⁻⁴³ Five studies were conducted in Australia⁴⁴⁻⁴⁸, 4 in Norway^{15,49-51}, 3 in New Zealand⁵²⁻⁵⁴, 2 each in Taiwan^{55,56}, Japan^{57,58} and Portugal^{59,60}, 1 each in Hong Kong⁶¹, Mexico⁶², the Philippines⁶³, Canada⁴⁴, Belgium⁶⁴, Hungary⁶⁵, Spain⁶⁶ and France⁶⁷. Only one study collected data across multiple countries⁴⁴. Of the 53 studies, four studies collected data from morning recess^{17,30,58,66}, seven from lunchtime^{23,32,57,52,53,63,67} and 12 from daily recess (i.e. no time of day reported).^{16,22,24,26-28,46,49-51,61,62} The remaining 30 studies collected data across multiple recess periods (e.g. morning recess and lunchtime).

[INSERT FIGURE 1 ABOUT HERE]

Correlates of school recess physical activity

Table 2 summarizes associations between identified correlates and recess physical activity.

Correlates that have been examined four or more times are reported below.

[INSERT TABLE 2 ABOUT HERE]

Individual variables

Sixteen individual variables were identified from 47 studies, with five of these being investigated four or more times. The most frequently studied variable was sex (38 studies), with boys consistently found to be significantly more active than girls.^{15-18,21,23,25,28,29,31,32,34,35,37-40,42,43,45-48,51,52,54,59,66-66} No associations were found for grade level^{15,18,20,23}, or body mass index/central adiposity^{18,30,39,40,42,54}, while the evidence for differences in physical activity between children with and without special education needs^{21,27,55-58} and age^{17,30,33,34,35,37,39,52,54,55,59,63,67} were inconclusive.

Social variables

Associations between social variables and physical activity were examined in ten studies. Five social variables were identified, with associations between perceived encouragement, socio-economic status, supervision and physical activity investigated four or more times. A positive association was found between physical activity and perceived encouragement from friends, family and schools.⁵³ Higher levels of perceived encouragement were associated with higher physical activity levels.⁵³ There was inconclusive evidence for associations between socio-economic status^{45,61,63} and adult supervision^{25,38,40,46-48} and physical activity.

Physical environmental variables

Twelve physical environmental variables were identified from 16 studies, six of which were studied four or more times. The most frequently studied variables were the availability of facilities in the outdoor school grounds, and fixed equipment and markings within the school grounds. The availability of separate facilities in outdoor physical activity spaces/areas (e.g. ball areas, sports fields, green spaces) has been mainly investigated in adolescents, and no association with physical activity was found.^{15,47,50} However, when overall facility provision was considered (that is, the sum of facilities available), positive associations with physical activity were identified.^{15,32,50,51} Fixed equipment and markings have been investigated in children and adolescents, and the association with physical activity was inconclusive.^{15,32,38,44,46,47,50} In comparison, unfixed equipment (e.g. loose equipment, balls, skipping ropes) has only been investigated in children, with positive associations found.^{25,38,40,46-48} An indeterminate result was obtained for weather/seasonal differences.^{34,36,38,40,46,48} No association was found between available outdoor space and physical activity.^{38,46}

Organizational/policy variables

Eleven variables were identified from 19 studies, two of which had been studied four or more times. An inconclusive association was found between recess duration (which is typically determined by school policy) and physical activity.^{24,37,39,45,46} Differences in physical activity levels between recess periods and physical education have primarily been investigated in children with special education needs, and the findings were inconclusive.^{20,26,28,41,56,61,62}

Discussion

This review provides evidence for factors to include in a social-ecological model of recess physical activity behavior but also highlights areas where evidence is lacking. Forty-four variables were identified across the four levels of the framework, though only 36% had been investigated four or more times. Only three studies examined correlates across all levels of the social-ecological model simultaneously.^{38,46,48} The majority of the variables identified were at the individual and physical environmental levels of the model, though only three consistent associations were found for sex, unfixed equipment and facility provision. Few studies examined social influences, despite recess providing an opportunity for youth to engage in social interactions with their peers relatively free of adult constraints.⁶⁸ Potentially important factors such as composition of social groups (e.g. single sex, mixed sex) have not been investigated, though this may be somewhat explained by the large number of child studies reviewed where the focus has typically been on teacher supervision and management.^{25,38,40,46-48} In addition, further research should investigate correlates of children's and adolescents' recess physical activity for variables that were not frequently investigated but indicated positive associations with physical activity. Identifying recess physical activity correlates may help inform future intervention development and school programming, as although recess interventions have generated positive results, reported changes are small and often focus on short-term changes.^{30,39,64}

Some limitations of included studies warrant attention. The majority of the research in this area has examined small samples, utilized cross-sectional study designs. A meta-analysis of the data may provide more information though such an analysis is difficult given the limited number of studies that report effect sizes and the lack of consistency in the correlates assessed between studies. A range of physical activity measures have been also used, which may influence the associations identified. While the majority of child studies have used

objective measures of physical activity, in particular accelerometry and direct observation, different accelerometer cut-points and observation systems may have influenced the strength of the associations observed. The majority of adolescent studies used self-report measures and while face and content validity of the measures were documented in some of the studies^{15,53,54}, further research using objective measures are needed to determine adolescents' recess physical activity levels.

Arguably the main weakness in the evidence to date is at the organizational/policy level, where only 2 variables have been investigated four or more times and the associations for recess duration and differences between recess periods and physical education were inconclusive. This weakness in the literature base is concerning, especially as there is an increasing trend to reduce the frequency and/or duration of recess.⁶⁹ The way in which recess is defined may explain the lack of consistent findings, as several studies included time spent eating at lunchtime and thus a substantial component of compulsory sedentary behavior.^{37,40} Moreover, some studies reported total daily recess duration (i.e. morning recess and lunchtime combined)^{35,60} whilst others examined the impact of individual recess periods^{18,31}, which may have also influenced the results.

Little research has documented the effect of written recess policies on recess provision and physical activity levels, particularly in countries where daily recess is not mandatory (e.g. the US).⁷⁰ However, mandatory recess periods may be one approach to impact on children's physical activity at the organizational/policy level.⁵¹ Schools are encouraged to have a written school physical activity programming policy that includes recess, which may contribute to stronger policies and programs at the school level.⁷⁰ This may also encourage schools that are contemplating the removal of recess to consider the benefits of retaining recess in the school

day not only on youth social, emotional, physical and cognitive development^{68,71}, but also their classroom behavior and academic performance.⁷¹ Furthermore, adopting written school physical activity programming policies may not only help promote more consistent scheduling of recess⁷⁰, but help to contribute to both children and adolescents' daily physical activity levels as a significant proportion of their recommended daily physical activity can be accumulated during this time.⁷²

Higher perceived encouragement from parents, peers and the school as a whole was associated with higher self-reported physical activity levels during recess periods, particularly in adolescents.⁵³ It was surprising, however, that perceived parental encouragement influenced physical activity levels in the younger adolescents given the lack of involvement parents have during this period of time. This result may be somewhat explained by the measures used, as the adolescents were not specifically asked about parental encouragement for physical activity and sport during school recess.⁵³ Interestingly, parental encouragement may benefit physical activity levels across a range of contexts, even in those where parental involvement is minimal.

Overall facility provision (i.e. sum of facilities available) was positively associated with physical activity.^{15,32,50,51} Interestingly, the provision of outdoor (e.g. sports fields, ball areas)¹⁵ and indoor spaces (e.g. gyms)⁵⁰ during recess periods were not associated with physical activity when they were investigated as individual facilities. These facilities may be associated with physical activity during other non-curricular periods, such as before or after school, though this was not examined in this review. Increasing access to different facilities during recess and/or lunchtime at school may benefit youth physical activity though single spaces may not be effective on their own. This supports research that has demonstrated that providing access to a range of spaces and facilities may stimulate physical activity by

increasing a sense of choice and providing supportive environments that facilitate active behaviors.^{37,73,74} Future research should examine whether increasing access to school facilities during recess periods increases physical activity levels in children and adolescents.

A positive association was found between unfixed equipment and recess physical activity in children. No studies examined this association in adolescents. These results contrast with previous correlates reviews that reported no or inconclusive associations between the availability of toys/equipment and overall physical activity.^{10,11} It is possible that unfixed equipment may affect physical activity participation in specific contexts, rather than overall daily physical activity. In the present study, there was no clear indication as to whether unfixed equipment increased MPA, VPA or MVPA, which may be dependent on the type and number of equipment provided. For example, Zask and colleagues⁴⁸ reported that the ratio of balls to children impacted on VPA, whilst others used a dichotomous variable to identify the presence or absence of equipment and found positive associations with both MPA and VPA.^{25,38,40,47} Further research is needed to determine whether specific types of equipment, or the overall availability of unfixed equipment, are associated with higher levels of physical activity. In elementary school settings, the provision of games equipment has been found to increase children's physical activity levels.⁶⁴ Examining whether the provision of unfixed equipment is a suitable strategy for increasing adolescents' recess physical activity is warranted.

This review found that boys are more physically active than girls during recess, supporting previous reviews of preschool, childhood and adolescence correlates.¹⁰⁻¹² Boys view recess as an opportunity to play competitive games that often dominate the available space.^{75,76} In comparison, girls may view recess as an opportunity to socialize with friends.^{75,77} However,

behaviors during recess have been primarily studied among children, and there are limited data concerning adolescents. There is a need to establish why adolescent males are more active than females during recess, as this will inform strategies targeted at promoting physical activity in this age group. Future research should examine the correlates of boys and girls physical activity separately. Identifying modifiable variables that differ by sex could be critical in the development of activity promotion strategies, and may help to inform evidence-based practice and policies designed to increase physical activity in primary and secondary school children.

Conclusion

In summary, there is currently a dearth of literature concerning correlates of physical activity during recess periods, particularly in adolescents. Despite the paucity of associations identified, it is recommended that schools should increase overall facility provision, provide unfixed equipment, and identify methods to increase social support, particularly by peers, to benefit children and adolescents' physical activity levels during recess.

Acknowledgements

Nicola Ridgers is supported by an Alfred Deakin Postdoctoral Research Fellowship. Jo Salmon is supported by a National Heart Foundation of Australia and sanofi-aventis Career Development Award. Rebecca Stanley is supported by an Australian Postgraduate Award Scholarship and a University of South Australia Top Up Scholarship.

No financial disclosures were reported by the authors of this paper.

References

1. Andersen L, Harro M, Sardinha L, et al. Physical activity and clustered cardiovascular risk in children: A cross-sectional study (The European Youth Heart Study). *Lancet* 2006;368:299-304.
2. Lubans DR, Morgan PJ, Cliff DP, Barnett LM, Okely AD. Fundamental movement skills in children and adolescents: review of associated health benefits. *Sports Med* 2010;40(12):1019-35.
3. Rethon C, Edwards P, Bhui K, Viner RM, Taylor S, Stansfeld SA. Physical activity and depressive symptoms in adolescents: a prospective study. *BMC Medicine*. 2010;8:32.
4. Tobias JH, Steer CD, Mattocks CG, Riddoch C, Ness AR. Habitual levels of physical activity influence bone mass in 11 year-old children from the UK: Findings from a large population-based cohort. *J Bone Miner Res* 2007;22(1):101-9.
5. World Health Organisation Global recommendations on physical activity for health. Geneva, Switzerland: World Health Organisation, 2010.
6. Jago R, Baranowski T. Non-curricular approaches for increasing physical activity in youth: a review. *Prev Med* 2004;39(1):157-63.
7. Robert Wood Johnson Foundation. Recess Rules - Why the undervalued playtime may be America's best investment for healthy kids and healthy schools. Robert Wood Johnson Foundation. 2007. <http://www.rwjf.org/files/research/sports4kidsrecessreport.pdf>.
8. Huberty JL, Siahpush M, Beighle A, Fuhrmeister E, Silva P, Welk GJ. Ready for Recess: A pilot study to increase physical activity in elementary school children. *J Sch Health* 2011;81:251-7.
9. National Association for Sport and Physical Education. Recess for Elementary School Students [Position paper]. Reston, VA, 2006.

10. Hinkley T, Crawford D, Salmon J, Okely AD, Hesketh K. Preschool children and physical activity: A review of correlates. *Am J Prev Med* 2008;34(5):435-41.
11. Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* 2000;32:963-75.
12. van der Horst K, Chin A Paw MJ, Twisk JWR, van Mechelen W. A brief review on correlates of physical activity and sedentariness in youth. *Med Sci Sports Exerc* 2007;39:1241-50.
13. Ridgers ND, Stratton G, Fairclough SJ. Physical activity levels of children during school playtime. *Sports Med* 2006;36(4):359-71.
14. Sallis JF, Owen N. Ecological models of health behaviour. In: Glanz K, Rimer BK, Marcus Lewis F, editors. *Health Behavior and Health Education: Theory, Research and Practice*. 3rd ed. San Francisco: Jossey-Bass; 2002: p. 462-84.
15. Haug E, Torsheim T, Sallis JF, Samdal O. The characteristics of the outdoor school environment associated with physical activity. *Health Educ Res.* 2010;25(2):248-56.
16. Barfield JP, Rowe DA, Michael TJ. Interinstrument consistency of the Yamax Digi-Walker pedometer in elementary school-aged children. *Meas Phys Educ Exerc Sci* 2004;8(2):109-16.
17. Beighle A, Morgan CF, Le Masurier G, Pangrazi RP. Children's physical activity during recess and outside of school. *J Sch Health* 2006;76(10):516-20.
18. Brusseau TA, Kulinna PH, Tudor-Locke C, Ferry M, van der Mars H, Darst PW. Pedometer-determined segmented physical activity patterns of fourth- and fifth-grade children. *J Phys Act Health* 2011;8(2):279-86.
19. Dale D, Corbin CB, Dale KS. Restricting opportunities to be active during school time: Do children compensate by increasing physical activity levels after school? *Res Q Exerc Sport* 2000;71:240-8.

20. Faison-Hodge J, Porretta DL. Physical activity levels of students with mental retardation and students without disabilities. *Adapt Phys Act Q* 2004;21:139-52.
21. Foley JT, Bryan RR, McCubbin JA. Daily physical activity levels of elementary school-aged children with and without mental retardation. *J Dev Phys Disabil* 2008;20(4):365-78.
22. Horvat M, Franklin C. The effects of the environment on physical activity patterns of children with mental retardation. *Res Q Exerc Sport* 2001;72(2):189-95.
23. Joens-Matre RR, Welk GJ, Calabro MA, Russell DW, Nicklay E, Hensley LD. Rural-urban differences in physical activity, physical fitness, and overweight prevalence of children. *J Rural Health* 2008;24(1):49-54.
24. McKenzie TL, Sallis JF, Elder JP, et al. Physical activity levels and prompts in young children at recess: A two-year study of a bi-ethnic sample. *Res Q Exerc Sport* 1997;68(3):195-202.
25. McKenzie TL, Crespo NC, Baquero B, Elder JP. Leisure-time physical activity in elementary schools: Analysis of contextual conditions. *J Sch Health* 2010;80(10):470-7.
26. Pitetti KH, Beets MW, Combs C. Physical activity levels of children with intellectual disabilities during school. *Med Sci Sports Exerc* 2009;41(8):1580-6.
27. Rosser Sandt DD, Frey GC. Comparison of physical activity levels between children with and without autistic spectrum disorders. *Adapt Phys Act Q* 2005;22:146-59.
28. Sarkin JA, McKenzie TL, Sallis JF. Gender differences in physical activity during fifth-grade physical education and recess periods. *J Teach Phys Educ* 1997;17:99-106.
29. Scruggs PW, Beveridge SK, Watson DL. Increasing children's school time physical activity using structured fitness breaks. *Pediatr Exerc Sci* 2003;15:156-69.

30. Stellino MB, Sinclair CD, Partridge JA, King KM. Differences in children's recess physical activity: recess activity of the week intervention. *J Sch Health* 2010;80(9):436-44.
31. Tudor-Locke C, Lee SM, Morgan CF, Beighle A, Pangrazi RP. Children's pedometer-determined physical activity during the segmented school day. *Med Sci Sports Exerc* 2006;38(10):1732-8.
32. Jones NR, Jones A, van Sluijs EMF, Panter J, Harrison F, Griffin SJ. School environments and physical activity: The development and testing of an audit tool. *Health Place*. 2010;16:776-83.
33. McLaughlin E, O'Donoghue PG. Activity profile of primary school children in the playground. *J Hum Movement Studies* 2002;42(2):91-108.
34. Ridgers ND, Stratton G. Physical activity during school recess: The Liverpool Sporting Playgrounds Project. *Pediatr Exerc Sci* 2005;17(3):281-90.
35. Ridgers ND, Stratton G, Fairclough SJ. Assessing physical activity during recess using accelerometry. *Prev Med* 2005;41(1):102-7.
36. Ridgers ND, Stratton G, Clark E, Fairclough SJ, Richardson DJ. Day-to-day and seasonal variability of physical activity during school recess. *Prev Med* 2006;42(5):372-4.
37. Ridgers ND, Stratton G, Fairclough SJ, Twisk JWR. Children's physical activity levels during school recess: a quasi-experimental intervention study. *Int J Behav Nutr Phys Act* 2007;4:19.
38. Ridgers ND, Fairclough SJ, Stratton G. Variables associated with children's physical activity levels during recess: the A-CLASS project. *Int J Behav Nutr Phys Act* 2010;7:74.
39. Ridgers ND, Fairclough SJ, Stratton G. Twelve-month effects of a playground intervention on children's morning and lunchtime recess physical activity levels. *J Phys Act Health* 2010;7(2):167-75.

40. Ridgers ND, Stratton G, McKenzie TL. Reliability and validity of the System for Observing Children's Activity and Relationships during Play (SOCARP). *J Phys Act Health* 2010;7(1):17-25.
41. Sleaf M, Warburton P. Physical activity levels of 5-11 year-old children in England as determined by continuous observation. *Res Q Exerc Sport* 1992;63(3):238-45.
42. Stratton G, Ridgers ND, Fairclough SJ, Richardson DJ. Physical activity levels of normal-weight and overweight girls and boys during primary school recess. *Obesity (Silver Spring)* 2007;15(6):1513-9.
43. Waring M, Warburton P, Coy M. Observation of children's physical activity levels in primary school: Is the school an ideal setting for meeting government activity targets? *Eur Phy Educ Rev* 2007;13(1):25-40.
44. Dymont JE, Bell AC, Lucas AJ. The relationship between school ground design and intensity of physical activity. *Child Geog* 2009;7(3):261-76.
45. Parrish AM, Iverson D, Russell K, Yeatman H. Observing children's playground activity levels at 13 Illawarra primary schools using CAST2. *J Phys Act Health* 2009;6(Suppl 1):S89-96.
46. Parrish AM, Russell K, Yeatman H, Iverson D. What factors influence children's activity? *Br J Sch Nurs* 2009;4(1):6-9.
47. Willenberg LJ, Ashbolt R, Holland D, et al. Increasing school playground physical activity: A mixed methods study combining environmental measures and children's perspectives. *J Sci Med Sport* 2010;13(2):210-6.
48. Zask A, van Beurden E, Barnett L, Brooks LO, Dietrich UC. Active school playgrounds - Myth or reality? Results of the "Move It Groove It" Project. *Prev Med* 2001;33:402-8.

49. Fjortoft I, Kristoffersen B, Sageie J. Children in schoolyards: Tracking movement patterns and physical activity in schoolyards using global positioning system and heart rate monitoring. *Landscape Urban Plan* 2009;93(3-4):210-7.
50. Haug E, Torsheim T, Samdal O. Physical environmental characteristics and individual interests as correlates of physical activity in Norwegian secondary schools: The health behaviour in school-aged children study. *Int J Behav Nutr Phys Act* 2008;5:47.
51. Haug E, Torsheim T, Samdal O. Local school policies increase physical activity in Norwegian secondary schools. *Health Promot Int* 2010;25(1):63-72.
52. Duncan JS, Badland HM, Schofield G. Combining GPS with heart rate monitoring to measure physical activity in children: A feasibility study. *J Sci Med Sport* 2009;12(5):583-5.
53. Hohepa M, Scragg R, Schofield G, Kolt GS, Schaaf D. Social support for youth physical activity: Importance of siblings, parents, friends and school support across a segmented school day. *Int J Behav Nutr Phys Act* 2007;4:54.
54. Hohepa M, Scragg R, Schofield G, Kolt GS, Schaaf D. Self-reported physical activity levels during a segmented school day in a large multiethnic sample of high school students. *J Sci Med Sport* 2009;12(2):284-92.
55. Pan CY. Objectively measured physical activity between children with autism spectrum disorders and children without disabilities during inclusive recess settings in Taiwan. *J Autism Dev Disord* 2008;38(7):1292-301.
56. Pan CY. School time physical activity of students with and without autism spectrum disorders during PE and recess. *Adapt Phys Act Q* 2008;25(4):308-21.
57. Tsujii N, Okada A, Kaku R, et al. Association between activity level and situational factors in children with attention deficit/hyperactivity disorder in elementary school. *Psychiatry Clin Neurosci* 2007;61(2):181-5.

58. Tsujii N, Okada A, Kaku R, Kuriki N, Hanada K, Shirakawa O. Differentiation between attention-deficit/hyperactivity disorder and pervasive developmental disorders with hyperactivity on objective activity levels using actigraphs. *Psychiatry Clin Neurosci* 2009;63(3):336-43.
59. Lopes V, Vasques CMS, Pereira MBFLO, Maia JAR, Malina RM. Physical activity patterns during school recess: A study in children 6 to 10 years old. *Int Electron J Health Educ.* 2006;9:192-201.
60. Mota J, Silva P, Santos MP, Ribeiro JC, Oliveira J, Duarte JA. Physical activity and school recess time: Differences between the sexes and the relationship between children's playground physical activity and habitual physical activity. *J Sports Sci* 2005;23(3):269-75.
61. Sit CH, McKenzie TL, Lian JM, McManus A. Activity levels during physical education and recess in two special schools for children with mild intellectual disabilities. *Adapt Phys Act Q* 2008;25(3):247-59.
62. Jennings-Aburto N, Nava F, Bonvecchio A, et al. Physical activity during the school day in public primary schools in Mexico City. *Salud Publica Mex* 2009;51(2):141-7.
63. Gonzalez-Suarez CB, Grimmer-Somers K. Physical activity pattern of prepubescent Filipino school children during school days. *J Sch Health* 2009;79(7):304-11.
64. Verstraete SJ, Cardon GM, DeClercq DL, De Bourdeaudhuij IM. Increasing children's physical activity levels during recess periods in elementary schools: the effects of providing game equipment. *Eur J Public Health.* 2006;16(4):415-9.
65. Ridgers ND, Toth M, Uvacek M. Physical activity levels of Hungarian children during school recess. *Prev Med* 2009;49(5):410-2.

66. Martinez-Gomez D, Calabro MA, Welk GJ, Marcos A, Veiga OL. Reliability and validity of a school recess physical activity recall in spanish youth. *Pediatr Exerc Sci* 2010;22(2):218-30.
67. Gavarry O, Bernard T, Giacomoni M, Seymat M, Euzet JP, Falgairette G. Continuous heart rate monitoring over 1 week in teenagers aged 11-16 years. *Eur J Appl Physiol* 1998;77:125-32.
68. Pellegrini AD, Bohn CM. The role of recess in children's cognitive performance and school adjustment. *Edu Res* 2005;34(1):13-19.
69. Ramstetter CL, Murray R, Garner AS. The crucial role of recess in schools. *J Sch Health* 2010;80(11):517-26.
70. Lee SM, Burgeson CR, Fulton JE, Spain CG. Physical education and physical activity: results from the School Health Policies and Programs Study 2006. *J Sch Health* 2007;77(8):435-63.
71. Ginsburg KR. The importance of play in promoting health child development and maintaining strong parent-child bonds. *Pediatrics* 2007;119:182-91.
72. Ridgers ND, Timperio A, Crawford D, Salmon J. Five-year changes in school recess and lunchtime and the contribution to children's daily physical activity. *Brit J Sports Med* doi:10.1136/bjism.2011.084921.
73. Sallis JF, Conway TL, Prochaska JJ, McKenzie TL, Marshall SJ, Brown M. The association of school environments with youth physical activity. *Am J Public Health* 2001;91:618-20.
74. Wechsler H, Devereaux AB, Davis M, Collins J. Using the school environment to promote physical activity and healthy eating. *Prev Med* 2000;31:S121-S37.

75. Blatchford P, Baines E, Pellegrini AD. The social context of school playground games: Sex and ethnic difference, and changes over time after entry to junior school. *Br J Dev Psychol* 2003;21:481-505.
76. Boyle DE, Marshall NL, Robeson WW. Gender at play - Fourth-grade girls and boys on the playground. *Am Behav Sci* 2003;46(10):1326-45.
77. Evans J. Children's attitudes to recess and changes taking place in Australian primary schools. *Res Education*. 1996;56:49-61.

Figure Legend:

Figure 1: Flow diagram of search results published between January 1990 and April 2011

Table 1: Rules for classifying variables regarding strength of associations with recess physical activity¹¹

Studies supporting association (%)	Summary code	Explanation of code
0-33	0	No association
34-59	?	Indeterminate/inconclusive association
60-100	+	Positive association
60-100	-	Negative association

When an outcome had been studied four or more times, it was coded as 00, ??, ++, or --

Table 2: Summary of reported outcomes of studies published January 1990 to April 2011

Correlate	Found association with PA during recess	Association (+/-)	Found no association with PA during recess	Summary coding			Summary code	
				n/N for row (%)	Association (-/+)			
	References	Association (+/-)	References				n/N for row (%)	Association (-/+)
Individual variables								
<i>Behavioral</i>								
Day-to-day variability			36				0/1 (0)	0
Physical conflict frequency	40	+					1/1 (100)	+
Sedentary activities (time)	40	-					1/1 (100)	-
Play ball games	46	+					1/1 (100)	+
Sports activities (time)	40	+					1/1 (100)	+
<i>Biological</i>								
Age	30, 33 ^b , 39, 59, 67	-	17, 34, 35, 37, 52, 55 ⁴				5/15 (33)	00
	33 ^a , 54, 55 ^c , 63	+						
Grade level	15	-	20, 23				1/4 (25)	00
	18	+						
BMI (overweight)	18 ^d , 39 ^{b,d} , 54 ^c	-	18 ^c , 30, 39 ^c , 40, 42, 54 ^d				3/10 (30)	00
	54 ⁿ	+						
Body mass	67	-					1/1	+
Maturation	67	-					1/1	+
Ethnicity (Caucasian)	54	-	18				1/2 (50)	?
Fitness			20				0/1 (0)	0
Sex (male)	15, 16, 17, 18, 21 ⁴ , 23, 25, 28, 29 ^{Step} , 31, 32, 34, 35, 37, 38 ^b , 39, 40, 42, 43 ^{a,b,d} , 45, 46, 47 ^b , 48, 51, 52, 54, 59, 64, 65, 66	+	22 ³ , 27, 29 ^{HR} , 30, 35 ^b , 38 ^a , 43 ^{b,c} , 47 ^a , 49, 62, 63, 66				32/43 (74)	++
	60	-						
Special educational needs	21, 55, 56, 58 ^{PDD}	+	20, 27, 57, 58 ^{ADHD}				4/8 (50)	??

(no disability)								
<i>Cognitive</i>								
Interest in physical activity	50, 51	+					2/2 (100)	+
Enjoyment			29				0/1 (0)	0
Social								
Group size (10 or more)	40	+					1/1 (100)	+
Perceived encouragement ¹	53 ^{d,e,i} , 53 ^{d,e,j} , 53 ^{d,g,i} , 53 ^{d,h,i}	+	53 ^{d,g,j} , 53 ^{d,h,j}				4/6 (67)	++
SES (low)	61 ³ , 63 ^d	-	45, 63 ^c				2/4 (50)	??
Supervision ²	25, 46 (observation, presence)	-	38, 40, 47 ^{a,d} , 48				3/8 (38)	??
Supervision ³	47 ^{b,d}	+						
Teacher management			46, 48				0/2 (0)	0
Physical environment								
Aesthetics			32				0/1 (0)	0
Unfixed equipment	25, 38 ^a , 40, 46, 47 ^b , 48 (ball/child ratio) ^b	+	38 ^b , 46 (# of balls), 47 ^a , 48				6/10 (60)	++
<i>Facility Availability and Accessibility</i>								
Outdoor physical activity areas	15 (sledding hill, soccer field ^k), 50 (green space)	+	15 (ball areas, green spaces, soccer field ^l), 47 (sports fields), 50 (ball areas, soccer fields, ski, water, woods)				3/12 (25)	00
Outdoor space	38 (space/child) ^b	+	38 (space/child ^a , total space), 46 (total space)				1/4 (25)	00
Indoor physical activity spaces/areas	50 (gym)	+	50 (ice skating, swimming)				1/3 (33)	0
Fixed equipment/markings	15 (equipment ^k , hopscotch ^k), 44 ^b , 46 (markings, shading), 47 ^a , 50 (equipment, obstacle course)	+	15 (boarding area, fixed equipment ^l , hopscotch ^l , obstacle course), 32 (design), 38 (fixed equipment, markings), 50				8/19 (42)	??

			(boarding area, climbing, fenced court yard)					
Fixed equipment/markings	47 ^b	-						
Overall facility provision	15 ¹ , 32 (sports provision), 50, 51	+	15 ^m , 32 (facility provision)				4/6 (67)	++
Play location (indoor)	19	-	65				1/2 (50)	?
Play surface (grass)	44, 46	+	47				2/3 (67)	+
School location (rural)	23	+					1/1 (100)	+
Seasons/temperature/ weather	38 (temp) ^b , 40 (temp), 46 (raining, temp)	-	34 (season), 36 (season), 38 (temp) ^a , 46 (humidity), 48 (heat stress)				4/9 (44)	??
Organizational/Policy								
Written policy	51	+					1/1 (100)	+
PA program involvement			51				0/1 (0)	0
Organized activities	51	+					1/2 (50)	?
	25	-						
Number on roll	48	-	38				1/2 (50)	?
Recess duration	39 ^{b,c HR} , 45	+	37 ^a , 39 ^{a,c} , 39 ^{a,b,d}				3/7 (43)	??
	24, 39 ^{b,c ACC} , 46	-						
Class time vs. recess	22 ³	-					1/1 (100)	-
Fitness break vs. recess	29	+					1/1 (100)	+
Morning vs. Lunchtime recess	25	+					1/2 (50)	?
	48	-						
PE vs. recess	20, 26 ^{b,3} , 41, 61 ³	-					4/7 (57)	??
	26 ^{a,3} , 28 ^o , 56, 62	+	28 ^p					
Provision of PE (5x/wk - Yes)			51				0/1 (0)	0
Number of daily recess periods			34				0/1 (0)	0

Note: The summary code is an overall summary of the findings for each variable: 0 = No association; ? = Indeterminate/inconclusive association; + = Positive association; - = Negative association.

¹ Perceived encouragement includes that from friends, parents, school and teachers.

² Supervision from adults (e.g. lunchtime assistants) and teachers - no direct management or organization of games/activities during recess/lunch

³ Data collected from special population only

⁴ Data from children without disorders in studies that included special populations

^a Association for MPA (moderate physical activity)

^b Association for VPA (vigorous physical activity)

^c Association for recess

^d Association for lunchtime

^e Association for friend support

f association for overall support

^g Association for parent support

h Association for school support

ⁱ Association for junior high adolescents

^j Association for senior high adolescents

^k Association for adolescent males

^l Association for adolescent females

^m Association for primary school children

ⁿ Association for central adiposity

^o Association for girls

^p Association for boys

HR = heart rate; ACC = accelerometry; Step = Pedometer; ADHD = Attention-deficit/hyperactivity disorder; PDD = Pervasive developmental disorders