

2008

Oxidative coupling of indoles using thallium(III) trifluoroacetate

Paul A. Keller

University of Wollongong, keller@uow.edu.au

N. R. Yepuri

University of Wollongong

M. J. Kelso

University of Wollongong, mkelso@uow.edu.au

M. Mariani

University of Wollongong

B. W. Skelton

University of Western Australia

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/scipapers>



Part of the [Life Sciences Commons](#), [Organic Chemistry Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Keller, Paul A.; Yepuri, N. R.; Kelso, M. J.; Mariani, M.; Skelton, B. W.; and White, A. H.: Oxidative coupling of indoles using thallium(III) trifluoroacetate 2008.
<https://ro.uow.edu.au/scipapers/151>

Oxidative coupling of indoles using thallium(III) trifluoroacetate

Abstract

The oxidative coupling of polysubstituted electron-rich indoles mediated by thallium trifluoroacetate was found to be a facile, clean, and high yielding reaction. Indolic coupling sites were determined by the nature of the substituents present, with dimerisation at the indole 2-position being the dominant outcome. Indoles bearing two potential reaction sites with similar reactivity were additionally found to undergo heterocoupling.

Keywords

biindole, oxidative dimerisation, thallium oxidative coupling, CMMB

Disciplines

Life Sciences | Organic Chemistry | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

This article was originally published as Keller, PA, Yepuri, NR, Kelso, MJ, Mariani, M, Skelton, BW & White, AH, Oxidative coupling of indoles using thallium(III) trifluoroacetate, *Tetrahedron*, 64(33), 2008, 7787-7795. Original journal article available [here](#)

Authors

Paul A. Keller, N. R. Yepuri, M. J. Kelso, M. Mariani, B. W. Skelton, and A. H. White

Oxidative coupling of indoles using thallium(III) trifluoroacetate

Paul A. Keller,^{a*} Nageshwar R. Yepuri,^a Michael J. Kelso,^a Michael Mariani,^a Brian W. Skelton,^b and Allan H. White^b

^aSchool of Chemistry, University of Wollongong, Wollongong, NSW 2522 Australia

^bSchool of Biomedical, Biomolecular and Chemical Sciences, University of Western Australia, WA 6009 Australia

Abstract—The oxidative coupling of polysubstituted electron-rich indoles mediated by thallium trifluoroacetate was found to be a facile, clean, and high yielding reaction. Indolic coupling sites were determined by the nature of the substituents present, with dimerisation at the indole 2-position being the dominant outcome. Indoles bearing two potential reaction sites with similar reactivity were additionally found to undergo heterocoupling.

Introduction

Numerous synthetic methods are available for introducing the biaryl linkage into organic compounds.^{1,2} The popular Suzuki reaction and its related organometallic cousins employ specific functionality to direct the sites of reaction and many examples have of course been reported.³ Also well known are oxidative coupling methods,⁴ where biaryl bonds are formed directly at unsubstituted aryl sites which have been activated either by the aryl units themselves or by ring substituents. The clear advantage of oxidative coupling over organometallic methods, particularly for the formation of biaryl dimers, is that it requires no prior functionalisation of the aromatic reaction site(s).¹

As part of our ongoing investigations into the biological activities of homo- and hetero-dimeric aromatic systems,⁵ we wished to identify simple methods for constructing focused libraries of symmetrical and non-symmetrical biindoles incorporating biaryl linkages at the indole 2 and 7 positions. While there have been many syntheses of 2,2'-biindoles,⁶ the procedures usually involve numerous steps and are impractical for preparing multiple analogues. In the case of 7,7'-biindoles, very few syntheses have been reported but one notable example⁷ describes the oxidative coupling of 4,6-dimethoxy-2,3-diphenylindole in the presence of quinone, chloranil or dichlorodicyanoquinone to yield the homodimeric 7,7'-coupled biindole in 100%, 70% and 60% yields respectively.


Attracted by the simplicity of oxidative coupling methods, we sought to identify a suitable oxidant for preparing our biindole libraries. Thallium (III) trifluoroacetate has previously been used for effecting

oxidative dimerisations of electron-rich naphthyl compounds,⁸ 2-substituted indoles⁹ and indolocarbazoles¹⁰ We now wish to report that thallium(III) trifluoroacetate, in combination with $\text{BF}_3 \cdot \text{Et}_2\text{O}$, is an effective reagent for the oxidative coupling of electron-rich polysubstituted indoles.

Results and Discussion

Synthesis of 7,7'-biindoles

Our preliminary investigations focussed on the 7,7'- dimerisation of a model compound 2,3-dicarbomethoxy-4,6-dimethoxyindole **1**. This indole monomer had its most reactive positions (C2 and C3) blocked, allowing the two methoxy substituents to activate the C7-position to oxidative dimerisation. The use of thallium trifluoroacetate/ $\text{BF}_3 \cdot \text{Et}_2\text{O}$ as oxidant afforded the 7,7'-dimer **2** as the sole product in excellent yield (Scheme 1). This rapid and selective reaction was superior to the previously reported oxidation, which employed ICl and produced **2** in 34% yield along with 31% of the iodinated monomer **3**.⁷




Scheme 1

Synthesis of 2,2'-biindoles

Formation of 2,2'-biindoles by oxidative dimerisation of 3-aryl-4,6-dimethoxyindoles using thallium trifluoroacetate / $\text{BF}_3 \cdot \text{Et}_2\text{O}$ was explored next. These monomers had two potential dimerisation sites (C2 and C7) but initial experiments showed that indole N-protection limited dimerisation at the 7-position (data not shown). The substituted indole monomers **4-7**¹¹ were NH protected using sodium hydride and benzenesulfonyl chloride to give **8-11** after recrystallisation from methanol (Scheme 2). Clean formation of the 2,2'-dimers was achieved by treating an acetonitrile solution of the protected indole monomers with a half-mole equivalent of thallium(III) trifluoroacetate, followed by addition of excess $\text{BF}_3 \cdot \text{Et}_2\text{O}$. After one hour of stirring at room temperature, the dimers were isolated and recrystallised from

methanol /dichloromethane. Compound **11** bearing the strongly electron withdrawing $p\text{-NO}_2\text{C}_6\text{H}_4$ at the indole 3-position, did not couple under these conditions, even after prolonged reaction.



Scheme 2

As a representative example, the structure of **12** was supported by analysis of the mass spectrum (ES), which showed a peak at m/z 845 assigned as the $[\text{M}+1]$ ion. Further support was provided by the ^1H NMR spectra, which showed the disappearance of a peak at 7.31 ppm assigned to H2 in **8**. The presence of only 'one set' of peaks in the ^1H NMR spectrum was consistent with the formation of a symmetrical dimer. Comparison of the ^1H NMR spectrum of **12** with that of the monomer **8** showed a shift in the aromatic peaks assigned to the benzenesulfonyl group from 7.89-7.56 to 7.53-7.30 ppm, and those assigned to the 3-aryl ring from 7.47-6.92 to 7.03-6.65 ppm. The structure of **12** was eventually confirmed by a single crystal X-ray study (Figure 1) which indicated that both phenyl rings are aligned to one side of the molecule, accounting for the downfield shifts of the aromatic signals in the NMR spectra of the dimer **12** relative to the monomer **8**. The 2,2'-dimeric indoles **12-15** can exist as

atropoisomers due to the presence of four *ortho* substituents. Presumably both of these isomers were formed in each of the products as racemic mixtures.





Figure 1. Molecular (ORTEP) projection of **12** down the quasi-two-fold axis.


Based on a mechanism previously proposed for thallium(III) promoted oxidative dimerisations⁸, we postulate that oxidative 2,2'-dimerisation of these indoles proceeds via an initial one-electron transfer from the electron-rich indole substrate to thallium(III) to form an indole radical cation (Scheme 3). Subsequent electrophilic substitution with a second indole substrate, followed by oxidative aromatization, gives the symmetrical 2,2'-dimers.



Scheme 3

Initial attempts to remove the benzenesulfonyl protecting group from the dimers using thionyl chloride at room temperature yielded complex mixtures. Deprotection was eventually achieved, albeit in disappointing yields of 10-15%, by heating the dimers at reflux for four hours in a 20% solution of sodium hydroxide. Dimer **17** was spectroscopically identical to the reported compound that had previously been synthesised in only 1.5% yield.¹²

The poor outcome in the final deprotection step led us to investigate indole 2,2'-dimerisations using alternative protecting groups. A selection of 3-arylsubstituted indoles were N-protected with various aryl or alkyl sulfonates and then subjected to the dimerisation/deprotection sequence. These results are summarised in Scheme 4. All of the protection and dimerisation reactions proceeded smoothly and in good yield with the exception of *N*-4-nitrobenzenesulfonyl-3-(*p*-nitrophenyl)indole **21**, which produced no dimer and quantitatively returned the starting material even after prolonged reaction times. This was not unexpected given the failure of compound **11** to undergo dimerisation. Confirmation of the structure of **28** was provided from a single crystal X-ray study (Figure 2).



R	X	Yield		
		monomer	dimer	deprotected dimer
<i>p</i> -Me C ₆ H ₄ SO ₂	<i>p</i> -BrC ₆ H ₄	70% 19	70% 25	Complex mixture
<i>p</i> -NO ₂ C ₆ H ₄ SO ₂	<i>p</i> -BrC ₆ H ₄	65% 20	70% 26	Complex mixture
<i>p</i> -NO ₂ C ₆ H ₄ SO ₂	<i>p</i> -NO ₂ C ₆ H ₄	65% 21	NR	-
MeSO ₂	Ph	75% 22	70% 27	Complex mixture
MeSO ₂	<i>p</i> -BrC ₆ H ₄	70% 23	75% 28	Complex mixture
MeSO ₂	<i>p</i> -OMeC ₆ H ₄	75% 24	70% 29	25% 16

Scheme 4




Figure 2: Molecular (ORTEP) projection of **28** down the crystallographic two-fold axis.


Different reaction conditions were trialled for deprotection of the dimers using thiophenol, DMF and K_2CO_3 at room temperature. Only **29** gave the deprotected 2,2'-biindole **16** in 25% yield, a slight improvement over the 15% obtained from **12** in NaOH at reflux (Scheme 2). All other deprotection reactions again resulted in complex mixtures.

Oxidative dimerisation with N-acyl protected indoles

The effects of indole N-protection were further explored using acetyl, trifluoroacetyl and Boc groups. The results from the protection, dimerisation and deprotection reactions are summarized in Scheme 5.

Indole acetylation was achieved using tetrabutylammonium hexafluorophosphate with potassium hydroxide and CH_2Cl_2 under phase transfer conditions. Trifluoroacetyl protection of the indoles was carried out using a modified literature protocol.¹³ Oxidative dimerisation was attempted on compounds **30-34** using the conditions described above. In stark contrast to monomers **8** and **24** (which contain 3-*p*-methoxyphenyl indolic substituents and N-phenylsulfonyl and N-methylsulfonyl groups respectively) the analogous N-acetyl and N-trifluoroacetyl-protected indoles **30** and **32** yielded no 2,2'-biindoles. Inexplicably, the corresponding N-Boc derivative **34** produced a complex mixture of products. Furthermore, given that compounds **11** and **21** (which carry 3-*p*-nitrophenyl indolic substituents) failed to undergo dimerisation, it was very surprising to find that the N-acetyl protected 3-(*p*-nitrophenyl)indole **31** dimerised to **35** in 30% yield! What was perhaps even more surprising was the finding that the 3-(*p*-nitrophenyl)indole **33** gave no trace of the 2,2-dimer but instead produced the

2,7',7,7''-terindole **36** in 35% yield (Scheme 5). The remaining mass was identified as the starting indole **33** and its N-deprotected precursor.



R	X	Yield		
		monomer	dimer	trimer
COCH ₃	<i>p</i> -OMePh	65% 30	NR	NR
COCH ₃	<i>p</i> -NO ₂ Ph	57% 31	NR	NR
COCH ₃	<i>p</i> -NO ₂ Ph		2,2'- dimer 30% 35	-
COCF ₃	<i>p</i> -OMePh	60% 32	NR	NR
COCF ₃	<i>p</i> -NO ₂ Ph	60% 33	-	2,7',7,7''- trimer 35% 36
Boc	<i>p</i> -OMePh	99% 34	-	-


Scheme 5

Support for the structure of the trimer **36** was provided by its mass spectrum (ES) which showed a peak at 1179 corresponding to the M+1 ion. Analysis of the ^1H NMR spectrum showed singlet peaks at 6.73, 6.66, and 6.44 ppm corresponding to the H5', H5 and H5'' protons. These three protons showed no *meta*-coupling to their respective H7 protons (usually observed), suggesting the molecule contains 2,7'- and 7,7''-linkages. Additionally, one H2 signal was absent from the ^1H NMR spectrum while there were clearly two other singlets at 7.21 and 6.95 ppm corresponding to the H2' and H2'' protons. The proposed structure was fully supported by gHMBC, gCOSY and NOESY correlations.

The mechanism for the formation of **36** is not clear, although on the basis of current knowledge about TTFA promoted oxidations and dimerisation of indoles,^{8,14} it may be due to the *p*-nitro group and trifluoroacetate group at the N1 position not activating the 2-position to the same level as occurs in the examples that underwent coupling. This would allow reaction at the C7 position to become a significant process. The 2,7'-dimer could then undergo further oxidative coupling between the 7 and 7' positions yielding the 2,7',7,7''-trimer **36**.

2.4. *Thallium(III) trifluoroacetate mediated coupling of 4,6-dimethoxy-3-aryl indoles without N-protection*

The oxidative dimerisation of indoles **4**, **6** and **7** using thallium (III) trifluoroacetate was attempted in the absence of NH protecting groups. After preliminary optimisation reactions, it was found that 3 to 5 equivalents of the Lewis acid were best added at 0 °C before continued stirring at room temperature for three hours. The results are summarised in Scheme 6. Indole **4** gave the 2,2'-dimer **16** in 35% yield, the remainder being starting material and other baseline products. Indole **6** gave 20% of 2,2'-dimer **18** with 50% of the starting material recovered. Interestingly, 4,6-dimethoxy-3-(*p*-nitrophenyl)indole **6** gave only the 2,7'-dimer **38** (20%) with no traces of formation of 2,2'-dimer.



X	Yield	
	2,2'-dimer	2,7'-dimer
<i>p</i> -OMePh	35% 16	-
<i>p</i> -BrPh	20% 18	-
<i>p</i> -NO ₂ Ph	-	15% 37

Scheme 6

For all oxidative coupling reactions attempted without *N*-protecting groups the reactions took longer and returned significant quantities of starting material, suggesting that dimerisation is facilitated by *N*-protection. The results also suggest that the alkyl or arylsulfonyl groups may help in the regioselectivity of the 2,2'-dimerisation provided the 3-aryl substituent is mildly electron-withdrawing or electron-donating.

Conclusions

Oxidative coupling of indoles with thallium(III) trifluoroacetate provides good to excellent yields of 2,2'-dimers. Protecting groups such as benzenesulfonyl, tosyl or methanesulfonyl aid formation of the 2,2'-dimer, provided that the *para* substituent on the 3-aryl ring is mildly deactivating or an electron-donating group. When the 3-aryl ring bears a *para* substituent which is a strong electron-withdrawing group, e.g. NO₂, no reaction is observed unless an *N*-acetyl protecting group is present which leads to

2,7'-dimerisation. Changing the indole N-protecting group to a trifluoroacetate group deactivates C2 and activates C7 such that trimeric-indoles are formed.

Experimental

Melting point determinations were carried out on a Gallenkamp melting point apparatus. Chemical ionization (CI) and electron impact (EI) mass spectra were obtained on a Shimadzu QP-5000 mass spectrometer by a direct insertion technique with an electron beam energy of 70 eV. Electrospray (ESI) mass spectra were obtained on a VG Autospec spectrometer. High-resolution mass spectra (HRMS) were determined on a micromass QToF2 spectrometer using polyethylene glycol or polypropylene glycol as the internal standard. The m/z values are stated with their peak intensity as a percentage in parentheses. Proton and carbon nuclear magnetic resonance (NMR) spectra were obtained as specified on a Varian Mercury 300 MHz or Varian Inova 500 MHz spectrometer. Spectra were recorded in the specified deuterated solvent, and referenced to the residual non-deuterated solvent signal. Chemical shifts (δ) in ppm were measured relative to the internal standard. Analytical thin layer chromatography (TLC) was carried out on Merck silica gel 60 F254 pre-coated aluminium plates with a thickness of 0.2 mm. All column chromatography was performed under 'flash' conditions on Merck silica gel 60 (230-400 mesh). Chromatography solvent mixtures were measured by volume. Sodium hydride was used as a dispersion in oil. Organic solvent extracts were dried with anhydrous magnesium sulfate. All compounds were judged to be of greater than 95% purity based upon ^1H NMR and TLC analysis.

7,7'-Bi(4,6-dimethoxyindolyl)-2,2',3,3'-tetracarboxylate (2)

To a solution of dimethyl 4,6-dimethoxyindole-2,3-dicarboxylate **1** (460 mg, 1.57 mmol) and thallium trifluoroacetate (432 mg, 0.79 mmol) in dry acetonitrile (40 mL) was added $\text{BF}_3 \cdot (\text{CH}_3\text{CH}_2)_2\text{O}$ (785 μL , 0.79 mmol) and the reaction allowed to stir at rt for 40 min. The reaction was then extracted with water (100 mL) and CH_2Cl_2 (4 x 50 mL). The combined organic extracts were dried (MgSO_4) and concentrated and the residue subjected to flash silica gel column chromatography (5:1 CH_2Cl_2 :EtOAc) to yield the dimer **2** (420 mg, 83%) as a pale yellow powder which was spectroscopically identical to that reported.⁷

General procedure (A) for the synthesis of protected indole starting materials:

To a stirred solution of 4,6-dimethoxy-3-phenylindole (1 mol) in CH_2Cl_2 under a nitrogen atmosphere, was added sodium hydride (1 mol, 60% dispersion in oil) and the reaction then stirred for 25 min. The

sulfonyl chloride (1 mol) was then added and the reaction left to stir at 25 °C for an additional 2 h. The solvent was then removed under reduced pressure and the resultant residue was suspended in water (50 mL) and extracted with CH₂Cl₂ (3 x 100 mL). The organic fractions were combined, dried (MgSO₄), concentrated under reduced pressure and the residue recrystallised from methanol yielding the corresponding protected indole.

N-Benzenesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)indole (**8**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-methoxyphenyl)indole **4** (2.00 g, 7.0 mmol), sodium hydride (0.28 g, 7.0 mmol in 60% oil dispersion) and benzenesulfonyl chloride (1.23 g, 7.0 mmol) to give the indole **8** (2.11 g, 73%) as a white solid, mp 131-133 °C. ¹H NMR δ: 7.89 (2H, d, *J* = 7.5 Hz, sulfonyl ArH2'' and 6''), 7.59-7.41 (3H, m, sulfonyl ArH3'', 4'', and 5''), 7.47 (2H, d, *J* = 8.7 Hz, ArH2' and 6'), 7.31 (1H, s, H2), 7.20 (1H, d, *J* = 1.8 Hz, H7), 6.92 (2H, d, *J* = 8.7 Hz, ArH3' and 5'), 6.32 (1H, d, *J* = 1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.84 (3H, s, C6 OCH₃), 3.71 (3H, s, ArC4 OCH₃). ¹³C NMR δ: 159.5 (ArC4'), 159.0 (C6), 155.0 (C4), 138.5 (C7a), 137.5 (sulfonyl ArC1''), 134.0 (sulfonyl ArC4''), 130.8 (sulfonyl ArC2'' and 6''), 129.4, (sulfonyl ArC3'' and 5''), 126.9 (C3), 124.4 (ArC1'), 121.1 (C2), 113.3 (ArC3' and 5'), 110.0 (C3a), 96.0 (C7), 90.0 (C5), 55.5 (C4 OCH₃), 54.5 (C6 OCH₃), 54.3 (ArC4 OCH₃). CI-MS *m/z*: 424 ([M+1]⁺, 10%), 391 (15), 340 (10), 284 (30), 149 (20), 125 (15), 113 (50), 97 (70), 85 (100); ES-HRMS *m/z*: calcd for [M+1]⁺ C₂₃H₂₂NO₅S 424.1219; found 424.1230.

N-Benzenesulfonyl-4,6-dimethoxy-3-phenylindole (**9**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-phenylindole **5** (2.00 g, 7.9 mmol), sodium hydride (0.32 g, 8.0 mmol) and benzenesulfonyl chloride (1.40 g, 7.9 mmol) to give the indole **9** (2.14 g, 69%) as a white solid, mp 125-128 °C. ¹H NMR δ: 7.90-7.54 (5H, m, sulfonyl ArH''), 7.46-7.36 (5H, m, ArH'), 7.36 (1H, s, H2), 7.20 (1H, d, *J* = 3.0 Hz, H7), 6.33 (1H, d, *J* = 3.0 Hz, H5), 3.95 (1H, s, C4 OCH₃), 3.65 (1H, s, C6 OCH₃). ¹³C NMR δ: 159.5 (C6), 154.5 (C4), 139.0 (C7a), 138.0 (ArC4'), 134.0 (sulfonylArC4''), 129.5 (sulfonylArC2'' and 6''), 129.2 (sulfonylArC1''), 129.0 (sulfonylArC3'' and 5''), 128.0 (ArC4'), 127.5 (ArC2' and 6'), 127.0 (ArC3' and 5'), 124.7 (C3) 122.0 (C2), 112.5 (C3a), 95.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.7 (C6 OCH₃). ES-MS *m/z*: 394 ([M+1]⁺, 80%), 310, (25), 254, (100), 187, (50), 141, (20); ES-HRMS *m/z*: calcd for [M+1]⁺ C₂₂H₂₀NO₄S 394.1113; found 394.1108.

N-Benzenesulfonyl-3-(4-bromophenyl)-4,6-dimethoxyindole (10)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (2.50 g, 7.5 mmol), sodium hydride (0.30 g, 7.5 mmol, 60% dispersion in oil) and benzenesulfonyl chloride (1.33 g, 7.5 mmol) to give the indole **10** (2.49 g, 70%) as a white solid, mp 137-139 °C. ¹H NMR δ : 7.96 (2H, d, J = 8.7 Hz, sulfonyl ArH2'' and 6''), 7.56 (2H, m, sulfonyl ArH3'' and 5''), 7.49-7.54 (4H, m, ArH2' 3' 5' and 6'), 7.25-7.48 (1H, m, sulfonyl ArH4''), 7.34 (1H, s, H2), 7.19 (1H, d, J = 1.8 Hz, H7), 6.33 (1H, d, J = 1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.71 (3H, s, C6 OCH₃). ¹³C NMR δ : 160.0 (C6), 154.0 (C4), 138.3 (C7a), 138.0 (ArC4'), 134.0 (sulfonyl ArC1''), 133.0 (sulfonyl ArC4''), 131.3 (ArC2' and 6'), 130.9 (ArC3' and 5'), 129.5 (sulfonyl ArC2'' and 6''), 127.0 (sulfonyl ArC3'' and 5''), 123.7 (ArC1''), 121.7 (C2), 121.4 (C3), 112.9 (C3a), 96.0 (C7), 90.0 (C5), 58.5 (C4 OCH₃), 58.0 (C6 OCH₃). CI-MS m/z : 474 ([M+1]⁺ ⁸¹Br 20%), 332 (15), 173 (30), 159 (75), 143 (100); EI-HRMS m/z : calcd for C₂₂H₁₈NO₄S⁷⁹Br 471.0139; found 471.0126.

N-Benzenesulfonyl-4,6-dimethoxy-3-(4-nitrophenyl)indole (11)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-nitrophenyl)indole **7** (2.00 g, 6.7 mmol), sodium hydride (0.27 g 6.7 mmol) and benzenesulfonyl chloride (1.18 g, 6.7 mmol) to give the the indole **10** (1.78 g, 61%) as a white solid, mp 150-152 °C. ¹H NMR δ : 8.22 (2H, d, J = 9.0 Hz, ArH2' and 6'), 7.93 (2H, d, J = 9.0 Hz, ArH3' and 5'), 7.69 (2H, d, J = 8.7 Hz, sulfonyl ArH2'' and 6''), 7.59-7.45 (2H, m, sulfonyl ArH3'', 4'', and 5''), 7.19 (1H, d, J = 2.1 Hz, H7), 6.36 (1H, d, J = 2.1 Hz, H5), 3.90 (3H, s, C4 OCH₃), 3.79 (3H, s, C6 OCH₃); ¹³C NMR δ : 159.8 (C6), 154.6 (C4), 146.9 (sulfonylArC1''), 141.1 (sulfonylArC1''), 138.1 (C7a) 137.7 (ArC4), 134.4 (sulfonylArC4''), 130.3 (ArC2' and 6'), 129.6, (sulfonylArC3' and 5'), 127.1 (ArC3' and 5'), 123.1 (sulfonylArC2'' and 6''), 122.7 (C2), 112.3 (C3a), 95.6 (C7), 89.9 (C5), 56.6 (C4 OCH₃), 55.4 (C6 OCH₃). ES-MS m/z 439 ([M+1]⁺, 100%); EI-HRMS m/z : calcd for C₂₂H₁₈N₂O₆S 438.0885; found 438.0907.

3-(4-Bromophenyl)-4,6-dimethoxy-N-(4'-methylphenylsulfonyl)indole (19)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (2.00 g, 6.0 mmol), sodium hydride (0.24 g, 6.0 mmol) and *p*-toluenesulfonyl chloride (0.19 g, 6.0 mmol) to give the indole **19** (2.20 g, 76%) as white solid, mp 138-140 °C. ¹H NMR δ : 7.78 (2H, d, J = 7.8 Hz, ArH2' and H6'), 7.49 (2H, d, J = 9.0 Hz, tosylArH3'' and 5''), 7.39 (2H, d, J = 9.0 Hz, tosylArH2'' and 6''), 7.34 (1H, s, H2), 7.25 (2H, d, J = 7.8 Hz, Ar3H' and 5'), 7.18 (1H, d, J = 1.8 Hz, H7), 6.32 (1H, d, J = 1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.70 (3H, s, C6 OCH₃), 2.35 (3H, s, CH₃). ¹³C NMR δ : 159.4 (C6), 154.6

(C4), 145.2 (tosylC1''), 137.5 (C7a), 135.1 (ArC4'), 133.0 (ArC4'), 131.3 (ArC2' and 6'), 130.9 (ArC3' and 5'), 130.1 (tosyl ArC2'' and 6''), 127.0 (tosyl ArC3'' and 5''), 125.0 (tosyl ArC4''), 123.4 (C3), 121.3 (ArC1'), 121.7 (C2), 112.8 (C3a), 95.4 (C7), 90.0 (C5), 56.1 (C4 OCH₃), 55.5, (C-6 OCH₃), 22.0 CH₃. EI-HRMS *m/z*: calcd for C₂₃H₂₀NO₄S⁷⁹Br 485.0296; found 485.0293.

3-(4-Bromophenyl)-4,6-dimethoxy-N-(p-nitrobenzenesulfonyl)indole (20)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (2.00 g, 6.0 mmol), sodium hydride (0.29 g, 6.0 mmol, 50% dispersion in oil) and *p*-nitrobenzenesulfonyl chloride (1.33 g, 6.0 mmol) to give the indole **20** (2.25 g, 73%) as yellow solid, mp 220-222 °C. ¹H NMR δ: 8.29 (2H, d, *J* = 8.7 Hz, Ar2H' and 6'), 8.05 2H, d, *J* = 8.7 Hz, ArH3' and 5'), 7.49 (2H, d, *J* = 6.6 Hz, sulfonylArH3'' and 5''), 7.37 (2H, d, *J* = 6.6 Hz, sulfonylAr2H'' and 6''), 7.3 (1H, s, H2), 7.17 (1H, d, *J* = 1.8 Hz, H7), 6.36 (1H, d, *J* = 1.8 Hz, H5), 3.89 (3H, s, C4 OCH₃), 3.71 (3H, s, C6 OCH₃). ¹³C NMR δ: 160.0 (C6), 155.0 (C4), 151.5 (sulfonylArC1'') 143.5 (ArC4'), 138.0 (C7a), 133.0 (ArC1'), 131.3 (ArC2' and 6'), 131.0 (ArC3' and 5'), 128.2 (sulfonylArC2'' and 6''), 125.0, (sulfonylArC4''), 124.8 (sulfonylArC3'' and 5''), 122.0 (C3), 121.1 (C2), 113.4 (C3a), 95.9 (C7), 90.1 (C5), 56.1 (C4 OCH₃), 55.4 (C6 OCH₃). CI-MS *m/z*: 519 ([M+1]⁺, ⁸¹Br, 100%), 455 (10), 332 (50), 236 (10); EI-HRMS *m/z*: calcd for C₂₂H₁₇⁷⁹BrN₂O₆S 515.9990; found 515.9982.

4,6-Dimethoxy-N-(p-nitrobenzenesulfonyl)-3-(4-nitrophenyl)indole (21)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4'-nitrophenyl)indole **7** (2.00 g, 6.7 mmol) sodium hydride (0.32 g, 6.7 mmol) and *p*-nitrobenzenesulfonyl chloride (1.48 g, 6.7 mmol) to give the indole **21** (2.17 g, 67%) as a pale yellow solid, mp 190-193 °C. ¹H NMR δ: 8.34 (2H, d, *J* = 9.0 Hz, ArH2 and 6), 8.24 (2H, d, *J* = 8.7 Hz, sulfonylArH3 and 5), 8.10 (2H, d, *J* = 9.0 Hz, ArH3 and 5), 7.68 (2H, d, *J* = 8.7 Hz, sulfonylArH2 and 6), 7.26 (1H, s, H2), 7.18 (1H, d, *J* = 1.8 Hz, H7), 6.39 (1H, d, *J* = 1.8 Hz, H5), 3.91 (3H, s, C4 OCH₃), 3.73 (3H, s, C6 OCH₃). ¹³C NMR δ: 160.1 (C6), 156.0 (C4), 146.1 (ArC4), 144.0 (ArC1), 141.0 (C7a), 148.0 (sulfonylArC4), 131.0 (ArC2 and 6), 128.1 (sulfonylArC2 and 6), 125.5 (ArC3 and 5), 125.0 (sulfonylArC1), 123.0 (C2), 122.0 (ArC2 and 6), 121.0 (C3), 113.0 (C3a), 96.0 (C7), 90.0 (C5), 58.0 (C4 OCH₃), 56.0 (C6 OCH₃). CI-MS *m/z* 484.0, ([M+1]⁺, 100%), 420 (10), 299 (30); EI-HRMS *m/z*: calcd for C₂₂H₁₇N₃O₈S 483.0736; found 483.0746.

N-Methanesulfonyl-4,6-dimethoxy-3-phenylindole (22)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-phenylindole **5** (2.00 g, 7.9 mmol) sodium hydride (0.32 g, 8.0 mmol) and methanesulfonyl chloride (0.90 g, 7.9 mmol) to give the indole **23** (1.96 g, 75%) as white solid, mp 140-142 °C. ¹H NMR δ : 7.57 (2H, d, J = 8.4 Hz, ArH2' and 6'), 7.38-7.34 (3H, m, ArH3', 4', and 5'), 7.23 (1H, s, H2), 7.10 (1H, d, J = 2.1 Hz, H5), 6.41 (1H, d, J = 2.1 Hz, H7), 3.89 (3H, s, C4 OCH₃), 3.76, (3H, s, C6 OCH₃), 3.11 (3H, s, CH₃). ¹³C NMR δ : 160.0 (C6), 155.0 (C4), 138.0 (C7a), 134.0 (ArC1') 130.0 (ArC2' and 6'), 128.0 (ArC3' and 5'), 127.6 (ArC4'), 125.0 (C3), 122.0 (C2), 113.0 (C3a), 96.0, (C7), 90.0 (C5), 56.0, (C4 OCH₃), 55.6 (C6 OCH₃), 40.0, CH₃. CI-MS m/z : 332 ([M+1]⁺, 45%), 270 (15), 132 (15), 103 (20), 71 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₁₇H₁₈NO₄S 332.0957 found; 332.0959.

3-(4-Bromophenyl)-N-methanesulfonyl-4,6-dimethoxyindole (23)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4'-bromophenyl)indole **6** (2.00 g, 6.0 mmol), sodium hydride (0.24 g, 6.0 mmol) and methanesulfonyl chloride (0.69 g, 6.0 mmol) to give the the indole **25** (1.72 g, 70%) as a white solid, mp 147-150 °C. ¹H NMR δ : 7.51 (2H, d, J = 9.0 Hz, ArH2' and 4'), 7.43 (2H, d, J = 9.0 ArH3' and 5'), 7.22 (1H, s, H2), 7.02, (1H, d, J = 2.1 Hz, H7), 6.40 (1H, d, J = 2.1 Hz, H5), 3.89 (3H, s, C4 OCH₃), 3.77, (3H, s, C6 OCH₃), 3.13, (s, Me). ¹³C NMR δ : 160.0 (C6), 156.5 (C4), 138.0 (C7a), 134.0 (ArC4') 130.5 (ArC2' and 6'), 130.0 (ArC3' and 5'), 124.0 (ArC1'), 123.0 (C3), 122.0 (C2), 112.0 (C3a), 96.0 (C7), 90.0 (C5), 55.0 (C4 OCH₃), 54.5 (C6 OCH₃), 42.0 CH₃. CI-MS m/z : 412 ([M+1]⁺, ⁸¹Br, 20%), 332 (15), 254(10), 113 (25), 97(30), 81 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₁₇H₁₇NO₄S⁷⁹Br 410.0062; found 410.0061.

N-Methanesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)indole (24)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4'-methoxyphenyl)indole **4** (2.00 g, 7.0 mmol), sodium hydride (0.28 g, 7.0 mmol) and methane sulfonyl chloride (0.81 g, 7.1 mmol) to give the indole **27** (1.99 g, 79%) as white solid, mp 145-148 °C. ¹H NMR δ : 7.50 (2H, d, J = 9.0 Hz, ArH2' and 6'), 7.17 (1H, s, H2), 7.10 (1H, d, J = 2.1 Hz, H5), 6.93 (2H, d, J = 9.0 Hz, ArH3' and 5'), 6.40 (1H, d, J = 2.1 Hz, H7), 3.88 (3H, s, C4 OCH₃), 3.85 (3H, s, C6 OCH₃), 3.77, (3H, s, ArC4 OCH₃), 3.09 (3H, s, CH₃). ¹³C NMR δ : 159.9 (ArC4'), 158.5 (C6), 155.0 (C4), 138.0 (C7a), 132.0 (ArC2' and 6') 126.0 (ArC1') 124.0 (C3), 121.0 (C2), 114.0 (ArC3' and 5'), 110.0 (C3a) 96.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.6 (C6 OCH₃) 55.7, (ArC4 OCH₃), 40.0, (CH₃). CI-MS m/z : 362 ([M+1]⁺, 90%), 340 (15),

284 (30), 149 (15), 113 (20), 97 (25), 81 (100); EI-HRMS m/z : calcd for $C_{18}H_{19}NO_5S$ 361.0983; found 361.0989.

N-Acetyl-4,6-dimethoxy-3-(4-methoxyphenyl)indole (**30**)

To a solution of 4,6-dimethoxy-3-(4-methoxyphenyl)indole **4** (0.50 g, 1.80 mmol), acetyl chloride (0.14 g, 1.80 mmol), and potassium hydroxide (0.10 g, 1.80 mmol) in CH_2Cl_2 (15 mL), was added a catalytic amount of the phase transfer catalyst tetrabutylammonium hexafluorophosphate (20 mol%) in CH_2Cl_2 (25 mL) under a nitrogen atmosphere. The reaction mixture was stirred for 12 h before water (30 mL) was added and the organic layer separated. The solvent was then removed under reduced pressure, and the oily liquid subjected to flash chromatography (ethyl acetate:hexane 2:8) to yield the indole **31** (0.37 g, 65%) as a white solid, mp 145-158 °C. 1H NMR δ : 7.77 (1H, d, J = 1.8 Hz, H7), 7.51 (2H, d, J = 9.0 Hz, Ar2' and 6'), 7.10 (1H, s, H2), 6.94, (2H, d, J = 9.0 Hz, Ar3' and 5'), 6.40 (1H, d, J = 1.8 Hz, H5), (3.89, 3.85, and 3.75, 3 x OCH_3), 2.61 (3H, s, $COCH_3$). ^{13}C NMR δ : 169.1 (C=O), 159.8 (ArC4'), 159.0 (C6), 154.4 (C4), 138.4 (C7a), 130.7 (ArC2' and 6'), 127.0 (ArC1'), 124.2 (C3), 120.4 (C2), 113.3 (ArC3' and 5'), 112.8 (C3a), 95.9 (C7), 93.1 (C5), (55.9, 55.5, and 55.4 3 x OCH_3), 24.4 ($COCH_3$). CI-MS m/z : 326 ($[M+1]^+$, 100%), 283 (20), 268 (10). EI-HRMS m/z : calcd for $C_{19}H_{19}NO_4$ 325.1314; found 325.1398.

N-Acetyl-4,6-dimethoxy-3-(4'-nitrophenyl)indole (**31**)

To a solution of 4,6-dimethoxy-3-(4-nitrophenyl)indole **7** (0.50 g, 1.70 mmol), acetyl chloride (0.13 g, 1.70 mmol), and potassium hydroxide (0.10 g, 1.8 mmol) in CH_2Cl_2 (15 mL) was added a catalytic amount of phase transfer catalyst tetrabutylammonium hexafluorophosphate (20 mol%) in CH_2Cl_2 (25 mL) under a nitrogen atmosphere. The reaction mixture was stirred for 12 h. To the reaction mixture water was added, and the organic layer separated. The solvent was removed under reduced pressure, and the oily liquid then purified by flash chromatography (ethyl acetate:hexane, 2:8) to yield the indole **32** (0.32 g, 57%) as a white solid, mp 180-182 °C. 1H NMR δ : 8.27 (2H, d, J = 9.0 Hz, ArH2' and 6'), 7.87 (2H, d, J = 9.0 Hz, ArH3' and 5'), (1H, s, H2), 7.70 (1H, d, J = 1.8 Hz, H7), 6.53 (1H, d, J = 1.8 Hz, H5), 3.84 (3H, s, C4 OCH_3), 3.79 (3H, s, C6 OCH_3), 2.68 (3H, s, acetyl CH_3). ^{13}C NMR δ : 160.0 (C6), 154.0 (C4), 149.0 (C=O), 142.0 (ArC4'), 139.0 (C7a), 131.0 (ArC2' and 6'), 123.0 (ArC3' and 5'), 122.0 (C2), 121.8 (C3), 112.0 (C3a), 97.0 (C7), 93.0 (C5), 56.0 (C4 OCH_3), 55.0 (C6 OCH_3), 26.0 ($COCH_3$). CI-MS m/z : 341 ($[M+1]^+$, 100%); EI-HRMS m/z : calcd for $C_{18}H_{16}N_2O_5$ 340.1059; found 340.1072.

N-tert-Butylcarbamate-4,6-dimethoxy-3-(4-methoxyphenyl)indole (34)

To a solution of 4,6-dimethoxy-3-(4'-methoxyphenyl)indole **4** (0.50 g, 1.80 mmol) and di-*tert*-butyldicarbonate (0.37 g, 1.80 mmol) in CH₂Cl₂ (25 mL) was added a catalytic amount of DMAP and the reaction stirred under a nitrogen atmosphere for thirty min. To the reaction mixture was added water, the organic layer separated, the solvent removed under reduced pressure, and the solid then recrystallised from methanol to yield the indole **35** (0.68 g, 99%), as a white solid, mp 151-153 °C. ¹H NMR δ: 8.22 (1H, d, *J* = 1.8 Hz, H7), 7.51 (2H, d, *J* = 9.0 Hz, ArH2' and 6'), 6.92 (2H, d, *J* = 9.0 Hz, ArH3' and 5'), 6.47 (1H, d, *J* = 1.8 Hz, H5), 6.35 (d, *J* = 2.1 Hz), 3.88 (3H, s, ArC4 OCH₃), 3.83 (3H, s, C4 OCH₃), 3.74 (3H, s, C6 OCH₃). ¹³C NMR δ: 159.9 (C=O), 159.8 (C6), 154.3 (C4), 150.0 (C7a), 130.8 (ArC2' and 6'), 127.4 (ArC1'), 121.1 (C2), 113.2 (ArC3' and 5'), 106.8 (C3a), 95.0 (C7), 91.7 (C5), 84.0 (CO₂C(CH₃)), (55.8, 55.4, and 55.4, 3 x OCH₃), 28.4, (3 x C, *t*-BOC). CI-MS *m/z*: 383 ([M⁺]⁺, 90%), 370 (15), 327 (100), 283 (30), 268 (10); ES-HRMS *m/z*: calcd for [M+1]⁺ C₂₂H₂₆NO₅ 384.1811; found 384.1807.

General procedure (B) for the synthesis of bisindoles: oxidative dimerisation:

Solid thallium(III) trifluoroacetate (0.53 mmol) was mixed with the solid protected indole (1.06 mmol) under a nitrogen atmosphere. Dry acetonitrile (7 mL) was added and the resultant mixture cooled to 0 °C. Boron trifluoride diethyletherate (8.0 mmol) was then added slowly into the reaction mixture, and allowed to stir for 1h. The reaction mixture was poured into water and extracted with CH₂Cl₂ (5 x 50 mL). The combined organic layers were concentrated to give a dark residue.

2,2'-Bi-[N-benzenesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)]indole (12)

This was prepared by general procedure (B) using *N*-benzenesulfonyl-3-(4-methoxyphenyl)-4,6-dimethoxyindole **8** (0.50 g, 1.18 mmol), thallium(III) trifluoroacetate (0.32 g, 0.55 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol) The residue was recrystallised from CH₂Cl₂/ methanol to give the indole **12** (0.35 g, 70%) as a white solid, mp 242-246 °C. ¹H NMR δ: 7.53 (2H, d, *J* = 6.0 Hz, sulfonylArH2'' and 6''), 7.46 (1H, t, sulfonylArH4''), 7.30 (2H, t, sulfonylArH3'' and 5''), 7.03 (2H, d, *J* = 12.0 Hz, ArH2' and 6'), 7.02 (1H, d, *J* = 1.8 Hz, H7), 6.65 (2H, d, *J* = 12.0 Hz, ArH3' and 5'), 6.25 (1H, d, *J* = 1.8 Hz, H5), 3.76 (3H, s, C4 OCH₃), 3.75 (3H, s, C6 OCH₃), 3.56 (1H, s, ArC4 OCH₃). ¹³C NMR δ: 159.3 (ArC4'), 159.0 (C6), 154.5 (C4), 139.9 (C7a), 139.0 (sulfonylArC1''), 133.9 (ArC4), 131.8 (sulfonylArC2'' and 6''), 129.5 (sulfonylArC3'' and 5''), 127.0 (C2), 126.7 (ArC2' and 6'), 126.3 (C3), 126.5 (ArC1'), 124.0 (C3a), 122.0 (C2), 112.5 (C3a), 112.0 (ArC3' and 5'), 94.5 (C5), 92.0 (C7), 56.0

(C4 OCH₃), 55.0 (C6 OCH₃). ES-MS m/z : 845 ([M+1]⁺, 35%), 797.2 (10), 704.2 (15), 625 (5), 552.3 (15), 475.4 (10), 411.2 (15), 332.3 (50%), 304.3 (100), 272.3 (50), 244.2 (80), 214.0 (40), 157.9 (75), 141.8 (45), 105.9 (90); ES-HRMS m/z : calcd for [M+1]⁺ C₄₆H₄₁N₂O₁₀S₂ 845.2203; found 845.2209.

2,2'-Bi-(N-benzenesulfonyl-4,6-dimethoxy-3-phenyl)indole (13)

This was prepared by general procedure (B) using *N*-benzenesulfonyl-3-phenyl-4,6-dimethoxyindole **9** (0.50 g, 1.3 mmol), thallium(III) trifluoroacetate (0.35 g, 0.65 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **13** (0.34 g, 69% yield) as a white solid, mp 224-226 °C. ¹H NMR δ: 7.57-7.01 (10H, m, sulfonylArH2'', 3'', 4'' and 6'', and ArH2', 3', 4', 5', and 6'), 7.00 (1H, d, *J* = 1.8 Hz, H7), 6.22 (1H, d, *J* = 1.8 Hz, H5), 3.76 (3H, s, C4 OCH₃), 3.53 (3H, s, C6 OCH₃). ¹³C NMR δ: 159.3 (C6), 154.9 (C4), 139.0 (C7a), 139.5 (sulfonylArC1''), 133.8 (C3), 133.4 (sulfonylArC4''), 130.7 (sulfonylArC2'' and 6''), 129.1 (sulfonylArC3'' and 5''), 129.0 (ArC4'), 128.5 (ArC4'), 128.0 (C2), 127.5 (ArC2' and 6'), 127.2 (ArC3' and 5'), 126.9 (ArC1'), 122.3 (C3a), 95.5 (C7), 90.8 (C5), 55.9 (C4 OCH₃), 55.4 (C6 OCH₃). ES-MS m/z : 785.0 ([M]⁺, 30%), 746.2 (20), 649.4 (25), 552.2 (20), 413, 304, 272, 244 (100%); ES-HRMS m/z : calcd for [M+1]⁺ C₄₄H₃₇N₂O₈S₂ 785.1991; found 785.1978.

2,2'-Bi-[N-benzenesulfonyl-3-(4-bromophenyl)-4,6-dimethoxy]indole (14)

This was prepared using general procedure (B) using thallium(III) trifluoroacetate (0.29 g, 0.53 mmol), *N*-benzenesulfonyl-3-(4-bromophenyl)-4,6-dimethoxyindole **10** (0.50 g, 1.06 mmol) and boron trifluoride diethyletherate (1.12, 7.90 mmol). The residue was recrystallised from CH₂Cl₂/methanol to give the indole **14** (0.37 g, 74%) as a white solid, mp 263-265 °C. ¹H NMR δ: 7.90 (2H, d, *J* = 9.0 Hz, sulfonylArH2'' and 6''), 7.55-7.34 (7H, m, sulfonylArH3'' and 4'', 5'' and ArH2', 3', 5', and 6'), 7.19 (1H, d, *J* = 1.8 Hz, H7), 6.34 (1H, d, *J* = 1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.72 (3H, s, C6 OCH₃). ¹³C NMR δ: 159.3 (C6), 154.5 (C4), 137.8 (C7a), 137.4 (ArC4'), 134.1 (sulfonylArC4'), 132.8 (C2), 131.2 (sulfonylArC2'' and 6''), 130.7 (sulfonylArC3'' and 5''), 129.9 (ArC1'), 129.4 (ArC1'), 129.2 (ArC2' and 6'), 126.8 (ArC3' and 5'), 123.5 (C3), 113.0 (C3a), 95.3 (C7), 89.9 (C5), 56.0 (C4 OCH₃), 55.3 (C6 OCH₃). ES-MS m/z : 942.7 ([M+1]⁺, ⁸¹Br, 30%), 761.3 (10), 488.0 (20), 392.2 (30), 338.4 (30), 316.3 (35), 314.2 (70), 288.3 (60), 270.2 (50), 244.2 (45), 174.8 (30), 166.9 (40), 144.9 (70), 125.8 (80), 103.9 (100); ES-HRMS m/z : calc for [M+1]⁺ C₄₄H₃₅N₂O₈⁷⁹Br₂S₂ 941.0202; found 941.0254.

2,2'-Bi-[4,6-dimethoxy-3-(4-methoxyphenyl)]indole (16)

A solution of 2,2'-bi[*N*-benzenesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)]indole **12** (0.30 g, 0.3 mmol) in dioxane (20 mL) with 20% aqueous sodium hydroxide (5 mL) was heated at reflux for 4 h. The reaction mixture was poured onto cold water (60 mL) and partitioned with CH₂Cl₂ (3 x 50 mL). The solvent was evaporated under vacuum and the residue recrystallised from CH₂Cl₂/methanol to give the indole **16** (0.03 g, 15%) as a pale yellow solid, mp 197-200 °C. ¹H NMR δ: 11.2 (1H, s, NH), 6.93 (2H, d, *J* = 9.0 Hz, ArH2' and 6'), 6.59 (2H, d, *J* = 9.0 Hz, ArH3' and 5'), 6.41 (1H, d, *J* = 1.8 Hz, H5), 6.10 (1H, d, *J* = 1.8 Hz, H7), 3.74 (3H, s, C4 OCH₃), 3.66 (3H, s, C6 OCH₃), 3.60 (3H, s, ArC4' OCH₃). ¹³C NMR (DMSO *d*₆) δ: 158.3 (ArC4'), 158.2 (C6'), 154.3 (C4'), 138.0 (C7a), 132.0 (ArC2' and 6'), 128.5 (ArC1'), 124.6 (C2), 116.3 (C3), 112.2 (ArC3' and 5'), 112.0 (C3a), 92.0 (C5), 87.2 (C7), 55.9 (C4 OCH₃), 55.6 (C6 OCH₃), 55.4 (ArC4 OCH₃); CI-MS *m/z*: 564([M+1]⁺, 60%), 338 (100), 285 (10), 197 (5), 153 (10), 137 (25), 121 (30), 97 (30); ES-HRMS *m/z*: calcd [M+1]⁺ C₃₄H₃₃N₂O₆ 564.2260; found 564.2259.

2,2'-Bi-(4,6-dimethoxy-3-phenyl)indole (17)

A solution of 2,2'-bi(*N*-benzenesulfonyl-4,6-dimethoxy-3-phenyl)indole **13** (0.30 g, 0.4 mmol) in dioxane (20 mL) with 20% aqueous sodium hydroxide (5 mL) was heated at reflux for 4 h. The reaction mixture was poured into cold water and partitioned with CH₂Cl₂ (3 x 50 mL). The solvent was evaporated and the residue recrystallised from CH₂Cl₂/methanol yielding the indole **17** (0.03 g, 15%) as a pale yellow solid mp 174-178 °C. ¹H NMR δ: 10.00 (1H, s, NH), 7.21-7.18 (2H, m, ArH2' and 6'), 7.09-7.06 (3H, m, ArH3', 4', and 5'), 6.49 (1H, d, *J* = 3.0 Hz, H7), 6.18 (1H, d, *J* = 3.0 Hz, H5), 3.77 (3H, s, C4 OCH₃), 3.66 (3H, s, C6 OCH₃); ¹³C NMR δ: 158.0 (C6), 155.0 (C4), 138.0 (C7a), 136.0 (C3), 131.0 (ArC4'), 126.7 (ArC2' and 6'), 125.2 (ArC3' and 5'), 125.0 (ArC1'), 118.0 (C2), 112.0 (C3a), 92.0 (C7), 86.2 (C5), 55.0 (C4 OCH₃), 54.5 (C6 OCH₃). ES-MS *m/z*: 504.2 ([M+1]⁺, 60%), 338.4 (25), 144.9 (20), 105.9 (45), 103.9 (100), 71.0 (25); ES-HRMS *m/z*: calcd for [M+1]⁺ C₃₂H₂₉N₂O₄ 504.2049; found 504.2054.

2',2-Bi-[3-(4-bromophenyl)-4,6-dimethoxy]indole (18)

This was prepared by general procedure (B) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (0.20 g, 0.6 mmol), thallium(III) trifluoroacetate (0.16 g, 0.3 mmol) and boron trifluoride diethyletherate (0.43 g, 3.0 mmol). The residue was subjected to flash silica gel column chromatography (50:50 hexane:ethyl acetate) yielding the indole **18** (0.04 g, 20%) as a pale yellow solid, mp 195-197 °C. ¹H NMR δ: 7.85

(2H, d, $J = 7.5$ Hz, ArH3' and 5'), 7.21 (2H, d, $J = 7.5$ Hz, ArH2' and 6'), 7.28 (1H, d, $J = 1.8$ Hz, H7), 6.32 (1H, s, H5), 3.84 (3H, s, C4 OCH₃), 3.49 (3H, s, C6 OCH₃). ¹³C NMR (CDCl₃) δ : 153.5 (C-6 and C-4), 140.0 (C-7a), 133.0 (C-2), 132.8 (C-3), 130.4 (ArC-4'), 128.0 (ArC2' and 6'), 127.8 (ArC2' and 5'), 126.0 (C-3a), 121.6 (ArC1'), 94.3 (C-7), 92.0 (C-5), 57.8 (C-4-OCH₃), 56.0 (C-6-OCH₃). ES-MS m/z : 663 ([M+1]⁺, ⁸⁰Br 60%), 607 (10), 468.5 (15), 332 (40), 316 (50), 304 (50), 288 (70) 244 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₃₂H₂₇N₂O₄⁷⁹Br₂ 661.0338; found 661.0331.

2,2'-Bi-[3-(4-bromophenyl)-(N-p-methylbenzenesulfonyl)-4,6-dimethoxy-]indole (**25**)

This was prepared by general procedure (**B**) using *N*-(4-methylphenylsulfonyl)-3-(4-bromophenyl)-4,6-dimethoxyindole **19** (0.50 g, 1.0 mmol), thallium(III) trifluoroacetate (0.28 g 0.50 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **25** (0.36 g 73%) as a yellow solid, mp 219 °C. ¹H NMR δ : 7.45 (2H, d, $J = 8.4$ Hz, sulfonylArH3'' and 5''), 7.17 (2H, d, $J = 8.7$ Hz, ArH2' and 6'), 7.14, (2H, d, $J = 8.7$ Hz, ArH3' and 5'), 7.04 (1H, d, $J = 2.1$ Hz, H7), 6.86, (2H, d, $J = 8.4$ Hz, sulfonylArH2'' and 6''), 6.26 (1H, d, $J = 2.1$ Hz, H5), 3.79 (3H, s, C6 OCH₃), 3.57 (3H, s, C4 OCH₃), 2.36 (3H, s, CH₃). ¹³C NMR δ : 159.6 (C6), 154.8 (C4), 144.7 (sulfonylArC4'), 138.7 (C7a), 136.7 (ArC4'), 132.9 (ArC1'), 132.2 (sulfonylArC2'' and 6''), 130.1 (ArC2' and 6'), 129.8 (ArC3' and 5'), 127.2 (sulfonylArC3'' and 5''), 126.0 (sulfonylArC1'), 124.0 (C2), 121.4 (C3), 112.3 (C3a), 95.4 (C7), 90.8 (C5), 55.8 (C4 OCH₃), 55.3 (C6 OCH₃), 21.8 (Me). ES-MS m/z : 970.7 ([M⁺1]⁺, ⁸¹Br, 20%), 501.7 (20), 304.3 (30), 288.2 (35), 272.2 (40), 244.3 (100), 225.1 (45), 103.7 (80). ES-HRMS m/z : calcd for [M+1]⁺ C₄₆H₃₇N₂O₂S₂⁷⁹Br₂ 969.0515; found 969.0524.

2,2'-Bi-[3-(4-bromophenyl)-4,6-dimethoxy-(N-p-nitrobenzenesulfonyl)]indole (**26**)

This was prepared by general procedure (**B**) using *N*-(p-nitrobenzenesulfonyl)-3-(4-bromophenyl)-4,6-dimethoxyindole **20** (0.50 g, 1.1 mmol), thallium(III) trifluoroacetate (0.30 g, 0.55 mmol), and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **26** (0.33 g, 67%) as a yellow solid, mp 232-235 °C. ¹H NMR δ : 8.14 (2H, d, $J=8.7$ Hz, ArH3 and 5), 7.48 (2H, d, $J=9.0$ Hz, ArH2 and 6), 7.20 (2H, d, $J = 8.7$ Hz, sulfonylArH2'' and 6''), 7.02 (1H, d, $J = 1.8$ Hz, 7H), 7.03 (2H, d, $J=9.0$ Hz, sulfonylArH3'' and 5''), 6.39 (1H, d, $J = 1.8$ Hz, H4), 3.82 (3H, s, C4 OCH₃), 3.63 (s, C6 OCH₃); ¹³C NMR δ : 160.6 (C6), 155.2 (C4), 150.3 (sulfonylArC4), 144.4 (sulfonylArC1), 139.1 (C7a), 132.3 (ArC1), 132.2 (sulfonyl ArC3and 5), 130.4 (ArC2 and 6), 130.0 (C2), 127.9 (sulfonylArC2 and 6), 124.51 (ArC3 and 5), 122.9 (ArC4), 122.0 (C3), 112.4 (C3a), 96.0 (C7), 90.8 (C5), 56.0 (C4 OCH₃), 55.4 (C6 OCH₃ and ArC4 OCH₃). ES-MS m/z : 1031 ([M+1]⁺, ⁸¹Br,

100%) 887.3 (10), 757.9 (30), 705.9 (60), 662.0 (50), 629.8, (50), 585.9 (80), 564.0 (95), 498.8 (70), 451.6, (45), 391.6 (100); ES-HRMS m/z : calcd for $[M+1]^+$ $C_{44}H_{33}N_4O_{12}S_2^{79}Br_2$ 1030.9903; found 1030.9919.

2,2'-Bi-(N-methanesulfonyl-4,6-dimethoxy-3-phenyl)indole (27)

This was prepared by general procedure (B) using *N*-methanesulfonyl-3-phenyl-4,6-dimethoxyindole **22** (0.50 g, 1.5 mmol) thallium(III) trifluoroacetate (0.41 g, 0.75 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH_2Cl_2 /methanol gave the indole **27** (0.33 g, 67%) as a white solid, mp 229-231 °C. 1H NMR δ : 7.25-7.16 (4H, m, ArH2', 3', 4', 5', and 6'), 7.11 (1H, d, J = 3.0 Hz, H5), 6.32 (1H, d, J = 3.0 Hz, H7), 3.87 (3H, s, C4 OCH₃), 3.59 (3H, s, C6 OCH₃), 2.90 (3H, s, CH₃). ^{13}C NMR δ : 160.0 (C7), 155.0 (C4), 148.0 (C7a), 134.0 (ArC1'), 130.0 (ArC2' and 6'), 127.3 (C3), 127.1 (ArC3' and 5'), 127.0 (C2), 126.5 (ArC4'), 113.0 (C3a), 95.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.6 (C6 OCH₃), 40.0 (Me). ES-MS m/z 661.1 ($[M+1]^+$, 50%), 348.1 (20), 132.8 (18), 105.9 (30), 103.9 (40), 71.0 (100); ES-HRMS m/z : calcd for $[M+1]^+$ $C_{34}H_{33}N_2O_8S_2$ 661.1678; found 661.1674.

2,2'-Bi-[3-(4-bromophenyl)-N-methanesulfonyl-4,6-dimethoxy]indole (28)

This was prepared by general procedure (B) using *N*-methanesulfonyl-3-(4-bromophenyl)-4,6-dimethoxyindole **23** (0.50 g, 1.2 mmol), thallium(III) trifluoroacetate (0.33 g, 0.6 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH_2Cl_2 /methanol gave the indole **28** (0.40 g, 80%) as a white solid, mp 179-182 °C. 1H NMR δ : 7.31 (2H, d, J = 9.0 Hz, ArH2 and 6), 7.08 (1H, d, J = 3.0 Hz, H5), 7.02 (2H, d, J = 9.0 Hz, ArH3 and 5), 6.33 (1H, d, J = 3.0 Hz, H7), 3.89 (3H, s, C4 OCH₃), 3.61 (3H, s, C6 OCH₃), 3.06 (3H, s, CH₃). ^{13}C NMR δ : 160.2 (C6), 155.0 (C4), 138.0 (C7a), 132.7 (ArC4'), 132.0 (ArC3' and 5'), 130.3 (ArC2' and 6'), 126.3 (ArC1'), 123.3 (C2), 121.4 (C3), 112.6 (C3a), 95.6 (C7), 89.9 (C5), 56.8 (C4 OCH₃), 56.0 (C6 OCH₃), 40.0 (CH₃). ES-MS m/z : 817.9 ($[M+1]^+$, ^{81}Br , 15%), 466.8 (25), 464.7 (10), 343.8 (10), 341.4 (15), 338.4 (30), 275.9 (15), 204.9 (20), 144.9 (20), 132.8 (25), 105.9 (40), 103.9 (100), 71.0 (35), 60.1 (50); ES-HRMS m/z : calcd for $[M+1]^+$ $C_{34}H_{31}N_2O_8S_2^{79}Br_2$ 816.9889; found 816.9863.

2,2'-Bi-[N-methanesulfonyl-4,6-dimethoxy-(4-methoxyphenyl)]indole (29)

This was prepared by general procedure (B) using *N*-methanesulfonyl-3-(4-methoxyphenyl)-4,6-dimethoxyindole **24** (0.50 g, 1.3 mmol), thallium(III) trifluoroacetate (0.36 g 0.65 mmol), boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH_2Cl_2 /methanol gave the indole

29 (0.38 g, 76%) as a white solid, mp 219-222 °C. ^1H NMR δ : 7.18 (2H, d, J = 9.0 Hz, ArH2' and 6'), 7.12 (1H, d, J = 1.8 Hz, H7), 6.74, (2H, d, J = 9.0 Hz, ArH3' and 5'), 6.33, (1H, d, J = 1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.74 (3H, s, C6 OCH₃), 3.62 (3H, s, ArC4 OCH₃), 2.93 (3H, s, CH₃); ^{13}C NMR δ : 159.9 (ArC4'), 158.5 (C6), 155.0 (C4), 138.0 (C7a), 132.0 (ArC2' and 6'), 126.0 (ArC1'), 123.8 (C3), 123.4 (C2), 113.5 (C3a), 113.0 (Ar3' and 5'), 96.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.7 (C6 OCH₃), 55.6 (ArC4 OCH₃), 40.0 (CH₃); ES-MS m/z : 721.2 ([M+1]⁺, 40%), 547.4 (15), 503.9 (10), 376.1 (20), 360.4 (25), 338.4 (70), 304.2, 244.1, 103.9, (50), 71.0 (50), 60.1 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₃₆H₃₇N₂O₁₀S₂ 721.1889; found 721.1883.

2,2'-Bi[N-acetyl-4,6-dimethoxy-3-(4-nitrophenyl)indole (35)]

This was prepared by general procedure (B) using *N*-acetyl-4,6-dimethoxy-3-(4-nitrophenyl)indole **32** (0.20 g, 0.6 mmol), thallium(III) trifluoroacetate (0.16 g, 0.3 mmol) and boron trifluoride diethyletherate (1.02 mL, 7.2 mmol). The residue was subjected to flash silica gel column chromatography (70:30 hexane:ethylacetate) yielding the indole **36** (0.06 g, 30%) as a pale yellow solid, mp 179-181 °C. ^1H NMR δ : 8.20 (2H, d, J = 9.0 Hz, ArH3' and 5'), 7.61 (1H, d, J = 2.1 Hz, H7), 7.57 (1H, d, J = 9.0 Hz, ArH2' and 6'), 6.36 (1H, d, J = 2.1 Hz, H5), 3.89 (3H, s, C4 OCH₃), 3.70 (3H, s, C6 OCH₃), 2.59 (3H, s, acetyl). ^{13}C NMR δ : 171.0 (C=O), 163.9 (C6), 157.8 (C4), 147.5 (ArC1'), 146.1 (ArC4'), 142.3 (C7a), 126.3 (ArC2' and 6'), 124.0 (ArC3' and C5'), 122.5 (C2), 104.0 (C3a), 96.5 (C7), 96.0 (C5), 56.0 (C6 OCH₃), 55.7 (C4 OCH₃), 26.7 (COCH₃). EI-MS m/z 689 ([M+1]⁺, 7%), 663 (100).

2-(4,6-Dimethoxy-3-(4-nitrophenyl)-N-trifluoroacetyl-1H-indol-7-yl)-4,4',6,6'-tetramethoxy-3,3'-bi(4-nitrophenyl)-N,N'-di(trifluoroacetyl)-1H,1'H-7,7'-biindole (36)]

This was prepared by general procedure (B) using thallium(III) trifluoroacetate (0.14 g, 0.25 mmol), *N*-trifluoro-4,6-dimethoxy-3-(4'-nitrophenyl)-1-trifluoroacetateindole **34** (0.20 g, 0.50 mmol) and boron trifluoride diethyletherate (0.29 g, 2.0 mmol). The residue was subjected to flash silica gel column chromatography (60:40 hexane:ethyl acetate) yielding the trimer **37** (0.70 g, 35%) as a yellow solid, mp 165-168 °C. ^1H NMR δ : 8.24 (2H, d, J = 5.1 Hz, ArH2' and 6'), 8.21 (2H, d, J = 5.1 Hz, ArH2'' and 6''), 7.94 (2H, d, J = 5.1 Hz, ArH2 and 6), 7.66 (2H, d, J = 4.8 Hz, ArH3' and 5') 7.55 (2H, d, J = 4.8 Hz, ArH3'' and 5''), 7.21 (1H, s, H2''), 7.12 (2H, d, J = 4.8 Hz, ArH3 and 5), 6.95 (1H, s, H2'), 6.73, (1H, s, H5''), 6.66 (1H, s, H5), 6.44 (1H, s, H5'), 3.95 (C4 OCH₃''), 3.92 (C4 OCH₃), 3.84 (C6 OCH₃''), 3.81 (C4 OCH₃') 3.79 (C6 OCH₃), 3.72 (C6 OCH₃'). ^{13}C NMR δ : 160.2 (C6'), 157.4 (C6''), 156.8 (C6), 156.6 (C4'), 154.8 (C4''), 154.0 (C4), 147.4 (ArC4'), 147.3 (ArC4''), 146.4 (ArC4), 142.0 (ArC1'), 141.1

(ArC1''), 140.0 (ArC1), 137.2 (C7a''), 137.2 (C7a), 137.0 (C7a'), 130.4 (ArC2' and 6', ArC2' and 6'', and ArC2 and 6), 128.0 (C2), 124.6 (C3''), 124.3 (C3'), 123.0 (ArC3' and 5' and ArC3'' and 5''), 122.4 (ArC3 and 5), 121.9 (C2''), 121.3 (C2'), 121.0 (C3), 114.2 (C3a''), 113.1 (C3a'), 112.8 (C3a), 106.2 (C7''), 105.5 (C7), 100.2 (C7'), 94.5 (C5''), 92.6 (C5), 91.8 (C5'), 57.2 (C4 OCH₃''), 57.1 (C4 OCH₃), 56.0 (C6 OCH₃'), 55.6 (C4 OCH₃'), 55.5 (C6 OCH₃''), 55.3 (C6 OCH₃). ES-MS m/z : 1179.2 ([M+1]⁺, 25%), 1082.3 (20), 986.3 (30), 416.2 (20), 400.3 (20), 338.4 (30), 313.2 (10), 218.9 (30), 179.0 (50), 104.0 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₅₄H₃₆N₆O₁₅F₉ 1179.2095; found 1179.2089.

4,6-Dimethoxy-2-[4',6'-dimethoxy-3'-(4-nitrophenylindol)-7'-yl]-3-[(4-nitrophenyl)indole (37)

This was prepared by general procedure (**B**) using 4,6-dimethoxy-3-(4'-nitrophenyl)indole **6** (0.20 g, 0.67 mmol), thallium(III) trifluoroacetate (0.18 g, 0.34 mmol), and boron trifluoride diethyletherate (0.47 g, 3.3 mmol). The crude residue was subjected to flash silica gel column chromatography (70:30, hexane:ethylacetate) to give the indole **38** (0.04 g, 20%) as a yellow solid, mp 245-248 °C. ¹H NMR δ : 8.56 (1H, s, NH), 8.18 (2H, d, J = 5.1 Hz, Ar3H'' and 5''), 8.02 (2H, d, J = 5.4 Hz, ArH3' and 5'), 7.94 (1H, s, NH'), 7.67 (2H, d, J = 5.1 Hz, Ar2H'' and 6''), 7.55 (2H, d, J = 5.4 Hz, Ar2H' and 6'), 6.88 (1H, s, H2') 6.57 (1H, d, J = 1.8 Hz, H7), 6.41 (1H, s, H5'), 6.32 (1H, d, J = 1.8 Hz, H5), 3.91 (3H, s, C6 OCH₃), 3.88 (3H, s, C6' OCH₃), 3.79 (3H, s, C4 OCH₃), 3.75 (3H, s, C4' OCH₃). ¹³C NMR δ : 158.3 (C6), 155.6 (C6'), 154.5 (C4), 155.3 (C4'), 145.7 (ArC1''), 145.4 (ArC1'), 144.9 (ArC4'), 143.7 (ArC4''), 138.2 (C7a), 137.7 (C7a'), 130.9 (ArC2' and 6'), 129.8 (ArC2'' and 6''), 127.5 (C3), 122.8 (ArC3' and 5'), 123.3 (ArC3'' and 5''), 122.9 (C2'), 117.3 (C3'), 114.9 (C2), 111.2 (C3a), 110.1 (C3a'), 97.1 (C7), 92.7 (C5), 89.7 (C5'), 86.9 (C7), 55.0 (C4 OCH₃), 53.5 (C4' OCH₃), 53.1 (C6' OCH₃), 52.0 C6. ES-MS m/z : 594.2 [M+1]⁺, (15%), 338.5 (50), 336.2 (100), 147.1 (90). ES-HRMS m/z : calcd for [M+1]⁺ C₃₂H₂₆N₄O₈ 595.1829; found 595.1832.

Structure determinations

Full spheres of CCD diffractometer data were measured (monochromatic Mo K α radiation, λ = 0.71073 Å; ω -scans; T ca. 153 K) yielding $N_{\text{(total)}}$ reflections, these merging to N unique (R_{int} cited) after 'empirical'/multiscan absorption correction (proprietary software), N_o with $F > 4\sigma(F)$ being considered 'observed'. All were used in the full matrix least squares refinement on F^2 , refining anisotropic displacement parameters for the non-hydrogen atoms, hydrogen atom treatment following a riding model; reflection weights were $(\sigma^2(F^2) + (aP)^2)^{-1}$ ($P = (F_o^2 + 2F_c^2)/3$). Neutral atom complex scattering factors were used within the SHELXL 97 program.¹⁵ Pertinent results are given in Figures 1, 2 and reference

16; full .cif depositions (without structure factor amplitudes) are deposited with the Cambridge Crystallographic Data Centre, CCDC 680107, 680108.

Acknowledgments

NRV would like to thank the University of Wollongong for a scholarship.

References

1. For general reviews of biaryl synthesis, see (a) Bringmann, G.; Walter, R.; Weirich, R. *Angew. Chem.* **1990**, *29*, 977-991; (b) Hassan, J.; Se'vignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, *102*, 1359-1469.
2. For reviews on the atropselective synthesis of biaryls, see (a) Bringmann, G.; Price Mortimer, A. J.; Keller, P. A.; Garner, J.; Gresser, M. J.; Breuning, M. *Angew. Chem.* **2005**, *44*, 5384-5427; (b) Wallace, T. *Org. Bio Chem.* **2006**, *4*, 3197-3210; (c) Lloyd-Williams, P.; Giralt, E. *Chem. Soc. Rev.* **2001**, *30*, 1454-157.
3. (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457-2483; (b) Suzuki, A. *J. Organomet. Chem.* **1999**, *576*, 147-168; (c) Kotha, S.; Lahiri, K.; Kashinath, D. *Tetrahedron* **2002**, *58*, 9633-9695; (d) Daudin, O. *Eur. J. Org. Chem.* **2005**, 4223-4229.
4. See reference 1 and (a) Gu, R.; Van Hecke, K.; Van Meervelt, L.; Toppet, S.; Dehaen, W. *Org. Biomol. Chem.* **2006**, *4*, 3785-3789; (b) Pezzella, A.; Panzella, L.; Natangelo, A.; Arzillo, M.; Napolitano, A.; d'Ischia, M. *J. Org. Chem.* **2007**, *72*, 9225-9230 and refs therein.
5. (a) Black, D. S.; Keller, P. A.; Kumar, N., *Tetrahedron Lett.* **1989**, *30*, 5807-5808; (b) Keller, P. A.; Birch, C.; Leach, S. P.; Tyssen, D.; Griffith, R. *J. Mol. Graph.* **2003**, *21*, 365-373. (c) Bremner, J. B.; Coates, J. A.; Coghlan, D. R.; David, D. M.; Keller, P. A.; Pyne, S. G. *N. J. Chem.* **2002**, *26*, 1549-1552.
6. See reference 4a and (a) Abbiati, G.; Arcadi, A.; Beccalli, E.; Bianchi, G.; Marinelli, F.; Rossi, E.; *Tetrahedron* **2006**, *62*, 3033-3039; (b) Wada, Y.; Nagasaki, H.; Tokuda, M.; Orito, K. *J. Org. Chem.* **2007**, *72*, 2008-2014; (c) Merlic, C. A.; You, Y.; McInnes, D. M.; Zechman, A. L.; Miller, M. M.; Deng, Q. *Tetrahedron* **2001**, 5199-5212 and refs therein; (d) Abbiati, G.; Arcadi, A.; Beccalli, E.; Bianchi, G.; Marinelli, F.; Rossi, E. *Tetrahedron* **2006**, *62*, 3033-3039; (e) Koza, D. J.; Euler, W. B.; *Heterocyclic Commun.* **1999**, *5*, 399-402

7. (a) Black, D. StC.; Bowyer, M. C.; Catalano, M. M.; Ivory, A. J.; Keller, P. A.; Kumar, N.; Nugent, S. J. *Tetrahedron* **1994**, *50*, 10497-10508; (b) Keawin, T.; Rajviroongit, S.; Black, D. StC. *Tetrahedron* **2005**, 853-861.
8. McKillop, A.; Turrell, A. G.; Young, D. W.; Taylor, E. C.; *J. Am. Chem. Soc.* **1980**, *102*, 6504-6512.
9. Banerji, A.; Ray, R.; Pal, S. C.; Banerji, D.; Maiti, K. K. *J. Indian Chem. Soc.* **1998**, *75*, 698-704.
10. Tholander, J.; Bergman, J. *Tetrahedron* **1999**, *55*, 12595-12602.
11. Black, D. S. C.; Bowyer, M. C.; Bowyer, P. K.; Ivory, A. J.; Kim, M.; Kumar, N.; McConnell, D. B.; Popiolek, M. *Aust. J. Chem.* **1994**, *47*, 1741-1750.
12. Black, D. StC.; Ivory, A. J.; Kumar, N.; *Tetrahedron*, **1996**, *52*, 7003-7012
13. (a) D'Ischia, M.; Napolitano, A.; Tsiakas, K.; Protà, G. *Tetrahedron* **1990**, *46*, 5789-5796; (b) Napolitano, A.; D'Ischia, M.; Protà, G.; Schultz, T. M.; Wolfram, L. J. *Tetrahedron* **1989**, *45*, 6749-6760.
14. Sheldrick, G.M.S. SHELXL 97 – A Program for Crystal Structure Refinement, University of Göttingen, 1997.
15. **12.** $\text{C}_{46}\text{H}_{40}\text{N}_2\text{O}_{10}\text{S}_2$, $M = 845.0$. Monoclinic, space group $P2_1/n$, $a = 12.725(1)$, $b = 17.671(1)$, $c = 18.158(1)$ Å, $\beta = 91.449(2)^\circ$, $V = 4082$ Å³. D_c ($Z = 4$) = 1.37_5 g cm⁻³. $\mu_{\text{Mo}} = 0.19$ mm⁻¹; specimen: 0.10 x 0.09 x 0.08 mm; $T'_{\text{min/max}} = 0.82$. $2\theta_{\text{max}} = 70^\circ$; $N_t = 67457$, $N = 18054$ ($R_{\text{int}} = 0.077$), $N_o = 8927$; $R1 = 0.055$, $wR2 = 0.14$ ($a = 0.067$). $|\Delta\rho_{\text{max}}| = 1.38$ e Å⁻³.
28. $\text{C}_{34}\text{H}_{30}\text{Br}_2\text{N}_2\text{O}_8\text{S}_2 \cdot 2(\text{CH}_3)_2\text{CO}$, $M = 934.7$. Monoclinic, space group $C2/c$, $a = 20.766(2)$, $b = 9.7298(8)$, $c = 22.543(2)$ Å, $\beta = 117.101(2)^\circ$, $V = 4055$ Å³. D_c ($Z = 4$) = 1.53_1 g cm⁻³. $\mu_{\text{Mo}} = 2.2$ mm⁻¹; specimen: 0.28 x 0.15 x 0.12 mm; $T'_{\text{min/max}} = 0.72$. $2\theta_{\text{max}} = 75^\circ$; $N_t = 42853$, $N = 10294$ ($R_{\text{int}} = 0.045$), $N_o = 5440$; $R1 = 0.037$, $wR2 = 0.077$ ($a = 0.034$). $|\Delta\rho_{\text{max}}| = 0.95$ e Å⁻³