UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers:
Part B

Faculty of Science, Medicine and Health

2019

Palaeolatitudinal distribution of lithologic indicators of climate in a palaeogeographic framework

Wenchao Cao
University of Sydney

Simon E. Williams
University of Sydney

Nicolas Flament
University of Wollongong, nflament@uow.edu.au

Sabin Zahirovic
University of Sydney

Christopher Scotese
Northwestern University

See next page for additional authors

Publication Details

Cao, W., Williams, S., Flament, N., Zahirovic, S., Scotese, C. & Muller, R. (2019). Palaeolatitudinal distribution of lithologic indicators of climate in a palaeogeographic framework. *Geological Magazine*, 156 (2), 331-354.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Palaeolatitudinal distribution of lithologic indicators of climate in a palaeogeographic framework

Abstract

Whether the latitudinal distribution of climate-sensitive lithologies is stable through greenhouse and icehouse regimes remains unclear. Previous studies suggest that the palaeolatitudinal distribution of palaeoclimate indicators, including coals, evaporites, reefs and carbonates, has remained broadly similar since the Permian period, leading to the conclusion that atmospheric and oceanic circulation control their distribution rather than the latitudinal temperature gradient. Here we revisit a global-scale compilation of lithologic indicators of climate, including coals, evaporites and glacial deposits, back to the Devonian period. We test the sensitivity of their latitudinal distributions to the uneven distribution of continental areas through time and to global tectonic models, correct the latitudinal distributions of lithologies for sampling- and continental area-bias, and use statistical methods to fit these distributions with probability density functions and estimate their high-density latitudinal ranges with 50% and 95% confidence intervals. The results suggest that the palaeolatitudinal distributions of lithologies have changed through deep geological time, notably a pronounced poleward shift in the distribution of coals at the beginning of the Permian. The distribution of evaporites indicates a clearly bimodal distribution over the past ~400 Ma, except for Early Devonian, Early Carboniferous, the earliest Permian and Middle and Late Jurassic times. We discuss how the patterns indicated by these lithologies change through time in response to plate motion, orography, evolution and greenhouse/icehouse conditions. This study highlights that combining tectonic reconstructions with a comprehensive lithologic database and novel data analysis approaches provide insights into the nature and causes of shifting climatic zones through deep time.

Publication Details

Cao, W., Williams, S., Flament, N., Zahirovic, S., Scotese, C. & Muller, R. (2019). Palaeolatitudinal distribution of lithologic indicators of climate in a palaeogeographic framework. *Geological Magazine*, 156 (2), 331-354.

Authors

Wenchao Cao, Simon E. Williams, Nicolas Flament, Sabin Zahirovic, Christopher Scotese, and R. Dietmar Muller

Paleolatitudinal distribution of lithologic indicators of climate in a paleogeographic framework

WENCHAO CAO^{*†}, SIMON WILLIAMS^{*}, NICOLAS FLAMENT[‡], SABIN ZAHIROVIC^{*}, CHRISTOPHER SCOTese[§] & R. DIETMAR MÜLLER^{*¶}

^{*}EarthByte Group and Basin GENESIS Hub, School of Geosciences, The University of Sydney, NSW 2006, Australia

[‡]School of Earth and Environmental Sciences, University of Wollongong, NSW 2522, Australia

[§]Paleomap Program, 134 Dodge Avenue, Evanston, Illinois 60202, USA

[¶]Sydney Informatics Hub, The University of Sydney, Sydney, NSW 2006, Australia

[†]Author for correspondence: wenchao.cao@sydney.edu.au

Whether the latitudinal distribution of climate-sensitive lithologies is stable through greenhouse and icehouse regimes remains unclear. Previous studies suggest that the paleolatitudinal distribution of paleoclimate indicators, including coals, evaporites, reefs and carbonates, have remained broadly similar since the Permian period, leading to the conclusion that atmospheric and oceanic circulation control their distribution rather than the latitudinal temperature gradient. Here we revisit a global-scale compilation of lithologic indicators of climate, including coals, evaporites and glacial deposits, back to the Devonian period. We test the sensitivity of their latitudinal distributions to the uneven distribution of continental areas through time and to global tectonic models, correct the latitudinal distributions of lithologies for sampling- and continental area-bias, and use statistical methods to fit these distributions with probability density functions and estimate their high-density latitudinal ranges with 50% and 95% confidence intervals. The results suggest that the paleolatitudinal distributions of lithologies have changed through deep geological time, notably a pronounced poleward shift in the distribution of coals at the beginning of the Permian. The distribution of evaporites indicate a clearly bimodal distribution over the past ~400 Ma, except for Early Devonian, Early Carboniferous, the earliest Permian and Middle and Late Jurassic times. We discuss how the patterns indicated from these lithologies change through time in response to plate motion, orography, evolution, and greenhouse/icehouse conditions. This study highlights that plate reconstructions, combined with a comprehensive lithologic database, and novel data analysis approaches provide insights into the nature and causes of shifting climatic zones through deep time.

Keywords: paleolatitudinal distribution, paleoclimate lithologic indicators, atmospheric CO₂, paleogeography, plate tectonic reconstructions, statistical analysis

1. Introduction

The paleolatitudinal distribution of climate-sensitive lithologic deposits is important to understand past climates (Gorden, 1975; Evans, 2006; Boucot *et al.* 2013), to reconstruct paleogeographies (Ronov *et al.* 1984, 1989; Scotese 2001, 2004; Golonka *et al.* 2006; Blakey, 2008), and to provide constraints for climate modelling (Boucot & Gray, 2001; Craggs *et al.* 2011) and plate motion histories (Scotese & Barrett, 1990; Witzke, 1990). However, the past distribution of climate-sensitive lithologies remains poorly understood, especially in deep geological times. Latitudinal distribution patterns of paleoclimate indicators have been considered broadly stable since the Permian, leading to an interpretation that atmospheric and oceanic circulation control their distribution rather than equator-to-pole temperature gradients (Ziegler *et al.* 2003). A global paleomagnetic compilation of Earth's basin-scale evaporite records suggests that evaporite paleolatitudes have varied to some degree over the past 2 billion years, with a relatively large difference between Cenozoic–Mesozoic times (a mean paleolatitude of $23 \pm 4^\circ$) and Devonian–Ediacaran times (a mean paleolatitude of $14 \pm 2^\circ$) (Evans, 2006). In addition, previous studies rarely considered the effect of the uneven distribution of continental areas through time (Vilhena & Smith, 2013) and the implications of applying different global tectonic reconstruction models (Scotese & Barrett, 1990; Ziegler *et al.* 2003) on latitudinal distribution of climate indicators.

In order to better understand the latitudinal distribution patterns of climate-sensitive lithologies and the associated climate change, we revisit a global-scale lithologic compilation of coals, evaporites and glacial deposits, with our analysis extending back to the Devonian period (~400 Ma ago). We test the sensitivity of their latitudinal distribution to uneven distribution of continental areas through time and to reconstruction models, and then quantify their distribution patterns with corrections of sampling- and continental area-bias, using Significant Zero crossings of the derivative (SiZer) of Chaudhuri & Marron (1999) to obtain the best fitting probability density functions and Highest Density Regions (HDR) of Hyndman (1996) to compute 50% and 95% confidence intervals. The shifting latitudinal distribution patterns reflect climate change associated with humid/arid, precipitation/evaporation or cold/warm climatic conditions. To understand the mechanism, we compare the trends of latitudinal distribution patterns of these paleoclimate deposits with the atmospheric CO₂ evolution (Foster *et al.* 2017) and tropic mountain range areas derived from the paleogeographic maps (Golonka *et al.* 2006; Cao *et al.* 2017) over time, as increased silicate weathering rates associated with tropical mountain uplift leads to a decrease in atmospheric CO₂ concentration and this may initiate glaciation (Tabor & Poulsen, 2008; Montañez & Poulsen, 2013; Goddérís *et al.* 2017; Swanson-Hysell & Macdonald, 2017).

2. Data and global plate tectonic models

We use a dataset of climate-sensitive lithologies, including coals, evaporites and glacial deposits (tillites, dropstones and glendonites), based on a comprehensive global-scale

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

86 compilation (Boucot *et al.* 2013). This compilation records and evaluates various types of
87 climatically sensitive lithologic deposits, including coal, cyclothems, laterite, bauxite,
88 lateritic manganese, oolitic ironstone, kaolinite, glendonite, tillites, dropstones, calcretes,
89 evaporites, clay minerals, palms, mangroves, and crocodilians, based on a comprehensive
90 literature review. It covers geological times from the Cambrian period to the Miocene epoch.
91 Each data point includes a reference source and geographic location. In order to analyse the
92 effect of global climate change on the latitudinal distribution patterns of climate-sensitive
93 lithologies, we use recorded occurrences of coals, (mostly marine) evaporites, and glacial
94 deposits (a combination of tillites, dropstones and glendonites) in this study, as distributions
95 of other lithologies included in the database are less reliable latitude indicators. For instance,
96 the record of palms, mangroves, and crocodilians suffers from sampling bias as they are
97 mostly collected from middle-high latitudes, while sample sizes are small for laterites and
98 oolitic ironstones.

100 The dataset used in this study covers Devonian to Miocene times and is divided into 23 time
101 intervals at the scale of geological stages (Table 1). The time range that we assign to each
102 data point is entirely derived from the time ranges that Boucot *et al.* (2013) has used to draw
103 the maps, and we use the middle ages of these time intervals to determine the reconstruction
104 times (Table 1). We use modern peats (as they may eventually become coals), evaporites and
105 glacial sediments (tillites and ‘glacio-marine’ beds) from Ziegler *et al.* (2003) for reference.
106 Coals and peats are generally considered as indicators of terrestrial humidity, reflecting
107 climatic regimes in which precipitation exceeds or is equal to evaporation (Parrish *et al.*
108 1982; Hallam, 1985; McCabe & Parrish, 1992; Price *et al.* 1995). Evaporites are formed in
109 climatic conditions where evaporation exceeds the combined effects of precipitation, marine
110 or river influx, and runoff (Craggs *et al.* 2011), yet, they may occur in temperate regions
111 (Boucot *et al.* 2013). Tillites, dropstones and glendonites in the dataset are glacial or glacial-
112 origin deposits indicating cold climate, generally due to high latitudes, but some could be
113 related to high elevations (Boucot *et al.* 2013).

[Insert Table 1]

117 The climatically sensitive lithologic deposits are distributed unevenly over different time
118 intervals (Figs. 1, 2). There are very few records of coals in the Devonian period (Figs. 1, 2),
119 with coals largely occurring in the Late Devonian stratigraphies (Boucot *et al.* 2013). Yet a
120 large number of coal records appear in Carboniferous, Permian, Late Triassic, Early Jurassic
121 and Neogene times (Figs. 1, 2a). Evaporites have relatively uniform richness over most of the
122 time intervals, except in Devonian, Late Carboniferous, Early Permian and Paleogene times
123 (Figs. 1, 2b), whereas, glacial deposit records are relatively rare and sparse over time (Figs. 1,
124 2c) due to the dominance of greenhouse conditions in the Phanerozoic (Hay, 2016).

[Insert Figures 1, 2]

128 In this study, we primarily use the global plate tectonic model of Matthews *et al.* (2016) to
129 reconstruct the lithologic data points back in time from present-day geographic locations.

This tectonic model represents a combination of a tectonic model for Mesozoic and Cenozoic times (230-0 Ma) (Müller *et al.* 2016) and a tectonic model for Paleozoic times (410-250 Ma) (Domeier & Torsvik, 2014), with a number of changes required to enable a smooth merging of the models as described by Matthews *et al.* (2016). It is a relative plate motion model that is ultimately tied to Earth's spin axis through an absolute reference frame (Matthews *et al.*, 2016). We removed the true polar wander correction from the original absolute plate motion model (Domeier & Torsvik, 2014) to tie the relative rotations to a paleomagnetic reference frame. Therefore, the tectonic reconstruction better reflects the true paleolatitudinal distribution of paleoclimate indicators (van Hinsbergen *et al.* 2015).

In order to test the sensitivity of the paleolatitudinal distribution of climate-sensitive deposits to the choice of reconstruction model, except for the plate motion model of Matthews *et al.* (2016), we use two alternative reconstruction models of Scotese (2008) and Golonka (2007) which are both based on the reconstruction of Scotese (2004). The reconstruction model of Golonka (2007) uses paleomagnetic data to constrain the paleolatitudinal positions of continents and rotation of plates, and hot spots, where applicable, are used as reference points to calculate paleolongitudes. A version of this model that includes an updated absolute reference frame is described in Wright *et al.* (2013) and the model is available in the supplementary materials of Wright *et al.* (2013). The reconstruction model of Scotese (2008) uses paleomagnetic reference frames for 410-250 Ma (Ziegler *et al.* 1979) and 250-100 Ma (Ziegler *et al.* 1983), and a hotspot reference frame for 100-0 Ma (Müller *et al.* 1993).

3. Methods

We first reconstructed the lithologic data for coals, evaporites and glacial deposits from their modern locations to ancient locations in geological time (Fig. 1) using the global plate motion model of Matthews *et al.* (2016). Due to the inconsistency of age assignments between some lithologic data and “underlying” terrane polygons, leading to unexpected reconstructed locations of these lithologic data, we only considered the data points whose time ranges of existence were within the time span and present-day spatial extent of the underlying terrane. We next binned the data points using a 5°x5° mesh of the global surface, following Ziegler *et al.* (2003) (Fig. 3b). Each bin contained a variable number of data points or none, for example, a bin could contain 0-24 data point(s) for modern peats at present-day coordinates (Fig. 3b). The recorded data points were unevenly distributed spatially (Fig. 3a, b), largely due to sampling bias. To remove the bias, we resampled the data by collapsing all the samples into a single value for each bin: 1 where data existed (Fig. 3c, red bins), and 0 where there were no data points (Fig. 3c, blue bins). We subsequently summed up the numbers of resampled data in each 5° latitudinal strip (Fig. 4b). These steps significantly reduced the numbers of data points considered per latitudinal strip (compare the y-axis of Fig. 4a, b).

[Insert Figures 3, 4]

Plate tectonic motion (Golonka, 2007; Scotese, 2008; Matthews *et al.* 2016) leads to a biased distribution of lithologic deposits on Earth. Latitude bands containing more continental area

are more likely to have more lithologic deposits preserved. In an attempt to remove this bias, we first calculated the continental area covered by each 5° latitudinal strip at each time interval since the Devonian period (Fig. 4c), using the HEALPix pixelization method that results in equal sampling of data on a sphere (Górski *et al.* 2005) and therefore equal sampling of surface areas. We next corrected the number of resampled lithologic data in a 5° latitudinal strip using the continental area in the same strip to remove the bias due to the uneven distribution of continental areas through time. Subsequently, the results were scaled to represent probabilities of data appearing in a 5° latitudinal strip (Fig. 4d). In order to investigate the paleolatitudinal zonal patterns of these climate indicators, we combined the data corrected for continental area bias from both hemispheres to form symmetric and composite zonal patterns, and normalised the results in a strip into probabilities with respect to the sum of all the values in the same strip (Fig. 4e). The symmetric zonal patterns erased the differences between the two hemispheres and can be used as indicating the influence of climate change on the distribution of climate-sensitive lithologic deposits (Scotese & Barrett, 1990; Ziegler *et al.* 2003).

We used SiZer (Significant ZERO crossing of the derivatives; by Chaudhuri & Marron, 1999), a data analysis method to identify significant peaks from kernel density estimation to obtain probability density functions of zonal patterns (Fig. 4f, black lines). Instead of trying to find the one bandwidth that provides the closest match to the unknown true density, SiZer looks at the whole range of bandwidths. Peaks and troughs are identified by finding the regions of significant gradient (zero crossings of the derivative; Chaudhuri & Marron, 1999). In order to identify high density intervals of distribution of these paleoclimate indicators, we used the Highest Density Regions (HDR) method of Hyndman (1996) to compute the high-density latitude ranges with 50% (Fig. 4f, grey area) and 95% (Fig. 4f, light grey area) confidence intervals. HDR is a statistical method for calculating confidence intervals when the sampling distribution has multiple peaks. For discrete-valued distribution, HDR simply consists of those elements of the sample space with highest probability (Hyndman, 1996).

4. Results

4.a. Paleolatitudinal distribution of coals, evaporites and glacial deposits

We counted the original data points for coals (Fig. 5a), evaporites (Fig. 6a) and glacial deposits (Fig. 7a) reconstructed using the plate tectonic model of Matthews *et al.* (2016) in each 5° latitudinal strip at each time interval since the Devonian period. Coals are distributed in a wide latitude range from the equator to the poles (Fig. 5a), while evaporites are concentrated between 60°N and 60°S (Fig. 6a). Glacial deposits generally occur middle-high latitudes except for a record of glendonites appearing near the equator in the Early Carboniferous (Tournaisian-Visean) from an outer shelf environment with near zero-degree seawater (Brandley & Krause, 1993a, b; 1994), a record of tillites in the Late Carboniferous (Kasimovian-Gzhelian) and a record of tillites in the Early Permian both from high elevation periglacial environment (Becq-Giraudon *et al.* 1996) (Fig. 7a). Therefore, these three records (highlighted with question marks in Figure 7) may not be reliable enough to indicate cold climate relevant to latitudinal change.

[Insert Figures 5-7]

Generally, more data points of coals (Fig. 5a), evaporites (Fig. 6a) and glacial deposits (Fig. 7a) are sampled from the northern than the southern hemisphere since Pangea breakup in the Early Jurassic (Ronov, *et al.* 1984, 1989; Scotese, 2001, 2004; Golonka *et al.* 2006; Blakey, 2008). During Early Devonian-Late Triassic time, the numbers of data points of these lithologies in the southern hemisphere exceed those in the northern hemisphere in general, with glacial deposits only occurring in the southern hemisphere in Early Devonian and Carboniferous times (Fig. 7a). This mainly reflects that plate tectonic motions modify the continental configuration through time, which has led to more continental areas in the northern hemisphere since the Early Jurassic (Figs. 5c, 8). Continental areas in the northern hemisphere have consistently increased since the Devonian period, while the areas in the southern hemisphere have decreased accordingly (Figs. 5c, 8).

[Insert Figure 8]

After removing the sampling- (Figs. 5b, 6b, 7b) and continental area-biases (Figs. 5d, 6d, 7d) over the data in each 5° strip at each time interval, the differences among strips at a time interval are evened out to some degree (for example, compare the y-axis of Figure 5a, b, d). The zonal patterns for coals (Fig. 5e), evaporites (Fig. 6e) and glacial deposits (Fig. 7e) considerably vary over time.

We compute the high-density latitudinal ranges with 50% and 95% confidence intervals for coals (Fig. 5f), evaporites (Fig. 6f) and glacial deposits (Fig. 7f), respectively, at each time interval. Considering the extent of these 95% confidence ranges, coals are relatively widely distributed on Earth's surface over the time period of interest, extending from the equator to the poles for many time periods (Fig. 5f, light grey area). Evaporites (96% confidence) are generally concentrated within low-middle latitudes (0-60°) in two hemispheres, with a mean paleolatitude of $26 \pm 3^\circ$ in each hemisphere since the Devonian (Fig. 6f, light grey area). This result is consistent with the mean paleolatitudes of $23 \pm 4^\circ$ during Cenozoic-Mesozoic time and $21 \pm 4^\circ$ during Permian-Carboniferous time (both with 95% confidence), recalculated from paleolatitudes of a global compilation of large evaporite basins extending back through Proterozoic time (Evans, 2006). Glacial deposits (95% confidence) occur in Late Devonian, Carboniferous, Permian, Jurassic, Early Cretaceous, and Late Paleogene-modern times (Fig. 7f, light grey area). This is generally consistent with three main Phanerozoic ice ages during Carboniferous-Permian, Late Jurassic-early Cretaceous and Early Eocene-modern times (Frakes *et al.* 1992).

Focussing on 50% confidence intervals, coal latitudinal belts show strong fluctuation over time (Fig. 5f, dark grey area). They move from the middle latitudes towards the equator during the Devonian, but the original coal deposits in the Devonian are too few to provide much climatological insight (Boucot *et al.* 2013). During the Carboniferous, they are strictly limited within 0-25°N and S, however, they separate into two latitudinal belts of 0-10° and

40-60° during the earliest Permian (Asselian-Sakmarian) (Fig. 5f). Subsequently, they undergo a poleward shift until reaching higher latitudes (60-90°) during the Late Jurassic, with a sharp equatorward movement in the Early Triassic and a rapid poleward shift in the Middle Triassic but these may be not reliable as they are based on very few data points (Figs. 1, 2a). Starting from the Early Cretaceous, the distributions of coals contract towards the equator until the Miocene when there are two latitudinal belts at ~30-50° and ~0-15° on two hemispheres, with a rapid shift towards the equator during the Early Eocene. Modern peats are mainly distributed within ~40-65° latitudes. Overall, the latitudinal belts determined for coals from probability density curves with 50% confidence generally occur at low latitudes during Paleozoic time, middle latitudes during Permian, Triassic and Cenozoic times, and high latitudes during Jurassic and Cretaceous times.

Evaporite latitudinal belts with 50% confidence (Fig. 6f, grey area) mostly occur at low latitudes (~0-30° N and S). They move poleward from the Early Devonian until the Early Carboniferous and then towards the equator until the Late Triassic. Starting from the Middle Triassic, they start to shift poleward until modern times at a mean latitude of ~35° N and S, with a short and rapid poleward movement during early Eocene (Lutetian).

The trends in how the highest density latitudes of glacial deposits with 50% confidence (Fig. 7f, grey area) shift northwards or southwards are similar to coals, but at higher latitude. They generally concentrate within middle-high latitudes through time (Fig. 7, grey area). They occur at ~50-70°N and S latitudes during the Late Devonian and then move equatorward to middle latitudes (~40-60°) during most of the Carboniferous, with a rapid poleward movement in the Late Mississippian. The Late Pennsylvanian-Early Permian transition is marked by a rapid shift of glacial deposits towards high latitudes. They remain at high latitudes during Jurassic and Cretaceous times, and then move to middle-high latitudes (~50-75°) during Paleogene and Miocene times. Modern glacial deposits mainly lie near the poles.

4.b. Sensitivity tests

4.b.1. Assumption of climate symmetry (zonal patterns)

In order to test the assumption that past climates were zonal and hemispherically symmetric, we investigated the latitudinal distributions of lithologies for each hemisphere independently through time, and estimated their high-density latitudinal belts using the same statistical methods (Supplementary Figs S1 and S2). At the first order, more lithologic data are distributed on the northern hemisphere over the timeframe, except for late Paleozoic and Cenozoic times when more glacial deposits formed in the southern hemisphere. The distributions of coals and evaporites in the northern hemisphere (Supplementary Figs S1 and S2) are similar to the results with data combined over two hemispheres (Figs 12a-c). This is largely because of the uneven distribution of continental areas through time, leading to a biased distribution of lithologic deposits on the northern hemisphere. Overall, the data distribution warrants combining the two hemispheres in zonal patterns.

4.b.2. Continental area bias

In order to examine sensitivity of the zonal patterns of these lithologic deposits to continental area bias, we compute the zonal patterns for coals (Fig. 9a), evaporites (Fig. 9b) and glacial deposits (Fig. 9c) for each time interval since the Devonian using the plate motion model of Matthews *et al.* (2016) with (Fig. 9, blue area) and without (Fig. 9, red) continental area correction, respectively. The distribution patterns of coals using the two approaches are significantly different in Middle Devonian, most of Carboniferous, Early and Middle Triassic, Jurassic, Cretaceous and Early Paleogene times. During these time periods except the Middle Devonian, the zonal patterns of coals without continental area correction are closer to the equator (Fig. 9, red). This mainly results from the uneven latitudinal distribution of continental areas, with very little to no continental coverage near the poles during these periods (Fig. 5c). Due to relatively small continental coverage near the equator (Fig. 5c), the zonal patterns of coals without area correction are closer to the poles during the Middle Devonian (Fig. 9, red).

[Insert Figure 9]

The uneven distribution of continental areas through time has relatively less impact on the zonal patterns of evaporites (Fig. 9b) and glacial deposits (Fig. 9c). Evaporite zonal patterns are generally identical over time, except during Late Mississippian, Late Pennsylvanian, the earliest Permian, Middle Triassic and late Cretaceous times (Fig. 9b). Due to different continental coverage at different latitudes (Fig. 5c), the area correction leads to more poleward distributions of evaporites during Late Mississippian, Late Pennsylvanian and earliest Permian times but more equatorward during Middle Triassic and late Cretaceous times. With and without area correction, glacial deposit zonal patterns are mostly identical over time, except for most of Permian, Late Paleogene and modern times when the area correction results in more poleward distributions and for Early and Late Mississippian and Late Pennsylvanian times, more equatorward distributions (Fig. 9c). Overall, latitudinal distributions of coals show stronger sensitivity to uneven distributions of continents than evaporites and glacial deposits, which could be mainly due to coals having a wider latitudinal distribution than evaporites and glacial deposits over time.

4.b.3. Reconstruction model bias

We use alternative reconstruction models of Scotese (2008) and Golonka (2007) with modifications as described in Wright *et al.* (2013), in addition to the model of Matthews *et al.* (2016), to analyze the sensitivity of latitudinal distributions of coals, evaporites and glacial deposits to reconstruction model. We use the same method described in the Methods section to obtain the probability density functions of the latitudinal distributions for these lithologies over time. The results indicate that the three reconstructions result in only second-order differences through time (Fig. 10). Differences in the distributions are most pronounced where reconstructed paleolatitudes differ for blocks that host a large proportion of data points for a given time interval.

[Insert Figure 10]

The latitudinal distributions of coals reconstructed using the three tectonic models are generally similar over time, except during Early Devonian, Early Carboniferous, the earliest Permian, Middle and Late Paleogene and Neogene times. In the Early Devonian, the latitudinal distributions of coals using the reconstruction of Golonka (2007) are more northward than those using the other two reconstructions, largely because North America and Eurasia in the model of Golonka (2007) are located further northward than in the other two models during this period (see Supplementary material). The more poleward location of Africa during the Early Carboniferous in the reconstruction of Matthews *et al.* (2016) leads to a more poleward distribution of coals. During the earliest Permian, numerous coal deposits occur in North and South China blocks. Their more northward locations in the model of Matthews *et al.* (2016), taken from the work of Domeier & Torsvik (2014), result in more occurrences at middle-latitudes (see Supplementary material). The terranes in the reconstruction of Scotese (2008) are generally further north relative to the other two models during Middle and Late Paleogene and Neogene times (see Supplementary material). This results in more poleward distributions of overall paleoclimate lithologies (Fig. 10a).

The latitudinal distributions of evaporites among the three reconstructions are very similar since the Devonian (Fig. 10b), except for Early Devonian and Carboniferous times. In the Early Devonian, evaporites are spread further from the equator using the reconstruction of Golonka (2007) than the other two reconstructions, which is mainly due to more northward locations of North America and Eurasia in the model of Golonka (2007). More southward locations of numerous plates in the reconstruction of Golonka (2007) (see Supplementary material) result in more equatorial distributions of evaporites during the Carboniferous (Fig. 10b).

The latitudinal distributions of glacial deposits are also similar through time among the three considered reconstructions (Fig. 9c), except for the Late Devonian, Early and Middle Mississippian and most of the Permian. During the Late Devonian, more glacial deposits are distributed at high latitudes using the reconstruction of Scotese (2008), mainly due to the more southerly location of South America in this reconstruction than that in the other two reconstructions (see Supplementary material). The more equatorial locations of South America and Northwest Africa in the model of Scotese (2008) leads to more equatorial distributions of glacial deposits during Early and Middle Mississippian times. During most of the Permian period, Antarctica and Australia are closer to the south pole in the reconstruction of Golonka (2007) (see Supplementary material), resulting in more poleward distribution of glacial deposits (Fig. 10c).

4.b.4. Spatial sampling resolution bias

[Insert Figure 11]

We compare the use of three different spatial resolutions to bin the original data points to remove the effect of sampling bias on the distribution patterns of these climatically lithologic deposits, using the plate motion model of Matthews *et al.* (2016), for each time interval since the Devonian period: 10° (Fig. 11a), 5° (Fig. 11b), and 2° (Fig. 11c). The results indicate that there are too few strips containing data points, especially during the Devonian, to be able to obtain much statistical significance using 10° as the binning size (Fig. 11a). Using 2° as the binning size results in noisier distributions (Fig. 11c) and a more difficult analysis of distribution patterns. A spatial resolution of 5° is a good compromise that captures the data distribution yet results in less noise, and follows previous work that aids comparison with our work (Fig. 11b).

5. Discussion

5.a. Paleolatitudinal distribution patterns of lithologic climate indicators over time

We present the high-density latitudinal ranges in one hemisphere, due to the assumed symmetry of two hemispheres, with 50% and 95% confidence intervals for coals (Fig. 12a), evaporites (Fig. 12b) and glacial deposits (Fig. 12c), respectively, at each time interval. We also present the cumulative high-density latitudinal ranges over the entire time period for these lithologies (rightmost panels in Fig. 12a, b, c). Overall, the latitudinal belts determined from probability density curves with 50% confidence for coals have changed since the Devonian. Some previous studies concluded that the paleolatitudinal distribution of coals has remained broadly similar since the Permian (Ziegler *et al.* 2003), whereas we have shown that results for earlier in the Paleozoic are significantly different from results from the Permian to present (Fig. 12a, b, c). In addition, the temporal division at the scale of geological periods in previous studies did not attempt to resolve variations over shorter time periods, whereas the dataset used in this study is based on a time division at the scale of geological stages. The temporal resolution based on the scale of geological periods does not allow us to capture variations of latitudinal distribution patterns of climate-sensitive lithologic deposits within geological periods, such as the significant movement towards the equator from probability density curves with 50% confidence for coals between Early Paleogene and Middle Paleogene times in Figure 12a.

[Insert Figure 12]

Highest density latitudinal belts of evaporites with 50% confidence exhibit relatively subtle shifts within a narrow range of latitudes over time (Fig. 12b), although they are very similar to the cumulative result covering the entire temporal range (right panel in Fig. 12b). Evans (2006) suggested that evaporite paleolatitudes remain stable during Cenozoic–Mesozoic times (a mean paleolatitude of $23 \pm 4^\circ$) and during Devonian–Ediacaran times (a mean paleolatitude of $14 \pm 2^\circ$). This might be because the time division, weighting method or source dataset of Evans (2006) does not capture variations of mean latitudes of evaporites at finer-resolution time divisions. In addition, Evans (2006) argued that the distribution of evaporites over the past two billion years has remained bimodal about the equator rather than being centred on the equator as expected from general-circulation climate simulations (Hunt,

1982; Jenkins, 2000; Jenkins, 2001). These considerations are important to test predictions from climate models with differing obliquities. Our results generally indicate a bimodal distribution of evaporites over the past ~400 Ma, except for Early Devonian, Early Carboniferous (Serpukhovian), earliest Permian (Asselian-Sakmarian) and Middle and Late Jurassic times during which the distribution is unimodal (Fig. 6f). These unimodal distributions should be treated with caution, since comparison with the unsmoothed distributions (Fig. 6e) suggests that the smoothing of the SiZer algorithm contributes to the existence of unimodal distributions in the final results. However, it is also apparent in both the smoothed and unsmoothed results that the clear bimodal pattern observed for the most recent time intervals is not clearly resolved within many older temporal intervals.

5.b. Contributors to the paleolatitudinal distribution of climate-sensitive lithologies

The evolution of latitudinal belts of coal samples with 50% confidence interval (Fig. 12a) may reflect the complex Phanerozoic fluctuations in the latitudinal extent of continental humid zones. The latitudinal distribution of coals with 50% confidence change over long time scales (Figs. 5, 12a), being concentrated at low latitudes during the Carboniferous period, middle latitudes during Permian and Triassic times, high latitudes during Jurassic-Early Paleogene times, before returning to middle latitudes during Middle Paleogene-modern times. The distribution during the Devonian period is unlikely to be reliable due to a lack of data samples (Figs. 1, 2). The Carboniferous period is significantly different from post-Paleozoic times (Fig. 12a). Abundant coal samples were accumulated in tropical regions (~0-25°N and S) during the Carboniferous period, while they were mostly deposited at middle or high latitudes during Mesozoic and Cenozoic times (Fig. 12a). This shift could indicate a climatic transition from humid to arid conditions from the Late Pennsylvanian through the Early Permian on equatorial Pangaea (Rowley *et al.* 1985; Ziegler, 1990; Parrish, 1993; Ziegler *et al.* 1997; Tabor & Montañez, 2004; Montañez *et al.* 2007; Tabor *et al.* 2008; Tabor & Poulsen, 2008). Tabor & Poulsen (2008) evaluated the climate factors possibly explaining this climate change, including due to changing continental configurations from tectonics, time-varying land-sea distribution, supercontinentality, monsoon, uplift/collapse of the Central Pangean Mountains (CPMs) (Fig. 13), waxing and waning of ice sheets in Gondwanaland and atmospheric CO₂. They suggested that the northward drift of most of Pangaea across climate zones, increasing atmospheric CO₂ level and Gondwanaland deglaciation could have been the main factors controlling the low-latitude climate of Pangea during Late Pennsylvanian-Early Permian times. The deglaciation of Gondwanaland is considered to be a possible cause of climate change over low-latitude Pangaea through their influence on large-scale atmospheric circulation (Ziegler *et al.* 1987; Perlmutter & Matthews, 1989; Cecil, 1990; Miller & West, 1993; Miller *et al.* 1996; Soreghan *et al.* 1997; Cecil *et al.* 2003; Perlmutter & Plotnick, 2003; Poulsen *et al.* 2007). Late Paleozoic climate simulations suggest that the deglaciation of Gondwana and atmospheric CO₂ rise could explain much of the Late Paleozoic climate change over low-latitude Pangaea and regional uplift/erosion of the CPMs has a secondary effect on tropical precipitation (Peyser & Poulsen, 2008). However, the change from dominantly humid paleoclimate indicators in Pennsylvanian strata

to arid and seasonal indicators in Permian strata is considered possibly related to the evolution of the CPMs (Goddris *et al.* 2017).

[Insert Figure 13]

We compared our results to a recent estimate of atmospheric CO₂ since the Devonian period (Foster *et al.* 2017; Fig. 12d), and to additional aspects of Earth's paleogeographic history extracted from digital paleogeographic reconstructions (Golonka *et al.* 2006; Cao *et al.* 2017; Fig. 12e-g). Atmospheric CO₂ levels do not indicate significant changes at the beginning of the Permian period that would directly coincide with the poleward shift in coal distributions. The areas of tropical mountain ranges close to the equator (Fig. 12e), thought to be regions of significantly enhanced weathering rates, similarly does not record a significant change from the Late Pennsylvanian through the Early Permian (Fig. 12e). In addition, we computed Phanerozoic ice sheets areas from paleogeographic maps (Golonka *et al.* 2006; Cao *et al.* 2017), and the results indicate that ice sheets indeed considerably reduced during the Late Pennsylvanian-Early Permian transition (Fig. 12g). Plate tectonic configurations (Golonka, 2007; Scotese, 2008; Matthews *et al.* 2016) indicate that most of Pangaea moved rapidly northward during the Late Pennsylvanian-Early Permian transition. Walker *et al.* (2002) argued that these motions influenced patterns of shallow-marine carbonates during Phanerozoic times through a long-term decrease in areas of tropical shelf based on the paleogeographic atlas of Scotese & Golonka (1992) - although, a persistent decrease in the area of tropical shallow marine environments over the past ~400 Ma is less apparent in paleogeographic maps of Golonka *et al.* (2006) and Cao *et al.* (2017) (Fig. 12f). The resulting migration of tropical basins from humid climate zones associated with the Intertropical Convergence Zone into the adjacent northern subtropical zone could be attributed to long-term tropical aridification (Witzke, 1990; Ziegler *et al.* 1977; Gibbs *et al.* 2002; Rees *et al.* 2002; Tabor *et al.* 2008).

It is important to consider some non-climatic factors that may influence the latitudinal distributions of these lithologies. Carboniferous coal-forming flora were dominated by pteridophytes with a low tolerance to groundwater fluctuations due to their shallow root system, making them less likely to form thick preserved coal seams even where favourable peat-forming conditions existed (Diessel, 1992). In contrast, post-Carboniferous coal measures were dominated by Gymnosperms from the Permian to the Cretaceous and angiosperms since the Triassic. They can occupy a wider range of latitudes due to their greater tolerance to groundwater fluctuations and adaptability in more fully marginal peat-forming conditions (Diessel, 1992). The evolution of plant groups through space and time could contribute to the changing distributions of coal deposits through time, especially its significant poleward shift in the Early Permian. Additionally, a further contributor to the Carboniferous low-latitude peak in coal distribution could be the relatively rare occurrence of extensive foreland-basin systems within the tropics during Pangea assembly, ideal for the formation and preservation of coal (Nelsen *et al.* 2016). In summary, the major poleward shift in coal distributions beginning in the Early Permian is likely to be driven by a

combination of processes, resulting from the interplay of tectonics, deglaciation, evolution of plants and paleoclimate.

The changing latitudinal belts of evaporites with 50% confidence (Fig. 12b) indicate the evolution of low-latitudinal arid zones over time but the fluctuations are subtle relative to other indicators. During the Early Paleogene, they underwent a relatively rapid shift away from the equator, contemporaneous with climate-driven shifts in coals and glacial deposits. In older times, the origins of changes are less clear (Fig. 12b). Ancient marine evaporite formation is not only related to dry climate but also tectonic setting (Warren, 2010). For instance, during the Early Devonian, many evaporites occurred in the northern margins of tropical Laurentia and low-latitude Siberia (Fig. 1), which could explain the equatorial distribution of evaporites during the time (Fig. 12b). The latitudinal belt of evaporites shift poleward to mean latitudes of ~25° N and S during Middle Devonian-Early Carboniferous time (Fig. 12b). This could be mostly due to that evaporites accumulate in the southern, western or eastern margins of Laurentia during the Middle Devonian-Early Carboniferous time (Fig. 1). From the assembly of Pangea in the Late Carboniferous until the Middle Triassic (Ziegler *et al.* 1979), abundant evaporites occur in the western and eastern margins of low-latitude Pangea (Fig. 1). Starting from the Late Triassic (~240 Ma) the pattern of rift basins formed during Pangea breakup (notably the Atlantic margins and the Gulf of Mexico) will have exerted an additional control on evaporite distributions (Fig. 1).

Atmospheric CO₂ is thought to be a primary driver of both ice sheet and climate variability (Montañez & Poulsen, 2013; Foster *et al.* 2017, Lowry *et al.*, 2004) and solar luminosity secondarily influences Paleozoic glaciation (Lowry *et al.*, 2004). Mountain uplift enhances silicate rock weathering, resulting in a decrease in atmospheric CO₂ concentrations (Kump & Arthur, 1997; Berner, 2004; Montañez & Poulsen, 2013; Foster *et al.* 2017). The latitudinal distribution patterns of glacial deposits reported in this study (Fig. 12c) record three major Phanerozoic glacial periods, during Carboniferous-Permian, Late Jurassic-early Cretaceous and Cenozoic times (Frakes *et al.* 1992). The late Paleozoic icehouse is the longest-lived ice age of the Phanerozoic (Montañez & Poulsen, 2013). Ice sheets were abundant over the South pole during this period. This ice age corresponds to intense mountain uplift around tropical Pangea (Fig. 12e) which could have contributed to late Paleozoic cooling through enhanced silicate weathering (Kump *et al.* 1999; Goddérís *et al.* 2017). The high temperature gradients during the time, derived from the comprehensive global-scale compilation of lithologies (Boucot & Gray, 2001; Boucot *et al.* 2013), also indicate a cooling (Fig. 12h). The Cenozoic glaciation corresponds to remarkable topographic uplift (Fig. 12e) of the Himalaya-Tibetan Plateau (Molnar & England, 1990) contributing to atmospheric CO₂ decrease through enhanced silicate weathering and resulting high temperature gradients (Fig. 12h). Therefore, our results support that mountain uplift could have been an important factor of the two major ice ages in late Paleozoic and Cenozoic times through decreased atmospheric CO₂ and enhanced temperature gradients. Yet, other factors including organic carbon burial and outgassing through volcanism and metamorphism (e.g. Pearson *et al.*, 2000) might also contribute to the drawdown of atmospheric CO₂.

5.c. Workflow limitations

The presented workflow uses a $5^{\circ} \times 5^{\circ}$ grid of Earth's surface to bin lithology data points. By resampling the data points, we removed the spatial sampling bias. However, latitude strips at or near the equator have larger areas than the polar ones (see Figure 1 in Vihena & Smith, 2013), which leads to over-binning in high latitude strips. Ideally, an equal area gridding scheme would be created to correct spatial sampling bias in the dataset (Vilhena & Smith, 2013), and this could be considered in future work. However, we used the simpler $5^{\circ} \times 5^{\circ}$ binning so that our results are comparable to the previous study of Ziegler *et al.* (2003).

There is also a temporal sampling bias in the workflow. The reconstruction ages used to construct the climatically sensitive lithologic data to ancient geographic locations represent discrete time periods of several millions of years. Because the reconstructed paleogeographic locations of the lithologic data points change within the time range for which their age is assigned in the source data, sample locations are not truly defined by one point paleolocation, and could be represented by an average location over a given time period in future work. Providing a continuous representation of paleolatitudinal distribution patterns indicated from paleoclimate lithologic data remains an important challenge. The temporal resolution in this study does not allow us to capture some short term fluctuations which may be interpreted from more detailed records (e.g. Horton *et al.* 2010; Montañez *et al.* 2016). For instance, our workflow considers the time interval of 290-252 Ma during the Permian period as a single stage, and cannot reflect the fluctuations indicated from atmospheric CO_2 level during that time (Fig. 12a, d). Higher temporal resolution would make it possible to refine climate reconstructions.

A further improvement to our analysis could be to use paleogeographic maps rather than continental polygons to correct for the bias due to the uneven distribution of areas through time over the lithologic data. Such maps have the potential to provide a clearer definition of which areas within the continental polygons used here are likely to host certain types of lithologies associated specifically with shallow marine or terrestrial environments. However, such paleogeographic maps, such as Ronov *et al.* (1984, 1989), Scotese (2001, 2004), Golonka *et al.* (2006), Blakey (2008) and Cao *et al.* (2017), are more interpretive than continental polygons and typically do not use divisions of the geological timescale that are exactly consistent either with each other or with the time range definitions of the lithologic data considered here, and remains an outstanding problem in terms of how to obtain paleogeographic maps and lithologies at the same temporal resolution.

6. Conclusions

This study provides a framework to investigate the shifting climatic zones through deep geological time by using plate reconstructions, a comprehensive database and new data analysis approaches. We quantified the paleolatitudinal zonal patterns of climate-lithologic deposits of coals, evaporites and glacial deposits since the Devonian, with corrections for sampling- and area-biases, and used statistical methods (SiZer and HDR) to obtain

probability density functions and estimate high probability latitudinal belts. The sensitivity test of the latitudinal distributions of climate-sensitive lithologic deposits on uneven distribution of continental areas through time indicate that these climate indicators are sensitive to the correction of continental areas, and that the distribution of coals are more sensitive than evaporites and glacial deposits to this correction, particularly in Middle Devonian, Carboniferous, Early and Middle Triassic, Jurassic, Cretaceous and Early Paleogene times. The latitudinal distributions of these lithologic indicators do not show strong sensitivity to reconstruction model.

The distribution of coal paleolatitudes herein with higher temporal resolution than previous studies shifted to significantly higher latitudes at the beginning of the Permian, contrasting with previous suggestion that the shift started in the Carboniferous (Diessel 1992). The changing distribution of coals over the period from the Permian to present also cannot be considered to have been constant as proposed by Ziegler et al. (2003). Our results indicate a clearly bimodal distribution of evaporites over the past ~400 Ma, except for the Early Devonian, Early Carboniferous, the earliest Permian and Middle and Late Jurassic (Fig. 6f). This suggests that the previously proposed bimodal or unimodal evaporite patterns could have alternated over geological times. The distribution of glacial deposits is consistent with previous interpretations of the main icehouse and greenhouse periods during the last ~400 Ma.

We considered some the main factors controlling the latitudinal distributions of the lithologic deposits over time. There is no single factor that dominates the changing distributions from the Early Devonian to present. As many previous studies have noted, tectonic factors associated with Pangaea assembly and breakup and the waxing and waning of ice sheets in Gondwanaland could have been significant factors influencing the distributions of the lithologies, including the extent of tropical continental humid zones from the Late Pennsylvanian through the Early Permian. The evolution of plant groups through space and time has influenced coal distributions, especially in a significant poleward shift during the Early Permian due to the rise of Gymnosperms. Due to these many factors, care should be taken when using the latitudinal distribution of these lithologies to constrain both past global climate and the past positions of continents.

Acknowledgements

This work is supported by Australian Research Council grants ARC grants IH130200012 (RDM, SZ, SW) and ARC DE160101020 (NF). W. Cao was also supported by a University of Sydney International Scholarship (USydIS). We thank John Cannon and Michael Chin for help with GPlates and pyGPlates, Sally Cripps, Rohitash Chandra, Rob Hyndman, Stephen Marron and Maelis Arnould for assistance with programming.

Supplementary material

We provide the lithologic data of coals, evaporites and glacial deposits in shapefile format, three reconstruction models used in this study and an animation demonstrating these three models since the Devonian period, and a PDF file containing two supplementary figures. All the files can be downloaded from the link https://www.dropbox.com/sh/x1d00ooxu5tikrh/AADq_LsQcX9khzW16EPUbeO6a?dl=0. The original lithologic data in tabular format can be found from the link https://www.researchgate.net/publication/263450893_Phanerozoic_Paleoclimate_An_Atlas_of_Lithologic_Indicators_of_Climate.

Appendix. Software availability

SiZer analysis is performed in MatLab using the ‘Smoothing’ package (Chaudhuri & Marron, 1999) available at http://www.stat.unc.edu/faculty/marron/marron_software.html. SiZer also can be found in the package ‘feature’ (version 1.2.13) in ‘R’ statistical software (version 3.3.2). HDR from Hyndman (1996) is performed using the ‘hdrcde’ package (version 3.1) in ‘R’ statistical software (version 3.3.2). HEALPix method is conducted using the library ‘healpy’ (version 1.10.3) in Python 2.0 and its documentation is available at <http://healpy.readthedocs.io/en/latest/index.html>.

References

- BECQ-GIRAUDON, J.F., MONTENAT, V. & VAN DEN DRIESCHE, J. 1996. Hercynian high-altitude phenomena in the French Massif Central: tectonic implications. *Palaeogeography, Palaeoclimatology, Palaeoecology* **122**, p. 227-240.
- BERNER, R.A. 2004. *The Phanerozoic Carbon Cycle: CO₂ and O₂*. Oxford Univ Press: New York pp. 150.
- BLAKEY, R.C. 2008. Gondwana paleogeography from assembly to breakup—A 500 m.y. odyssey. *Geological Society of America Special Papers* **441**(1), 1–28.
- BOUCOT, A.J., CHEN, X. & SCOTese, C.R. 2013. *Phanerozoic paleoclimate: An atlas of lithologic indicators of climate*. SEPM Concepts in Sedimentology and Paleontology No. 11, Tulsa, Oklahoma, 1-478p.
- BOUCOT, A.J. & GRAY, J. 2001. A critique of phanerozoic climatic models involving changes in the CO₂ content of the atmosphere. *Earth-Science Reviews* **56**(1–4), 1–159.
- BRANDLEY, R.T. & KRAUSE, F.F. 1993a. Carbonate and siliciclastic deposition on a wave swept, cold-water, low-latitude ramp: Lower Carboniferous Mount Head Formation, southwestern Alberta, Canada. *Canadian Society of Petroleum Geologists, Annual Convention, Program and Abstracts, Carboniferous to Jurassic Pangea*, p. 35.
- BRANDLEY, R.T. & KRAUSE, F.F. 1993b. Thinolite-type pseudomorphs of ikaite: cold water indicators in the Mt. Head (Early Carboniferous- Viséan), Rocky Mountains, Canada. *Canadian Society of Petroleum Geologists, Annual Convention, Program and Abstracts, Carboniferous to Jurassic Pangea*, p. 34.
- BRANDLEY, R.T. & KRAUSE, F.F. 1994. Thinolite-type pseudomorphs after ikaite: indicators of cold water on the subequatorial western margin of Lower Carboniferous North America. *Canadian Society of Petroleum Geologists Memoir* **17**, p. 333-344.

- 699 CAO, W., ZAHIROVIC, S., FLAMENT, N., WILLIAMS, S., GOLONKA, J. & MÜLLER, R.D. 2017.
700 Improving global paleogeography since the late Paleozoic using paleobiology.
701 *Biogeosciences* **14**, 5425–5439.
- 702 CECIL, C.B., 1990. Palaeoclimate controls on stratigraphic repetition of chemical and
703 siliciclastic rocks. *Geology* **18**, 533–536.
- 704 CECIL, C.B., DULONG, F.T., WEST, R.R., STAMM, R., WARDLAW, B. & EDGAR, N.T. 2003.
705 Climate controls on the stratigraphy of a middle Pennsylvanian cyclothem in North
706 America. In: Cecil, C.B., Edgar, T.N. (Eds.), *Climate Controls on Stratigraphy*. Soc.
707 Econ. Paleon. Min. Spec. Pub., vol. **77**, pp. 151–180.
- 708 CHAUDHURI, P. & MARRON, J.S. 1999. SiZer for exploration of structure in curves. *Journal of*
709 *the American Statistical Association* **94**(447), 807–823.
- 710 CLEVELAND, W.S. & DEVLIN, S.J. 1988. Locally weighted regression: an approach to
711 regression analysis by local fitting. *J. Am. Stat. Assoc.*, **83**, pp. 596–610.
- 712 GORDON, W.A. 1975. Distribution by latitude of Phanerozoic evaporite deposits. *The Journal*
713 *of Geology* **83**(6), 671–684.
- 714 CRAGGS, H.J., VALDES, P.J. & WIDDOWSON, M. 2011. Climate model predictions for the
715 latest Cretaceous: An evaluation using climatically sensitive sediments as proxy
716 indicators. *Palaeogeography, Palaeoclimatology, Palaeoecology* **315–316**, 12–23.
- 717 DIESSEL, C.F.K. 1992. Coal-bearing depositional systems. *Berlin, Springer-Verlag*, 72p.
- 718 DOMEIER, M. & TORSVIK, T.H. 2014. Plate tectonics in the late Paleozoic. *Geoscience*
719 *Frontiers* **5**(3), 303–350.
- 720 EVANS, D.A.D. 2006. Proterozoic low orbital obliquity and axial-dipolar geomagnetic field
721 from evaporite palaeolatitudes. *Nature* **444**(7115), 51–55.
- 722 Frakes, L. A., Francis, J. E. & Syktus, J. I. 1992. Climate Modes of the Phanerozoic.
723 *Cambridge Univ. Press, Cambridge*.
- 724 FOSTER, G.L., ROYER, D.L. & LUNT, D.J. 2017. Future climate forcing potentially without
725 precedent in the last 420 million years. *Nature Communications* **8**, 14845.
- 726 GIBBS, M.T., REES, P.M., KUTZBACH, J.E., ZIEGLER, A.M., BEHLING, P.J. & ROWLEY, D.B.
727 2002. Simulations of Permian climate and comparisons with climate-sensitive sediments.
728 *J. Geol.* **110**, 33–55.
- 729 GODDERIS, Y., DONNADIEU, Y., CARRETIER, S., ARETZ, M., DERA, G., MACQUIN, M. &
730 REGARD, V. 2017. Onset and ending of the late Palaeozoic ice age triggered by
731 tectonically paced rock weathering. *Nature Geoscience* **10**, 382–386.
- 732 GOLONKA, J. 2007. Late Triassic and Early Jurassic palaeogeography of the world.
733 *Palaeogeography, Palaeoclimatology, Palaeoecology* **244**(1–4), 297–307.
- 734 GOLONKA, J., KROBICKI, M., PAJAK, J., GIANG, N. V. & ZUCHIEWICZ, W. 2006. *Global Plate*
735 *Tectonics and Paleogeography of Southeast Asia*, Faculty of Geology, Geophysics and
736 Environmental Protection, AGH University of Science and Technology, Arkadia,
737 Krakow, Poland, 1-128p.
- 738 GORSKI, K.M., HIVON, E., BANDAY, A.J., WANDELT, B.D., HANSEN, F.K., REINECKE, M. &
739 BARTELMAN, M. 2005. HEALPix -- a Framework for High Resolution Discretization,
740 and Fast Analysis of Data Distributed on the Sphere. *The Astrophysical Journal* **622**(2),
741 759–771.
- 742 HALLAM, A. 1985. A review of Mesozoic climates. *Journal of the Geological Society* **142**(8),

- 433–445.
- HAY, W.W. 2016. *Experimenting on a Small Planet: A History of Scientific Discoveries, a Future of Climate Change and Global Warming*. Second Edition. Springer International Publishing Switzerland, 1-819p.
- Horton, D.E., Poulsen, C.J. & Pollard, D. 2010. Influence of high-latitude vegetation feedbacks on late Paleozoic glacial cycles. *Nat. Geosci.* **3**, 572–77.
- HUNT, B.G. 1982. The impact of large variations of the Earth's obliquity on the climate. *J. Meteorol. Soc. Jpn* **60**, 309–318.
- HYNDMAN, R.J. 2009. Computing and Graphing Highest Density. *The American Statistician* **50**(2), 120–126.
- JENKINS, G.S. 2000. Global climate model high-obliquity solutions to the ancient climate puzzles of the Faint Young Sun Paradox and low-altitude Proterozoic Glaciation. *J. Geophys. Res.* **105**, 7357–7370.
- JENKINS, G.S. 2001. High-obliquity simulations for the Archean Earth: Implications for climatic conditions on early Mars. *J. Geophys. Res.* **106**, 32903–32913.
- KUMP, L.R., & ARTHUR, M.A. 1997, Global chemical erosion during the Cenozoic: weatherability balances the budgets, in Ruddiman, W.F., editor, *Tectonic Uplift and Climate Change*; New York, Plenum Press, p. 399–426.
- KUMP, L.R., ARTHUR, M.A., PATZKOWSKY, M.E., GIBBS, M.T., PINKUS, D.S. & SHEEHAN, P.M. 1999, A weathering hypothesis for glaciation at high atmospheric pCO₂ during the Late Ordovician. *Palaeogeography, Palaeoclimatology, Palaeoecology* **152**, p. 173–187.
- LOWRY, D.P., POULSEN, C.J., HORTON, D.E., TORSVIK, T.H. & POLLARD, D. 2014. Thresholds for Paleozoic ice sheet initiation. *Geology* **42**, 627–630.
- MILLER, K.B. & WEST, R.R. 1993. A Reevaluation of Wolfcampian Cyclothems in Northeastern Kansas: Significance of Aerial Exposure and Flooding Surfaces. *Kansas Geol. Surv. Bull.*, vol. **235**, pp. 1–26.
- MILLER, K.B., MCCAHOON, T.J. & WEST, R.R. 1996. Lower Permian (Wolfcampian) palaeosol-bearing cycles of the U.S. Midcontinent: evidence of climatic cyclicity. *J. Sediment. Res.* **66**, 71–84.
- MATTHEWS, K.J., MALONEY, K.T., ZAHIROVIC, S., WILLIAMS, S.E., SETON, M. & MÜLLER, R.D. 2016. Global plate boundary evolution and kinematics since the late Paleozoic. *Global and Planetary Change* **146**, 226–250.
- MCCABE, P.J. & PARRISH, J.T. 1992, Tectonic and climatic controls on the distribution and quality of Cretaceous coals, in McCabe, P.J., and Parrish, J.T., eds., *Controls on the distribution and quality of Cretaceous coals*: Geological Society of America Special Paper **267**, p. 1–15.
- Molnar, P. & England, P. 1990. Late Cenozoic uplift of mountain ranges and global climatic change: Chicken or egg? *Nature* **346**, 29–34.
- MONTAÑEZ, I.P. & POULSEN, C.J. 2013. The Late Paleozoic Ice Age: An Evolving Paradigm. *Annual Review of Earth and Planetary Sciences* **41**(1), 629–656.
- Montañez, I.P., McElwain, J.C., Poulsen, C.J., White, J.D., DiMichele, W.A., Wilson, J.P., Griggs, G. & Hren, M.T. 2016. Climate, pCO₂ and terrestrial carbon cycle linkages during late Palaeozoic glacial–interglacial cycles. *Nat. Geosci.* **9**, 824–828.

- 786 MONTAÑEZ, I.P., TABOR, N.J., NIEMEIER, D., DiMICHELE, W.A., FRANK, T.D., FIELDING, C.R.
 787 & ISBELL, J.L. 2007. CO₂-forced climate and vegetation instability during Late
 788 Palaeozoic deglaciation. *Science* **315**, 87–91.
- 789 MÜLLER, R.D., ROYER, J.Y. & LAWVER, L.A. 1993. Revised plate motions relative to the
 790 hotspots from combined Atlantic and Indian Ocean hotspot tracks. *Geology* **21**(3), 275–
 791 278.
- 792 MÜLLER, R.D., SETON, M., ZAHIROVIC, S., WILLIAMS, S.E., MATTHEWS, K.J., WRIGHT, N.M.,
 793 SHEPHARD, G.E., MALONEY, K.T., BARNETT-MOORE, N., BOWER, D.J. & CANNON, J.S.
 794 2016. Ocean basin evolution and global-scale reorganization events since Pangea
 795 breakup. *Annual Review of Earth and Planetary Science Letters* **44**, 107–138.
- 796 NELSEN, M.P., DiMICHELE, W.A., PETERS, S.E. & BOYCE, C.K. 2016. Delayed fungal
 797 evolution did not cause the Paleozoic peak in coal production. *Proceedings of the*
 798 *National Academy of Sciences* **113**(9), 2442–2447.
- 799 PARRISH, J.T. 1993. Climate of the supercontinent Pangaea. *J. Geol.* **101**, 215–233.
- 800 PARRISH, J.T., ZIEGLER, A.M. & SCOTese, C.R. 1982. RAINFALL PATTERNS AND THE
 801 DISTRIBUTION OF COALS AND EVAPORITES IN THE MESOZOIC AND
 802 CENOZOIC JUDITH. *Palaeogeography, Palaeoclimatology, Palaeoecology* **40**(1–3),
 803 31–66.
- 804 PEARSON, P. N. & PALMER, M.R. 2000. Atmospheric carbon dioxide concentrations over the
 805 past 60 million years. *Nature* **406**, 695–699.
- 806 PERLMUTTER, M.A. & MATTHEWS, M.D. 1989. Global cyclostratigraphy — a model. In:
 807 Cross, T.A. (Ed.), *Quantitative Dynamic Stratigraphy*. Prentice Hall, Englewood Cliffs,
 808 New Jersey, pp. 223–260.
- 809 PERLMUTTER, M.A. & PLOTNICK, R.E. 2003. Hemispheric asymmetry of the marine
 810 stratigraphic record: conceptual proof of a unipolar ice cap. In: Cecil, C.B., Edgar, T.N.
 811 (Eds.), *Climate Controls on Stratigraphy*. Soc. Econ. Paleont. Min. Spec. Pub., vol. 77,
 812 pp. 51–66.
- 813 PEYSER, C.E. & POULSEN, C.J. 2008. Controls on Permo-Carboniferous precipitation over
 814 tropical Pangaea: a GCM sensitivity study. *Palaeogeography, Palaeoclimatology,*
 815 *Palaeoecology* **268**(3–4), 181–192.
- 816 POULSEN, C.J., POLLARD, D., MONTAÑEZ, I.P. & ROWLEY, D. 2007. Late Palaeozoic tropical
 817 climate response to Gondwana deglaciation. *Geology* **35**, 771–774.
- 818 PRICE, G.D., SELLWOOD, B.W. & VALDES, P.J. 1995. Sedimentological evaluation of general
 819 circulation model calculations for the ‘greenhouse’ Earth: Cretaceous and Jurassic case
 820 studies. *Sedimentary Geology* **100**, 159–180.
- 821 REES, P.M., ZIEGLER, A.M., GIBBS, M.T., KUTZBACH, J.E., BEHLING, P.J. & ROWLEY, D.B.
 822 2002. Permian phytogeographic patterns and climate data/model comparisons. *J. Geol.*
 823 **110**, 1–31.
- 824 RONO, A., KHAIN, V. & SESLAVINSKY, K. 1984. Atlas of Lithological-Paleogeographical
 825 Maps of the World, Late Precambrian and Paleozoic of Continents. *U.S.S.R. Academy of*
 826 *Sciences, Leningrad*, 70 pp.
- 827 RONO, A., KHAIN, V. & BALUKHOVSKY, A. 1989. Atlas of Lithological-Paleogeographical
 828 Maps of the World, Mesozoic and Cenozoic of Continents and Oceans, *U.S.S.R.*
 829 *Academy of Sciences, Leningrad*, 77 pp.

- 830 ROWLEY, D.B., RAYMOND, A., PARRISH, J.T., LOTTES, A.L., SCOTese, C.R. & ZIEGLER, A.M.
831 1985. Carboniferous palaeogeographic, phytogeographic, and palaeoclimatic
832 reconstructions. *Int. J. Coal Geol.* **5**, 7–42.
- 833 SCOTese, C.R. 2001. Atlas of Earth History, Volume 1, Paleogeography. *PALEOMAP*
834 *Project, Arlington, Texas*, 52 pp.
- 835 SCOTese, C.R. 2004. A Continental Drift Flipbook. *The Journal of Geology* **112**(6), 729–741.
- 836 SCOTese, C.R. 2008. *The PALEOMAP Project PaleoAtlas for ArcGIS, Volume 2, Cretaceous*
837 *paleogeographic and plate tectonic reconstructions*. PALEOMAP Project, Arlington,
838 Texas.
- 839 SCOTese, C.R. & BARRETT, S.F. 1990. Gondwana's movement over the South Pole during the
840 Palaeozoic: evidence from lithological indicators of climate. *Geological Society, London,*
841 *Memoirs* **12**, 75–85.
- 842 SCOTese, C.R. & GOLONKA, J. 1992. *Paleogeographic atlas. PALEOMAP Progress Report*
843 *20-0692*. Arlington, Department of Geology, University of Texas, 34 p.
- 844 Soreghan, G.S., 1997. Walther's Law, climate change, and Upper Paleozoic cyclostratigraphy
845 in the Ancestral Rocky Mountains. *Geology* **67**, 1001–1004.
- 846 SWANSON-HYSELL, N.L. & MACDONALD, F.A. 2017. Tropical weathering of the Taconic
847 orogeny as a driver for Ordovician cooling. *Geology* **45**(8), 719–722.
- 848 TABOR, N.J. & POULSEN, C.J. 2008. Palaeoclimate across the Late Pennsylvanian-Early
849 Permian tropical palaeolatitudes: A review of climate indicators, their distribution, and
850 relation to palaeophysiographic climate factors. *Palaeogeography, Palaeoclimatology,*
851 *Palaeoecology* **268**(3–4), 293–310.
- 852 TABOR, N.J. & MONTAÑEZ, I.P. 2004. Permo-Pennsylvanian alluvial palaeosols (north-
853 central Texas): high-resolution indicator records of the evolution of early Pangaeon
854 palaeoclimate. *Sedimentology* **51**, 851–884.
- 855 TABOR, N.J., MONTAÑEZ, I.P., SCOTese, C.R., POULSEN, C.J. & MACK, G.H. 2008. Paleosol
856 archives of environmental and climatic history in paleotropical western Pangea during
857 the latest Pennsylvanian through Early Permian, in Fielding, C.R., Frank, T.D., and Isbell,
858 J.L., eds., *Resolving the Late Paleozoic Ice Age in Time and Space*: Geological Society of
859 America Special Paper 441, p. 291–303.
- 860 VAN HINSBERGEN, D.J.J., DE GROOT, L. V., VAN SCHAIK, S.J., SPAKMAN, W., BIJL, P.K.,
861 SLUIJS, A., LANGEREIS, C.G. & BRINKHUIS, H. 2015. A paleolatitude calculator for
862 paleoclimate studies. *PLoS ONE* **10**(6), 1–21.
- 863 VILHENA, D.A. & SMITH, A.B. 2013. Spatial Bias in the Marine Fossil Record. *PLoS ONE*
864 **8**(10), e74470.
- 865 Walker, L.J., Wilkinson, B.H. & Ivany, L.C. 2002, Continental drift and Phanerozoic
866 carbonate accumulation in shallow shelf and deep marine settings. *The Journal of*
867 *Geology* **110**, 75–88.
- 868 Warren, J.K. 2010. Evaporites through time: Tectonic, climatic and eustatic controls in
869 marine and nonmarine deposits. *Earth-Science Reviews* **98**, 217–268.
- 870 WITZKE, B.J. 1990. Palaeoclimatic constraints for palaeozoic Palaeolatitudes of Laurentia and
871 Euramerica. *Geological Society, London, Memoirs* **12**(1), 57–73.

WRIGHT, N., ZAHIROVIC, S., MÜLLER, R.D. & SETON, M. 2013. Towards community-driven paleogeographic reconstructions: Integrating open-access paleogeographic and paleobiology data with plate tectonics. *Biogeosciences* **10**(3), 1529–1541.

ZIEGLER, A.M. 1990. Phytogeographic patterns and continental configurations during the Permian Period. In: McKerrow, W.S., Scotese, C.R. (Eds.), *Palaeozoic Palaeogeography and Biogeography*. Geol. Soc. Mem., vol. 12, pp. 363–379.

ZIEGLER, A.M., ESHEL, G., MCALLISTER REES, P., ROTHFUS, T.A., ROWLEY, D.B. & SUNDERLIN, D. 2003. Tracing the tropics across land and sea: Permian to present. *Lethaia* **36**, 227–254.

ZIEGLER, A.M., HANSEN, K.S., JOHNSON, M.E., KELLY, M.A., SCOTese, C.R. & VAN DER VOO, R. 1977. Silurian continental distribution, palaeogeography, climatology, and biogeography. *Tectonophysics* **40**, 13–51.

ZIEGLER, A.M., HULVER, M.L. & ROWLEY, D.B. 1997. Permian world topography and climate. In: Martini, I.P. (Ed.), *Late Glacial and Post-Glacial Environmental Changes—Quaternary, Carboniferous–Permian and Proterozoic*. Oxford University Press, Oxford, pp. 111–146.

ZIEGLER, A.M., RAYMOND, A., GEIRLOWSKI, T.C., HORRELL, M.A., ROWLEY, D.B. & LOTTES, A.L. 1987. Coal, climate and terrestrial productivity: the present and early Cretaceous compared. In: Scott, A.C. (Ed.), *Coal and Coal-bearing Strata: Recent Advances*. Geol. Soc. London Spec. Pap., vol. 32, pp. 25–49.

ZIEGLER, A.M., SCOTese, C.R. & BARRETT, S.F. 1983. Mesozoic and Cenozoic Paleogeographic Maps. *Tidal Friction and the Earth's Rotation II*, 240–252.

ZIEGLER, A. M., SCOTese, C. R., MCKERROW, W. S., JOHNSON, M. E. & BAMBACH, R.K. 1979. Paleozoic paleogeography. *Annual Review of Earth and Planetary Sciences* **7**, 473–502.

Table 1. Time division of the climate-sensitive lithologic indicators of Boucot *et al.* (2013) using the 2016 time scale of the International Commission on Stratigraphy (ICS2016). The reconstruction time is the rounded-off middle age of each time interval.

916

Era	Period	Epoch/Age	Start (Ma)	End (Ma)	Reconstruction Time (Ma)
Cenozoic	Neogene	Miocene	23.0	5.3	14
		Paleogene	33.9	23.0	28
	Paleogene	Middle and Late Eocene (Lutetian-Priabonian)	47.8	33.9	41
		Early Eocene (Ypresian)	56.0	47.8	52
		Paleocene	66.0	56.0	61
Mesozoic	Cretaceous	Late Cretaceous (Coniacian-Maastrichtian)	89.8	66.0	78
		Late Cretaceous (Albian-Turonian)	113.0	89.8	101
		Early Cretaceous (Berriasian-Aptian)	145.0	113.0	129
	Jurassic	Late Jurassic	164	145	155
		Early and Middle Jurassic	201	164	183
	Triassic	Late Triassic	237	201	219
		Middle Triassic	247	237	242
		Early Triassic	252	247	250
	Permian	Middle-Late Permian (Artinskian-Lopingian)	290	252	271
		Early Permian (Asselian-Sakmarian)	299	290	295
Late Paleozoic	Carboniferous	Late Carboniferous (Kasimovian-Gzhelian)	307	299	303
		Late Carboniferous (Bashkirian-Moscovian)	323	307	315
		Early Carboniferous (Serpukhovian)	331	323	327
		Early Carboniferous (Tournaisian-Visean)	359	331	345
	Devonian	Late Devonian	383	359	371
		Middle Devonian (Givetian)	388	383	386
		Middle Devonian (Eifelian)	393	388	391
		Early Devonian	419	393	406

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

Captions:

941

942

Figure 1. (Colour online) Locations of climate-sensitive lithologic data reconstructed using the plate motion model of Matthews et al. (2016): coals (green), evaporites (red) and glacial

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

deposits (blue) since the Early Devonian (Boucot *et al.* 2013; Ziegler *et al.* 2003). Ages on the top of each subplot are reconstruction times according to the time scale described in Table 1.

Figure 2. (a, b, c) Sum of data points of coals, evaporites and glacial deposits at each time interval since the Devonian period. (d, e, f) Sum of resampled data points for coals, evaporites and glacial deposits at each time interval since the Devonian period.

Figure 3. (Colour online) (a) Present-day locations of modern global peats (green points) from Ziegler *et al.* (2003). Grey polygons are modern terrane geometries. (b) Distribution of modern peat points binned using a 5°x5° mesh of the global surface. (c) Distribution of resampled modern peat points (red: bins containing at least one data point; blue: bins containing no data points).

Figure 4. (a) Number of original modern peat points from Ziegler *et al.* (2003) in each 5° latitudinal strip. (b) Number of resampled modern peat points in each 5° latitudinal strip. (c) Modern continental area in each 5° latitudinal strip as a percentage of total modern continental area. (d) Distribution of resampled modern peat points with continental area bias corrected. (e) Distribution of modern peat points with sampling- and area-biases corrected. The two hemispheres are combined to form a symmetric distribution pattern (also known as zonal pattern; Scotese & Barrett, 1990; Ziegler *et al.* 2003). (f) Fitted probability density function (black line) using the method of SiZer from Chaudhuri & Marron (1999). High-density latitude ranges at confidence intervals of 50% (grey area) and 95% (light grey area) using the method of Highest Density Regions from Hyndman (1996).

Figure 5. (a) Numbers of original coal points of Boucot *et al.* (2013) and Ziegler *et al.* (2003) reconstructed using the tectonic model of Matthews *et al.* (2016) in each 5° latitudinal strip for each time interval since the Devonian period. (b) Number of resampled coal points in each 5° latitudinal strip for each time interval. (c) Continental area in each 5° latitudinal strip as a percentage of Earth's surface for each time interval. (d) Distribution of coals with continental area corrected in each 5° latitudinal strip for each time interval. (e) Symmetric zonal pattern of coals with sampling- and continental area-biases removed. (f) Probability density function of symmetric zonal pattern of coals for each time interval. The cumulative results for all times are presented on the rightmost side with bold borders.

Figure 6. (a) Number of original evaporate points of Boucot *et al.* (2013) and Ziegler *et al.* (2003), reconstructed using the tectonic model of Matthews *et al.* (2016), in each 5° latitudinal strip for each time interval since the Devonian period. (b) Number of resampled evaporate points in each 5° latitudinal strip for each time interval. (c) Continental area in each 5° latitudinal strip as a percentage of total continental area for each time interval. (d) Distribution of evaporites with continental area corrected in each 5° latitudinal strip for each time interval. (e) Zonal pattern of evaporites with sampling- and continental area-biases corrected. (f) Probability density function of zonal pattern of evaporites for each time

interval. The cumulative results for all times are presented on the rightmost side with bold borders.

Figure 7. (a) Numbers of original glacial deposit points of Boucot *et al.* (2013) and Ziegler *et al.* (2003) reconstructed using the tectonic model of Matthews *et al.* (2016) in each 5° latitudinal strip for each time interval since the Devonian period. (b) Number of resampled glacial deposit points in each 5° latitudinal strip for each time interval. (c) Continental area in each 5° latitudinal strip as a percentage of total continental area for each time interval. (d) Distribution of glacial deposit with continental area corrected in each 5° latitudinal strip for each time interval. (e) Zonal pattern of glacial deposit with sampling- and continental area-biases corrected. (f) Probability density function of zonal pattern of glacial deposits for each time interval. The troughs with a question mark indicate records unlikely to represent latitudinal changes in climate (see text). The cumulative results for all times are presented on the rightmost side with bold borders.

Figure 8. (Colour online) Continental area in the northern and southern hemispheres respectively as a percentage of total continental area at each time interval since the Devonian period.

Figure 9. (Colour online) Zonal patterns of lithologic deposits of coals (a), evaporites (b) and glacial deposits (c), with continental area bias correction (blue) and without continental area bias correction (red), for each time interval since the Devonian period using the plate motion model of Matthews *et al.* (2016). The number of original data points at each time interval are reported at the bottom of each subplot.

Figure 10. (Colour online) Latitudinal distribution of coals (a), evaporites (b) and glacial deposits (c), for each time interval since the Devonian period, using three reconstruction models (blue: Matthews *et al.* (2016); red: Scotese (2008); green: Golonka (2007)). The number of original data points at each time interval are present on the bottom of each subplot.

Figure 11. Distribution patterns of coals, reconstructed using the plate motion model of Matthews *et al.* (2016), for each time interval since the Devonian period using three different bin sizes (red: 10° (a); blue: 5° (b); green: 2° (c)) to bin data points in order to remove the sampling bias. The number of original data points at each time interval are reported at the bottom of each subplot.

Figure 12. (Colour online) (a, b, c) Paleolatitudinal high-density latitudinal belts of coals, evaporites and glacial deposits, with 50% (blue area) and 95% (light grey area) confidence intervals for each time interval since the Devonian period. The blue lines are the most significant peaks of latitudinal distribution. The areas marked with a question mark indicate records of limited use for climate interpretation. Cumulative high-density probabilities are presented on the rightmost side of Figure 12a, b, c with bold borders. (d) Atmospheric CO₂ concentration curve since the Devonian period derived from Foster *et al.* (2017). 68% and 95% confidence intervals are shown as dark and light grey bands. The blue line represents a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

locally weighted scatterplot smoothing (LOESS) fit, a nonparametric regression fit from Cleveland & Devlin (1988), through the data. Icehouse intervals and greenhouse intervals (Hay, 2016) are indicated by a blue band and a white band at the top of the panel, respectively on the top. (e) Global paleolatitudinal distribution of mountain ranges since the Devonian period calculated from the set of paleogeographic maps (Golonka *et al.* 2006; Cao *et al.* 2017). (f) Global paleolatitudinal distribution of shallow marine environments (Golonka *et al.* 2006; Cao *et al.* 2017). (g) Global paleolatitudinal distribution of ice sheets from the paleogeographic maps (Golonka *et al.* 2006; Cao *et al.* 2017). (h) Climate temperature gradients since the Early Devonian derived from Boucot *et al.* (2013).

Figure 13. (Colour online) Global paleogeography in the Late Carboniferous from Cao et al. (2017) showing the location of the Central Pangaeian Mountains (CPMs). Black toothed lines indicate subduction zones, and other black lines denote middle-ocean ridges and transforms. Grey outlines delineate reconstructed present-day coastlines and terranes. Mollweide projection with 0°E central meridian.

Figure 1. (Colour online) Locations of climate-sensitive lithologic data reconstructed using the plate motion model of Matthews et al. (2016): coals (green), evaporites (red) and glacial deposits (blue) since the Early Devonian (Boucot et al. 2013; Ziegler et al. 2003). Ages on the top of each subplot are reconstruction times according to the time scale described in Table 1.

154x142mm (600 x 600 DPI)

Figure 2. (a, b, c) Sum of data points of coals, evaporites and glacial deposits at each time interval since the Devonian period. (d, e, f) Sum of resampled data points for coals, evaporites and glacial deposits at each time interval since the Devonian period.

72x30mm (300 x 300 DPI)

Figure 3. (Colour online) (a) Present-day locations of modern global peats (green points) from Ziegler et al. (2003). Grey polygons are modern terrane geometries. (b) Distribution of modern peat points binned using a 5°x5° mesh of the global surface. (c) Distribution of resampled modern peat points (red: bins containing at least one data point; blue: bins containing no data points).

123x191mm (600 x 600 DPI)

Figure 4. (a) Number of original modern peat points from Ziegler et al. (2003) in each 5° latitudinal strip. (b) Number of resampled modern peat points in each 5° latitudinal strip. (c) Modern continental area in each 5° latitudinal strip as a percentage of total modern continental area. (d) Distribution of resampled modern peat points with continental area bias corrected. (e) Distribution of modern peat points with sampling- and area-biases corrected. The two hemispheres are combined to form a symmetric distribution pattern (also known as zonal pattern; Scotese & Barrett, 1990; Ziegler et al. 2003). (f) Fitted probability density function (black line) using the method of SiZer from Chaudhuri & Marron (1999). High-density latitude ranges at confidence intervals of 50% (grey area) and 95% (light grey area) using the method of Highest Density Regions from Hyndman (1996).

178x400mm (300 x 300 DPI)

Figure 5. (a) Numbers of original coal points of Boucot et al. (2013) and Ziegler et al. (2003) reconstructed using the tectonic model of Matthews et al. (2016) in each 5° latitudinal strip for each time interval since the Devonian period. (b) Number of resampled coal points in each 5° latitudinal strip for each time interval. (c) Continental area in each 5° latitudinal strip as a percentage of Earth's surface for each time interval. (d) Distribution of coals with continental area corrected in each 5° latitudinal strip for each time interval. (e) Symmetric zonal pattern of coals with sampling- and continental area-biases removed. (f) Probability density function of symmetric zonal pattern of coals for each time interval. The cumulative results for all times are presented on the rightmost side with bold borders.

138x114mm (600 x 600 DPI)

Figure 6. (a) Number of original evaporate points of Boucot et al. (2013) and Ziegler et al. (2003), reconstructed using the tectonic model of Matthews et al. (2016), in each 5° latitudinal strip for each time interval since the Devonian period. (b) Number of resampled evaporate points in each 5° latitudinal strip for each time interval. (c) Continental area in each 5° latitudinal strip as a percentage of total continental area for each time interval. (d) Distribution of evaporites with continental area corrected in each 5° latitudinal strip for each time interval. (e) Zonal pattern of evaporites with sampling- and continental area-biases corrected. (f) Probability density function of zonal pattern of evaporites for each time interval. The cumulative results for all times are presented on the rightmost side with bold borders.

138x114mm (600 x 600 DPI)

Figure 7. (a) Numbers of original glacial deposit points of Boucot et al. (2013) and Ziegler et al. (2003) reconstructed using the tectonic model of Matthews et al. (2016) in each 5° latitudinal strip for each time interval since the Devonian period. (b) Number of resampled glacial deposit points in each 5° latitudinal strip for each time interval. (c) Continental area in each 5° latitudinal strip as a percentage of total continental area for each time interval. (d) Distribution of glacial deposit with continental area corrected in each 5° latitudinal strip for each time interval. (e) Zonal pattern of glacial deposit with sampling- and continental area-biases corrected. (f) Probability density function of zonal pattern of glacial deposits for each time interval. The troughs with a question mark indicate records unlikely to represent latitudinal changes in climate (see text). The cumulative results for all times are presented on the rightmost side with bold borders.

137x112mm (600 x 600 DPI)

Figure 8. (Colour online) Continental area in the northern and southern hemispheres respectively as a percentage of total continental area at each time interval since the Devonian period.

43x24mm (600 x 600 DPI)

Figure 9. (Colour online) Zonal patterns of lithologic deposits of coals (a), evaporites (b) and glacial deposits (c), with continental area bias correction (blue) and without continental area bias correction (red), for each time interval since the Devonian period using the plate motion model of Matthews et al. (2016). The number of original data points at each time interval are reported at the bottom of each subplot.

78x36mm (600 x 600 DPI)

Figure 10. (Colour online) Latitudinal distribution of coals (a), evaporites (b) and glacial deposits (c), for each time interval since the Devonian period, using three reconstruction models (blue: Matthews et al. (2016); red: Scotese (2008); green: Golonka (2007)). The number of original data points at each time interval are present on the bottom of each subplot.

79x37mm (600 x 600 DPI)

Figure 11. Distribution patterns of coals, reconstructed using the plate motion model of Matthews et al. (2016), for each time interval since the Devonian period using three different bin sizes (red: 10° (a); blue: 5° (b); green: 2° (c)) to bin data points to in order to remove the sampling bias. The number of original data points at each time interval are reported at the bottom of each subplot.

77x35mm (600 x 600 DPI)

Figure 12. (Colour online) (a, b, c) Paleolatitudinal high-density latitudinal belts of coals, evaporites and glacial deposits, with 50% (blue area) and 95% (light grey area) confidence intervals for each time interval since the Devonian period. The blue lines are the most significant peaks of latitudinal distribution. The areas marked with a question mark indicate records of limited use for climate interpretation. Cumulative high-density probabilities are presented on the rightmost side of Figure 12a, b, c with bold borders. (d) Atmospheric CO₂ concentration curve since the Devonian period derived from Foster et al. (2017). 68% and 95% confidence intervals are shown as dark and light grey bands. The blue line represents a locally weighted scatterplot smoothing (LOESS) fit, a nonparametric regression fit from Cleveland & Devlin (1988), through the data. Icehouse intervals and greenhouse intervals (Hay, 2016) are indicated by a blue band and a white band at the top of the panel, respectively on the top. (e) Global paleolatitudinal distribution of mountain ranges since the Devonian period calculated from the set of paleogeographic maps (Golonka et al. 2006; Cao et al. 2017). (f) Global paleolatitudinal distribution of shallow marine environments (Golonka et al. 2006; Cao et al. 2017). (g) Global paleolatitudinal distribution of ice sheets from the paleogeographic maps (Golonka et al. 2006; Cao et al. 2017). (h) Climate temperature gradients since the Early Devonian derived from Boucot et al. (2013).

145x125mm (600 x 600 DPI)

Figure 13. (Colour online) Global paleogeography in the Late Carboniferous from Cao et al. (2017) showing the location of the Central Pangaeian Mountains (CPMs). Black toothed lines indicate subduction zones, and other black lines denote middle-ocean ridges and transforms. Grey outlines delineate reconstructed present-day coastlines and terranes. Mollweide projection with 0°E central meridian.

42x22mm (600 x 600 DPI)