UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers:
Part B

Faculty of Science, Medicine and Health

2019

Alkaloids and styryllactones from *Goniothalamus cheliensis*

Wuttichai Jaidee

Mae Fah Luang University

Raymond J. Andersen

University of British Columbia

Brian O. Patrick

University of British Columbia

Stephen G. Pyne

University of Wollongong, spyne@uow.edu.au

Chatchai Muanprasat

Mahidol University

See next page for additional authors

Publication Details

Jaidee, W., Andersen, R. J., Patrick, B. O., Pyne, S. G., Muanprasat, C., Borwornpinyo, S. & Laphookhieo, S. (2019). Alkaloids and styryllactones from *Goniothalamus cheliensis*. *Phytochemistry*, 157 8-20.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Alkaloids and styryllactones from *Goniiothalamus cheliensis*

Abstract

Eight previously undescribed compounds, including four alkaloids and five styryllactones together with 36 known compounds were isolated from the twig and leaf extracts of *Goniiothalamus cheliensis*. Their structures were elucidated by extensive analysis of their spectroscopic data. The absolute configuration of (–)-(4S,5S,6R,7S,8S)-goniochelenlactone and (–)-(4S,5S,6R,7S,8S)-7-acetylgoniochelenlactone were established from single crystal X-ray analysis using Cu K α radiation. The absolute configurations of the other related compounds were identified by comparisons of their ECD spectra with those of related known compounds. Most of the isolated compounds were evaluated for their cytotoxicities against human colorectal cancer cells (HCT-116). Griffithazanone A was the most potent with an IC₅₀ value of 2.39 μ M.

Publication Details

Jaidee, W., Andersen, R. J., Patrick, B. O., Pyne, S. G., Muanprasat, C., Borwornpinyo, S. & Laphookhieo, S. (2019). Alkaloids and styryllactones from *Goniiothalamus cheliensis*. *Phytochemistry*, 157 8-20.

Authors

Wuttichai Jaidee, Raymond J. Andersen, Brian O. Patrick, Stephen G. Pyne, Chatchai Muanprasat, Suparerk Borwornpinyo, and Surat Laphookhieo

Alkaloids and styryllactones from *Goniothalamus cheliensis*

Wuttichai Jaidee^{a,b}, Raymond J. Andersen^c, Brian O. Patrick^c, Stephen G. Pyne^d,
Chatchai Muanprasat^{e,f}, Suparerk Borwornpinyo^g, Surat Laphookhieo^{a,b,*}

^a Center of Chemical Innovation for Sustainability (CIS), Mae Fah Luang University,
Chiang Rai 57100, Thailand

^b School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

^c Department of Chemistry, University of British Columbia, 2036 Main Mall,
Vancouver, BC, Canada V6T 1Z1

^d School of Chemistry, University of Wollongong, Wollongong, New South Wales,
2522, Australia

^e Department of Physiology, Faculty of Science, Mahidol University, Rajathevi,
Bangkok 10400, Thailand

^f Excellent Center for Drug Discovery, Faculty of Science, Mahidol University,
Rajathevi, Bangkok 10400, Thailand

^g Department of Biotechnology, Faculty of Science, Mahidol University, Rajathevi,
Bangkok 10400, Thailand

* Corresponding author.

E-mail address: surat.lap@mfu.ac.th (S. Laphookhieo)

ABSTRACT

Eight previously undescribed compounds, including four alkaloids and five styryllactones together with 36 known compounds were isolated from the twig and leaf extracts of *Goniothalamus cheliensis*. Their structures were elucidated by extensive analysis of their spectroscopic data. The absolute configuration of (–)-(4*S*,5*S*,6*R*,7*S*,8*S*)-goniochelienlactone and (–)-(4*S*,5*S*,6*R*,7*S*,8*S*)-7-acetylgoniochelienlactone were established from single crystal X-ray analysis using Cu K α radiation. The absolute configurations of the other related compounds were identified by comparisons of their ECD spectra with those of related known compounds. Most of the isolated compounds were evaluated for their cytotoxicities against human colorectal cancer cells (HCT-116). Griffithazanone A was the most potent with an IC₅₀ value of 2.39 μ M.

Keywords: *Goniothalamus cheliensis*, Annonaceae, Alkaloids, Styryllactones

1. Introduction

Goniothalamus is the one of the largest genera of the family Annonaceae. It is distributed throughout the world and is found predominately in Southeast Asia (Duc et al., 2016). This genus includes about 160 species, of which 25 are found in Thailand (Saunders et al., 2008). Some species have been used as traditional medicines for the treatment of fever (Surivet and Vatele, 1999), scabies (Heyne, 1950), edema, rheumatism (Jiang et al., 2011), tympanites (Quisumbing, 1951), and typhoid fever (Efendi et al., 2010). *Goniothalamus cheliensis* Hu is a large green tree which is distributed throughout the forested slopes at 1500 m, in South Yunnan, China (Saunders et al., 2008). This plant is found in the Northern parts of Thailand, especially in Chiang Rai and Nan Provinces. Previous phytochemical investigations of *G. cheliensis*, resulted in the isolation and identification of various types of compounds, including styryllactones (Jiang et al., 2011; Wang et al., 2002; Zhu et al., 2012, Duc et al., 2016; Wang et al., 2003, Si et al., 2002), alkaloids (Jiang et al., 2008; Duc et al., 2016), flavonoids (Wang et al., 2003, Sun et al., 2014), and acetogenins (Duc et al., 2016). Many of these isolated compounds exhibited a variety of biological activities such as antiproliferative (Sun et al., 2014), antifungal (Duc et al., 2016), antimalarial (Duc et al., 2016) and cytotoxic activities (Wang et al., 2002; Zhu et al., 2012). In this study, we describe the isolation and identification of eight previously undescribed compounds (**1–3** and **5–9**) and 36 known compounds (**4** and **10–44**) (Fig. 1) from the twig and leaf extracts of *G. cheliensis* collected from Doi Tung, Chiang Rai Province, Thailand. Their cytotoxicities against the colon cancer cell line HCT-116 are also reported.

2. Results and Discussion

The acetone extracts of the leaves and twigs of *G. cheliensis* were individually subjected to separation by silica gel and Sephadex LH-20 column chromatography (CC) followed by semipreparative RP-HPLC to afford eight previously undescribed compounds (**1–3** and **5–9**) and 36 known compounds (**4** and **10–44**). The known compounds were identified as 3-methyl-1*H*-benz[*f*]indole-4,9-dione (**4**) (Efendi et al., 2010), (–)-goniobutenolide B (**10**) (Xu et al., 1994), 7-*epi*-(–)-goniobutenolide B (**11**) (Popsavin et al., 2012), (+)-goniodiol (**12**) (Fang et al., 1991), goniodiol-8-monoacetate (**13**) (Wu et al., 1992), (+)-7-*O*-acetylgoniodiol (**14**) (Kampong et al., 2013), 8-acetoxy goniofufurone (**15**) (Zhang et al., 1998), isoaltholactone (**16**) (Colegate et al., 1990), (+)-glaberide I (**17**) (Wu et al., 2013), (–)-glaberide I (**18**) (Chang et al., 2015), (+)-syringaresinol (**19**) (Rahman et al., 2007; Yamauchi et al., 2015), (+)-medioresinol (**20**) (Nakasone et al., 1996; Yamauchi et al., 2015), (+)-episyringaresinol (**21**) (Chen et al., 1998), (–)-syringaresinol (**22**) (Rahman et al., 2007; Yamauchi et al., 2015), (–)-episyringaresinol (**23**) (Lui et al., 2015), (–)-pinoresinol (**24**) (Ming et al., 2001; Yamauchi et al., 2015), griffithazanone A (**25**) (Zhang et al., 1999), cleistopholine (**26**) (Waterman and Muhammad, 1985), vanillic acid (**27**) (Wu et al., 2000), *p*-hydroxybenzoic acid (**28**) (Rukachaisirikul et al., 2010), *p*-methylbenzoic acid (**29**) (Zhou et al., 2013), 4-hydroxy-3-methoxypropiophenone (**30**) (Ito et al., 2001), 3,5-dimethoxy-4-hydroxypropiophenone (**31**) (Zdero et al., 1986), *p*-hydroxybenzaldehyde (**32**) (Zhou et al., 2013), ethyl-4-hydroxybenzoate (**33**)

(Fan et al., 2016), *trans*-ferulic acid (**34**) (Salum et al., 2010), (–)-(3*R*)-mellein methyl ether (**35**) (Li et al., 2017), derrusin (**36**) (East et al., 1969), 5-hydroxy-7-methoxy-3',4'-methylene dioxy isoflavone (**37**) (Liping et al., 1996), derrustone (**38**) (Singh and Muthukrishnan, 2005), robustone methyl ether (**39**) (East et al., 1969), derrugenin (**40**) (Ortega et al., 2015), robustigenin (**41**) (Pancharoen et al., 2008), methyl-BRM-5 (**42**) (Tahara et al., 1986), *trans*-*N*-coumaroyltyramine (**43**) (Holzbach and Lopes, 2010), and *N*-*trans*-feruloyl tyramine (**44**) (Holzbach and Lopes, 2010) by comparison of their spectroscopic data with those published.

Compound **1**, goniochelienic acid A, was obtained as yellow needles. The core structure of compound **1** was a benzo[*f*]indole-4,9-dione (Efendi et al., 2010) which was confirmed by a single-crystal X-ray diffraction analysis using Mo K α radiation (Fig. 5). The HRESIMS spectrum showed a [M + Na]⁺ ion peak at *m/z* 294.0379 (calcd. for C₁₄H₉NO₅Na, 294.0378) which also supported this structural assignment. The ¹H NMR spectrum of **1** (Table 1) showed resonances for a hydroxy proton of a carboxylic acid [δ_{H} 13.68 (COOH)], a H-bonded hydroxy proton [δ_{H} 12.20 (1H, s, 8-OH)], aromatic protons of an ABC spin system [δ_{H} 7.82 (1H, d, *J* = 7.4 Hz, H-5)/ δ_{C} 120.7, 7.64 (1H, dd, *J* = 7.8, 7.4 Hz, H-6)/ δ_{C} 136.0, 7.32 (1H, d, *J* = 7.8 Hz, H-7)/ δ_{C} 126.1], an olefinic proton [δ_{H} 7.74 (1H, s, H-2)/ δ_{C} 138.0], and a *N*-methyl group [δ_{H} 4.15 (3H, s, *N*-CH₃)/ δ_{C} 37.5]. The structure of **1** was further supported by the following key HMBC correlations (Fig. 3): δ_{H} 12.20 (8-OH) with C-7 (δ_{C} 126.1), C-8 (δ_{C} 162.5), and C-8a (δ_{C} 114.5); δ_{H} 7.82 (H-5) with C-4 (δ_{C} 183.1) and C-6 (δ_{C} 136.0); δ_{H} 4.15 (*N*-Me) with C-9a (δ_{C} 130.9), C-2 (δ_{C} 138.0); δ_{H} 7.74 (H-2) with C-3 (δ_{C} 114.5), C-3a (δ_{C} 123.5), C-9a (δ_{C} 130.9), and C-10 (δ_{C} 161.5). From these data, the structure of compound **1** was assigned as 8-hydroxyl-1-methyl-4,9-dioxy-1*H*-benzo[*f*]indole-3-carboxylic acid. Full assignments of the ¹H and ¹³C NMR spectroscopic data are shown in Table 1.

Compound **2**, goniochelienic acid B, was isolated as a yellow powder. Its molecular formula, C₁₄H₉NO₄, was established by the HRESIMS which showed a [M + Na]⁺ ion peak at *m/z* 278.0436 (calcd. for C₁₄H₉NO₄Na, 278.0429). The ¹H and ¹³C NMR spectroscopic data (Table 1) of **2** were similar to those of **1**. The main difference between these two compounds was that the H-bonded proton at δ_{H} 12.22 (OH-8) was not observed in the ¹H NMR spectrum of **2**. Compound **2** showed ¹H NMR resonances for four aromatic protons, that formed part of a 1,2-disubstituted aromatic ring, at δ_{H} 8.26 (1H, dd, *J* = 1.3, 7.4 Hz, H-5), 7.79 (1H, td, *J* = 1.5, 7.5 Hz, H-6), 7.81 (1H, td, *J* = 1.5, 7.5 Hz, H-7), 8.21 (1H, dd, *J* = 7.4, 1.3 Hz, H-8) instead of the ABC spin system observed in the ¹H NMR spectrum of compound **1**. Finally, the structure of **2** was supported with the following key HMBC correlations (Fig. 3): δ_{H} 8.26 (H-5) with C-4 (δ_{C} 184.0) and C-6 (δ_{C} 134.5); δ_{H} 4.16 (*N*-Me) with C-9a (δ_{C} 131.8), C-2 (δ_{C} 137.5); δ_{H} 7.73 (H-2) with C-3 (δ_{C} 116.4), C-3a (δ_{C} 123.1), C-9a (δ_{C} 131.8), and C-10 (δ_{C} 162.1). Therefore, compound **2** was assigned as 1-methyl-4,9-dioxy-1*H*-benzo[*f*]indole-3-carboxylic acid.

Compound **3**, methyl goniochelienate, was isolated as a yellow amorphous powder. The molecular formula C₁₄H₉NO₄ was deduced from HRESIMS analysis which showed a [M + Na]⁺ at *m/z* 278.0435 (calcd for C₁₄H₉NO₄Na, 278.0429). The

¹H NMR spectroscopic data of **3** (Table 1) were similar to those of **4**; whose structure was determined by single crystal X-ray structural analysis (Fig. 6). The difference between compounds **3** and **4** is that **3** has a C-3 methoxy carbonyl substituent (δ_{H} 3.89) and **4** has a C-3 methyl group. The position of the methoxy group in **3** was confirmed by the HMBC correlation from δ_{H} 3.89 (3-CO₂Me) with C-10 (δ_{C} 163.2) (Fig. 3). The other key HMBC correlations were shown in Fig. 3. Therefore, the structure of **3** was identified as methyl 4,9-dioxo-1*H*-benz[*f*]indole-3-carboxylate.

Compound **5** was obtained as a yellow solid. The molecular formula of **5** was established as C₁₃H₁₁NO₅ based on the HRESIMS (m/z 284.0530 [M + Na]⁺, calcd for C₁₃H₁₁NO₅Na 284.0535). The ¹H and ¹³C NMR spectra of **5** in CDCl₃ solution shown resonances characteristic of a naphthoquinone derivative (Table 2), which were similar to those of goniothalamine B (**45**), previously isolated from *Goniothalamus scortechinii* (Prawat et al., 2012). The significant difference between these two compounds was observed for the chemical shift of H-7/C-7 and H-8/C-8. Compound **5** displayed these resonances at δ_{H} 7.05 (1H, d, J = 8.4 Hz, H-7)/ δ_{C} 113.2 and 7.75 (1H, d, J = 8.4 Hz, H-8)/ δ_{C} 121.0, whereas for goniothalamine B these appeared at δ_{H} 7.22 (1H, d, J = 8.4 Hz, H-7)/ δ_{C} 117.8 and δ_{H} 7.76 (1H, d, J = 8.4 Hz, H-8)/ δ_{C} 120.7 (Table 2). In addition, the chemical shifts for C-2, C-3 and C-6 were also different; δ_{C} 106.9 (C-2) and δ_{C} 158.4 (C-3) 155.8 (C-6) for compound **5** and δ_{C} 109.4 (C-2), δ_{C} 152.5 (C-3) and δ_{C} 152.1 (C-6) for goniothalamine B. These data indicated that compound **5** was a structural isomer of goniothalamine B. To help further assign all ¹H NMR resonances as well as to confirm the location of amino group, the NMR spectra of compound **5** were also recorded in DMSO-*d*₆. The ¹H NMR spectrum in DMSO-*d*₆ (Table 2) showed additional downfield resonances at δ_{H} 9.12 (1H, br s) and δ_{H} 10.68 (1H, br s) assignable to the primary amine protons. Finally, the structure was further confirmed from the following HMBC correlations: δ_{H} 10.68 (2-NH₂) with C-1 (δ_{C} 178.1); δ_{H} 3.91 (6-OMe) with C-6 (δ_{C} 155.8); δ_{H} 13.94 (5-OH) with C-6 (δ_{C} 155.8), C-5 (δ_{C} 151.6), C-4a (δ_{C} 114.3); δ_{H} 7.62 (H-8) with C-1 (δ_{C} 178.1), C-8a (δ_{C} 121.8), and C-7 (δ_{C} 114.7) (Fig. 3 and S27). Thus, the structure of **5** was identified as goniochelinone, a structural isomer of goniothalamine B. Full assignments of the ¹H and ¹³C NMR spectroscopic data are shown in Table 2 and the COSY and selected HMBC correlations in Fig. 3.

Plausible biosynthetic pathways of compounds **1–5** are shown in Fig. 11. Compounds **1–4** and **5** might arise from different polyketide units (Wagoner et al., 2008; King et al., 2009; Bauer et al., 2010). The benzo[*f*]indole-4,9-diones **1–4** could be obtained from a hexaketide generated from an isobutyryl-CoA starter and a five acetyl-CoA extender (Zhou et al., 2016) involving multi-steps including the incorporation of nitrogen (Fig. 11a) (Wagoner et al., 2008; Bauer et al., 2010). Compound **4** undergoes oxidation of the methyl group to give the carboxylic acid derivative (**4a**), esterification of **4a** would yield compound **3**. Alternatively, **4a** was regioselectively *N*-methylated to obtain compound **2** which was then oxidized at C-8 to give compound **1**. The naphthoquinone **5** might arise from heptaketide (Fig. 11b) with the similar process of **4**.

Compound **6** was isolated as a colorless prism. Analysis of its HRESIMS showed a $[M + Na]^+$ ion peak at m/z 273.0749 (calcd for $C_{13}H_{14}O_5Na$, 273.0739) which established the molecular formula $C_{13}H_{14}O_5$. The 5,7-dihydroxy-8-phenylbicyclo[3.3.1]nonan-2-one based structure and the absolute configuration (4*S*,5*S*,6*R*,7*S*,8*S*) of compound **6** were established by single-crystal X-ray diffraction analysis using Cu $K\alpha$ radiation (Fig. 7) with the flack parameter of -0.04(7). The ECD spectrum of **6** (Fig. 9) displayed a negative Cotton effect at 227 nm, similar to that of parvistone D and parvistone E (Liuo et al., 2014) which also supported the identification of its absolute configuration. The 1H NMR spectrum further supported this structure and showed resonances for the mono substituted aromatic ring at δ_H 7.38 (2H, d, $J = 7.9$ Hz, H-2'/6'), 7.30 (2H, d, $J = 7.6$ Hz, H-3'/5') and 7.26 (1H, d, $J = 7.2$ Hz, H-4'). The structure of compound **6** was almost identical to that of (+)-goniopypyrone except for the configuration at C-7 which was 7*S* for compound **6** and 7*R* for (+)-goniopypyrone (Tsubuki et al., 1999; Zhang et al., 1998). The 1H NMR spectroscopic data of these two compounds were significantly different, except for the aromatic proton chemical shifts, as shown in Table S1. Thus, compound **6** was assigned as (-)-(4*S*,5*S*,6*R*,7*S*,8*S*)-goniocheliolactone, the C-7 epimer of (+)-goniopypyrone.

Compound **7** was isolated as a colorless prism. Its molecular formula $C_{15}H_{16}O_6$ was established the HRESIMS analysis which showed a $[M + Na]^+$ ion peak at m/z 315.0854 (calcd for $C_{15}H_{16}O_6Na$, 315.0845). The core structure of compound **7** was a [3.3.1] bicyclic, similar to that found in compound **6**. A single-crystal X-ray diffraction analysis of compound **7** using Cu $K\alpha$ radiation established the structure of compound **7** and its absolute configuration as 4*S*,5*S*,6*R*,7*S*,8*S* (Fig. 8) with the Flack parameter 0.00 (11). Its ECD spectrum (Fig. 9) was similar to that of compound **6** which showed the Cotton effect at 232 nm to support the 4*S*,5*S*,6*R*,7*S*,8*S* configuration. The full assignments of its 1H and ^{13}C NMR spectroscopic data are summarized in Table 3. It should be noted that compound **7** is the C-7 is epimer of 7-acetylgoniopypyrone (Zhang et al., 1998). Differences were also observed between these epimers from their 1H NMR spectra, especially the chemical shifts for the [3.3.1] bicyclic protons as shown in Table S2. Thus, the structure of **7** was identified as (-)-(4*S*,5*S*,6*R*,7*S*,8*S*)-7-acetylgoniocheliolactone.

Compound **8**, (+)-(7*S*,8*S*)-goniocheliolbutenolide A, was obtained as a colorless viscous oil. Its molecular formula $C_{15}H_{14}O_5$ was deduced from its HRESIMS (m/z 297.0730 $[M + Na]^+$, calcd for $C_{15}H_{14}O_5Na$, 297.0739). The 1H and ^{13}C NMR spectroscopic data of **8** were similar to those of goniobutenolide A (Fang et al., 1991). The main difference between compounds **8** and goniobutenolide A was that compound **8** displayed an additional 1H NMR resonance for acetoxy protons at δ_H 2.12. Additionally, the resonances for H-7 (δ_H 5.10) and H-8 (δ_H 5.88) of compound **8** were shifted downfield compared to the same protons of goniobutenolide A, [δ_H 4.98 (H-7) and 4.94 (H-8)], (SI Table S3 and S4). The HMBC correlations between H-8 (δ_H 5.85) with the acetate carbonyl carbon (δ_C 168.0), H-2'/H-6' (δ_H 7.40) with C-8 (δ_C 76.5) and H-7 (δ_H 4.98) with C-8 (δ_C 76.5) supported the position of the acetoxy group

at C-8 (Fig. 3). The relative stereochemistry of **8** was proposed to be *ethyro* based on the coupling constant of H-7/H-8 ($J_{7,8} = 6.0$ Hz) indicating their *syn* orientation ($J_{7,8} = 6-8$ Hz) (Fang et al., 1991, Shing et al., 1994; Xu et al., 1994). The *Z*-geometry of C-5/C-6 was determined from a comparison of the ^1H NMR spectroscopic data to those of 7-*epi*-goniobutenolide A and B (SI Table S3 and S4). The absolute configuration of **8** (7*S*,8*S*) was proposed by the comparison of its ECD spectrum to that of compound **11** which showed a negative Cotton effect at 230 and 274 nm (Fig. 10). Thus, the structure of compound **8** was deduced as (7*S*,8*S*,*Z*)-5-(3-acetyl-2-hydroxy-3-phenylpropylidene)-2(5*H*)-furanone.

Compound **9**, (-)-(7*S*,8*R*)-goniochelienbutenolide B, was obtained as a colorless viscous oil. Its molecular formula $\text{C}_{15}\text{H}_{14}\text{O}_5$ was established by HRESIMS (m/z 297.0746 [$\text{M} + \text{Na}$] $^+$, calcd for $\text{C}_{15}\text{H}_{14}\text{O}_5\text{Na}$, 297.0739). The ^1H NMR spectroscopic data of **9** were similar to that of **8** (Table 4 and SI Fig.S3), but H-3 (δ_{H} 6.19) of compound **9** showed long range coupling with that of H-6 (δ_{H} 5.82) with the J value of 1.5 Hz indicating the *E*-geometry at the exo-cyclic alkene isomer of compound **9** (Fang et al., 1991; Popsavin et al., 2012). The oxymethine protons of C-7 and C-8 were assigned the *threo* configuration on the basis of their coupling constant $J_{7,8} = 4.2$ Hz (Fang et al., 1991; Shing et al., 1995; Popsavin et al., 2012). The absolute configuration of **9** as 7*S*,8*R* was proposed from the comparison of its ECD spectrum to that of **10** (Fig. 10). Therefore, the structure was characterized as (7*S*, 8*R*, *E*)-5-(3-acetyl-2-hydroxy-3-phenylpropylidene)-2(5*H*)-furanone.

Annonaceae is a large family containing over 120 genera (Saripalli and Dixit, 2013). Each genus produces different major phytochemicals. For example, polyoxygenated cyclohexenes are commonly found in many species of *Uvaria* (Awale et al., 2017; Lekphrom et al., 2018; Hsu et al., 2016; Kaweetripob et al., 2015; Auranwiwat et al., 2017; Macabeo et al., 2017), but they are also found in some species of *Artabotrys* (Liu et al., 2018), *Cleistochlamys* (Nyandoro et al., 2017), *Dasymaschalon* (Hongthong et al., 2015), *Ellipeiopsis* (Kijjoa et al., 2002), and *Monanthotaxis* (Starks et al., 2012) genera. Flavonoids are major compounds from many genera including *Desmos* (Clement et al., 2017; Tuntipaleepun et al., 2012; Kuo et al., 2015; Prachyawarakorn et al., 2013; Bajgai et al., 2011; Rittiwong et al., 2011), and *Friesodielsia* (Meesakul et al., 2017; Prawat et al., 2012; Fleischer et al., 1997; Joseph et al., 2007), while alkaloids, including aristololactams, aporphines and oxoprotoberberine, are commonly found in the *Miliusa* (Promchai et al., 2018; Promchai et al., 2016; Hasan et al., 2000; Chen et al., 2003; Harrigan et al., 1994) and *Fissistigma* (Ge et al., 2013; Zhou et al., 2016; Zhou et al., 2018; Chia et al., 2000) genera. Styryllactones are the major compounds found in *Goniiothalamus* (Moharam et al., 2012; Macabeo et al., 2013; Zhang et al., 1998; Fang et al., 1991., Suchaichit et al., 2015, Hisham et al., 2000; Uma et al., 2012) and *Polyalthia* (Nantapap et al., 2015; Nurunajah et al., 2001; Samoath and Vasanthi, 2013) whereas acetogenin (Moghadamtousi et al., 2015; Dang et al., 2011; Melot et al., 2009; Liaw et al., 2008; Sun et al., 2017; Guzmán et al., 2009; Liaw et al., 2004; Queiroz et al., 2003) and diterpenoids (Supudompol et al., 2004; Li et al., 2005; Li et al., 2009; Deepalard et al., 2007; Meng et al., 2007) were found in *Annona* and *Mitrephora*.

Most of the isolated compounds (**1**, **2**, **4**, **6**, **7**, **10**, **12–14**, **16**, **19**, **22**, **25**, **27**, **28**, **30**, **31**, **34**, **36–39**) were evaluated for their cytotoxicities against the colon cancer cell line HCT-116. Only compound **25** (griffithazanone A) showed potent cytotoxicity with an IC₅₀ value of 2.39 μ M (doxorubicin as a positive control had an IC₅₀ value of 9.74 μ M). All remaining compounds were inactive (IC₅₀ >50 μ M). The IC₅₀ value of compound **25** against HCT-116 is less than those against human cervical cancer cells (20.2 μ M and 11.7 μ M for KB cells and HeLa cells, respectively) (Tip-pyang et al., 2010). Indicating that the cytotoxic effect of compound **25** is not cell-line specific and that compound **25** may be more effective against colorectal cancer cell lines than cervical cancer cell lines.

3. Conclusions

The phytochemical investigation of the extracts of the *G. cheliensis* led to the identification of 44 compounds, of which eight were previously undescribed compounds (**1–3**, **5**, and **6–9**). The discovery of styryllactones as the major compounds was in good agreement with the previous report of this genus (Moharam et al., 2012; Macobeo et al., 2013; Zhang et al., 1998; Fang et al., 1991., Suchaichit et al., 2015, Hisham et al., 2000; Uma et al., 2012). To the best of our knowledge, the indole alkaloids **1–3** and alkaloid **5** were found in this species for the first time. In a previous paper, the [3.3.1] bicyclic core structure of compounds **6** and **7** was reported as having the 7*R* configuration (Fang et al., 1991; Tsubuki et al., 1999; Zhang et al., 1998), however, we report herein related compounds having the 7*S* configuration with their absolute configurations being confirmed by single-crystal X-ray diffraction analysis. The isolated compounds were found to have no cytotoxicities against the colon cancer cell line HCT-116 at 50 μ M except for griffithazanone A (**25**) which had potent cytotoxicities. This compound may have potential as a lead compound for the development of new anti-cancer agents.

4. Experiment section

4.1. General experiment procedures

The optical rotation [α]_D values were determined with a Bellingham and Stanley ADP400 polarimeter. Melting points were measured with a Buchi B-540 melting point apparatus. UV-vis spectra were recorded with Varian Cary 5000 UV-vis-NIR spectrophotometer. The IR spectra were recorded using a Perkin-Elmer FTS FT-IR spectrophotometer. Electronic circular dichroism spectra were recorded on a JASCO J-815 CD spectropolarimeter. The NMR spectra were recorded using Bruker Avance 600 MHz spectrometers. Chemical shifts are reported in parts per million (δ), and coupling constants (*J*) are expressed in hertz. HRESIMS spectra were obtained on a Bruker-Hewlett-Packard 1100 Esquire-LC system mass spectrometer. Single-crystal X-ray diffraction measurements were made on a Bruker APEX DUO diffractometer with cross-coupled multilayer optics Cu-K α radiation. HPLC was performed on a Waters 1525 HPLC pump system, equipped with a Waters 2487 dual λ absorbance detector using the following column: Phenomenex Luna 5u C₈ column (10 $\times$ 250 mm)

and CHIRALCEL OD-H column (4.6 × 250 mm). Quick column chromatography (QCC) and column chromatography (CC) were carried out on Si gel 60 H (5–40 μm, Merck), Si gel 100 (63–200 μm, Merck) or silica gel 100 (63–200 μm, SiliCycle[®] Inc.), respectively. Waters 10 g Sep-Pak's were used for reversed-phase flash chromatography. Sephadex LH-20 was also used for CC. Precoated plates of silica gel 60 F₂₅₄ were used for analytical purposes.

4.2 Plant material

The twigs and leaves of *Goniothalamus cheliensis* Hu (Annonaceae) were collected from Doi Tung, Chiang Rai Province, Thailand (GPS coordinates: 20°17.351' N, 99°48.661' E), in October 2015, winter. The plant material was authenticated by Mr. Martin van de Bult (Doi Tung Development Project, Chiang Rai, Thailand), and a voucher specimen (MFU-NPR0146) was deposited at the Natural Products Research Laboratory, School of Science, Mae Fah Luang University.

4.3 Extraction and isolation

The dried twigs of *G. cheliensis* (5.1 kg) were extracted with acetone. The acetone extract (41.2 g) was subjected to QCC eluting with a gradient solvent system of *n*-hexanes-acetone (1:0 to 0:1, v/v) to afford 11 fractions (A–K). Fraction D (2.3 g) was subjected to Sephadex LH-20 (100% MeOH) to yield seven subfractions (D₁–D₇). Fraction D₄ (95.8 mg) was subjected to Sep-Pak C₁₈ CC, eluted with MeOH-H₂O (3:2, v/v) to yield subfraction D₁A. Fraction D₁A (61.6 mg) was separated over Sephadex LH-20 to give 4 subfractions (D₁A-1–D₁A-4). Fraction D₁A-2 (40.4 mg) was fractionated using silica gel CC eluting with *n*-hexanes-acetone (7:3, v/v) to obtained 6 subfractions (D₁A-2-1–D₁A-2-6). Fraction D₁A-2-2 (4.5 mg) was chromatographed using RP C₁₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 1:1, 2 mL/min) to give compounds **29** and **32** (0.3 mg, *t*_R = 11.6 min), **30** (1.0 mg, *t*_R = 15.6 min), and **33** (0.9 mg, *t*_R = 18.5 min). Compound **31** (2.1 mg, *t*_R = 15.6 min) was obtained from D₁A-2-3 by RP C₁₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 1:1, 2 mL/min). Fraction D₅ (205.0 mg) was subjected to Sep-Pak C₁₈ CC, eluted with MeOH-H₂O (3:2, v/v) to give subfraction D₅A (172.6 mg) following separated by Sephadex LH-20 (100% MeOH) to yield 4 subfractions (D₅A-1–D₅A-4). Compound **35** (3.9 mg) and **28** (1.4 mg) were purified from D₅A-2 (47.5 mg) by silica gel CC eluting with *n*-hexanes-acetone (7:3, v/v). Fraction D₇ (29.4 mg) was subjected to Sep-Pak C₁₈ CC eluting with MeOH-H₂O (3:2, v/v) to give D₇A (26.1 mg) and further separation on a silica gel CC eluting with CH₂Cl₂-MeOH (97:3, v/v) to yield compound **5** (0.8 mg). Fraction E (14.9 mg) was subjected to RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:2, 2 mL/min) to give compound **2** (1.0 mg, *t*_R = 12.8 min) and compound **1** (2.1 mg, *t*_R = 13.6 min). Fraction F (8.9 g) was partition with CH₂Cl₂ and water, the CH₂Cl₂ soluble fraction (7.9 g) was subjected to silica gel CC, eluting with a gradient of CH₂Cl₂-MeOH (99:1 to 19:1, v/v) to afford 7 subfractions (F1–F7). Fraction F2 (912.8 mg) was chromatographed on a silica gel CC eluting with CH₂Cl₂-MeOH (99:1, v/v) to afford compound **4** (1.1 mg) and 4 subfractions (F2-1–F2-4). Fraction F2-2 (588.0 mg) was subjected to Sephadex LH-

20 CC (100% MeOH) to afford 5 subfractions (F2-2-1–F2-2-5). Fraction F2-2-1 (114.5 mg) was purified using RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:7, 2 mL/min) to afford compounds **8** (1.1 mg, *t_R* = 34.9 min), **9** (0.3 mg, *t_R* = 48.7 min), and seven subfractions (F2-2-1-1–F2-2-1-7). Fraction F2-2-1-3 and F2-2-1-6 were subjected to RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:7, 2 mL/min) to yield compounds **16** (7.7 mg, *t_R* = 40.5 min), **13** (1.3 mg, *t_R* = 62.9 min), **7** (0.9 mg, *t_R* = 70.0 min), **14** (1.7 mg, *t_R* = 81.9 min), and **15** (1.1 mg, *t_R* = 91.4 min). Fraction F2-2-2 (257.7 mg) was chromatographed on silica CC eluting with *n*-hexanes-acetone (7:3, v/v) to give 5 subfractions (F2-2-2-1–F2-2-2-5). Fraction F2-2-2-2 (11.8 mg) was subjected to RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:7, 2 mL/min) to yield compound **26** (0.7 mg, *t_R* = 20.6 min). Compound **20** (3.9 mg, *t_R* = 29.1 min), enantiomeric mixtures of **17** and **18** (1.9 mg, *t_R* = 15.2 min), **23** and **21** (0.7 mg, *t_R* = 29.1 min) and **19** and **22** (15.7 mg) were obtained from fraction F2-2-2-4 (28.5 mg). The chiral column OD-H (eluted with *n*-hexanes/MeOH, 3:2, 2 mL/min) was used to yield compounds **17** (0.2 mg, *t_R* = 10.9 min) and **18** (0.4 mg, *t_R* = 11.4 min), **23** (0.3 mg, *t_R* = 9.2 min) and **21** (0.2 mg, *t_R* = 10.9 min), and **19** (5.2 mg, *t_R* = 6.3 min) and **22** (6.9 mg, *t_R* = 7.4 min). Fraction F2-2-4 (13.7 mg) was subjected to RP C₁₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 2:3, 2 mL/min) to give compounds **24** (0.5 mg, *t_R* = 14.1 min) and **25** (1.9 mg, *t_R* = 11.8 min). Compound **3** (1.8 mg, *t_R* = 15.9 min) was purified by RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:7, 2 mL/min) from fraction F2-2-5 (8.4 mg). Compound **34** (1.0 mg, *t_R* = 9.9 min) was purified from fraction F3 (435.7 mg) by QCC over silica gel, eluting with a gradient of CH₂Cl₂-MeOH gradient (99:1 to 97:3, v/v) to give 4 subfractions (F3-1–F3-4), with each of these purified with RP C₁₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 2:3, 2 mL/min) and Sephadex LH-20 (100% MeOH). Fraction G (10.8 g) was chromatographed on a silica gel CC eluting with CH₂Cl₂-MeOH (95:5, v/v) to give 10 subfractions (G1–G10). Fraction G3 (15.7 mg) was purified using RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:7, 2 mL/min) to give compound **12** (1.7 mg, *t_R* = 7.4 min). Fraction G4 (142.3 mg) was subjected to silica gel CC, eluted with CH₂Cl₂-MeOH (97:3, v/v) following RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 1:3, 2 mL/min) to give compounds **27** (5.2 mg, *t_R* = 14.2 min), **10** (0.4 mg, *t_R* = 23.3 min), and **11** (0.3 mg, *t_R* = 24.6 min). Compound **6** (36.9 mg) was recrystallized from *n*-hexanes-acetone (1:1, v/v) from subfraction G6 (234.7 mg).

The dried leaves of *G. cheliensis* (412.1 g) were extracted with acetone (3 × 3 L) at room temperature. The acetone extract (33.8 g) was subjected to QCC over silica gel, eluting with a gradient of *n*-hexanes-acetone gradient (1:0 to 0:1, v/v) to afford fourteen fractions (A–N). The combined of fraction E–H were further separated on a Sep-Pak C₁₈ CC eluting with MeOH-H₂O (1:1, v/v) to give fraction B1 (819.4 mg) and further separation by silica gel CC, eluting with *n*-hexanes-acetone (9:1, v/v) gave 5 subfractions (B1-1–B1-5). Fraction B1-3 (73.2 mg) was purified by RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:2, 2 mL/min) to yield compounds **37** (2.0 mg, *t_R* = 24.7 min) and **42** (1.3 mg, *t_R* = 37.3 min). The combination of fraction I–K was subjected to Sep-Pak C₁₈ CC eluting with MeOH-H₂O (1:1, v/v) to give

fraction C1 (806.4 mg) and further separation by Sephadex LH-20 CC (100% MeOH) to yield six subfractions (C1-1–C1-6). Fraction C1-3 (8.7 mg) was purified by silica gel CC, eluting with *n*-hexanes-acetone (7:3, v/v) to give compound **39** (1.7 mg) and 5 subfractions (C1-3-1–C1-3-5). Compounds **36** (1.4 mg, t_R = 32.8 min) and **41** (0.9 mg, t_R = 34.0 min) were obtained from subfraction C1-3-5 (4.5 mg), by a RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 1:1, 2 mL/min). After further purification by RP C₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 1:1, 2 mL/min), compounds **38** (2.1 mg, t_R = 20.7 min) and **40** (0.4 mg, t_R = 23.5 min) were obtained from subfraction C1-4 (6.1 mg). The combined of fractions L–N were further separated by Sephadex LH-20 CC (100% MeOH) to give nine subfractions (D1–D9). Fraction D6 (18.2 mg) was isolated by RP C₁₈ HPLC (eluted with CH₃CN/H₂O with 0.05% TFA, 3:7, 2 mL/min) to afford compounds **43** (0.3 mg, t_R = 28.9 min) and **44** (0.8 mg, t_R = 30.5 min).

4.3.1 *Goniochelienic acid A (1)*

Yellow needle; mp 215–217 °C; UV (MeOH) $\lambda_{\max}$ (log ϵ) 253 (3.47), 287 (3.08), 328 (2.51) nm; IR (neat) $\nu_{\max}$ 2920, 2850, 1721, 1451, 1236, 747 cm⁻¹; For ¹H-NMR (600 MHz, CDCl₃) and ¹³C-NMR (125 MHz, CDCl₃), see Table 1; HRESIMS (positive-ion mode) m/z 294.0379 [M + Na]⁺ (calcd for C₁₄H₉NO₅Na, 294.0378).

4.3.2 *Goniochelienic acid B (2)*

Yellow amorphous solid; mp 220–222 °C; UV (MeOH) $\lambda_{\max}$ (log ϵ) 253 (3.74), 278 (3.33), 331 (2.95) nm; IR (neat) $\nu_{\max}$ 2918, 2850, 1727, 1459, 1228, 717 cm⁻¹; For ¹H-NMR (600 MHz, CDCl₃) and ¹³C-NMR (125 MHz, CDCl₃), see Table 1; HRESIMS (positive-ion mode) m/z 278.0436 [M + Na]⁺ (calcd for C₁₄H₉NO₄Na, 278.0429).

4.3.3 *Methyl goniochelienate (3)*

Yellow solid; mp 214–216 °C; UV (MeOH) $\lambda_{\max}$ (log ϵ) 251 (4.37), 276 (3.95), 309 (3.53), 374 (3.13) nm; IR (neat) $\nu_{\max}$ 2924, 2853, 1678, 1208, 1135 cm⁻¹; For ¹H-NMR (600 MHz, MeOH-*d*₄) and ¹³C-NMR (125 MHz, MeOH-*d*₄), see Table 1; HRESIMS (positive-ion mode) m/z 278.0435 [M + Na]⁺ (calcd for C₁₄H₉NO₄Na, 278.0429).

4.3.4 *Goniochelieninone (5)*

Yellow solid; mp 162–165 °C; UV (MeOH) $\lambda_{\max}$ (log ϵ) 215 (3.53), 282 (3.20), 417 (2.59) nm; IR (neat) $\nu_{\max}$ 3335, 3215, 2920, 2851, 1615, 1452, 1371, 1260 cm⁻¹; For ¹H-NMR (600 MHz, CDCl₃) and ¹³C-NMR (125 MHz, CDCl₃), see Table 2; HRESIMS (positive-ion mode) m/z 284.0530 [M + Na]⁺ (calcd for C₁₃H₁₁NO₅Na, 284.0535).

4.3.5 (–)-(4*S*,5*S*,6*R*,7*S*,8*S*)-*Goniochelienlactone (6)*

Colorless solid; mp 215–217 °C; [α]_D²⁵ –47.0 (*c* 0.1, EtOH); UV (EtOH) $\lambda_{\max}$ (log ϵ) 215 (2.76), 255 (1.51) nm; CD (EtOH) $\lambda_{\max}$ ($\Delta\epsilon$) 227 (–0.89) nm; IR (neat)

$\nu_{\max}$ 3395, 2925, 2873, 2853, 1713, 1177, 1081 cm^{-1} ; ^1H NMR (600 MHz, $\text{DMSO}-d_6$) and ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$) data, see Table 3; HRESIMS (positive-ion mode) m/z 273.0749 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{13}\text{H}_{14}\text{O}_5\text{Na}$, 273.0739).

4.3.6 (–)-(4*S*,5*S*,6*R*,7*S*,8*S*)-7-acetylgoniocheliolactone (**7**)

Colorless solid; mp 152–155 °C; $[\alpha]_D^{25}$ –58.0 (c 0.1, EtOH); UV (EtOH) $\lambda_{\max}$ (log ϵ) 212 (2.69), 249 (0.97) nm; CD (EtOH) $\lambda_{\max}$ ($\Delta\epsilon$) 232 (–0.26) nm; IR (neat) $\nu_{\max}$ 3422, 2924, 1731, 1220, 1054 cm^{-1} ; ^1H NMR (600 MHz, CDCl_3) and ^{13}C NMR (125 MHz, CDCl_3) data, see Table 3; HRESIMS (positive-ion mode) m/z 315.0854 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{16}\text{O}_6\text{Na}$, 315.0845).

4.3.7 (+)-(7*S*,8*S*)-Goniocheliolbutenolide A (**8**)

Colorless viscous; $[\alpha]_D^{25}$ +55.0 (c 0.1 EtOH); UV (EtOH) $\lambda_{\max}$ (log ϵ) 206 (2.9), 273 (2.8) nm; CD (EtOH) $\lambda_{\max}$ ($\Delta\epsilon$) 227 (–0.30), 274 (–0.59) nm; IR (neat) $\nu_{\max}$ 3419, 2926, 1682, 1750, 1208, 1135 cm^{-1} ; ^1H NMR (600 MHz, acetone- d_6) and ^{13}C NMR (125 MHz, acetone- d_6) data, see Table 4; HRESIMS (positive-ion mode) m/z 297.0730 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{14}\text{O}_5\text{Na}$, 297.0739).

4.3.8 (–)-(7*S*,8*R*)-Goniocheliolbutenolide B (**9**)

Colorless viscous; $[\alpha]_D^{25}$ –34.0 (c 0.1 CHCl_3); UV (CHCl_3) $\lambda_{\max}$ (log ϵ) 225 (2.9), 274 (2.8) nm; CD (CHCl_3) $\lambda_{\max}$ ($\Delta\epsilon$) 228 (+0.04), 281 (+0.76) nm; IR (neat) $\nu_{\max}$ 3472, 2922, 2852, 1743, 1236 cm^{-1} ; ^1H NMR (600 MHz, CDCl_3) and ^{13}C NMR (125 MHz, CDCl_3) data, see Table 4; HRESIMS (positive-ion mode) m/z 297.0746 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{15}\text{H}_{14}\text{O}_5\text{Na}$, 297.0739).

4.4 X-ray crystal analysis of compounds **1**, **4**, **6**, and **7**

X-ray diffraction data were collected using Mo $\text{K}\alpha$ radiation ($\lambda = 0.71073$ Å) and Cu $\text{K}\alpha$ radiation ($\lambda = 1.54178$ Å) on a Bruker APEX-II CCD diffractometer. Using Olex2 (Dolomanov et al., 2009), the structure was solved with the XT (Sheldrick, 2015) structure solution program, using the Intrinsic Phasing solution method. The model was refined with version 2018/1 of XL (Sheldrick, 2015) using Least Squares minimisation. Crystallographic data of compounds **1**, **4**, **6**, and **7** have been deposited at the Cambridge Crystallographic Data Center under the reference number CCDC 1846457 (**1**), 1846454 (**4**), 1846455 (**6**), and 1846456 (**7**). These data can be obtained, free of charge, from the Cambridge Crystallographic Data Centre via http://www.ccdc.cam.ac.uk/data_request/cif.

4.4.1 Crystal data of **1**

Crystals of **1** were obtained as orange irregular-shaped crystals from Me_2CO -hexanes. $\text{C}_{31}\text{H}_{24}\text{N}_2\text{O}_{11}$, $M_r = 600.52$, monoclinic, space group $\text{C}2/c$ (No. 15), $a = 13.549(6)$ Å, $b = 26.964(12)$ Å, $c = 8.522(4)$ Å, $\beta = 122.191(10)^\circ$, $\alpha = \gamma = 90^\circ$, $V = 2635(2)$ Å³, $T = 90(2)$ K, $Z = 4$, $Z' = 0.5$, $\mu(\text{Mo } \text{K}\alpha) = 0.117$, 9693 reflections

measured, 1731 unique ($R_{int} = 0.0813$) which were used in all calculations. The final wR_2 was 0.1937 (all data) and R_I was 0.0692 ($I > 2(I)$).

4.4.2 Crystal data of **4**

Crystals of **4** were obtained as orange needle-shaped crystals from Me₂CO-hexanes. C₁₃H₉NO₂, $M_r = 211.21$, monoclinic, space group P2₁/c (No. 14), $a = 15.1793(14)$ Å, $b = 3.8477(5)$ Å, $c = 17.6197(18)$ Å, $\beta = 109.578(5)^\circ$, $\alpha = \gamma = 90^\circ$, $V = 969.59(19)$ Å³, $T = 100(2)$ K, $Z = 4$, $Z' = 1$, $\mu(\text{Mo } K_\alpha) = 0.099$, 9656 reflections measured, 2298 unique ($R_{int} = 0.0343$) which were used in all calculations. The final wR_2 was 0.1790 (all data) and R_I was 0.0602 ($I > 2(I)$).

4.4.3 Crystal data of **6**

Crystals of **6** were obtained as colourless irregular-shaped crystals from Me₂CO-hexanes. C₁₃H₁₄O₅, $M_r = 250.24$, orthorhombic, space group P2₁2₁2₁ (No. 19), $a = 9.4413(5)$ Å, $b = 10.3844(6)$ Å, $c = 11.6131(6)$ Å, $\alpha = \beta = \gamma = 90^\circ$, $V = 1138.57(11)$ Å³, $T = 90(2)$ K, $Z = 4$, $Z' = 1$, $\mu(\text{Cu } K_\alpha) = 0.948$, 9069 reflections measured, 2012 unique ($R_{int} = 0.0325$) which were used in all calculations. The final wR_2 was 0.0615 (all data) and R_I was 0.0241 ($I > 2(I)$). Flack parameter = -0.04(7).

4.4.4 Crystal data of **7**

Crystals of **7** were obtained as colourless irregular-shaped crystals from Me₂CO-hexanes. C₁₅H₁₆O₆, $M_r = 292.28$, monoclinic, P2₁ (No. 4), $a = 10.9487(4)$ Å, $b = 6.8358(3)$ Å, $c = 18.6909(7)$ Å, $\beta = 90.243(2)^\circ$, $\alpha = \gamma = 90^\circ$, $V = 1398.87(10)$ Å³, $T = 90(2)$ K, $Z = 4$, $Z' = 2$, $\mu(\text{Cu } K_\alpha) = 0.910$, 20213 reflections measured, 4945 unique ($R_{int} = 0.0351$) which were used in all calculations. The final wR_2 was 0.0900 (all data) and R_I was 0.0336 ($I > 2(I)$). Flack parameter = 0.00(11).

4.5 Biological assay

4.5.1 Bioassay for Cytotoxicity

The cytotoxicities of compounds on colon cancer cell were performed following by the previous method (Sriyatep et al., 2017). Doxorubicin used as a positive control (IC₅₀ = 9.74 µM).

Acknowledgements

This research was financially supported by the Thailand Research Fund and Mae Fah Luang University through the Basic Research Grant (BRG5980012), the Thailand Research Fund through the Royal Golden Jubilee Ph.D. Program (PHD/0019/2557) and Direct Basic Research Grant (DBG5980001) and the Mae Fah Luang University Graduate Student Research Grant. University of British Columbia is also acknowledged for laboratory facilities. We would like to thank Mr. Martin van de Bult, Doi Tung Development Project, Chiang Rai, Thailand for plant collection and identification.

Appendix A. Supplementary data

Supplementary data related to this article can be found at

References

- Auranwiwat, C., Wongsomboon, P., Thaima, T., Rattanajak, R., Kamchonwongpaisan, S., Willis, A.C., Lie, W., Pyne, S.G., Limtharakul, T., 2017. 2-Phenyl-naphthalenes and a polyoxygenated cyclohexene from the stem and root extracts of *Uvaria cherrevensis* (Annonaceae). *Fitoterapia* 120, 103–107.
- Awale, S., Tawia, A.M., Dibwe, D.F., Ueda, J., Sun, S., Athikomkulchai, S., Balachandran, C., Saiki, I., Matsumoto, K., Esumi, H., 2017. Highly oxygenated antiausterity agents from the leaves of *Uvaria dac*. *Bioorg. Med. Chem. Lett.* 27, 1967–1971.
- Bajgai, S.P., Prachyawarakorn, V., Mahidol, C., Ruchirawat, S., Kittakoop, P., 2011. Hybrid flavan-chalcones, aromatase and lipooxygenase inhibitors, from *Desmos choichinchinensis*. *Phytochemistry* 72, 2062–2067.
- Bauer, J.D., King, R.W., Brady, S.F., 2010. Utahmycins A and B, azaquinones produced by an environmental DNA clone. *J. Nat. Prod.* 73, 976–979.
- Chang, C.W., Chang, H.S., Cheng, M.J., Peng, C.F., Chen, I.S., 2015. Identification of five new minor constituents from the whole plant of *Amischotolype hispida*. *Helv. Chim. Acta.* 98, 347–358.
- Chen, B., Feng, C., Li, B.G., Zhang, G.L., 2003. Two new alkaloids from *Milusa cuneate*. *Nat. Prod. Res.* 17, 397–402.
- Chia, Y.C., Chang, F.R., Teng, C.M., Wu, Y.C., 2000. Aristolactams and dioxoaporphines from *Fissistigma balansae* and *Fissistigma oldhamii*. *J. Nat. Prod.* 63, 1160–1163.
- Chen, C.Y., Wu, T.Y., Chang, F.R., Wu, Y.C., 1998. Lignans and kauranes from the stems of *Annona cherimola*. *J. Chin. Chem. Soc.* 45, 6294–634.
- Clement, J.A., Ondeyka, J.G., Goetz, M.A., 2017. Benzoate esters and flavonoids from *Desmos pedunculatus*. *Phytochem. Lett.* 22, 117–121.
- Colegate, S.M., Din, L.B., Latiff, A., Salleh, K.M., Samsudin, M.W., Skelton, B.W., Tadano, K.I., White, A.H., Zakaria, Z., 1990. (+)-Isoalcoholactone: a furanopyrone isolated from *Goniiothalamus* species. *Phytochemistry* 29, 1701–1704.
- Dang, Q.L., Kim, W.K., Nguyen, C.M., Choi, Y.H., Choi, G.J., Jang, K.S., Park, M.S., Lim, C.H., Luu, N.H., Kim, J.C., 2011. Nematicidal and antifungal activities of Annonaceous acetogenins from *Annona squamosa* against various plant pathogens. *J. Chem. Food Chem.* 59, 11160–11167.
- Deepralard, K., Pengsuparp, T., Moriyasu, M., Kawanishi, K., Suttisri, R., 2007. Chemical constituents of *Mitrephora maingayi*. *Biochem. Syst. Ecol.* 35, 695–699.
- Dolomanov, O.V., Bourhis, L.J., Gildea, R.J., Howard, J.A.K., Puschmann, H., 2009. Olex2: A complete structure solution, refinement and analysis program. *J. Appl. Cryst.* 42, 339–341.
- Duc, L.V., Thanh, T.B., Thanh, H.N., Tien, V.N., 2016. Chemical constituents and cytotoxic effect from the barks of *Goniiothalamus cheliensis* Merr. & Chun. growing in Vietnam. *J. Appl. Pharm. Sci.* 6, 1–5.

606 East, A.J., Ollis, W.D., Wheeler, R.E., 1969. Natural occurrence of 3-aryl-4-
 607 hydroxycoumarins. Part I. Phytochemical examination of *Derris robusta*
 608 (Roxb.) Benth. J. Chem. Soc. 365–374.
 609 Efdi, M., Fujita, S., Inuzuka, T., Koketsu., 2010. Chemical studies on *Goniothalamus*
 610 *tapis* Miq. Nat. Prod. Res. 24, 657–662.
 611 Fang, X.P., Andersen, J. E., Chang, C.J., Mclaighlin, J.L., 1991. Two new styryl
 612 lactones, 9-deoxygoniopypyrone and 7-*epi*-goniofufurone, from
 613 *Goniothalamus giganteus*. J. Nat. Prod. 54, 1034–1043.
 614 Fan, C.M., Chou, G.X., Zhu, E.Y., 2016. Chemical constituents from *Crotalaria*
 615 *sessiliflora* L. Acta. Pharmacol. Sin. 51, 775–779.
 616 Fleischer, T.C., Waigh, R.D., Waterman, P.G., 1997. Bisabolene sesquiterpenes and
 617 flavonoids from *Friesodielsia enghiana*. Phytochemistry 44, 315–318.
 618 Ge, Y.W., Zhu, S., Shang, M.Y., Zang, X.Y., Wang, X., Bai, Y.J., Li. L., Komatsu,
 619 K., Cai, S.Q., 2013. Aristololactams and aprophines from the stems of
 620 *Fissistigma oldhamii* (Annonaceae). Phytochemistry 86, 201–207.
 621 Schlie-Guzmán, M.A., García-Carrancá, A., González-Esquinca., 2009. *In vitro* and
 622 *in vivo* antiproliferative activity of Laherradurin and cherimolin-2 of *Annona*
 623 *diverifolia* Saff. Phytoter. Res. 24, 1128–1133.
 624 Harrigan, G.G., Gunatilaka, A.A.L., Kingston, D.G., 1994. Ioslotion of bioactive and
 625 other oxoaporphine alkaloids from two annonaceous plants, *Xylopia*
 626 *aethiopica* and *Miliusa* CF. *banacea*. J. Nat. Prod. 57, 68–73.
 627 Hasan, C.M., Jumana, S., Rashid, M.A., 2000. (+)-Isocorydine α -N-oxide: a new
 628 aporphine alkaloid from *Miliusa velutina*. Nat. Prod. Lett. 14, 393–397.
 629 Heyne, K., 1950. De nuttige planten van Indonesie, 3rd ed., Wageningen: H.
 630 Veenman and Konen.
 631 Hishman, A., Harassi, A., Shuaily, W., Echigo, S., Fujimoto, Y., 2000.
 632 Cardiopetalactone: A novel styryllactones from *Goniothalamus*
 633 *cardiopetalus*. Tetrahedron 56, 9985–9989.
 634 Holzbach, J.C., Lope, L.M.X., 2010. Aristolactams and alkaloids of *Aristolochia*
 635 *gigantean*. Molecules 15, 9462–9472.
 636 Hongthong, S., Kuhakarn, C., Jaipetch, T., Prabpai, S., Kongsaree, P.,
 637 Piyachaturawat, P., Jariyawat, S., Suksen, K., Limthongkul, J., Panthong, A.,
 638 Nuntasaeen, N., Reutrakul, V., 2015. Polyoxzygenated cyclohexene derivatives
 639 isolated from *Dasymaschalon sootepense* and their biological activities.
 640 Fitoterapia 106, 158–166.
 641 Hsu., Y.M., Wu, T.Y., Du, Y.C., Shazly, M.E., Beerhues, L., Thang, T.D., Luu, H.V.,
 642 Hwang, T.L., Chang, F.R., Wu, Y.C., 2016. 3-Methyl-4,5,dihydro-oxepine,
 643 polyoxygenated seco-cyclohexenes and cyclohexenes from *Uvaria flexuosa*
 644 and their anti-inflammatory activity. Phytochemistry 122, 184–192.
 645 Ito, J., Chang, F.R., Wang, H.K., Park, Y.K., Ikegaki, M., Kilgore, N., Lee, K.H.,
 646 2001. Anti-AIDS agents. 48.¹ anti-HIV activity of moronic acid derivatives
 647 and the new melliferone-related triterpenoid isolated from Brazilian propolis.
 648 J. Nat. Prod. 64, 1278–1281.

- 649 Ichimaru, M., Nakatani, N., Moriyasu, M., Nishiyama, Y., Kato, A., Mathenge, S.G.,
650 Juma, F.D., Chalomotiso, P.B., 2010. Hydroxyespintanol and
651 schefflerichalcone: two new compounds from *Uvaria scheffleri*. J. Nat. Med.
652 64, 75–79.
- 653 Jiang, M.M., Feng, Y.F., Gao, H., Zhang, X., Tang, J.S., Yao, X.S., 2011. Three new
654 bis-styryllactones from *Goniiothalamus cheliensis*. Fitoterapia 82, 524–527.
- 655 Jiang, M.M., Feng, Y.F., Zhang, X., Yao, X.S., 2011. Chemical constituents from
656 roots of *Goniiothalamus cheliensis* (II). Chin. Tradit. Herbal Drugs. 12, 2386–
657 2388.
- 658 Jiang, M.M., Wen, R., Rui, W., Fu, F.F., Zhao, L., Yao, X., Feng, Y., 2011. Analysis
659 of styryllactones from *Goniiothalamus cheliensis* by UPLC-Q-TOF-MS.
660 China J. Chin. Mat. Med. 36, 1323–1326.
- 661 Jiang, M.M., Zhang, X., Dai, Y., Gao, H., Lui, H.W., Wang, N.L., Ye, W.C., Yao,
662 X.S., 2008. Alkaloids from the root barks of *Goniiothalamus cheliensis*. Chin.
663 Chem. Lett. 19, 302–304.
- 664 Joseph, C.C., Magadula, J.J., Nkunya, M.H.H., 2007. A novel antiplasmodial 3',5'-
665 diformylchalcone and other constituents of *Friesodielsia obovate*. Nat. Prod.
666 Res. 21, 1009–1015.
- 667 Kampong, R., Pompimon, W., Meepowpan, P., Sukdee, S., Sombutsiri, P., Nantasaen,
668 N., Krachodnok, S., 2013. (–)-7-*O*-acetylgoniodiol as cancer chemopreventive
669 agent from *Goniiothalamus griffithii*. Int. J. Chem. Sci. 11, 1234–1246.
- 670 Kawetripob, W., Mahidol, C., Prawat, H., Ruchirawat, S., 2015. Cyclohexene long-
671 chain fatty acid esters from *Uvaria dulcis* (Dunal). Phytochem. Lett. 12,
672 248–251.
- 673 Khallouki, F., Haubner, R., Ulrich, C.M., Owen, R.W., 2011. Ethnobotanical survey,
674 chemical composition, and antioxidant capacity of methanolic extract of the
675 root bark of *Annona cuneata* Oliv. J. Med. Food. 14, 1397–1402.
- 676 Kijjoa, A., Bessa, J., Pinto, M.M.M., Anatachoke, C., Silva, A.M.S., Eaton, G., Herz,
677 W., 2002. Polyoxygenated cyclohexene derivatives from *Ellipeiopsis*
678 *cherrevensis*. Phytochemistry 59, 543–549.
- 679 King, R.W., Bauer, J.D., Brady, S.F., 2009. An environmental DNA-derived type II
680 polyketide biosynthetic pathway encodes the biosynthesis of the pentacyclic
681 polyketide erdacin. Angew. Chem. Int. Ed. 48, 6257–6261.
- 682 Kuo, P.C., Thang, T.D., Huang, G.J., Huang, B.S., Hoa, L.T.M., Yang, M.L., Wu,
683 T.S., 2015. Flavonoids from the fruits of *Desmos cochinchinesis* var. *fulvecens*
684 and their inhibitory effect on NO production. Chem. Nat. Compd. 51,
685 152–155.
- 686 Lekphrom, R., Kanokmedhakul, K., Schevenels, F., Kanokmedhakul, S., 2018.
687 Antimalaria polyoxygenated cyclohexene derivatives from the roots of
688 *Uvaria cherrevensis*. Fitoterapia 127, 420–424.
- 689 Li, Chen., Lee, D., Graf, T.N., Phifer, S.S., Nakanishi, Y., Burgess, J.P., Riswan, S.,
690 Setyowati, F.M., Saribi, A.M., Soejarto, D.D., Farnsworth, N.R., Falkinham,
691 J.O., Kroll, D.J., Kinghorn, A.D., Wani, C.C., Oberlies, N.H., 2005. A

692 hexacyclic *enttrachylobane* diterpenoid possessing an oxetane ring from
 693 *Mitrephora glabra*. Org. Lett. 7, 5709–5712.

694 Li, Chen., Lee, D., Graf, T.N., Phifer, S.S., Nakanishi, Y., Riswan, S., Setyowati,
 695 F.M., Saribi, A.M., Soejarto, D.D., Farnsworth, N.R., Falkinham, J.O., Kroll, D.J.,
 696 Kinghorn, A.D., Wani, M.C., Oberlies, N.H., 2009. Bioactive constituents of
 697 the stem bark of *Mitrephora glabra*. J. Nat. Prod. 72, 1949–1953.

698 Liaw, C.C., Chang, F.R., Wu, Y.C., Wang, H.K., Nakanishi, Y., Bastow, K.F., Lee,
 699 K.H., 2004. Montacin and *cis*-montacin, two new cytotoxic
 700 monotetrahydrofuran Annonaceae acetogenins from *Annona montana*. J. Nat.
 701 Prod. 67, 1804–1808.

702 Liaw, C.C., Yang, Y.L., Chen, M., Chang, F.R., Chen, S.L., Wu, S.H., Wu, Y.C.,
 703 2008. Mono-tetrahydrofuran Annonaceae acetogenins from *Annona*
 704 *squamosa* as cytotoxic agents and calcium ion chelators. J. Nat. Prod. 71,
 705 764–771.

706 Li, W., Wiesenfeldt, M.P., Glorius, F., 2017. Ruthenium-NHC-diamine catalyzed
 707 enantioselective hydrogenation of isocoumarins. 139, 2585–2588.

708 Liping, Z., Meigua, J., Changqi, H., 1996. Studies on isoflavones from Chinese
 709 peashrub (*Caragana sinica*). Zhongcaoyao 27, 134–136.

710 Liou, J.R., Wu, T.Y., Thang, T.D., Hwang, T.L., Wu, C.C., Cheng, Y.B., Chiang,
 711 M.Y., Lan, Y.H., El-Shazly, M., Wu, S.L., Beerhues, L., Yuan, S.S., Hou,
 712 M.F., Chen, S.L., Chang, F.R., Wu, Y.C., 2014. Bioactive 6*S*-styryllactone
 713 constituents of *Polyalthia parviflora*. J. Nat. Prod. 77, 2626–2632.

714 Lui, Y.P., Tang, J.Y., Hua, Y., Lai, L., Luo, X.L., Zhang, Z.J., Yin, W.Q., Chen,
 715 G.Y., Fu, Y.H., 2018. Bioactive polyoxygenated seco-cyclohexenes from
 716 *Artabotrys hongkongensis*. Bioorg. Chem. 76, 386–391.

717 Lui, Y.P., Wang, X.C., Li, X.B., Li, K.K., Huang, L.G., Wen, C.Q. Fu, Y.H., 2015.
 718 Studies on non-alkaloid constituents from *Ochrosia elliptica*. Chin. J. Chin.
 719 Mat. Med. 40, 1508–1513.

720 Macabeo, A.P.G., Lopez, A.D.A., Schmidt, S., Heilmann, J., Dahse, H.M., Alejandro,
 721 G.J.D., Franzblau, S.G., 2013. Antitubercular and cytotoxic constituents from
 722 *Goniathalamus gitingensis*. Rac. Nat. Prod. 8:1, 41–45.

723 Macabeo, A.P.G., Rubio, P.Y.M., Higuchi, T., Umezawa, N., Federl, C., Budde, S.,
 724 Bangcaya, P., Alejandro, G.J.D., 2017. Polyoxygenated *seco*-cyclohexenes
 725 and other constituents from *Uvaria valderramensis*. Biochem. Syst. Ecol. 71,
 726 200–204.

727 Macabeo, A.P.G., Letada, A.G., Budde, S., Federl, C., Dahse, H.M., Franzblau, S.G.,
 728 Alejandro, G.J.D., Pierens, G.K., Garson, M.J., 2017. Antitubercular and
 729 cytotoxic chlorinated *seco*-cyclohexenes from *Uvaria alba*. J. Nat. Prod. 80,
 730 3319–3323.

731 Meesakul, P., Pudhom, K., Pyne, S.G., Laphookhieo, S., 2017. Hydride flavan-
 732 flavanones from *Friesodielsia desmodies* and their inhibitory activities against
 733 nitric oxide production. RSC Adv. 7, 17545–17550.

734 Melot, A., Fall, D., Gleye, C., Champy, P., 2009. Apolar Annonaceae acetogenins
 735 from the fruit pulp of *Annona muricata*. Molecules 14, 4387–4395.

736 Meng, D.H., Xu, Y.P., Chen, W.L., Zou, J., Lou, L.G., Zhao, W.M., 2007. Anti-tumor
 737 clerodane-type diterpenes from *Mitrephora thorelii*. J. Asian Nat. Prod. Res. 9,
 738 679–684.
 739 Ming, H., Jinghua, Z., Changqi, H., 2001. Studies on the chemical components of
 740 *Clematis chinensis*. J. Chin. Pharm. Sci. 10, 180–182.
 741 Moghadamtousi, S.Z., Fadaeinasab, M., Nikzad, S., Mohan, G., Ali, H.M., Kadir,
 742 H.A., 2015. *Annona muricata* (Annonaceae): A review of its traditional uses,
 743 isolated acetogenins and biological activities. Int. J. Mol. Sci. 16,
 744 15625–15658.
 745 Moharam, B.A., Jantan, I., Jalil, I., Ahmad, F., 2012. Inhibitory effect of compounds
 746 from *Goniiothalamus tapis* Miq. and *Goniiothalamus uvaroides* king on
 747 platelet-activating factor receptor binding. Phyother. Res. 26, 687–691.
 748 Nakasone, Y., Takara, K., Wada, K., Tanaka, J., Yogi, S., Nakatani, N., 1996.
 749 Antioxidative compounds isolated from kakuto, non-centrifugal cane sugar.
 750 Biosci. Biotech. Biochem. 60, 1714–1716.
 751 Nantapap, S., Sangrueng, K., Nuntasen, N., Meepowpan, P., Pompimon, W., 2015.
 752 Chemical constituents from aerial parts of *Polyalthia evecta* (Pierre) Finet &
 753 Gagnep. Var. *attopeuensis*. Int. J. Chem. Sci. 13, 1705–1712.
 754 Nurunajah, G., Ahmat, N., Hadiani, I.N., Ishak, Z., 2011. Flavonoid constituents from
 755 the stem bark of *Polyalthia cauliflora* var. *cauliflora*. Aust. J. Basic Appl. Sci.
 756 5, 154–158.
 757 Nyandoro, S.S., Munissi, J.J., Gruhonjic, A., Duffy, S., Pan, F., Puttreddy, R.,
 758 Holleran, J.P., Fitzpatrick, P.A., Pelletier, J., Avery, V.M., Rissanen, K.,
 759 Erdélyi, M., 2017. Polyoxygenated cyclohexenes and other constituents of
 760 *Cleistochlamys kirkii* leaves. J. Nat. Prod. 80, 114–125.
 761 Ortega, A.R., Toscano, R.A., Barragán, A.H., Cisneros, C.A., Nathan, P.J., 2015.
 762 Structure elucidation of a new isoflavone by exclusive use of ¹H NMR
 763 measurements. Magn. Reson. Chem. 53, 860–865.
 764 Pancharoen, O., Athipornchai, A., Panthong, A., Taylor, W.C., 2008. Isoflavone and
 765 rotenoids from the leaves of *Millettia brandisiana*. Chem. Pharm. Bull. 56,
 766 835–838.
 767 Popsavin, V., Kavačević, I., Benedeković, G., Popsavin, M., Kojoć, V., Bogdanović.,
 768 2012. Divergent synthesis of cytotoxic styryl lactones related to
 769 goniobutenolides A and B, and to crassalactone D. Org. Lett. 14,
 770 5956–5959.
 771 Prachyawarakorn, V., Sangpetsiripan, S., Surawatanawong, P., Mahidol, C.,
 772 Ruchirawat, S., Kittakoop, P., 2013. Flavans from *Desmos choichinchinensis*
 773 as potent aromatase inhibitors. Med. Chem. Commun. 4, 1590–1596.
 774 Prawat, U., Chaimanee, S., Butsuri, A., Salae, A.W., Tuntiwachwuttikul, P., 2012.
 775 Bioactive styryllactones, two new naphthoquinones and one new
 776 styryllactones, and other constituents from *Goniiothalamus scortechinii*.
 777 Phytochem. Lett. 5, 529–534.

778 Prawat, U., Chairerk, O., Phupornprasert, U., Salae, A.W., Tuntiwachwuttikul, P.,
 779 2013. Two new C-benzylated dihydrochalcone derivatives from the leaves of
 780 *Melodorum siamensis*. *Planta. Med.* 79, 83–86.
 781 Prawat, U., Phupornprasert, D., Butsuri, A., Salae, A.W., Boonsri, S.,
 782 Tuntiwachwuttikul, P., 2012. Flavonoids from *Friesodielsia discolor*.
 783 *Phytochem. Lett.* 5, 809–813.
 784 Promchai, T., Jaidee, A., Cheenpracha, S., Trisuwas, K., Rattanajak, R.,
 785 Kamchonwongpaisan, S., Laphookhieo, S., Pyne, S.G., Ritthiwigrom, T.,
 786 2016. Antimalarial oxoprotoberberine alkaloids from the leaves of *Miliusa*
 787 *cuneate*. *J. Nat. Prod.* 79, 978–983.
 788 Promchai, T., Saesong, T., Ingkaninan, K., Laphookhieo, S., Pyne, S.G., Limtharakul,
 789 T., 2018. Acetylcholinesterase inhibitory activity of chemical constituents
 790 isolated from *Miliusa thorelii*. *Phytochem. Lett.* 23, 33–37.
 791 Queiroz, E.F., Roblot, F., Laprévote, O., Paulo, M.Q., Hocquemiller, R., 2003. Two
 792 unusual acetogenins from the roots of *Annona salzmanii*. *J. Nat. Prod.* 66,
 793 755–758.
 794 Quisumbing, E., 1951. Medicinal plants of the Philippines. Manila, Philippines.
 795 Rahman, A., Katayama, T., Suzuki, T., Nakagawa, T., 2007. Stereochemistry and
 796 biosynthesis of (+)-lyoniresinol, a syringyl tetrahydronaphthalene lignan in
 797 *Lyonia ovalifolia* var. *elliptical* I: isolation and stereochemistry of syringyl
 798 lignans and predicted precursors to (+)-lyoniresinol from wood. *J. Wood. Sci.*
 799 53, 161–167.
 800 Rittiwong, T., Mutarapat, T., Ponglimanont, C., Mahabusarakam, W., Chakthong, S.,
 801 2011. Saiyutones A–D: four new unusual biflavones from *Desmos chinensis*.
 802 *Tetrahedron* 67, 5444–5449.
 803 Rukachaisirikul, V., Khamthong, N., Sukpondma, Y., Phongpaichit, S., Towatana,
 804 N.H., Graidist, P., Sakayaroj, J., Kirtikara, K., 2010. Cyclohexene
 805 diketopiperazine, lactone and phenol derivatives from the sea fan-derived
 806 fungi *Nigrospora* sp. PSU-F11 and PSU-F12. *Arch. Pharm. Res.* 33, 375–380.
 807 Sampath, M., Vasanthi, M., 2013. Isolation, structural elucidation of flavonoids from
 808 *Polyalthia longifolia* (SONN.) *thawaites* and evaluation of antibacterial,
 809 antioxidant and anticancer potential. *Int. J. Pharm. Pharm. Sci.* 5, 336–341.
 810 Salum, M.L., Robles, C.J., Erra-Balsells, R., 2010. Photoisomerization of ionic liquid
 811 ammonium cinnamates: one-pot synthesis–isolation of Z-cinnamic acids. 12,
 812 4808–4811.
 813 Saripalli, H.R., Dixit, P.K., 2013. Studies on morphological features and biological
 814 activities of the genus *annonia* of Ethiopia, N. E. Africa with a special
 815 emphasis on *Graviola*: A review. *Int. J. Pharm. Sci. Res.* 821–827.
 816 Saunders, R.M.K., Chalermglin, P., 2008. A synopsis of *Goniothalamus* species
 817 (Annonaceae) in Thailand, with descriptions of three new species. *Bot. J.*
 818 *Linn. Soc.* 156, 355–384.
 819 Sheldrick, G.M., 2015. SHELXT-integrated space-group and crystal-structure
 820 determination. *Acta Cryst.* A71, 3–8

821 Sheldrick, G.M., 2015. Crystal structure refinement with SHELXL. *Acta Cryst.* C71,
822 3–8.

823 Shing, T.K.M., Tsui, H.C., Zhou, Z.H., 1995. Enantiospecific syntheses of (+)-
824 goniofufurone, (+)-7-*epi*-goniofufurone, (+)-goniobutenolide A, (–)-
825 goniobutenolide B, (+)-goniopypyrone, (+)-altholactone, (+)-goniotriol, and
826 (+)-7-acetylgoniotriol. *J. Org. Chem.* 60, 3121–3130.

827 Shing, T.K.M., Tai, V.W.F., Tsui, H.C., 1994. Goniobutenolides A and B;
828 serendipitous syntheses, relative and absolute configuration. *J. Chem. Soc.*
829 *Chem. Commun.* 1293–1294.

830 Si, W., Zhang, Y.J., Chen, R.Y., Yu, D.Q., 2002. Goniolactones A-F, six new
831 styrylpyrone derivatives from the roots of *Goniiothalamus cheliensis*. *J. Nat.*
832 *Prod.* 65, 835–841.

833 Singh, O., Muthukrishnan, M., 2005. Synthesis of isoflavones containing naturally
834 occurring substitution pattern by oxidative rearrangement of respective
835 flavanones using thallium(III) *p*-tosylate. *Indian J. Chem.* 44B, 2575–2581.

836 Sriyatep, T., Andersen, R.J., Patrick, B.O., Pyne, S.G., Muanprasat, C., Seemakham,
837 S., Borwornpinyo, S., Laphookhieo, S., 2017. Scalemic caged xanthon
838 isolated from the stem bark extract of *Garcinia propinqua*. *J. Nat. Prod.* 80,
839 1658–1667.

840 Starks, C.M., Williams, R.B., Rice, S.M., Norman, V.L., Lawrence, J.A., Goering,
841 M.G., Johnson, M.O.N., Hu, J.F., Eldridge, G.R., 2012. Polyoxygenated
842 cyclohexene derivatives from *Monanthotaxis congoensis*. *Phytochemistry* 74,
843 185–189.

844 Suchaichit, N., Kanokmedhakul, K., Panthama, N., Poopasit, K., Moosophon, P.,
845 Kanokmedhakul, S., 2015. A 2H-tetrahydropyran derivative and bioactive
846 constituents from the bark of *Goniiothalamus elegants* Ast. *Fitoterapia* 103,
847 206–212.

848 Sun, L., Zhao, R., Lan, X., Chen, R., Wang, S., Du, G., 2014. Goniolactone C, a styryl
849 lactone derivative, inhibits PDGF-BB-induced vascular smooth muscle cell
850 migration and proliferation via PDGFR/ERK signaling. *Molecules* 19, 19501–
851 19515.

852 Sun, S., Liu, J., Sun, X., Zhu, W., Yang, F., Felczak, L., Dou, Q.P., Zhou, K., 58.
853 Novel Annonaceous acetogenins from Graviola (*Annona muricata*) fruits with
854 strong anti-proliferative activity. *Tetrahedron Lett.* 58, 1895–1899.

855 Supudompol, B., Chaowasku, T., Kingfang, K., Burud, K., Wongseripipatana, S.,
856 Likhitwitayawuid, K., 2004. A new pimarane from *Mitrephora tomentosa*.
857 *Nat. Prod. Res.* 18, 387–390.

858 Surivet, J.P., Vatéle, J.M., 1999. Total synthesis of antitumor *Goniiothalamus*
859 styryllactones. *Tetrahedron* 55, 13011–13028.

860 Tahara, S., Hashidoko, Y., Ingham, J.L., Mizutani, J., 1986. New 5-*O*-
861 methylisoflavone in the roots of yellow lupin (*Lupinus luteus* L. cv. Barpine).
862 *Agric. Biol. Chem.* 50, 1809–1819.

- Tip-pyang, S., Limpipatwattana, Y., Khumkratok, S., Siripong, P., Sichaem, J. 2010. A new cytotoxic 1-azaanthraquinone from the stems of *G. laoticus*. *Fitoterapia* 81, 894–896.
- Tsubuki, M., Kanai, K., Nagase, H., Honda, T., 1999. Stereocontrolled syntheses of novel styryl lactones, (+)-goniodiol, (+)-goniotriol, (+)-8-acetylgoniotriol, (+)-goniofufurone, (+)-9-deoxygoniopypyrone, (+)-goniopypyrone, and (+)-altholactone from common intermediates and cytotoxicity of their congeners. *Tetrahedron* 55, 2493–2514.
- Tuntipaleepun, M., Chakthong, S., Ponglimanont, C., Plodpai, P., Voravuthikunchai, S.P., 2012. Antifungal and cytotoxic substances from the stem barks of *Desmos chinensis*. *Chin. Chem. Lett.* 23, 587–590.
- Wagoner, R.M.V., Mantle, P.G., Wright, J.L.C., 2008. Biotsynthesis of scorinone, a 2-aaanthraquinone from *Amorosia littoralis*, a fungus from marin sediment. *K. Nat. Prod.* 71, 426–430.
- Wang, S., Zhang, Y.J., Chen, R.Y., Yu, D.Q., 2002. Goniolactones A–F, six new styrylpyrone derivatives from the roots of *Goniothalamus cheliensis*. *J. Nat. Prod.* 65, 835–841.
- Wang, S., Dai, S.J., Zhang, P.C., Chen, R.Y., Zeper, A., Yu, S.S., Yu, D.Q., 2003. Studies on the chemical constituents of the roots of *Goniothalamus cheliensis* and *Uvaria macrophylla*. *Acta Chin. Sinica.* 61, 1090–1096.
- Waterman, P. G., Muhammad, I., 1985, Sesquiterpenes and alkaloids from *Cleistopholis patens*. *Phytochemistry* 24, 523–527.
- Wu, M.D., Cheng, M.J., Chen, I.S., Su, Y.S., Hsieh, S.Y., Chang, H.S., Chang, C.W., Yuan, G.F., 2013. Phytochemistry investigation of *Annulohypoxyton ilanense*, an endophytic fungus derived from *Cinnamomun* species. *Chem. Biodivers.* 10, 493–505.
- Wu, T.S., Chan, Y.Y., Leu, Y.L., 2000. The constituents of the root and stem of *Aristolochia cucurbitifolia* Hayata and their biological activity. *Chem. Pharm. Bull.* 48, 1006–1009.
- Wu, Y.C., Chang, F.R., Duh, C.Y., Wang, S.K., Wu, T.S., 1992. Cytotoxic Styrylpyrones of *Goniothalamus amuyon*. *Phytochemistry* 31, 2851–2853.
- Xu, D.X., Sharpless, K.B., 1994. Synthesis and stereochemical assignments for goniobutenolides A and B. *Tetrahedron Lett.* 35, 4685–4688.
- Yamauchi, S., Ichikawa, H., Nishiwaki, H., Shuto, Y., 2015. Evaluation of plant growth regulatory activity of furofuran lignan bearing a 7,9':7',9-diepoxy structure using optical pure (+)- and (–)-enantiomers. *J. Agric. Food Chem.* 63, 5224–5228.
- Zdero, C., Bohlmann, F., King, R.M., Robinson, H., 1986. Diterpene glycosides and other constituents from Argentinian *Baccharis* species. *Phytochemistry* 12, 2841–2855.
- Zhang, Y.J., Kong, M., Chen, R.Y., Yu, D.Q., 1999. Alkaloids from the roots of *Goniothalamus griffithii*. *J. Nat. Prod.* 62, 1050–1052.
- Zhang, Y.J., Zhou, G.X., Chen, R.Y., Yu, D.Q., 1998. Styryllactones from the rhizomes of *Goniothalamus griffithii*. *J. Asian Nat. Prod. Res.* 1, 189–197.

907 Zhou C.C., Lui, C.T., Huang, X.X., Wu, J., Li, L.Z., Li, D.M., Song, S.J., 2013.
 908 Isolation and identification of aromatic compounds from the leaves of
 909 *Crataegus pinnatifida* Bge. Chin. J. Med. Chem. 23, 213–217.
 910 Zhou, Q., Fu, Y.H., Zhang, Y.Q., Wu, S.Y., Song, X.P., Cao, C.K., Xu, W., Chen,
 911 G.Y., 2016. Fissitungfines A and B, two novel aporphine related alkaloids
 912 from *Fissistigma tungfangense*. Tetrahedron Lett. 57, 4162–4164.
 913 Zhou, Q., Fu, Y.H., Zhang, Y.Q., Wu, S.Y., Song, X.P., Cao, C.K., Xu, W., Chen,
 914 G.Y., 2018. Bioactive aporphine alkaloids from *Fissistigma tungfangense*.
 915 Phytochem. Lett. 25, 105–108.
 916 Zhou W., Zhuang, Y., Bai, Y., Bi, H., Liu, T., Ma, Y., 2016. Biosynthesis of
 917 phlorisovalerophenone and 4-hydroxy-6-isobutyl-2-pyrone in *Escherichia coli*
 918 from glucose. Microb. Cell Fact. 15, 149.
 919 Zhu, J.X., Yu, D.L., Huang, W.H., Sun, L., Lv, Y., Zheng, Q.T., Lee, K.H., Yu, J.G.,
 920 2012. Goniodilactone and gonioheptenolactone, two novel cytotoxic
 921 styryllactones from the leaves of *Goniotalamus cheliensis*. Chin. Chem. Lett.
 922 23, 583–586.
 923

Graphical abstract

Alkaloids and styryllactones from *Goniothalamus cheliensis*

Wuttichai Jaidee, Raymond J. Andersen, Brian O. Patrick, Stephen G. Pyne, Chatchai Muanprasat, Suparek Borwornpinyo, Surat Laphookhieo *

Goniothalamus cheliensis

Eight previously undescribed compounds were isolated and characterized from *Goniothalamus cheliensis*. Griffithazanone A showed potent cytotoxicities against human colorectal cancer cells (HCT-116).

Figure legends

Fig. 1. The structures of compounds from *G. cheliensis* twigs.

Fig. 2. The structures of compounds from *G. cheliensis* leaves.

Fig. 3. The key COSY and HMBC correlations of **1–3** and **5–9**.

Fig. 4. The key NOESY correlations of **6** and **7** (molecular structures generated using 3D structure ChemBioDraw Ultra 11.0).

Fig. 5. Single crystal X-ray structure of **1**.

Fig. 6. Single crystal X-ray structure of **4**.

Fig. 7. Single crystal X-ray structure of **6**.

Fig. 8. Single crystal X-ray structure of **7**.

Fig. 9. ECD spectra of **6** and **7**.

Fig. 10. ECD spectra of **8–11**.

Fig. 11. Plausible biosynthetic pathway of **1–4** (**a**) and **5** (**b**).

Fig. 1.

Fig. 2.

Fig. 3.

Fig.4.

Fig. 5.

Fig. 6.

Fig. 7.

Fig. 8.

Fig. 9.

Fig. 10.

Fig. 11.

Table 1¹H (600 MHz) and ¹³C (150 MHz) NMR Spectroscopic data of **1–3**.

position	1 (CDCl ₃)		2 (CDCl ₃)		3 (methanol- <i>d</i> ₄)	
	δ_{C} , type	δ_{H} (<i>J</i> in Hz)	δ_{C} , type	δ_{H} (<i>J</i> in Hz)	δ_{C} , type	δ_{H} (<i>J</i> in Hz)
2	138.0, CH	7.74, s	137.5, CH	7.73, s	130.2, CH	7.81, s
3	114.5, C		116.4, C		115.2, C	
3a	123.5, C		123.1, C		124.0, C	
4	183.1, C		184.0, C		179.1, C	
4a	132.0, C		132.6, C		133.7, C	
5	120.7, CH	7.82, d (7.4)	127.4, CH	8.26, dd (7.4, 1.3)	124.0, CH	8.14, dd (7.8, 1.2)
6	136.0, CH	7.64, dd (7.8, 7.4)	134.5, CH	7.79, td (7.5, 1.5)	132.1, CH	7.78, dd (7.2, 1.2)
7	126.1, CH	7.32, d (7.8)	133.6, CH	7.81, td (7.5, 1.5)	131.4, CH	7.77, ddd (7.8, 7.2, 1.2)
8	162.5, C		126.6, CH	8.21, dd (7.4, 1.3)	125.2, CH	8.19, dd (7.8, 1.2)
8a	114.5, C		132.6, C		133.0, C	
9	180.9, C		175.5, C		175.3, C	
9a	130.9, C		131.8, C		134.4, C	
10	161.5, C		162.1, C		163.2, C	
<i>N</i> -Me	37.5, CH ₃	4.15, s	37.3, CH ₃	4.16, s		
8-OH		12.20, s				
3-CO ₂ <i>Me</i>					49.3, CH ₃	3.89, s
3-COOH		13.68, s		13.70, s		

Table 2¹H (600 MHz) and ¹³C (150 MHz) NMR Spectroscopic data of **5** and goniiothalamine B.

position	5 (DMSO- <i>d</i> ₆)		5 (CDCl ₃)		goniiothalamine B (CDCl ₃)	
	δ_C , type	δ_H (<i>J</i> in Hz)	δ_C , type	δ_H (<i>J</i> in Hz)	δ_C , type	δ_H (<i>J</i> in Hz)
1	178.1, C		178.0, C		180.2, C	
2	107.5, C		106.9, C		109.4, C	
3	155.0, C		158.4, C		152.5, C	
4	187.7, C		186.3, C		185.5, C	
4a	114.3, C		113.4, C		114.0, C	
5	151.6, C		151.8, C		152.0, C	
6	155.8, C		155.8, C		152.1, C	
7	114.7, CH	7.29, d (8.4)	113.2, CH	7.05, d (8.4)	117.8, C	7.22, d (8.4)
8	120.7, CH	7.62, d (8.4)	121.0, CH	7.75, d (8.4)	120.7, C	7.76, d (8.4)
8a	121.8, C		126.7, CH		125.0, C	
5-OH		13.94, s		13.97, s		11.71, s
6-OMe	56.2, CH ₃	3.91, s	56.0, CH ₃	4.02, s	56.4, CH ₃	3.99, s
NH ₂		9.12, br s				7.07, br s
		10.68, br s				10.66, br s
COMe	199.9, C		200.7, C		202.4, C	
COMe	33.2, CH ₃	2.56, s	33.0, CH ₃	2.73, s	33.2, CH ₃	2.73, s

Table 3¹H (600 MHz) and ¹³C (150 MHz) NMR Spectroscopic data of **6** and **7**.

position	6 (DMSO- <i>d</i> ₆)		7 (CDCl ₃)	
	δ_C , type	δ_H (<i>J</i> in Hz)	δ_C , type	δ_H (<i>J</i> in Hz)
2	168.7, C		167.0, C	
3 _{ax}	35.2, CH ₂	2.78, d (19.3)	34.6, CH ₂	3.15, dd (19.6, 1.2)
3 _{eq}		3.07, d (19.3, 5.2)		3.02, dd (19.6, 5.2)
4	69.3, CH	4.07, d (4.6)	69.7, CH	4.38, m
5	64.7, CH	4.15, td (4.3, 1.6)	65.3, CH	4.33, dd (4.6, 1.6)
6	78.4, CH	4.45, m	74.5, CH	4.91, dt (4.6, 2.6)
7	67.2, CH	3.87, ddd (9.8, 7.2, 2.3)	69.1, CH	5.37, dd (10.3, 2.6)
8	72.9, CH	4.24, d (9.8)	70.6, CH	4.67, d (10.3)
1'	139.7, C		136.0, C	
2'	128.7, CH	7.38, d (7.9)	126.7, CH	7.41, m
3'	127.7, CH	7.30, t (7.6)	128.1, CH	7.37, m
4'	127.6, CH	7.26, t (7.2)	128.6, CH	7.34, m
5'	127.7, CH	7.30, t (7.6)		7.37, m
6'	128.7, CH	7.38, d (7.9)		7.41, m
5-OH		5.70, d (4.0)		
7-OH		5.16, d (6.9)		
7-OCOMe			169.1, C	
7-OCOMe			20.2, CH ₃	1.94, s

Table 4¹H (600 MHz) and ¹³C (150 MHz) NMR Spectroscopic data of **8** and **9**.

position	8 (acetone- <i>d</i> ₆)		9 (CDCl ₃)	
	δ_{C} , type	δ_{H} (<i>J</i> in Hz)	δ_{C} , type	δ_{H} (<i>J</i> in Hz)
2	168.2, C		169.6, C	
3	119.6, CH	6.31, d (5.5)	121.2, CH	6.19, dd (5.6, 1.5)
4	114.0, CH	7.74, d (5.5)	139.6, CH	7.50, d, (5.6)
5	149.8, C		152.4, C	
6	112.9, CH	5.46, d (9.2)	107.0, CH	5.82, dd (10.0, 1.5)
7	67.8, CH	4.98, dd (9.2, 5.3)	72.3, CH	5.69, dd (10.0, 4.2)
8	76.5, CH	5.85, d (5.3)	74.8, CH	5.06, d (4.2)
1'	135.8, C		137.4, C	
2'	126.7, CH	7.40, d (7.4)	125.9, CH	
3'	127.5, CH	7.33, t (7.4)	128.1, CH	7.34-7.38, m
4'	127.3, CH	7.27, t (7.3)	128.8, CH	
5'	127.5, CH	7.33, t (7.4)	128.1, CH	
6'	126.7, CH	7.40, d (7.4)	125.9, CH	
8-OCOMe	168.0, C		168.2, C	
8-OCOMe	19.5, CH ₃	2.00, overlapped	21.0, CH ₃	2.07, s