UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers:
Part B

Faculty of Science, Medicine and Health

2018

Antibacterial and cytotoxic activities of phenolic constituents from the stem extracts of *Spatholobus parviflorus*

Thanika Promchai
Chiang Mai University

Pawinee Janhom
Chiang Mai University

Wisanu Maneerat
Mae Fah Luang University

Roonglawan Rattanajak
National Science and Technology Development Agency

Sumalee Kamchonwongpaisan
National Science and Technology Development Agency

See next page for additional authors

Publication Details

Promchai, T., Janhom, P., Maneerat, W., Rattanajak, R., Kamchonwongpaisan, S., Pyne, S. G. & Limtharakul, T. (2018). Antibacterial and cytotoxic activities of phenolic constituents from the stem extracts of *Spatholobus parviflorus*. *Natural Product Research*, Online First 1-5.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Antibacterial and cytotoxic activities of phenolic constituents from the stem extracts of *Spatholobus parviflorus*

Abstract

A new 2-arylbenzofuran, spathobenzofuran (1), together with ten known compounds including a 2-arylbenzofuran, three pterocarpanes and six isoflavones were isolated from the acetone crude extract of the stems of *Spatholobus parviflorus*. All compounds were characterised by spectroscopic methods. Compound 4 was active (MIC 8 µg/mL) against Gram-negative *Pseudomonas aeruginosa* TISTR 781 while compound 2 had modest activity against Gram-positive *Staphylococcus aureus* TISTR 1466 with a MIC value of 16 µg/mL. All isolated compounds showed no cytotoxicity against Vero and KB cells.

Publication Details

Promchai, T., Janhom, P., Maneerat, W., Rattanajak, R., Kamchonwongpaisan, S., Pyne, S. G. & Limtharakul, T. (2018). Antibacterial and cytotoxic activities of phenolic constituents from the stem extracts of *Spatholobus parviflorus*. *Natural Product Research*, Online First 1-5.

Authors

Thanika Promchai, Pawinee Janhom, Wisanu Maneerat, Roonglawan Rattanajak, Sumalee Kamchonwongpaisan, Stephen G. Pyne, and Thunwadee Limtharakul

Antibacterial and cytotoxic activities of phenolic constituents from the stem extracts of *Spatholobus parviflorus*

Thanika Promchai^{a,b}, Pawinee Janhom^a, Wisanu Maneerat^c, Roonglawan Rattanajak^d, Sumalee Kamchonwongpaisan^d, Stephen G. Pyne^e and Thunwadee Limtharakul^{a,f} (née Ritthiwigrom)

^aDepartment of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Chiang Mai University, Chiang Mai, Thailand; ^bThe Graduate School, Chiang Mai University, Chiang Mai, Thailand; ^cSchool of Science, Mae Fah Luang University, Chiang Rai, Thailand; ^dMedical Molecular Biology Research Unit, Natural Center for Genetic Engineering and Biotechnology, National Science and Technology Development Agency, Pathumthani, Thailand; ^eSchool of Chemistry, University of Wollongong, Wollongong, NSW, Australia; ^fResearch Center on Chemistry for Development of Health Promoting Products from Northern Resources, Chiang Mai University, Chiang Mai, Thailand.

Contact: Thunwadee Limtharakul, *e-mail* address: thunwadee.r@cmu.ac.th

Antibacterial and cytotoxic activities of phenolic constituents from the stem extracts of *Spatholobus parviflorus*

ABSTRACT

A new 2-arylbenzofuran, spathobenzofuran (**1**), together with ten known compounds including a 2-arylbenzofuran, three pterocarpanes and six isoflavones were isolated from the acetone crude extract of the stems of *Spatholobus parviflorus*. All compounds were characterised by spectroscopic methods. Compound **4** was active (MIC 8 µg/mL) against Gram-negative *Pseudomonas aeruginosa* TISTR 781 while compound **2** had modest activity against Gram-positive *Staphylococcus aureus* TISTR 1466 with a MIC value of 16 µg/mL. All isolated compounds showed no cytotoxicity against Vero and KB cells.

Keywords: *Spatholobus parviflorus*; arylbenzofuran; antibacterial activity; cytotoxicity

1. Introduction

Spatholobus parviflorus (Fabaceae) is a woody climbing plant known as “Thao Pun Sai” and “Kwang Phu” in Thai. Four species from this genus have been found in Thailand (Smitinand 2014). A water decoction of the stems and leaves (Mahidol University 1996) of *S. parviflorus* is used in traditional medicine by tribal communities in the North of Thailand to treat indigestion, gastritis, menstrual pain, and other diseases. Plants of the genus *Spatholobus* are a major source of flavonoids (Yoon et al. 2004; Tang et al. 2012; Wang et al. 2017). Recently, Sichaem and co-workers reported the isolation of flavonoids and phenolic derivatives and their cholinesterase activity from the extracts of the stems of *S. parviflorus* (Sichaem et al. 2018). From our ongoing search for bioactive compounds from Thai medicinal plants, we report here the isolation, structural characterisation, and bioactivities of a new 2-arylbenzofuran (**1**) as well as ten known compounds (**2-11**) from the stem extracts of this plant. The isolation of 2-arylbenzofurans and compounds **3**, **5**, **9**, and **11** from plants of this genus are reported here for the first time.

2. Results and discussion

The dried stems of *S. parviflorus* were extracted with acetone and the crude extract was separated by column chromatography which led to the isolation of a new 2-

arylbenzofuran, spathobenzofuran (**1**), and ten known compounds; vignafuran (**2**) (Csékei et al. 2008), lyratin B (**3**) (Zhang et al. 2010), (6a*R*, 11a*R*)-medicarpin (**4**) (Yoon et al. 2016), phaseollidin (**5**) (Dagne et al. 1993), daidzein (**6**) (Takeya and Itokawa 1982), formononetin (**7**) (Granados-Covarrubias and Maldonado 2009), genistein (**8**) (Huang et al. 2013), 3'-methoxydaidzein (**9**) (Goto et al. 2009), 8-*O*-methylretusin (**10**) (Mizuno et al. 1990) and 7,3'-dihydroxy-8,4'-dimethoxyisoflavone (**11**) (Puebla et al. 2010) as shown in Figure 1. The structures of all isolated compounds were elucidated by spectroscopic methods, especially 1D and 2D NMR spectroscopy, mass spectrometry and physical data.

Compound **1** was obtained as a yellow gum with a molecular formula of C₁₇H₁₄O₆ from the HRESI-MS ion peak at m/z 315.0867 [M+H]⁺ (calculated for C₁₇H₁₅O₆, 315.0869). The UV spectrum showed maximum absorption bands at λ_{max} 247, 289 and 341 nm which corresponded to a 2-arylbenzofuran derivative (Abdel-Kader 2001; Belofsky et al. 2006; Wu et al. 2018). The IR spectrum show stretching bands for a hydroxy group and a carboxaldehyde group at 3362 and 1700 cm⁻¹, respectively. The ¹H NMR spectrum of compound **1** (Table S1) showed resonances for a carboxaldehyde group [δ_{H} 9.93 (*s*, 1H)], two isolated aromatic protons [δ_{H} 7.60, 7.00 (each *s*, 1H)], a 1,2,4-trisubstituted benzene [δ_{H} 7.42 (*d*, J = 8.4 Hz, 1H), 6.61 (*d*, J = 2.2 Hz, 1H), 6.55 (*dd*, J = 8.4 and 2.2 Hz, 1H)] and two methoxy groups [δ_{H} 3.94, 3.83 (each *s*, 3H)]. The ¹³C NMR spectrum (Table S1) indicated resonances for 17 carbons, including a carboxaldehyde group (δ_{C} 189.7), two methoxy groups (δ_{C} 56.9 and 56.1) and 14 aromatic carbons (9 ArC and 5 ArCH) which contributed further proof that compound **1** contained a core 2-arylbenzofuran structure. From 2D NMR spectral data, the low field singlet aromatic proton resonance at δ_{H} 7.60 was assigned to H-4 (δ_{C} 104.1) on the basis of its HMQC spectrum and its HMBC correlations with C-3 (δ_{C} 118.6), C-6 (δ_{C} 150.7) and C-8 (δ_{C} 147.3). The carboxaldehyde group (δ_{H} 9.93) was located at C-3 of the furan ring from the HMBC correlations of the aldehyde proton with C-3 and C-9 (δ_{C} 117.7). The other singlet aromatic proton, resonating at δ_{H} 7.00 was attached to C-7 (δ_{C} 98.9) on the basis of the HMQC spectrum and the HMBC correlations of H-7 with C-5 (δ_{C} 147.9), C-6 and C-9. The substituents at C-5 and C-6 were identified as a methoxy group (δ_{H} 3.94) from the HMBC spectrum and a hydroxy group based on the downfield ¹³C NMR chemical shift of C-6 (δ_{C} 150.7), respectively. While the C-8 position (δ_{C} 147.3) was connected to the oxygen atom that

formed part of the furan ring. The 1,2,4-trisubstituted benzene ring substituent was linked to C-2 (δ_c 163.1) of the furan ring on the basis of the HMBC correlations of H-6' (δ_H 7.42) with C-2, C-2' (δ_c 160.3) and C-4' (δ_c 164.7). Two aromatic protons resonating at δ_H 6.61 and 6.55 were then attributed to H-3' and H-5', respectively, according to their splitting patterns and coupling constants. The substituents at C-2' and C-4' were identified as a methoxy group (δ_H 3.83), on the basis of the HMBC spectrum, and a hydroxy group, based on its ^{13}C NMR chemical shift (δ_c 164.7), respectively. In support of these assignments for the positions of these methoxy groups, NOESY NMR experiments were performed. These showed cross peaks between the methoxy group at δ_H 3.94 and δ_H 7.60 (H-4), and the methoxy group at δ_H 3.83 and δ_H 6.61 (H-3'). Therefore, the structure of compound **1** was elucidated as 6-hydroxy-2-(4'-hydroxy-2'-methoxyphenyl)-5-methoxy-3-benzofuran-carboxaldehyde.

All compounds, except for **3**, **5**, and **9**, were evaluated for their antibacterial activities against Gram-positive bacteria (*Bacillus cereus*, *Staphylococcus aureus*, and *Staphylococcus epidermidis*) and Gram-negative bacteria (*Escherichia coli*, *Salmonella typhimurium*, *Pseudomonas aeruginosa*, and *Serratia marcescens*). Compound **4** presented the best activity against *P. aeruginosa* with a MIC value of 8 $\mu\text{g/mL}$ while its activity against *B. cereus* was moderate with a MIC value of 32 $\mu\text{g/mL}$. Compound **2** showed antibacterial activity against *S. aureus* with a MIC value of 16 $\mu\text{g/mL}$, while its activity against *E. coli* and *S. typhimurium* was modest with MIC values of 32 $\mu\text{g/mL}$. The other compounds exhibited weak antibacterial activities with MIC values ranging between 64 and 128 $\mu\text{g/mL}$ (Table S2). All isolated compounds were evaluated for their cytotoxicities against human mouth epidermal carcinoma cells (KB) and Vero cells. None of the tested compounds showed cytotoxicity in these assays.

3. Experimental

For the details of the isolation of all compounds see the Supplementary material.

Spathobenzofuran (**1**): yellow gum; UV (MeOH) λ_{max} (log ϵ): 247 (3.5), 289 (3.3), 341 (3.3) nm; IR (neat) ν : 3362, 2851, 2836, 1700 and 1620 cm^{-1} ; ^1H NMR (500 MHz, CD_3OD): δ_H 9.93 (s, 1H, H-10), 7.60 (s, 1H, H-4), 7.42 (d, J = 8.4 Hz, 1H, H-6'), 7.00 (s, 1H, H-7), 6.61 (d, J = 2.2 Hz, 1H, H-3'), 6.55 (dd, J = 8.4 and 2.2 Hz, 1H, H-5'), 3.94 (s, 3H, 5-OCH₃), 3.83 (s, 3H, 6'-OCH₃); ^{13}C NMR (125 MHz, CD_3OD): δ_c 189.7 (C-10),

164.7 (C-4'), 163.1 (C-2), 160.3 (C-2'), 150.7 (C-6), 147.9 (C-5), 147.3 (C-8), 118.6 (C-3), 117.7 (C-9), 110.3 (C-1'), 133.7 (C-6'), 109.1 (C-5'), 104.1 (C-4), 100.5 (C-3'), 98.9 (C-7), 56.9 (5-OCH₃), 56.1 (2'-OCH₃) (HRESI-MS m/z [M+H]⁺ 315.0867 (calcd for C₁₇H₁₅O₆, 315.0869)).

4. Conclusions

A phytochemical investigation of the stems extracts of *S. parviflorus* led to the isolation and identification of a new arylbenzofuran (**1**) and ten known compounds (**2-11**). The arylbenzofurans **1** and **2**, and compounds **3**, **5**, **9**, and **11** were found for the first time from plants in this genus, while the other compounds (**4**, **6**, **7**, **8** and **10**) were found in a very recent study of this plant (Sichaem et al. 2018). In addition, the arylbenzofuran (**2**) was moderately active against a Gram-positive strain (*S. aureus*) while pterocarpan (**4**) showed slightly better activity against a Gram-negative strain (*P. aeruginosa*). These compounds showed no cytotoxicity to KB and Vero cells.

Acknowledgements

This paper is dedicated to memory of Mr. James F. Maxwell (Herbarium of Chiang Mai University). TP thanks the Science Achievement Scholarship of Thailand (SAST) and the Center of Excellence for Innovation in Chemistry (PERCH-CIC). We would like to thank the bioassay laboratory, BIOTEC, NSTDA, Pathumthani, Thailand, for the KB and Vero cells, and financial support from NSTDA's Cluster Program Management to S.K.

Disclosure statement

The authors declare no conflict of interest

References

- Abdel-Kader MS. 2001. Phenolic constituents of *Ononis vaginalis* roots. *Planta Med.* 67:388–390.
- Belofsky G, Carreno R, Lewis K, Ball A, Casadei G, Teges GP. 2006. Metabolites of the “Smoke Tree”, *Dalea spinosa*, potentiate antibiotic activity against multidrug-resistant *Staphylococcus aureus*. *J Nat Prod.* 69:261–264.

- Csékei M, Novák Z, Kotschy A. 2008. Development of a one-pot sequential Sonogashira coupling for the synthesis of benzofurans. *Tetrahedron*. 64:8992-8996.
- CLSI, 2012. Reference methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Approved Standard ninth ed.; Clinical and Laboratory Standards Institute: Wayne, PA, USA.
- Dagne E, Gunatilaka AAL, Kingston DGI. 1993. Two bioactive pterocarpanes from *Erythrina burana*. *J Nat Prod*. 56:1831–1834.
- Granados-Covarrubias EH, Maldonado LA. 2009. A Wacker–Cook synthesis of isoflavones: formononetin. *Tetrahedron Lett*. 50:1542–1545.
- Goto H, Terao Y, Akai S. 2009. Synthesis of various kinds of isoflavones, isoflavanes, and biphenyl-ketones and their 1,1-diphenyl-2-picrylhydrazyl radical-scavenging activities. *Chem Pharm Bull*. 57:346–360.
- Huang R, Ding ZG, Long YF, Zhao JY, Li MG, Cui XL, Wen ML. 2013. A new isoflavone derivative from *Streptomyces* sp. YIM GS3536. *Chem Nat Comp*. 48:966–969.
- Mahidol University. 1996. Sa Moon Prai Peun Ban Lanna (in Thai). Bangkok: Amarin Printing & Publishing Public Company. p 203.
- Mizuno M, Matsuura N, Iinuma M, Tanaka T, Phengklai C. 1990. Isoflavones from stems of *Euchresta horsfieldii*. *Phytochemistry*. 29:2675–2677.
- Puebla P, Oshima-Franco Y, Franco LM, Santos MGD, Silva RVD, Rubem-Mauro L, Feliciano AS. 2010. Chemical constituents of the bark of *Dipteryx alata* Vogel, an active species against *Bothrops jararacussu* Venom. 15:8193–8204.
- Smitinand T. 2014. Thai plant names. Bangkok: The Forest Herbarium Royal Forest Department. p 523.
- Sichaem J, Ruksilp T, Sawasdee P, Khumkratok S, Tip-pyang S. 2018. Chemical constituents of the stems of *Spatholobus parviflorus* and their cholinesterase inhibitory activity. *Chem Nat Comp*. 54:356–357
- Takeya K, Itokawa H. 1982. Isoflavonoids and the other constituents in callus tissues of *Pueraria lobata*. *Chem Pharm Bull*. 30:1496–1499.
- Tang RN, Qu XB, Guan SH, Xu PP, Shi YY, Guo DA. 2012. Chemical constituents of *Spatholobus suberectus*. *Chin J Nat Med*. 10:32–35.

- Vichai V, Kirtikara K. 2006. Sulforhodamine B colorimetric assay for cytotoxicity. screening. Nat Protoc. 1:1112–1116.
- Wang LX, Zheng HR, Ren FC, Chen TG, Li XM, Jiang XJ, Wang F. 2017. Polysubstituted isoflavonoids from *Spatholobus suberectus*, *Flemingia macrophylla*, and *Cudrania cochinchinensis*. Nat Prod Bioprospect. 7:201–206.
- Wu X, Li M, Wang X, Shen T, Wang S, Ren D. 2018. Two new 2-arylbenzofuran derivatives from the leaves of *Morus alba*. Nat Prod Res. <https://doi.org/10.1080/14786419.2018.1443095>.
- Yoon KH, Park KJ, Yin J, Yoon KH, Lee JY, Hwang YJ, Lee DI, Choi, YW, Lee MW. 2016. Antioxidative and antitumor effects of isoflavones isolated from the leaves of *Maackia fauriei*. Rec Nat Prod. 10: 441–451.
- Yoon JS, Sung SH, Park JH, Kim YC. 2004. Flavonoids from *Spatholobus suberectus*. Arch Pharm Res. 27:589–592.
- Zhang DW, Li GH, Yu QY, Dai SJ. 2010. New anti-inflammatory 4-hydroxyisoflavans from *Solanum lyratum*. Chem Pharm Bull. 58:840–842.