



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers

Faculty of Science, Medicine and Health

2013

Gas-phase transformation of phosphatidylcholine cations to structurally informative anions via ion/ion chemistry

John R. Stutzman
Purdue University

Stephen J. Blanksby
University of Wollongong, blanksby@uow.edu.au

Scott A. Mcluckey
Purdue University

Publication Details

Stutzman, J. R., Blanksby, S. J. & Mcluckey, S. A. (2013). Gas-phase transformation of phosphatidylcholine cations to structurally informative anions via ion/ion chemistry. *Analytical Chemistry*, 85 (7), 3752-3757.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Gas-phase transformation of phosphatidylcholine cations to structurally informative anions via ion/ion chemistry

Abstract

Gas-phase transformation of synthetic phosphatidylcholine (PC) monocations to structurally informative anions is demonstrated via ion/ion reactions with doubly deprotonated 1,4-phenylenedipropionic acid (PDPA). Two synthetic PC isomers, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (PC16:0/18:1) and 1-oleoyl-2-palmitoyl-sn-glycero-3-phosphocholine (PC18:1/16:0), were subjected to this ion/ion chemistry. The product of the ion/ion reaction is a negatively charged complex, $[PC + PDPA - H]^-$. Collisional activation of the long-lived complex causes transfer of a proton and methyl cation to PDPA, generating $[PC - CH_3]^-$. Subsequent collisional activation of the demethylated PC anions produces abundant fatty acid carboxylate anions and low-abundance acyl neutral losses as free acids and ketenes. Product ion spectra of $[PC - CH_3]^-$ suggest favorable cleavage at the sn-2 position over the sn-1 due to distinct differences in the relative abundances. In contrast, collisional activation of PC cations is absent of abundant fatty acid chain-related product ions and typically indicates only the lipid class via formation of the phosphocholine cation. A solution phase method to produce the gas-phase adducted PC anion is also demonstrated. Product ion spectra derived from the solution phase method are similar to the results generated via ion/ion chemistry. This work demonstrates a gas-phase means to increase structural characterization of phosphatidylcholines via ion/ion chemistry.

Keywords

chemistry, transformation, gas, phosphatidylcholine, phase, cations, structurally, informative, anions, via, ion

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Stutzman, J. R., Blanksby, S. J. & Mcluckey, S. A. (2013). Gas-phase transformation of phosphatidylcholine cations to structurally informative anions via ion/ion chemistry. *Analytical Chemistry*, 85 (7), 3752-3757.

Gas-Phase Transformation of Phosphatidylcholine Cations to Structurally Informative Anions via Ion/Ion Chemistry

John R. Stutzman[†], Stephen J. Blanksby[‡], and Scott A. McLuckey^{†*}

[†]Department of Chemistry
Purdue University
West Lafayette, IN, USA 47907-2084

[‡]ARC Centre of Excellence for Free Radical Chemistry and Biotechnology, School of Chemistry
University of Wollongong
Wollongong, NSW, Australia 2522

Running title: Transformation of Phosphatidylcholine Cations

*Address reprint requests to:

Dr. S. A. McLuckey
560 Oval Drive
Department of Chemistry
Purdue University
West Lafayette, IN 47907-2084, USA
Phone: (765) 494-5270
Fax: (765) 494-0239
E-mail: mcluckey@purdue.edu

ABSTRACT

Gas-phase transformation of synthetic phosphatidylcholine (PC) monocations to structurally informative anions is demonstrated via ion/ion reactions with doubly deprotonated 1,4-phenylenedipropionic acid (PDPA). Two synthetic PC isomers, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (PC_{16:0/18:1}) and 1-oleoyl-2-palmitoyl-*sn*-glycero-3-phosphocholine (PC_{18:1/16:0}) were subjected to this ion/ion chemistry. The product of the ion/ion reaction is a negatively charged complex, [PC+PDPA-H]⁻. Collisional activation of the long-lived complex causes concomitant transfer of a proton and methyl cation to PDPA, generating [PC-CH₃]⁻. Subsequent collisional activation of the demethylated PC anions produces abundant fatty acid carboxylate anions and low abundance acyl neutral losses as free acids and ketenes. Product ion spectra of [PC-CH₃]⁻ suggest favorable cleavage at the *sn*-2 position over the *sn*-1 due to distinct differences in the relative abundances. In contrast, collisional activation of PC cations is absent of abundant fatty acid chain-related product ions and typically indicates only the lipid class via formation of the phosphocholine cation. A solution phase method to produce the gas-phase adducted PC anion is also demonstrated. Product ion spectra derived from the solution phase method are similar to the results generated via ion/ion chemistry. This work demonstrates a gas-phase means to increase structural characterization of phosphatidylcholines via ion/ion chemistry.

Key words: Ion/ion, phosphatidylcholines, charge inversion, lipids, lipidomics

INTRODUCTION

Mass spectrometry (MS) has become an essential analytical tool for the identification and structural characterization of biological macromolecules.^{1,2,3} In tandem MS (MS/MS or MSⁿ), the polarity in which the analyte ion is most efficiently formed does not always provide sufficient structural information upon collision-induced dissociation (CID). For instance, phosphatidylcholines (PC), a subclass of glycerophospholipids, are efficiently ionized in the positive polarity, but their cationic versions generate limited structural information upon CID. PCs compose thirty percent of cell membranes in mammalian cells⁴ and can act as secondary messengers in metabolic signaling/pathways.⁵ PCs are composed of a glycerol backbone with two fatty acid chains attached at the *sn*-1 and *sn*-2 positions while a phosphocholine moiety is attached at the *sn*-3 position. Due to the presence of a fixed positive charge quaternary amine, PCs readily generate gas-phase cations in positive ion mode electrospray ionization (ESI).⁶ However, collisional activation of PC cations generally yields the phosphocholine moiety (184 *m/z*) as the dominant and often exclusive product. The phosphocholine fragment ion provides insight on only the phospholipid class and does not provide structural information about the individual acyl chains.^{7,8} Elucidating the composition of the individual fatty acid chains is important due to the diverse physical properties of PCs (*i.e.*, the number of carbons and extent of unsaturation) as these properties relate to their lipid biochemistry.^{9,10,11} With the presence of the choline moiety, PCs cannot be ionized in the negative polarity by deprotonation⁶ and therefore structural information of the individual fatty acid chains cannot be easily accessed without solution phase additives (see below).

Efforts to better characterize PCs have included CID studies of adducted PCs in both the positive¹² and negative ion modes.^{8,13,14,15} Gas-phase lithium adducts of PCs generated by positive ESI of solution phase PC mixtures have shown neutral losses of lithiated fatty acid chains (*i.e.*, RCO₂Li, where R is the alkyl chain) upon CID. The difference in *m/z* between the product and precursor ion corresponds to the degree of unsaturation and chain length. However, the structurally informative peaks were of low relative abundance and the spectra were complicated by additional neutral losses.¹² In the negative ion mode approach, millimolar concentrations of salts were added to a PC solution to produce gas-phase adducted PC anions (*e.g.*, [PC+Cl]⁻ and [PC+CH₃CO₂]⁻). Collisional activation of the adducted PC anions produced a neutral loss comprised of the counter anion and a methyl cation from the quaternary ammonium (*i.e.*, CH₃Cl and CH₃CO₂CH₃), yielding [PC-CH₃]⁻. Subsequent CID of [PC-CH₃]⁻ generated fatty acid carboxylate anions (RCO₂⁻) and neutral losses associated with free acids (RCO₂H) or ketene derivatives (RCHCO), which are all indicative of the composition of the individual fatty acid chains.^{8,13-15} In both of the solution phase methods, the high salt concentrations can lead to poor ionization efficiency, spectral complexity upon ionization, and incompatibilities with liquid chromatography (LC) solvent systems.


The type of ion subjected to tandem MS is usually determined by the ionization method and conditions. An alternative approach is to transform in the gas-phase an ion-type that is readily generated by an ionization method to another ion-type that more readily yields structural information of interest, preferably mass-selection. Gas-phase ion/ion reactions have been shown to be particularly useful for many ion transformations,^{16,17} particularly when conducted within the context of MSⁿ, whereby the analyte and reagent ions are each mass-selected prior to reaction. The use of ion/ion reactions within the context of an MSⁿ experiment allows the

definition of the ion-type to be de-coupled from the initial ionization conditions.¹⁸ As a result, the ion-type subjected to activation can be independently optimized to generate the most informative fragmentation. Ion/ion reactions have been utilized in acid/base reactions (*i.e.*, proton transfer),^{19, 20} reduction/oxidation reactions (*i.e.*, electron transfer),^{21, 22} and covalent^{23,24,25,26} and/or electrostatic modification.^{27,28} Previous studies have demonstrated the significance of ion-type transformation and its relation to increased sequence coverage.^{29,30,31} For instance, MALDI-derived tryptic peptide monocations covalently and electrostatically modified with benzene disulfonic acid dianions via ion/ion reactions demonstrated increased sequence coverage upon collisional activation, especially in the form of b-type ions.³²

In the case of PC ions, an example of gas-phase ion-type transformation via ion/ion chemistry has been demonstrated in the electron transfer dissociation (ETD) of doubly sodiated PC cations, $[\text{PC}+2\text{Na}]^{2+}$ using azobenzene radical anions as the ETD reagent. ETD induced cleavages at each ester linkage (*i.e.*, neutral loss of a fatty acid chain) with additional neutral loss of the quaternary nitrogen moiety. CID of the charged reduced electron transfer product that did not undergo dissociation (*i.e.*, the ETnoD ion), $[\text{M}+2\text{Na}]^{+}$, generated results similar to the electron transfer reaction. Similar to the lithiated PC cation study, this method provided information on the number of carbons and the degree of unsaturation for each fatty acid chain via neutral losses.³³

In this study, we demonstrate the gas phase transformation of PC monocations into demethylated PC anions (*i.e.*, $[\text{PC}-\text{CH}_3]^-$) via ion/ion reactions, where pre- and post-transformation structural characterization is compared. PC monocations are subjected to ion/ion reactions with doubly deprotonated 1,4-phenylenedipropionic acid (PDPA). The product of the

ion/ion reaction is a negatively charged complex, comprised of PDPA and PC, *i.e.* [PC+PDPA-H]⁻. Collisional activation of the complex generates [PC-CH₃]⁻ as seen in process (1):



Gas-phase ion/ion reactions have been performed on synthetic PC isomers, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (16:0/18:1 PC) and 1-oleoyl-2-palmitoyl-*sn*-glycero-3-phosphocholine (18:1/16:0 PC). Collisional activation of [PC-CH₃]⁻ generates large relative contributions from fatty acid carboxylate anions (RCO₂⁻) and less abundant neutral losses of fatty acids (RCO₂H) and ketene derivatives (RCHCO), whereas collisional activation of [PC+H]⁺ produces exclusively the phosphocholine cation. Transformation of PC monocations via ion/ion reactions facilitates the production of structurally informative anions and the identification of fatty acid composition without recourse for solution phase salt additives.

EXPERIMENTAL SECTION

Materials. Methanol, chloroform, and ammonium hydroxide were purchased from Mallinckrodt (Phillipsburg, NJ). Acetonitrile and 1,4-phenylenedipropionic acid were purchased from Sigma-Aldrich (St. Louis, MO). 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine and 1-oleoyl-2-palmitoyl-*sn*-glycero-3-phosphocholine were purchased from Avanti Polar Lipids, Inc (Alabaster, AL). PC analytes were prepared at a concentration of ~30 μM in a solution of 50/50 (v/v) chloroform/methanol prior to positive nanoelectrospray ionization (nESI). The reagent anion, PDPA, was prepared at a concentration of ~ 220 μM in a solution of 49.5/49.5/1 (v/v/v) acetonitrile/methanol/ammonium hydroxide prior to negative nESI.


Solution Phase PDPA Spiking. A solution mixture of PC (*i.e.*, PC_{16:0/18:1} or PC_{18:1/16:0}) and PDPA was prepared at concentrations of 10 μM and 140 μM , respectively. The solution was subjected to negative mode nESI, where MSⁿ analyses and subsequent mass analysis were performed in Q3.

Mass Spectrometry. All experiments were performed on a QTRAP 4000 triple quadrupole/linear ion trap mass spectrometer (AB SCIEX, Concord, ON, Canada), which has been modified to perform ion/ion reactions.³⁴ Alternately pulsed nESI emitters sequentially inject mass-selected anions and cations into the q2 reaction cell.^{35,36} Doubly deprotonated PDPA anions were first ionized and accumulated in the q2 reaction cell. Next, the singly charged PC cations were ionized and transferred to the q2 cell to undergo a mutual storage reaction for 1000 milliseconds. Following accumulation of the doubly deprotonated PDPA anions and throughout the ion/ion reaction period in the transformation efficiency study, any singly charged PDPA anions that were formed from ion/molecule reactions due to adventitious basic vapors or ion/ion reactions were subjected to continuous resonance ejection to minimize the possibility of singly charged anions reacting with the singly charged cations. Product anions resulting from the ion/ion reaction were transferred to Q3 to perform MSⁿ analyses and subsequent mass analysis via mass-selective axial ejection (MSAE).³⁷

RESULTS AND DISCUSSION

Transformation of PC cations. Two PC isomers, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (16:0/18:1 PC) and 1-oleoyl-2-palmitoyl-*sn*-glycero-3-phosphocholine (18:1/16:0 PC), have been subjected to ion/ion reactions with doubly deprotonated PDPA. The ion/ion reaction produces a negatively charged complex (*i.e.*, [PC+PDPA-H]⁻), which is

stabilized by a strong electrostatic interaction between the fixed charge quaternary ammonium group of PC and a carboxylate of PDPA (Scheme 1).²⁴ Figure 1(a) illustrates the product ion spectrum following an ion/ion reaction between $[\text{PDPA-2H}]^{2-}$ anions and $[\text{PC}_{16:0/18:1}+\text{H}]^+$ cations. The post ion/ion reaction spectrum generates $[\text{PC}_{16:0/18:1}+\text{PDPA-H}]^-$ and a charge reduced reagent anion (*i.e.*, $[\text{PDPA-H}]^-$). Proton transfer to the reagent dianion can occur via two pathways: a fly-by mechanism or charge transfer upon separation of the long-lived complex.³⁸



Scheme 1: The sequence of generating $[\text{PC-CH}_3]^-$ via ion/ion reactions between PDPA dianions and PC monocations. Top, mass-selected precursor ions; middle, electrostatic complex generated via a mutual storage ion/ion reaction; bottom, the main products following collisional activation of the negative complex.

Subsequent isolation and collisional activation of the long-lived complex, $[\text{PC}_{16:0/18:1}+\text{PDPA-H}]^-$, produces a highly abundant $[\text{PC}_{16:0/18:1}-\text{CH}_3]^-$ ion along with consecutive fragments consistent with charged and neutral losses of the fatty acid chains (Figure 1(b)). Observation of the demethylated PC anions indicates concomitant methyl cation and proton transfers to the anionic reagent, thus producing a neutralized PDPA reagent (see Scheme 1). The observation of $[\text{PC}-\text{CH}_3]^-$ mirrors the results of solution phase acetate and chloride salt experiments, where collisional activation of $[\text{PC}+\text{CH}_3\text{CO}_2\text{H}]^-$ or $[\text{PC}+\text{Cl}]^-$ generated a demethylated PC anion.^{8,13-15} Gas-phase methyl cation transfer between carboxylates and quaternary ammonium groups has been observed previously within the context of an ion/ion reaction.²⁶ Alkyl (*i.e.*, ethyl, propyl, butyl, isobutyl) cation transfer has also been observed upon CID of ESI-generated complexes composed of sulfophenyl benzoic acid dianions and tetraalkylammonium monocations.³⁹ Collisional activation of the gas-phase anionic complex produced either proton or alkyl cation transfer to the sulfophenyl benzoic acid dianions, as similarly seen in this work with collisional activation of PDPA-adducted PCs. The relative abundance of the proton transfer product (*i.e.*, $[\text{PDPA-H}]^-$) in the dissociation spectrum is particularly noteworthy (Figure 1(b)). This low contribution of $[\text{PDPA-H}]^-$ compared to $[\text{PC}_{16:0/18:1}-\text{CH}_3]^-$ suggests that concomitant transfer of a proton and methyl cation upon collisional activation is a dominant pathway, while sole proton transfer without concomitant methyl cation transfer is a less favorable pathway. The other PC isomer, 18:1/16:0 PC, yields similar results to the 16:0/18:1 compound for both the ion/ion reaction and subsequent collisional activation of the long-lived complex (Supporting Information, S-1).




Figure 1: (a) Product ion spectrum following ion/ion reactions between PDPA dianions and 16:0/18:1 PC monocations, (b) Ion trap CID of the long-lived complex, $[\text{PC}_{16:0/18:1}+\text{PDPA-H}]^-$

The analytical utility of the gas-phase transformation of PC cations to demethylated PC anions is dependent upon the value of the information gained from changing one ion-type to another, as discussed below, as well as the reaction efficiency. A lower limit to the efficiency of converting $\text{PC}_{16:0/18:1}$ cations to the anion complex $[\text{PC}_{16:0/18:1}-\text{CH}_3]^-$ can be derived from the spectrum of Figure 1(a) from the ratio of $[\text{PDPA-H}]^-/([\text{PDPA-H}]^-+[\text{PC}_{16:0/18:1}-\text{CH}_3]^-)$. This yields in an efficiency of roughly 25%. However, a significant fraction of the observed $[\text{PDPA-H}]^-$ signal in Figure 1(a) arises from ion/molecule proton transfer from polar vapors present in the vacuum system. Furthermore, the accumulation of $[\text{PDPA-H}]^-$ due to ion/ion and ion/molecule reactions leads to an increase in the neutralization of the PC cations from ion/ion reactions with

the [PDPA-H]⁻ anions. We therefore chose to measure an “operational” efficiency of transforming PC_{16:0/18:1} cations to demethylated PC anions, which is defined as:^{40,41}

$$= \frac{\sum([PC - CH_3 - H]) + \sum(ConFragments)}{\sum PC_i - \sum PC_f} \times 100$$

where $\Sigma[PC-CH_3-H]$ is the abundance of demethylated PC anions produced upon CID of the long-lived complex ([PC+PDPA-H]⁻), $\Sigma(ConFragments)$ is the abundance of consecutive fragments from [PC-CH₃]⁻ following CID of the long-lived complex, ΣPC_i is the abundance of [PC+H]⁺ after isolation and before ion/ion transformation, and ΣPC_f is the abundance of residual [PC+H]⁺ after transformation. A resonance ejection voltage was applied throughout the reaction period to eject [PDPA-H]⁻ ions generated by ion/ion and ion/molecule reactions to minimize any neutralization of the PC cations associated with ion/ion reactions with [PDPA-H]⁻. By subtracting PC_i from PC_f , the transformation efficiency is limited to the fraction of [PC+H]⁺ that undergoes an ion/ion reaction with [PDPA-2H]²⁻. The transformation efficiency determined in this way represents the maximum possible efficiency for conversion of [PC+H]⁺ to [PC-CH₃]⁻ because it represents the value that would be obtained if all reactant cations undergo an ion/ion reaction. Complete depletion of analyte ions is achievable provided there is an excess of reagent ions, there is good overlap in the ion populations, and the reaction time is sufficient to allow for the reaction to go to completion. It is also emphasized here that the measurement yields an “operational” efficiency rather than an absolute efficiency due to possible differences in detection efficiencies for positive ions and negative ions. The absolute values for voltages applied to the conversion dynode and multiplier were the same in both detection polarities with the only difference being the polarity of the conversion dynode. In this study, the transformation

efficiency of PC cations to demethylated PC anions was measured to be $49 \pm 7\%$. The major loss process is expected to be neutralization of the cation by $[\text{PDPA-2H}]^{2-}$ as there are no major competing charge inversion channels and little resonance ejection of the $[\text{PC+PDPA-H}]^-$ complex is expected under the collisional activation conditions used to dissociate it. Overall, the results demonstrate that the ion/ion charge inversion/CID process used to generate the $[\text{PC-CH}_3]^-$ ions is clearly efficient enough for analytical use.

MS/MS of $[\text{PC-CH}_3]^-$ Anions. Ion trap CID of 16:0/18:1 PC prior to and after ion/ion transformation (*i.e.*, $[\text{PC}_{16:0/18:1}+\text{H}]^+$ and $[\text{PC}_{16:0/18:1}-\text{CH}_3]^-$) are compared in Figure 2. Collisional activation of $[\text{PC}_{16:0/18:1}-\text{CH}_3]^-$ produces a structurally informative product ion spectrum, where loss of neutral and charged fatty acid chains are observed (Figure 2(a)). These fragment ions allow for the diagnosis of chain length and the degree of unsaturation. The most favorable dissociation pathway appears to be the generation of a charged fatty acid chain from the *sn*-1 or *sn*-2 position, *i.e.*, $[\text{C}_{16}\text{H}_{31}\text{CH}_2\text{COO-H}]^-$ and $[\text{C}_{14}\text{H}_{29}\text{CH}_2\text{COO-H}]^-$. Neutral losses of fatty acids as ketenes (*e.g.*, $[\text{PC}_{16:0/18:1}-\text{CH}_3-\text{C}_{14}\text{H}_{29}\text{CH}=\text{C}=\text{O-H}]^-$) and free fatty acids (*e.g.*, $[\text{PC}_{16:0/18:1}-\text{CH}_3-\text{C}_{14}\text{H}_{29}\text{CH}_2\text{COOH-H}]^-$) are also observed in low abundance. Similar dissociation behavior of demethylated PC anions^{8,13-15} and other glycerophospholipids^{42,43,44,45} has been noted previously in the literature. In contrast, CID of the $[\text{PC}_{16:0/18:1}+\text{H}]^+$ ion yields the phosphocoline cation as the strongly dominant product (Figure 2(b)). Figure 2(c) serves to depict the relationship between bond cleavage and product ion-type (*e.g.*, cleavage of the ester bond leads to charged or neutral loss of a single fatty acid chain). In addition, a comparison of *sn*-cleavage position reveals a larger contribution from ions produced from the cleavage at the *sn*-2 position (*i.e.*, $[\text{C}_{16}\text{H}_{31}\text{CH}_2\text{COO-H}]^-$ and $[\text{PC}_{16:0/18:1}-\text{CH}_3-\text{C}_{16}\text{H}_{31}\text{CH}=\text{C}=\text{O-H}]^-$) compared to those produced from cleavage at the *sn*-1 position (*i.e.*, $[\text{C}_{14}\text{H}_{29}\text{CH}_2\text{COO-H}]^-$ and $[\text{PC}_{16:0/18:1}-\text{CH}_3-$

$C_{14}H_{29}CH=C=O-H]^+$). Ekroos *et al.* noted the relative favorability of *sn*-2 cleavage in synthetic PC isomers and biological samples;¹³ however, structural characterization via relative abundance comparisons can be misleading due to the likely isomeric mixtures present in synthetic and natural samples. Additionally, relative abundance ratios of the *sn*-1 and *sn*-2 position were previously demonstrated to be dependent on fatty acid chain length and degree of saturation, thus suggesting caution on assigning exact structures.⁴⁶ While the phosphocholine moiety in the product ion spectrum of Figure 2(b) allows for lipid class profiling, CID of the transformed PC anion produces more structurally informative product ions.




Figure 2: (a) Ion trap CID of $[PC_{16:0/18:1}-CH_3]^-$. (b) Ion trap CID of $[PC_{16:0/18:1}+H]^+$. (c) Relating bond cleavage and product ion type for $[PC_{16:0/18:1}-CH_3]^-$.

Figure 3(a) depicts the product ion spectrum following ion trap CID of $[PC_{18:1/16:0}-CH_3]^-$. Collisional activation of $[PC_{18:1/16:0}-CH_3]^-$ produces similar fragment ions to the 16:0/18:1 PC isomer, (*i.e.*, neutral losses of fatty acid ketenes, free fatty acids, and charged fatty acids), thus providing a diagnostic fragmentation spectrum. As seen with the 16:0/18:1 PC (Figure 2(a)), the

dissociation behavior of 18:1/16:0 is dominated by the production of fatty acid anions (Figure 3(a)). However, the most significant difference between the two isomers (*i.e.*, between 16:0/18:1 PC and 18:1/16:0 PC) is the reversal of the relative abundances of the $[C_{14}H_{29}CH_2COO-H]^-/[C_{16}H_{31}CH_2COO-H]^-$ and $[PC-CH_3-C_{14}H_{29}CH=C=O-H]^-/[PC-CH_3-C_{16}H_{31}CH=C=O-H]^-$ ions. This reversal in intensity may be attributed to the position of the fatty acid chain (*i.e.*, *sn*-1 or *sn*-2), where cleavage of *sn*-2 appears to be more favorable. Ekroos *et al.* reported a similar change in relative abundances between isomeric demethylated PC anions.¹³ Collisional activation of $[PC_{18:1/16:0}+H]^+$ is depicted in Figure 3(b). Ion trap CID of the PC cation produces a dominant phosphocholine moiety, as also seen with 16:0/18:1 PC (Figure 2(b)).




Figure 3: (a) Ion trap CID of $[PC_{18:1/16:0}-CH_3]^-$. (b) Ion trap CID of $[PC_{18:1/16:0}+H]^+$.

Solution Phase PDPA Mixture. The preceding results demonstrate the formation of a $[PC+PDPA-H]^-$ complex, which dissociates to the $[PC-CH_3]^-$ ion that can yield structurally

informative products upon CID in the gas-phase. We also sought to generate such a complex directly from solution for two reasons. First, generation of the complex directly from solution would enable access to the precursor ion of interest without recourse to gas-phase ion/ion reactions, in analogy with the work of Ekroos *et al.*¹³ Of course, such an approach might also require a separate positive ion experiment to confirm the phospholipid class information. Second, we were interested to see if a complex formed in solution would differ in its CID behavior from that of the complex formed via ion/ion reaction. Direct ionization of a solution phase mixture containing PDPA and 16:0/18:1 PC via negative mode nESI is illustrated in Figure 4(a). Ionization of the mixture produces low abundance PDPA-adducted PC anions (*i.e.*, $[\text{PC}_{16:0/18:1}+\text{PDPA-H}]^-$) among dimers of PDPA and substantial chemical noise. Collisional activation of the complex produces the demethylated PC anion (data not shown). Ion trap CID of the demethylated PC anion (Figure 4(b)) generates similar dissociation behavior compared to the analogous $[\text{PC}_{16:0/18:1}-\text{CH}_3]^-$ produced via ion/ion reactions (Figure 3(a)). The solution phase method clearly exhibits spectral complexity as well as limited ion signal from the adducted PC anion upon ionization, which could lead to complications in complex mixture analysis. The main advantages of the ion/ion chemistry are the high degree of control over the identities of the reactants via mass-selection prior to reaction, the avoidance of spectral complexity that can arise in solution phase reaction mixtures, the avoidance of ion suppression in the ionization step arising from the high salt concentration in the sample, and ready access to the original $[\text{PC}+\text{H}]^+$ ions and the product $[\text{PC}-\text{CH}_3]^-$ ion from a single solution and set of ionization conditions. Solution phase experiments with 18:0/16:0 PC (S-2) produced results similar to those observed with 16:0/18:1 PC. The dissociation behavior of $[\text{PC}_{18:0/16:0}-\text{CH}_3]^-$ derived from the solution phase is also similar to that observed with the ion/ion chemistry.




Figure 4: (a) Ionization of PDPA and 16:0/18:1 solution phase mixture via negative mode nESI
 (b) Ion trap CID of [PC_{16:0/18:1}-CH₃]⁻

CONCLUSIONS

Synthetic phosphatidylcholine monocations have been subjected to ion/ion reactions with doubly deprotonated PDPA. Collisional activation of the long-lived complex, comprised of PDPA and PC, generates a demethylated PC anion (*i.e.*, [PC-CH₃]⁻), indicating concomitant transfer of a proton from the phosphate group and methyl cation from the quaternary amine group to the carboxylate groups of PDPA. Ion trap CID of [PC-CH₃]⁻ generates product ions consistent with charged fatty acid chains and neutral losses of the acyl chains from the PC analyte as free acids and ketenes. Fragmentation spectra generate structurally informative product ions, which allows for the elucidation of fatty acid composition (*i.e.*, the number of

carbons and degree of unsaturation in each acyl chain). The dissociation behavior of demethylated PC anions also exhibits preferential cleavages at the *sn*-2 over the *sn*-1 position as reflected in the relative abundances. The dissociation behaviors in this study reflect previous solution phase experiments reported in the literature. Solution phase studies demonstrated here, where a mixture of PC and PDPA are directly ionized via negative mode nESI, show that the complex that leads to demethylated PC anions can also be formed directly in solution, although a relatively low abundance. The ion/ion reaction approach is attractive in that a single set of ionization conditions in one polarity, which can be optimized for $[\text{PC}+\text{H}]^+$ formation, can be used to generate two ion-types, the $[\text{PC}+\text{H}]^+$ and $[\text{PC}-\text{CH}_3]^-$ ions, that provide complementary information upon CID.

ACKNOWLEDGEMENTS

This research was supported by AB Sciex and the National Institutes of Health under Grant GM 45372. S.J.B. Also acknowledged support from the Australian Research Council through the Centres of Excellence (CE0561607) and Discovery (DP120102922) Programs.

References

- 1 Hunt, D. F.; Yates, J. R.; Shabanowitz, J.; Winston, S.; Hauer, C. R. *Proc. Natl. Acad. Sci. U.S.A.* **1986**, *83*, 6233–6237.
- 2 Biemann, K.; Martin, S. A. *Mass Spectrom. Rev.* **1987**, *6*, 1–75.
- 3 Aebersold, R.; Mann, M. *Nature.* **2003**, *422*, 198–207.
- 4 Kawai, K.; Fujita, M.; Nakao, M. *Biochim. Acta.* **1974**, *369*, 222-233.
- 5 Zeisel, S. H.; Blusztajn, J. K. *Annu. Rev. Nutr.* **1994**, *14*, 269–296.
- 6 Han, X.; Gross, R. *Proc. Natl. Acad. Sci. USA.* **1994**, *91*, 10635-10639.
- 7 Brugger, B.; Erben, G.; Sandhoff, R.; Wieland, F. T.; Lehmann, W. D. *Proc. Natl. Acad. Sci. USA.* **1997**, *94*, 2339-2344.
- 8 Han, X.; Gross, R. *J. Am. Soc. Mass Spectrom.* **1995**, *6*, 1202-1210.
- 9 Farooqui, A. A.; Horrocks, L. A. *Cell. Mol. Neurobiol.* **1998**, *18*, 599–608.
- 10 Mitchell, T. W.; Pham, H.; Thomas, M. C.; Blanksby, S. J. *J. Chromatog. B.* **2009**, *877*, 2722–2735.
- 11 Brown, S. H. J.; Mitchell, T. W.; Oakley, A. J.; Pham, H. T.; Blanksby, S. J. *J. Am. Soc. Mass Spectrom.* **2012**, *23*, 1441-1449.
- 12 Hsu, F. F.; Bohrer, A.; Turk, J. *J. Am. Soc. Mass Spectrom.* **1998**, *9*, 516–526.
- 13 Ekroos, K.; Ejsing, C.; Bahr, U.; Karas, M.; Simons, K.; Shevchenko, A. *J. Lipid Res.* **2003**, *2181-2192*.
- 14 Harrison, K.A.; Murphy, R.C. *J. Mass Spectrom.* **1995**, *30*, 1772–1773.
- 15 Kerwin, J. L.; Tuininga, A. R.; Ericsson, L. H. *J. Lipid Res.* **1994**, *35*, 1102–1114.
- 16 McLuckey, S.A.; Huang, T.-Y. *Anal. Chem.* **2009**, *81*, 8669-8676.
- 17 McLuckey, S.A.; Mentinova, M. *J. Am. Soc. Mass Spectrom.* **2011**, *22*, 3-12.
- 18 McLuckey, S. A. *Eur. J. Mass Spectrom.* **2010**, *16*, 429-436.
- 19 Stephenson, Jr, J. L.; McLuckey, S. A. *J. Am. Chem. Soc.* **1996**, *118*, 7390-7397.
- 20 Stephenson, Jr, J. L.; McLuckey, S. A. *Anal. Chem.* **1996**, *68*, 4026-4032.

-
- 21 Syka, J. E. P.; Coon, J. J.; Schroeder, M. J.; Shabanowitz, J.; Hunt, D. F. *Proc. Natl. Acad. Sci. USA*. **2004**, *101*, 9528-9533.
- 22 Coon, J. J.; Shabanowitz, J.; Hunt, D. F.; Syka, J. E. P. *J. Am. Soc. Mass Spectrom.* **2005**, *16*, 880-882.
- 23 Han, H.; McLuckey, S. A. *J. Am. Chem. Soc.* **2009**, *131*, 12884-12885.
- 24 Mentinova, M.; McLuckey, S. A. *J. Am. Chem. Soc.* **2010**, *132*, 18248–18257.
- 25 McGee, W. M.; Mentinova, M.; McLuckey, S. A. *J. Am. Chem. Soc.* **2012**, *134*, 11412-11414.
- 26 Prentice, B. M.; Gilbert, J. D.; Stutzman, J. R.; Forrest, W. P.; McLuckey, S. A. *J. Am. Soc. Mass Spectrom.* DOI: 10.1007/s13361-012-0506-8.
- 27 He, M.; Emory, J. F.; McLuckey, S. A. *Anal. Chem.* **2005**, *77*, 3173-3182.
- 28 Stutzman, J. R.; Luongo, C. A., McLuckey, S. A. *J. Mass Spectrom.* **2012**, *47*, 669-675.
- 29 Hassell, K. M.; Stutzman, J. R.; McLuckey S. A. *Anal. Chem.* **2010**, *82*, 1594-1597.
- 30 Mentinova, M. McLuckey, S.A.; *Int. J. Mass Spectrom.* **2011**, *308*, 133-136.
- 31 Stutzman, J. R.; Hassel, K. M.; McLuckey, S. A. *Int. J. Mass Spectrom.* **2012**, *312*, 195-200.
- 32 Stutzman, J. R.; McLuckey, S.A. *Anal. Chem.* **2012**, *84*, 10679-10685.
- 33 Liang, X.; Liu, J.; LeBlanc, Y.; Covey, T.; Ptak, A. C.; Brenna, J. T.; McLuckey, S. A. *J. Am. Soc. Mass Spectrom.* **2007**, *18*, 1783-1788.
- 34 Xia, Y.; Wu, J.; Londry, F. A.; Hager, J. W.; McLuckey, S.A. *J. Am. Soc. Mass Spectrom.* **2005**, *16*, 71–81.
- 35 Liang X.; Xia, Y.; McLuckey, S. A. *Anal. Chem.* **2006**, *78*, 3208–3212.
- 36 Liang X.; Han, H.; Xia, Y.; McLuckey, S. A. *J. Am. Soc. Mass Spectrom.* **2007**, *18*, 369–376.
- 37 Londry, F.A.; Hager, J. W. *J. Am. Soc. Mass Spectrom.* **2003**, *14*, 1130–1147.
- 38 Wells, J. M.; Chrisman, P. A.; McLuckey, S. A. *J. Am. Chem. Soc.* **2003**, *125*, 7238–7249.
- 39 Gronert, S. *Acc. Chem. Res.* **2003**, *36*, 848-857.
- 40 Hassell, K. M.; LeBlanc, Y.; McLuckey, S.A. *Anal. Chem.* **2011**, *83*, 3252-3255.
- 41 Hassell, K. M.; Hilger, R.T., McLuckey, S.A. *Phys. Chem. Chem. Phys.* **2011**, *13*, 18418-18427.
- 42 Hsu, F. F.; J. Turk. *J. Am. Soc. Mass Spectrom.* **2000**, *11*, 797–803.
- 43 Hsu, F. F.; J. Turk. *J. Am. Soc. Mass Spectrom.* **2000**, *11*, 892-899.

-
- 44 Hsu, F. F.; J. Turk. *J. Am. Soc. Mass Spectrom.* **2000**, *11*, 986-999.
- 45 Hsu, F. F.; J. Turk. *J. Am. Soc. Mass Spectrom.* **2000**, *11*, 1036-1043.
- 46 Huang, Z.; Gage, D. A.; Sweeley, C. C. *J. Am. Soc. Mass Spectrom.* **1992**, *3*, 71-78.