

University of Wollongong

## Research Online

---

Faculty of Health and Behavioural Sciences -  
Papers (Archive)

Faculty of Science, Medicine and Health

---

2010

### Genetic adult lactase persistence is associated with risk of Crohn's Disease in a New Zealand population

Deborah Nolan

*University of Wollongong, djn297@uow.edu.au*

Dug Yeo Han

*Uni of Auckland, NZ*

Wen Jiun Lam

*Uni of Auckland, NZ*

Angharad R. Morgan

*Uni of Auckland, NZ*

Alan G. Fraser

*Uni of Auckland, NZ*

*See next page for additional authors*

Follow this and additional works at: <https://ro.uow.edu.au/hbspapers>



Part of the [Arts and Humanities Commons](#), [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

---

#### Recommended Citation

Nolan, Deborah; Han, Dug Yeo; Lam, Wen Jiun; Morgan, Angharad R.; Fraser, Alan G.; Tapsell, Linda C.; and Ferguson, Lynnette R.: Genetic adult lactase persistence is associated with risk of Crohn's Disease in a New Zealand population 2010, 1-18.

<https://ro.uow.edu.au/hbspapers/616>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: [research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

# Genetic adult lactase persistence is associated with risk of Crohn's Disease in a New Zealand population

## Abstract

Background Mycobacterium avium subspecies paratuberculosis (MAP) is an infective agent found in ruminants and milk products, which has been suggested to increase the risk of gastrointestinal inflammation in genetically susceptible hosts. It is hypothesized that lactase persistence facilitates exposure to such milk products increasing the likelihood of adverse outcomes. Individuals either homozygous or heterozygous for the T allele of DNA variant, rs4988235, located 14kb upstream from the LCT locus, are associated with having lactase persistence. The aim of this study was to determine whether lactase persistence as evident by the T allele of rs4988235 is associated with Crohn's Disease (CD) in a New Zealand population. Findings Individuals homozygous for the T allele (T/T genotype) showed a significantly increased risk of having CD as compared with those homozygous for the C allele (OR=1.61, 95% CI=1.03-2.51). Additionally, a significant increase in the frequency of the T allele was observed in CD patients (OR=1.30, 95% CI=1.05-1.61, p=0.013), indicating that the T allele encoding lactase persistence was associated with an increased risk of CD. Conclusions Our findings indicate that lactase persistence as evident by the presence of the T allele of rs4988235 is associated with risk of CD in this New Zealand Caucasian population.

## Keywords

persistence, crohn, disease, lactase, zealand, population, adult, associated, genetic, risk

## Disciplines

Arts and Humanities | Life Sciences | Medicine and Health Sciences | Social and Behavioral Sciences

## Publication Details

Nolan, D., Han, D., Lam, W., Morgan, A. R., Fraser, A. G., Tapsell, L. C. & Ferguson, L. R. (2010). Genetic adult lactase persistence is associated with risk of Crohn's Disease in a New Zealand population. *B M C Research Notes*, 3 (339), 1-18.

## Authors

Deborah Nolan, Dug Yeo Han, Wen Jiun Lam, Angharad R. Morgan, Alan G. Fraser, Linda C. Tapsell, and Lynnette R. Ferguson

# **Genetic adult lactase persistence is associated with risk of Crohn's Disease in a New Zealand population.**

Deborah J Nolan<sup>1§</sup> Dug Yeo Han<sup>2,3</sup>, Wen Jiun Lam<sup>2,3</sup>, Angharad R Morgan<sup>2,3</sup>, Alan G Fraser<sup>2</sup> Linda C Tapsell<sup>1,4</sup>, Lynnette R Ferguson<sup>2</sup>

<sup>1</sup>Smart Foods Centre, University of Wollongong, Northfields Ave, Wollongong, New South Wales, Australia

<sup>2</sup>Nutrigenomics New Zealand, Discipline of Nutrition, University of Auckland, Park Rd, Auckland, New Zealand

<sup>3</sup>Discipline of Nutrition, University of Auckland, Park Rd, Auckland, New Zealand

<sup>4</sup>Illawarra Health and Medical Research Institute, University of Wollongong, Northfields Ave, Wollongong, New South Wales, Australia

<sup>§</sup>Corresponding author

Deborah J Nolan, Smart Foods Centre, Building 41.257, University of Wollongong, Northfields Ave, Wollongong, New South Wales, Australia, 2522

Telephone: +61 2 4221 5122

Fax: +61 2 4221 4844

Email addresses:

Deborah Nolan: [djn297@uow.edu.au](mailto:djn297@uow.edu.au)

Dug Yeo Han: [dy.han@auckland.ac.nz](mailto:dy.han@auckland.ac.nz)

Wen Jiun Lam: [w.lam@auckland.ac.nz](mailto:w.lam@auckland.ac.nz)

Angharad Morgan: [ang.morgan@auckland.ac.nz](mailto:ang.morgan@auckland.ac.nz)

Alan Fraser: [a.fraser@auckland.ac.nz](mailto:a.fraser@auckland.ac.nz)

Linda Tapsell: [ltapsell@uow.edu.au](mailto:ltapsell@uow.edu.au)

Lynnette Ferguson: [l.ferguson@auckland.ac.nz](mailto:l.ferguson@auckland.ac.nz)

## **Abstract**

### **Background**

*Mycobacterium avium* subspecies *paratuberculosis* (MAP) is an infective agent found in ruminants and milk products, which has been suggested to increase the risk of gastrointestinal inflammation in genetically susceptible hosts. It is hypothesized that lactase persistence facilitates exposure to such milk products increasing the likelihood of adverse outcomes. Individuals either homozygous or heterozygous for the T allele of DNA variant, rs4988235, located 14kb upstream from the *LCT* locus, are associated with having lactase persistence.

The aim of this study was to determine whether lactase persistence as evident by the T allele of rs4988235 is associated with Crohn's Disease (CD) in a New Zealand population.

### **Findings**

Individuals homozygous for the T allele (T/T genotype) showed a significantly increased risk of having CD as compared with those homozygous for the C allele (OR=1.61, 95% CI=1.03-2.51). Additionally, a significant increase in the frequency of the T allele was observed in CD patients (OR=1.30, 95% CI=1.05-1.61, p=0.013), indicating that the T allele encoding lactase persistence was associated with an increased risk of CD.

### **Conclusions**

Our findings indicate that lactase persistence as evident by the presence of the T allele of rs4988235 is associated with risk of CD in this New Zealand Caucasian population.

# Findings

## Background

Crohn's Disease (CD) is a form of inflammatory bowel disease of both genetic and environmental aetiology. Somewhat controversially *Mycobacterium avium* subsp. *Paratuberculosis* (MAP) exposure has resurfaced within the literature as a potential environmental variant that may be implicated in the development of CD [1]. MAP is an infective agent frequently found in ruminants such as cows and sheep that results in Johne's disease, an inflammatory bowel condition similar to that experienced by humans with CD [2]. MAP is secreted into the milk of infected ruminants [3] and is delivered to humans via the consumption of milk products [4] due to its reported resistance to commercial pasteurisation methods [5-6].

One of the most important factors that may influence the amount of milk consumed, and hence exposure to MAP in humans relates to the presence of Lactase-Phlorizin Hydrolase (lactase) in the small intestinal mucosa. Lactase is the enzyme required to hydrolyse lactose, the sugar found in milk products, into its constituent galactose and glucose [7]. Infants are born with the ability to digest lactose, but for the majority of the world's population this declines with age, a phenomenon known as lactase non persistence. However, lactase persistence is highly prevalent in descendants from populations with a rich pastoralist heritage [8]. Greater than 70% of individuals of white European heritage are reportedly lactose persistent compared to less than 40% throughout individuals from Asian countries [9].

Ennatah et al [10] identified a DNA variant, rs4988235 (c.1993+327 C>T, previously termed C/T-13910); 14kb upstream from the *LCT* locus that demonstrated complete agreement with clinically identified lactase non persistence in a Finnish

population. *LCT* is a gene which encodes Lactase Phlorizin Hydrolase (LPH) for the hydrolysis of lactose. Homozygotes for the C allele of rs4988235 were found to be lactase-non persistent, whilst those heterozygous or homozygous for the T allele were determined to be lactase persistent. This finding has been replicated in several other European populations [11, 12] and support for the functionality of this DNA variant has been demonstrated *in vitro* [13]. Thus the C/C genotype for rs4988235 is now a widely accepted marker of adult-type hypolactasia in Europeans.

Evaluating the association of lactase persistence with CD as a key factor influencing exposure to MAP via the consumption of milk products is particularly important in New Zealand, a country with one of the highest incidences of CD in the Western world at 16.5/100,000 per year [14] coupled with the highest herd prevalence estimate of Johne's disease at 60% [5].

The aim of this study was to determine whether lactase persistence as evident by the presence of the T allele of rs4988235 is associated with CD in a New Zealand population.

## **Methods**

### ***Samples***

The Auckland CD Project is a population-based study of genetic and environmental determinants of CD aetiology. CD patients were recruited between May 2005 and April 2007 through local doctors and surgeries in Auckland, New Zealand, and also other North Island centres, by media campaigns, e.g. local newspaper and television. Controls were recruited by the same means, and also recruited from non-affected family members and friends of CD patients. Participants consented to collection of peripheral blood for DNA extraction and genotyping.

A total of 945 subjects (333 CD patients and 612 controls) consented to take part. The cases in this study are a subset of the Caucasian participants of the Auckland CD Project. Their gastroenterologists were approached, clinical records scrutinised to confirm diagnosis, and CD was defined using standard diagnostic criteria [15]. Cases were phenotyped according to the Montreal Classification systems, allowing genotype-phenotype analysis to be performed. Phenotypic information was available for 323 CD patients. All participants included in this study were of self-reported European ancestry (subjects who self-reported having any Maori or other non-Caucasian ancestry are not included in the dataset).

The study was conducted under ethical protocol MEC/04/12/011, authorised through the New Zealand Multi-Region Human Ethics Committee. All study subjects gave informed consent.

DNA was extracted from blood samples or buccal swabs using Qiagen's DNA extraction kit (Qiagen, Valencia, CA, USA) and following the manufacturer's instructions.

### ***Genotyping***

Genotyping of rs4988235 was performed using the ABI TaqMan MGB diallelic discrimination system with forward and reverse primer sequences 5'-CTG CGC TGG CAA TAC AGA TAA G-3' and 5'AAA TGC AAC CTA AGG AGG AGA GTT C-3' respectively and VIC-- ATA ATG TAG TCC CTG GCC T and FAM--ATA ATG TAG CCC CTG GC probes (Applied Biosystems, Melbourne, Australia). The reactions were prepared using 2x TaqMan Universal Master Mix, 40x SNP Genotyping Assay Mix, DNase-free water, and 10 ng genomic DNA in a final volume of 5 ul per reaction. The PCR amplification was performed using the ABI



Prism 7900 HT sequence-detector machine under the following conditions: 50°C for 2 minutes, 10 min 95°C enzyme activation followed by 40 cycles at 95°C for 15 sec and 60°C for 1 min (annealing/extension).

To reduce the possibility of genotyping error, genotyping was repeated utilising different primers (5'-CTC TGC GCT GGC AAT ACA G and 5'-AAA TGC AAC CTA AGG AGG AGA) and probes (VIC-ATA AGA TAA TGT AGC CCC TGG C and FAM- ATA AGA TAA TGT AGT CCC TGG C) (Applied Biosystems, Melbourne, Australia) under the following conditions: 60°C for 1 minute, 95°C for 10 minutes, followed by 55 cycles of 92°C for 15s and 60°C for 1 minute and 60°C for 1 minute.

The allelic discrimination results for both sets of genotyping were determined after the amplification by performing an endpoint read.

Assays were optimized in 24 samples consisting of 20 reference Centre d'Etude du Polymorphisme Humain (CEPH) samples and 4 blanks. All sample plates contained cases, controls, blanks, CEPH and duplicate samples. Quality control measures included independent double genotyping, blind to sample identity and blind to the other caller, and comparison of our CEPH genotypes to those in the HapMap ([www.hapmap.org](http://www.hapmap.org)).

### **Statistical analysis**

The allelic trend test [16] was used to compare frequencies between case and control alleles. An exact test was used to test for departures from Hardy-Weinberg equilibrium (HWE) in the case and the control samples [17]. Lactase persistence was defined as having genotypes C/T and T/T of rs4988325 whilst C/C was defined as lactase non-persistence.

Genotypic analysis was carried out and the allelic odds ratio with confidence interval was calculated to test the effect of the T allele on CD risk.

SAS (V9.1 SAS Institute., Cary, NC, USA) and R (Ihaka and Gentleman, 1996) were used for statistical analyses.

## Results

The genotypes did not meet Hardy-Weinberg equilibrium (HWE) for both the cases and controls. To ensure genotyping errors did not exist in the data, genotyping was repeated using an alternative primer and probe set and the genotypes were confirmed to be correct. Also, the minor allele frequency was 0.273 which is similar to the HapMap CEU population frequency of 0.274. Deviations from HWE may have resulted by chance, as a result of population admixture or perhaps the rs4988235 SNP is associated with a currently unknown factor in controls.

For rs4988235, the T/T genotype showed a significantly increased risk of having CD as compared with the C/C genotype (OR=1.61, 95% CI=1.03-2.51) however, no significant difference was found between C/T and C/C. Additionally, a significant increase in the frequency of the T allele was observed in CD patients (OR=1.30, 95% CI=1.05-1.61,  $p=0.013$ ), indicating that the T allele encoding lactase persistence was associated with risk of CD (Table 1).

Genotype-phenotype analysis: of 323 cases, 88 (28.4%) had relatives with IBD, 98 (31.6%) had undergone a bowel resection, 44 (14.2%) had extra-intestinal manifestations of CD occurring and 112 (46.9%) smoked at diagnosis (Table 2).

Individuals carrying the T allele of rs4988235 had an increased probability of developing CD that was diagnosed between 17 and 40 years (OR=1.36, 95% CI=1.05-1.76,  $p=0.021$ ); and a significantly increased risk of developing ileal CD (OR=1.50, 95% CI=1.05-1.76,  $p=0.032$ ) or CD of an inflammatory nature (OR=1.36, 95%

CI=1.02-1.82,  $p=0.039$ ). In addition, a significantly increased risk of having other relatives with IBD and of undergoing bowel resection was observed in individuals with the T allele of rs4988235 (OR=1.71, 95% CI=1.17-2.51,  $p=0.006$  and OR=1.47, 95% CI=1.03-2.10,  $p=0.033$  respectively).

## Discussion

Results from this study suggest that genetic adult-type lactase persistence as evident by the T allele of rs4988235 may increase the risk of developing CD in this New Zealand population.

Our findings are in agreement with Juste et al [18], who reported an increased frequency of the T allele of rs4988235 in CD patients (61.9%) compared with controls (47.1%,  $p=0.0275$ ). Similarly a highly significant correlation between incidence of CD and lactase persistence ( $r= -0.655$ ,  $p= 0.0017$ ) was reported in an observational study encompassing data from 20 countries [19], whilst Shrier et al [20] reported a reduction in CD risk with increasing prevalence of lactase non-persistence in a review of global population data.

Conversely, Büning et al [21] failed to find an association between the C/C genotype of rs4988235 and the pathogenesis of CD in a German case-control study of 166 CD patients and 187 healthy controls. This discrepancy may relate to the smaller sample size of this study; however evaluation of this phenomenon on a regional basis, taking into consideration adult lactase status, dairy food consumption, and degree and management of MAP infection within ruminant populations may provide evidence to ensure that a consensus can be reached regarding the role of lactase persistence and MAP exposure in the aetiology of CD.

The finding that lactase persistence was associated with an increased risk of having a relative with inflammatory bowel disease may reflect the autosomal

dominant nature of lactase persistence, whilst the increased risk of CD diagnosis between 17-40 years of age may correspond with the onset of hypolactasia that typically occurs in adulthood in susceptible individuals. The T allele for rs4988235 was also associated with an increased risk of inflammation, having an ileal CD location and bowel resection amongst CD cases. This was an unexpected finding and warrants further investigation to conclusively determine whether lactase persistence may influence the phenotypic expression of CD.

As associations between MAP and inflammatory bowel disease have largely been limited to individuals with CD, this study did not evaluate lactase persistence in the ulcerative colitis context. However, the inclusion of this clinical population in future research may provide further insights into the potential association between lactase persistence and inflammatory bowel disease.

Whether lactase persistence is an independent risk factor for CD or merely a marker of the likelihood of exposure to the MAP pathogen, found predominantly in milk products, is beyond the scope of this particular study. However epidemiological evidence is conflicting, indicating an urgent need for further, well controlled dietary and genetic case-control studies to distinguish between these two hypotheses.

In Japan, Shoda et al [22] reported that incidence of CD was strongly correlated with consumption of milk protein and Shrier et al [18] identified a trend with estimated yearly per capita dairy food consumption and increased risk of CD. In contrast Abubakar et al [23] reported that consumption of pasteurized milk was associated with a decreased risk of CD in a case-control study of 218 individuals with CD in England.

Recent advances in technology have ensured that the presence of MAP *in vivo* is more accurately detected [24]. These advances have contributed to the resurgence

of the MAP and CD hypothesis. Viable MAP was recently identified in the peripheral blood of a high proportion of individuals with CD, with only MAP DNA isolated in a small proportion of controls [25]. This may reflect different physiological responses to the pathogen between cases and controls. These findings were supported locally in a New Zealand study, utilising a larger cohort of 361 CD patients and 200 controls. In this study MAP DNA in peripheral blood was found to be over-represented in CD patients in comparison to controls [26].

The Zoonotic potential of MAP was discussed in a review by Grant [5], who reported that hypotheses suggesting a causal role for MAP in the pathogenesis of CD were plausible, however insufficient available evidence can either prove or disprove this idea at present.

Results from this exploratory study do not prove that MAP exposure is responsible for the development of CD; or that milk products increase risk of the development of CD. However our findings indicate that there may be a complex interaction between exposure to this infectious agent as a consequence of an evolutionary adaptation to ensure tolerance to the milk products that may serve as the medium for its delivery to humans. As the worldwide incidence of CD is rapidly increasing, [27-29], there is an urgent need to determine whether our findings can be replicated in larger populations. If so, efforts to eradicate MAP through animal vaccination and greater prevention of its contamination into the food supply should become key public health priorities.

## **Conclusion**

Following a case-control evaluation of rs4988235 in a New Zealand CD population, we found that lactase persistence as evident by the presence of a T allele was

associated with increased CD risk. This finding may relate to the increased exposure to dairy products in individuals with lactase persistence, a food group that has been implicated in the aetiology of CD due to the presence of MAP species in milk. Further research is required to substantiate our findings and to conclusively determine the nature of the observed association between lactase persistence and CD.

## **Competing interests**

The author(s) declare that they have no competing financial interests.

## **Authors' contributions**

DJN conceived the study and conducted the genotyping of blood samples with WJL.

DJN assisted with data analysis, interpretation of data and drafted the manuscript.

DYH performed the statistical analyses and contributed to the manuscript. ARM made suggestions to the study and contributed to the manuscript. AGF provided the clinical data and many of the samples obtained. LRF made suggestions for the study and revised the manuscript. LCT revised the manuscript. All authors read and approved the final draft.

## **Acknowledgements**

Deborah Nolan is the recipient of an Australian Endeavour Research Fellowship.

Nutrigenomics New Zealand is a collaboration between AgResearch Ltd, Plant and Food research and The University of Auckland and is largely funded by the foundation for Research, Science and Technology.

## References

1. Chamberlin W, Graham DY, Hulten K, El-Zimaity HMT, Schwartz MR, Naser S, Shafran I, El-Zaatari FAK: *Mycobacterium avium subsp. Paratuberculosis* as one cause of Crohn's disease. *Aliment Pharmacol Ther* 2001, **15**:337-346
2. Greenstein, RJ: **Is Crohn's disease caused by a mycobacterium? Comparisons with leprosy, tuberculosis, and Johne's disease.** *Lancet Infect Dis* 2003, **3**:507-514
3. Sweeney RW, Whitlock RH, Rosenberger AE: *Mycobacterium paratuberculosis* cultured from milk and supramammary lymph nodes of infected asymptomatic cows. *J. Clin. Microbiol* 1992 **30**:166-171
4. Ayele WY, Svastova P, Roubal P, Bartos M, Pavlik I: *Mycobacterium avium Subspecies paratuberculosis* Cultured from Locally and Commercially Pasteurized Cow's Milk in the Czech Republic. *Appl. Environ. Microbiol* 2005 **71**:1210-1214
5. Grant IR: **Zoonotic potential of *Mycobacterium avium* ssp. paratuberculosis: the current position.** *J Appl Microbiol* 2005 **98**:1282-1293.
6. Grant IR, Ball HJ, Rowe MT: **Incidence of *Mycobacterium paratuberculosis* in Bulk Raw and Commercially Pasteurized Cows' Milk from Approved Dairy Processing Establishments in the United Kingdom.** *Appl. Environ. Microbiol* 2002 **68**:2428-2435
7. Vesa TH, Marteau P, Korpela R: **Lactose Intolerance.** *J Am Coll Nutr* 2000 **19**:165S-175.
8. Tishkoff SA, Reed FA, Ranciaro A, Voight BF, Babbitt CB, Silverman JS, Powell K, Mortensen HM, Hirbo JB, Osman M, Ibrahim M, Omar SA, Lema G, Nyambo TB, Gori J, Bumpstead S, Pritchard JK, Wray GA, Deloukas P: **Convergent adaptation of human lactase persistence in Africa and Europe.** *Nat Genet* 2007 **39**:31-40

9. Sahi, T: **Genetics and Epidemiology of Adult-type Hypolactasia.** Scand J Gastroenterol 1994 **29**:7-20
10. Enattah NS, Sahi T, Savilahti E, Terwilliger JD, Peltonen L, Järvelä I: **Identification of a variant associated with adult-type hypolactasia.** Nat Genet 2002 **30**:233 – 237
11. Kerber M, Oberkanins C, Kriegshäuser G, Kollertis B, Dossenbach-Glaninger A, Fuchs D, Ledochowski M: **Hydrogen breath testing versus LCT genotyping for the diagnosis of lactose intolerance: A matter of age.** Clinica Chimica Acta 2007 **383**:91-96
12. Ridefelt P, Håkansson LD: **Lactose intolerance: Lactose tolerance test versus genotyping.** Scand J Gastroenterol 2005 **40**:822-826
13. Lewinsky RH, Jensen TGK, Møller J, Stensballe A, Olsen J, Troelsen JT: **T-13910 DNA variant associated with lactase persistence interacts with Oct-1 and stimulates lactase promoter activity *in vitro*.** Hum Mol Genet 2005 **14**:3945-3953
14. Gearry RB, Richardson A, Frampton CMA, Collett JA, Burt MJ, Chapman BA, Barclay ML: **High incidence of Crohn's disease in Canterbury, New Zealand: Results of an epidemiologic study.** Inflamm Bowel Dis 2006 **12**:936-943
15. Lennard-Jones JE: **Classification of Inflammatory Bowel Disease.** Scand J Gastroenterol 1989 **24**:2-6
16. Sasieni P D: **From Genotypes to Genes: Doubling the Sample Size.** Biometrics 1997 **53**:1253-1261
17. Wigginton JE, Abecasis GR: **PEDSTATS: descriptive statistics, graphics and quality assessment for gene mapping data.** Bioinformatics 2005 **21**: 3445-3447(3)
18. Juste RA, Elguezabal N, Chamorro S, Molina E, Garrido JM, Riestra S, de Francisco R, Rodrigo L: **Associations between CARD15 polymorphisms, MAP DNA in blood and lactase persistence in a Crohn's disease case-control study in North Spain.** In *Proceedings of the 10<sup>th</sup> International Colloquium on paratuberculosis: 9-14 August 2009, Minnesota.* 2009:269
19. Juste R A: **Crohn's disease and ruminant farming. Got lactase?** Med Hypotheses 2010 **75**:7-13



20. Shrier I, Szilagyi A, Correa JA: **Impact of Lactose Containing Foods and the Genetics of Lactase on Diseases: An Analytical Review of Population Data.** Nutr Cancer 2008 **60**:292- 300
21. Büning C, Ockenga J, Krüger S, Jurga J, Baier P, Dignass A, Vogel A, Strassburg C, Weltrich R, Genschel J, Lochs H, Schmidt H: **The C/C (-13910) and G/G (-22018) Genotypes for Adult-type Hypolactasia are not Associated with Inflammatory Bowel Disease.** Scand J Gastroenterol 2003 **38**:538 – 542
22. Shoda R, Matsueda K, Yamato S, Umeda N: **Epidemiologic analysis of Crohn disease in Japan: increased dietary intake of n-6 polyunsaturated fatty acids and animal protein relates to the increased incidence of Crohn disease in Japan.** Am J Clin Nutr 1996 **63**:741-745
23. Abubakar I, Myhill DJ, Hart AR, Lake AR, Harvey I, Rhodes JM, Robinson R, Lobo AJ, Probert CSJ, Hunter PR: **A Case-Control Study of Drinking Water and Dairy Products in Crohn's Disease--Further Investigation of the Possible Role of *Mycobacterium avium paratuberculosis*.** Am. J. Epidemiol, 2007 **165**:776-783.
24. Bull TJ, McMinn EJ, Sidi-Boumedine K, Skull A, Durkin D, Neild P, Rhodes G, Pickup R, Hermon-Taylor J: **Detection and Verification of *Mycobacterium avium subsp. paratuberculosis* in Fresh Ileocolonic Mucosal Biopsy Specimens from Individuals with and without Crohn's Disease** J. Clin. Microbiol, 2003 **41**:2915-2923.
25. Naser SA, Ghobrial G, Romero C, Valentine JF: **Culture of *Mycobacterium avium subspecies paratuberculosis* from the blood of patients with Crohn's disease.** Lancet, 2004 **364**:1039-1044
26. Bentley RW, Keenan JI, Gearry RB, Kennedy MA, Barclay ML, Roberts RL: **Incidence of *Mycobacterium avium Subspecies paratuberculosis* in a Population-Based Cohort of Patients With Crohn's Disease and Control Subjects.** The American Journal of Gastroenterology, 2008 **103**:1168-1172
27. Munkholm P, Langholz E, Nielsen OH, Kreiner S, Binder V: **Incidence and Prevalence of Crohn's Disease in the County of Copenhagen, 1962-87: A Sixfold Increase in Incidence.** Scand J Gastroenterol, 1992 **27**:609-614

28. Phavichitr N, Cameron DJ, Catto-Smith AG: **Increasing incidence of Crohn's disease in Victorian children.** Journal of Gastroenterology and Hepatology, 2003 **18**:329-332
29. Leong RWL, Lau JY, Sung JJY: **The epidemiology and phenotype of Crohn's disease in the Chinese population.** Inflamm Bowel Dis, 2004 **10**:646-651

Table 1. Genotype and allele count frequencies for rs4988235 in CD patients and in controls

Genotype	CD	Control	OR (95%CI)	p
Allele	N (%)	N(%)		
C/C	32 (9.6)	82 (13.4)	1.00	
C/T	118 (35.4)	238 (38.9)	1.27 (0.80-2.02)	0.065
T/T	183 (55.0)	292 (47.7)	<b>1.61 (1.03-2.51)</b>	
<b>Z= 2.34, p &lt; Z 0.009*</b>				
T	484 (72.7)	822 (67.2)	<b>1.30 (1.05-1.61)</b>	<b>0.013</b>
C	182 (27.3)	402 (32.8)	1.00	

\*Cochran-Armitage Trend Test

Table 2. Summary of clinical data in subgroups of CD patients with a T allele for rs4988235

	N (%)	OR (95% CI)	p
Age at first diagnosis			
0-16 years	30 (10.6)	0.98 (0.55-1.75)	1.00
17-40 years	203 (71.5)	<b>1.36 (1.05-1.76)</b>	<b>0.021</b>
>40 years	51 (18.0)	1.56 (0.96-2.53)	0.087
CD location			
Ileal	97 (36.5)	<b>1.50 (1.05-2.15)</b>	<b>0.032</b>
Colonic	86 (32.3)	1.26 (0.88-1.81)	0.245
Ileocolonic	83 (31.2)	1.25 (0.86-1.81)	0.273
CD Behaviour			
Inflammatory	152 (57.6)	<b>1.36 (1.02-1.82)</b>	<b>0.039</b>
Strictureing	81 (30.7)	1.27 (0.87-1.85)	0.233
Penetrating	31 (11.7)	1.41 (0.77-2.57)	0.315
Relative with IBD	88 (28.4)	<b>1.71 (1.17-2.51)</b>	<b>0.006</b>
Bowel resection	98 (31.6)	<b>1.47 (1.03-2.10)</b>	<b>0.033</b>
Extra Intestinal Manifestations	44 (14.2)	1.38 (0.84-2.28)	0.228
Smoker at diagnosis	CD: 112 (46.9) Control: 130 (31.7)	1.04 (0.69-1.56)	0.918

\*IBD – Inflammatory Bowel Disease