2019

Phosphine-Scavenging Cationic Gold(I) Complexes: Alternative Applications of Gold Cocatalysis in Fundamental Palladium-Catalyzed Cross-Couplings

Curtis Ho

University of Tasmania

Alireza Ariaferd

Islamic Azad University, University of Tasmania (UTAS), Alireza.Ariaferd@utas.edu.au

Christopher J. T Hyland

University of Wollongong, chrhyl@uow.edu.au

Alex C. Bissember

University of Tasmania (UTAS)

Publication Details

Ho, C. C., Ariaferd, A., Hyland, C. J.T. & Bissember, A. C. (2019). Phosphine-Scavenging Cationic Gold(I) Complexes: Alternative Applications of Gold Cocatalysis in Fundamental Palladium-Catalyzed Cross-Couplings. *Organometallics*, 38 (13), 2683-2688.

Phosphine-Scavenging Cationic Gold(I) Complexes: Alternative Applications of Gold Cocatalysis in Fundamental Palladium-Catalyzed Cross-Couplings

Abstract

We have demonstrated that air-stable cationic gold(I) cocatalysts have the capacity to enhance the efficiency of palladium-catalyzed cross-couplings. Specifically, we determined that a 1:1 $[\text{Pd}\{\text{P}(\text{t-Bu})_3\}_2]/[\text{Au}\{\text{P}(\text{t-Bu})_3\}(\text{NTf}_2)]$ system provides superior reactivity relative to $[\text{Pd}\{\text{P}(\text{t-Bu})_3\}_2]$, across Suzuki-Miyaura, Stille, and Mizoroki-Heck reactions performed under mild conditions. Our results are consistent with cationic gold(I) species serving primarily as phosphine scavengers in this chemistry, as recently predicted by density functional theory (DFT).

Publication Details

Ho, C. C., Ariafard, A., Hyland, C. J.T. & Bissember, A. C. (2019). Phosphine-Scavenging Cationic Gold(I) Complexes: Alternative Applications of Gold Cocatalysis in Fundamental Palladium-Catalyzed Cross-Couplings. *Organometallics*, 38 (13), 2683-2688.

Phosphine-Scavenging Cationic Gold(I) Complexes: Alternative Applications of Gold Cocatalysis in Fundamental Palladium-Catalyzed Cross-Couplings

Curtis C. Ho,[†] Alireza Ariafard,[†] Christopher J. T. Hyland,[‡] and Alex C. Bissember^{*†}

[†]School of Natural Sciences – Chemistry, University of Tasmania, Hobart, Tasmania 7001, Australia

[‡]School of Chemistry and Molecular Bioscience, University of Wollongong, Wollongong, New South Wales 2522, Australia

ABSTRACT: We have demonstrated that air-stable cationic gold(I) cocatalysts have the capacity to enhance the efficiency of palladium-catalyzed cross-couplings. Specifically, we determined that a 1:1 [Pd{P(*t*-Bu)₃}₂]/ [Au{P(*t*-Bu)₃}(NTf₂)] system provides superior reactivity relative to [Pd{P(*t*-Bu)₃}₂], across Suzuki–Miyaura, Stille, and Mizoroki–Heck reactions performed under mild conditions. Our results are consistent with cationic gold(I) species serving primarily as phosphine scavengers in this chemistry, as recently predicted by density functional theory (DFT).

INTRODUCTION

Palladium-catalyzed cross-couplings represent a fundamentally important class of synthetic transformations of both academic and industrial significance.¹ Arylphosphine ligands have been employed to great effect as versatile supporting ligands in these reactions, as have bulky, electron-rich trialkylphosphines.¹ The latter family of ligands, particularly the venerable P(*t*-Bu)₃, have enabled efficient Pd-catalyzed reactions under mild conditions.² Indeed, in 2010, P(*t*-Bu)₃ was described as, “the most widely used single ligand in modern cross coupling chemistry”.^{2d}

In contrast to many other Pd/PR₃ catalyst systems, it is generally accepted that the oxidative addition of aryl electrophiles to P(*t*-Bu)₃-ligated Pd(0) species predominantly proceeds via the monophosphine-ligated intermediate [Pd{P(*t*-Bu)₃}].^{3,4} It has also been reported that catalytically-inactive [Pd{P(*t*-Bu)₃}₂] comprises the catalyst resting state in various Mizoroki–Heck and Suzuki–Miyaura couplings.^{3a,c} Thus, ~1:1 ratios of Pd : P(*t*-Bu)₃ are often employed in catalysis performed at lower temperatures, and the active species is often generated in situ from standard palladium precatalysts.^{2c,3a,5}

Beyond the development of new ligands that provide enhanced reactivity, a number of strategies have been used to augment various features of Pd-mediated couplings and the emergence of homo- and heterobimetallic systems in homogenous catalysis are examples of this.⁶ Gold complexes have been employed in the latter capacity.^{6a,b} Many of these studies describe Au(I) species serving as transmetalating intermediates derived from various alkyne/ allene activation processes characteristic of Au(I) complexes, or from preformed organogold complexes.⁷ Au(I) species have been employed as cocatalysts in Pd-catalyzed Stille reactions to

facilitate efficient cross-couplings with sterically hindered nucleophiles and tin/ gold transmetalation mechanisms are suggested to operate in this system.⁸ Other modes of reactivity have also been invoked. For example, Blum and co-workers reported a detailed study demonstrating the viability of a Pd/ Au cocatalyst system to promote the carbostannylation of alkynes.⁹ The mechanism is consistent with the coordination of a Lewis acidic Au(I) species to the alkyne, which is key to activation in this manifold.^{9,10}

In 2017, Ariafard’s density functional theory (DFT) computational study first predicted the phosphine-scavenging ability of cationic Au(I) complexes (Figure 1).¹¹ Specifically, this work suggested that this property could be exploited to lower the barrier leading to the catalytically active monophosphine-ligated [Pd{P(*t*-Bu)₃}] from its stable (catalytically-inactive) bis-ligated precursor. This prompted us to postulate that Pd/ Au cocatalysis could be exploited to enhance the reactivity of this (and perhaps other Pd/ PR₃) catalyst systems under mild conditions by sequestering free phosphine. Herein, we illustrate the validity of this postulate in practice.

Figure 1. Calculated Gibbs free energies (kcal/mol) for the formation of [Pd{P(*t*-Bu)₃}] from [Pd{P(*t*-Bu)₃}₂] in the presence or absence of [Au–L]⁺.¹¹

Our results provide evidence for the capacity of cationic Au(I) complexes to formally abstract $P(t\text{-Bu})_3$ from $[\text{Pd}\{P(t\text{-Bu})_3\}_2]$. Furthermore, we determined that $[\text{Pd}\{P(t\text{-Bu})_3\}_2]/[\text{Au}\{P(t\text{-Bu})_3\}(\text{NTf}_2)]$ systems can provide superior efficiency relative to $[\text{Pd}\{P(t\text{-Bu})_3\}_2]$ in key couplings, including: Suzuki–Miyaura, Stille, and Mizoroki–Heck reactions. These data are consistent with the gold cocatalyst playing a key role primarily as a phosphine scavenger.

RESULTS AND DISCUSSION

First, we investigated the capacity of $[\text{Au}\{P(t\text{-Bu})_3\}(\text{NTf}_2)]$ (**1**) to formally abstract $P(t\text{-Bu})_3$ from the bis(phosphine)-ligated Pd(0) complex $[\text{Pd}\{P(t\text{-Bu})_3\}_2]$ (**2**) (Scheme 1). After the addition of Au complex **1** to Pd complex **2**, a new signal (δ 97.0 ppm) appeared, consistent with the rapid formation of $[\text{Au}\{P(t\text{-Bu})_3\}]^+$, as judged by ^{31}P NMR spectroscopy.^{12,13} We also isolated and crystallographically characterized linear mononuclear $[\text{Au}\{P(t\text{-Bu})_3\}_2]\text{NTf}_2$ (**3**).¹⁴

Scheme 1. Reaction of Au complex **1** with Pd complex **2** to afford $[\text{Au}\{P(t\text{-Bu})_3\}_2]\text{NTf}_2$ (**3**). In the structural representation of Au complex **3**, thermal ellipsoids are drawn at the 50% probability level and the $[\text{NTf}_2]^-$ counteranion and hydrogen atoms are omitted for clarity.

Next, we explored the capacity of Au complex **1** to promote the oxidative addition of *p*-bromotoluene to a Pd species derived from $[\text{Pd}\{P(t\text{-Bu})_3\}_2]$ (**2**). In the absence of Au complex **1**, reaction of the aryl bromide with Pd complex **2** was not observed after 24 h, as judged by ^{31}P NMR spectroscopy (Scheme 2). This is in agreement with germane findings.^{3a} In stark contrast, the presence of equimolar quantities of Au complex **1** rapidly promoted oxidative addition in minutes. Specifically, ^{31}P NMR spectroscopy indicated the presence of a new signal (δ 61.5 ppm),¹⁵ which was consistent with the presence of Pd(II) species **4**. Conversion of Pd(0) complex **2** to Pd(II) complex **4** was complete within 0.25 h and we also observed the concomitant formation of $[\text{Au}\{P(t\text{-Bu})_3\}]^+$ by ^{31}P NMR spectroscopy.¹²

We obtained analogous results employing other commercially available, air-stable cationic Au(I) complexes **5–7** in equivalent experiments. In each case, oxidative addition to form Pd(II) complex **4** promptly occurred at ambient temperature. This property of species **1**, and **5–7** is notable as Fu and coworkers have suggested that (catalytically inactive) Pd complex **2** comprises the catalyst resting state in various Mizoroki–Heck and Suzuki–Miyaura couplings.^{3a,c} The presence of free phosphine has also been shown to reduce the efficiency of transmetalation in Suzuki–Miyaura

and Stille cross-coupling reactions mediated by Pd/ PPh_3 systems.^{16,17}

The capacity of substoichiometric quantities of copper salts to accelerate and enhance the efficiency of Pd-catalyzed Stille and Suzuki–Miyaura couplings is well established.^{18,19} Much of the pioneering work exploring the role of copper additives in Stille chemistry was undertaken by Liebeskind and excellent reviews discussing this “copper effect” within the context of Pd cross-coupling chemistry have been published more recently.^{6b,20} Although it can be difficult to unambiguously determine the mechanistic basis of this phenomenon in every instance, two primary effects have been proposed.^{6b} Specifically, the operation of possible tin/ copper transmetalation mechanisms have been invoked in these Stille processes (particularly in very polar solvents such as NMP, DMF, and DMSO),^{6b,17b,21} in addition to the phosphine-scavenging capacity of copper species.^{17b,c,18a,22} The latter effect is thought to be most pronounced when free, strong donor phosphine ligands are present in ethereal solutions as their presence can impede transmetalation.^{6b,17b,c,23}

Scheme 2. Investigating the capacity of Au complexes **1** and **5–7** and Cu complex **8** to promote the oxidative addition of a Pd species to *p*-bromotoluene.

Donnelly and Finet first suggested phosphine-scavenging processes may be operative in Suzuki–Miyaura reactions performed in the presence of CuI.^{19a} Furthermore, Hobbs noted a specific 1:2 Pd/ Cu cocatalyst ratio provided optimal results in Sonogashira reactions employing $[\text{Pd}(\text{PPh}_3)_4]$ and proposed that this type of behavior may also be present.²⁴ Ariafard’s DFT studies also support the phosphine-scavenging ability of cationic Cu(I) complexes.¹¹ Consequently, we also determined that $[\text{Cu}(\text{MeCN})_4]\text{PF}_6$ (**8**) could rapidly promote the oxidative addition of *p*-bromotoluene to an active species derived from $[\text{Pd}\{P(t\text{-Bu})_3\}_2]$ (Scheme 2). We also isolated and crystallographically characterized the mononuclear complex $[\text{Cu}\{P(t\text{-Bu})_3\}_2]\text{PF}_6$ (**9**).²⁵

To explore the viability of our proposed phosphine-scavenging strategy in synthesis, we investigated the effects of Pd complex **2**/ Au complex **1** (and Pd complex **2**/ Cu complex **8**) cocatalyst systems in Pd-catalyzed cross-coupling

processes under simple and mild reaction conditions. In each case, we compared the respective rates of product formation employing these two cocatalyst systems and benchmarked these data against equivalent experiments featuring Pd complex **2** exclusively.

The enhanced efficiency provided by our Pd/ Au cocatalyst system was clearly evident in a Suzuki–Miyaura cross-coupling involving *p*-bromotoluene and phenylboronic acid (Figure 2). In the absence of additive **1**, the reaction essentially featured an induction period (~0.5 h). In contrast, the process using 1% [Au(PPh₃)₃](NTf₂) (**1**) provided an enhanced rate of product formation.^{26,27} Employing 3% Au complex **1** afforded negligible differences in reactivity relative to 1% loading (see Supporting Information), which is consistent with the phosphine-scavenging capacity of Au complex **1** playing a key role in improving the efficiency of this transformation.²⁸ Similar results were also obtained when a 1% Pd(dba)₂/ 1% [HP(*t*-Bu)₃]BF₄ catalyst system was utilized in this transformation (see Supporting Information).

Figure 2. Effects of cocatalysts Au complex **1** (red line) or Cu complex **8** (blue line) on the progress of Pd-mediated Suzuki–Miyaura cross-couplings. The experiment employing only Pd catalyst **2** is represented with a black line.²⁹

In a recent study,^{16,30} we identified that the poor turnover of key on-cycle intermediate *trans*-[Pd(PPh₃)₂(Ar)(I)] in the presence of free PPh₃ was responsible for the surprisingly poor reactivity of aryl iodides in [Pd(PPh₃)₄]-catalyzed Suzuki–Miyaura cross-couplings performed at relatively low temperatures (Schemes 3A and 3B).³¹ Our report also illustrated the drastic difference between this process and the equivalent reaction utilizing aryl bromides (Scheme 3A). With this in mind, we investigated whether gold catalysis could ameliorate the aforementioned issues affecting catalyst turnover in this transformation. Although a 1% loading of [Au(PPh₃)(NTf₂)] (**5**) provided a negligible effect, 2% Au complex **5** facilitated a much more efficient process (Scheme 3C).

Employing 3% [Au(PPh₃)(NTf₂)] afforded essentially

the same results relative to 2% **5**. Significant reactivity differences were observed when 1:1 and 1:2 ratios of [Pd(PPh₃)₄] and [Au(PPh₃)(NTf₂)] were employed in these Suzuki–Miyaura couplings. In contrast, similar results were obtained using 1:2 and 1:3 ratios of [Pd(PPh₃)₄] and [Au(PPh₃)(NTf₂)]. These data are consistent with the phosphine-scavenging capacity of Au complex **5** playing a role in enhancing the efficiency of this process.^{28,32} We also investigated the ability of [Au(PPh₃)(NTf₂)] (**5**) to formally abstract PPh₃ from [Pd(PPh₃)₄] (Scheme 3D). After the addition of Au complex **5** to a solution containing this Pd(0) species, a new signal (δ 22.1 ppm) was observed, consistent with the formation of [Au(PPh₃)₃]⁺ (**10**), as judged by ³¹P NMR spectroscopy.³³ This demonstrates capacity for Au complex **5** to scavenge PPh₃.³⁴

Scheme 3. A) Reported inefficient Pd-catalyzed Suzuki–Miyaura couplings of aryl iodides.¹⁶ B) Reported poor turnover of *trans*-[Pd(PPh₃)₂(Ar)(I)] in the presence of free PPh₃.¹⁶ C) Investigating the effects of cocatalysts **5** or **8** on the efficiency of Pd-catalyzed Suzuki–Miyaura reactions with *p*-iodotoluene. D) Reaction of Au complex **5** with [Pd(PPh₃)₄] to afford [Au(PPh₃)₃]⁺ (**10**). ^aYields after 0.25 h are provided in parentheses.²⁹

Next, we demonstrated the superior results provided by our Pd/ Au cocatalyst system in a Stille cross-coupling involving *p*-iodotoluene and tributylphenylstannane (Figure 3). Specifically, the process featuring 1% Au complex **1** provided a more efficient process relative to the reaction

featuring only Pd complex **2**.^{26,27} Employing 3% Au complex **1** provided negligible improvements in reactivity relative to 1% Au complex **1** (see Supporting Information). However, we cannot exclude the possibility that tin/ gold transmetalation mechanisms may also be operative under these conditions. The enhanced efficiency we observed in each of the Pd-catalyzed Stille and Suzuki–Miyaura cross-couplings investigated in this study is consistent with the Au cocatalyst scavenging phosphine and, by doing so, accelerating the respective rates of transmetalation in these processes.

Figure 3. Effects of cocatalysts Au complex **1** (red line) or Cu complex **8** (blue line) on the progress of Pd-mediated Stille cross-couplings. The experiment employing only Pd catalyst **2** is represented with a black line.²⁹

Figure 4. Effects of cocatalysts Au complex **1** (red line) or Cu complex **8** (blue line) on the progress of Pd-mediated Mizoroki–Heck cross-couplings. The experiment employing only Pd catalyst **2** is represented with a black line.²⁹

The improved results provided by our Pd/ Au cocatalyst

system were also demonstrated in a Mizoroki–Heck cross-coupling involving *p*-iodotoluene and ethyl acrylate (Figure 4).^{26,27} Our results indicated that the process featuring 2% [Au{P(*t*-Bu)₃}(NTf₂)] (**1**) provided a more efficient process relative to the reaction featuring only Pd complex **2**. Employing 4% Au complex **1** provided minor improvements in reactivity relative to 2% Au complex **1** (see Supporting Information). Similar results were also obtained when a 2% Pd(dba)₂/ 2% [HP(*t*-Bu)₃]BF₄ catalyst system was employed to effect this transformation (see Supporting Information).

CONCLUSIONS

In summary, we have demonstrated that substoichiometric quantities of air-stable cationic Au(I) complexes can accelerate and enhance the efficiency of Pd-catalyzed cross-coupling reactions. Our observations are consistent with Au(I) cocatalysts serving primarily as phosphine scavengers in these processes and these findings reinforce the predictions of a recently published DFT study.¹¹ Our results also suggest that gold additives may be more effective than copper salts in this regard, which should be investigated further in the future.

Arguably, the viability and general utility of this Au(I) cocatalysis approach in augmenting the development of Pd-catalyzed reactions under mild conditions will be influenced by factors including: the donor-capacity of the ligand employed, binding affinity of the gold complex, choice of solvent, and additives present. Subsequent studies will seek to investigate the broader implications and further applications of this strategy in other metal-catalyzed transformations, including olefin polymerization processes,³⁵ and to utilize this approach to facilitate new reaction development under mild conditions.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures and compound characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail for A.C.B.: alex.bissember@utas.edu.au

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We gratefully acknowledge the University of Tasmania (UTAS) and the Australian Research Council (grant number DP180101332) for financial support, UTAS Central Science Laboratory for providing access to NMR spectroscopy services, and Dr. Mairi F. Haddow from Heriot-Watt University for assistance with X-ray crystallography. Part of this research was undertaken on the MX2 beamline at the Australian Synchrotron, Victoria, Australia.

REFERENCES

- (1) See for example: *Metal-Catalyzed Cross-Coupling Reactions and More*, 3rd Ed.; de Meijere, A., Braš, S., Oestreich, M., Eds.; Wiley-VCH: Weinheim, Germany, 2014.
- (2) For selected reviews on applications of alkylphosphines in Pd-catalyzed C–C cross-couplings, see: (a) Rossi, R.; Bellina, F.; Lessi, M.; Manzini, C.; Marianetti, G.; Perego, L. A. Recent Applications of Phosphane-based Palladium Catalysts in Suzuki–Miyaura Reactions Involved in Total Syntheses of Natural Products *Curr. Org. Chem.* **2015**, *19*, 1302–1409. (b) Li, H.; Johansson, Seechurn, C. C. C.; Colacot, T. J. Development of Preformed Pd Catalysts for Cross-Coupling Reactions, Beyond the 2010 Nobel Prize. *ACS Catal.* **2012**, *2*, 1147–1164. (c) Fu, G. C. The Development of Versatile Methods for Palladium-Catalyzed Coupling Reactions of Aryl Electrophiles through the Use of $P(t\text{-Bu})_3$ and PCy_3 as Ligands. *Acc. Chem. Res.* **2008**, *41*, 1555–1564. (d) Fleckenstein, C. A.; Pleino, H. Sterically demanding trialkylphosphines for palladium-catalyzed cross coupling reactions—alternatives to $Pt\text{Bus}$. *Chem. Soc. Rev.* **2010**, *39*, 694–711.
- (3) See, for example: (a) Littke, A. F.; Fu, G. C. A Versatile Catalyst for Heck Reactions of Aryl Chlorides and Aryl Bromides under Mild Conditions. *J. Am. Chem. Soc.* **2001**, *123*, 6989–7000. (b) Littke, A. F.; Schwarz, L.; Fu, G. C. $Pd/P(t\text{-Bu})_3$: A Mild and General Catalyst for Stille Reactions of Aryl Chlorides and Aryl Bromides. *J. Am. Chem. Soc.* **2003**, *124*, 6343–6348. (c) Littke, A. F.; Dai, C.; Fu, G. C. Versatile Catalysts for the Suzuki Cross-Coupling of Arylboronic Acids with Aryl and Vinyl Halides and Triflates under Mild Conditions. *J. Am. Chem. Soc.* **2000**, *122*, 4020–4028. (d) Beller, M.; Riermeier, T. H. Phosphapalladacycle-Catalyzed Heck Reactions for Efficient Synthesis of Trisubstituted Olefins: Evidence for Palladium(0) Intermediates. *Eur. J. Inorg. Chem.* **1998**, 29–35. (e) van Strijdonck, G. P. F.; Boele, M. D. K.; Kamer, P. C. J.; de Vries, J. G.; van Leeuwen, P. W. N. M. Fast Palladium Catalyzed Arylation of Alkenes Using Bulky Monodentate Phosphorus Ligands. *Eur. J. Inorg. Chem.* **1999**, 1073–1076. (f) Hartwig, J. F.; Paul, F. Oxidative Addition of Aryl Bromide after Dissociation of Phosphine from a Two-Coordinate Palladium(0) Complex, Bis(tri-*o*-tolylphosphine)Palladium(0). *J. Am. Chem. Soc.* **1995**, *117*, 5373–5374.
- (4) For a study of reductive elimination of aryl halides from $P(t\text{-Bu})_3$ -ligated Pd complexes, see: Roy, A. H.; Hartwig, J. F. Reductive Elimination of Aryl Halides from Palladium(II). *J. Am. Chem. Soc.* **2001**, *123*, 1232–1233.
- (5) Generating the active catalyst in situ from various Pd(0)- or Pd(II)-precatalysts and $[HP(t\text{-Bu})_3]BF_4$ allows for easier handling as free $P(t\text{-Bu})_3$ is susceptible to oxidation by oxygen (ref. 2c). $[Pd\{P(t\text{-Bu})_3\}_2]$ is an air-stable solid and can be handled in air (refs. 2c and 3c).
- (6) For selected reviews on homogeneous heterobimetallic catalysis featuring palladium, see: (a) Hirner, J. J.; Shi, Y.; Blum, S. A. Organogold Reactivity with Palladium, Nickel, and Rhodium: Transmetalation, Cross-Coupling, and Dual Catalysis. *Acc. Chem. Res.* **2011**, *44*, 603–613. (b) Pérez-Temprano, M. H.; Casares, J. A.; Espinet, P. Bimetallic Catalysis using Transition and Group 11 Metals: An Emerging Tool for C–C Coupling and Other Reactions. *Chem. Eur. J.* **2012**, *18*, 1864–1884. (c) Powers, D. C.; Ritter, T. Bimetallic Redox Synergy in Oxidative Palladium Catalysis. *Acc. Chem. Res.* **2012**, *45*, 840–850. (d) Bay, K. L.; Yang, Y.-F.; Houk, K. N. Multiple roles of silver salts in palladium-catalyzed C–H activations. *J. Organomet. Chem.* **2009**, *864*, 19–25.
- (7) For studies describing Au(I) transmetalating species in cross-couplings see, for example: (a) Shi, Y.; Ramgren, S. D.; Blum, S. A. Palladium-Catalyzed Carboauration of Alkynes and Palladium/Gold Cross-Coupling. *Organometallics* **2009**, *28*, 1275–1277. (b) Shi, Y.; Roth, K. E.; Ramgren, S. D.; Blum, S. A. Catalyzed Catalysis Using Carbophilic Lewis Acidic Gold and Lewis Basic Palladium: Synthesis of Substituted Butenolides and Isocoumarins. *J. Am. Chem. Soc.* **2009**, *131*, 18022–18023. (c) Al-Amin, M.; J. S. Johnson, J. S.; Blum, S. A. Selectivity, Compatibility, Downstream Functionalization, and Silver Effect in the Gold and Palladium Dual-Catalytic Synthesis of Lactones. *Organometallics* **2014**, *33*, 5448–5456. (d) Al-Amin, M.; Roth, K. E.; Blum, S. A. Mechanistic Studies of Gold and Palladium Cooperative Dual-Catalytic Cross-Coupling Systems. *ACS Catal.* **2014**, *4*, 622–629. (e) García-Domínguez, P.; Nevado, C. Au–Pd Bimetallic Catalysis: The Importance of Anionic Ligands in Catalyst Speciation. *J. Am. Chem. Soc.* **2016**, *138*, 3266–3269.
- (8) delPozo, J.; Carrasco, D.; Pérez-Temprano III, M. H.; García-Melchor, M.; Álvarez, R.; Casares, J. A.; Espinet, P. Stille Coupling Involving Bulky Groups Feasible with Gold Cocatalyst. *Angew. Chem. Int. Ed.* **2013**, *52*, 2189–2193.
- (9) Shi, Y.; Peterson, S. M.; Haberaecker III, W. W.; Blum, S. A. Alkynes as Stille Reaction Pseudohalides: Gold- and Palladium-Catalyzed Synthesis of Tri- and Tetra-Substituted Olefins. *J. Am. Chem. Soc.* **2008**, *130*, 2168–2196.
- (10) Ariafard, A.; Rajabi, N. A.; Atashgah, M. J.; Canty, A. J.; Yates, B. F. Computational Study of Carbostannylation Implicating Bimetallic Catalysis Involving “AuI–Vinyl–PdIII” Species. *ACS Catal.* **2014**, *4*, 860–869.
- (11) Khaledifard, Y.; Nsiri, B.; Javidy, S. A.; Sereshk, A. V.; Yates, B. F.; Ariafard, A. Phosphine-Scavenging Role of Gold(I) Complexes from $Pd(P\text{Bu}_3)_2$ in the Bimetallic Catalysis of Carbostannylation of Alkynes *Organometallics* **2017**, *36*, 2014–2019.
- (12) Diversi, P.; Cuzzola, A.; Ghiotto, F. Chemistry of Polynuclear Cationic Gold(I) Thiolates of Formula $[Au_2(S\text{tBu})(L_2)][BF_4]$. *Eur. J. Inorg. Chem.* **2009**, *4*, 545–553.
- (13) A black powder, presumed to be Pd black, precipitated after ~0.5 h.
- (14) CCDC 1875862 (**3**) contains the supplementary crystallographic data. These data are available free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (15) Stambuli, J. P.; Incarvito, C. D.; Buhl, M.; Hartwig, J. F. Synthesis, Structure, Theoretical Studies, and Ligand Exchange Reactions of Monomeric, T-Shaped Arylpalladium(II) Halide Complexes with an Additional, Weak Agostic Interaction. *J. Am. Chem. Soc.* **2004**, *126*, 1184–1194.
- (16) For studies demonstrating the effects of free PPh_3 on reducing the efficiency of transmetalation in Pd-catalyzed Suzuki–Miyaura cross-couplings see: Ho, C. C.; Olding, A.; Smith, J. A.; Bissember, A. C. Nuances in Fundamental Suzuki–Miyaura Cross-Couplings Employing $[Pd(PPh_3)_4]$: Poor Reactivity of Aryl Iodides at Lower Temperatures. *Organometallics*, **2018**, *37*, 1745–1750 and references cited therein.
- (17) For studies demonstrating the effects of free PPh_3 on reducing the efficiency of transmetalation in Pd-catalyzed Stille cross-couplings see, for example: (a) Farina, V.; Krishnan, B. Large rate accelerations in the stille reaction with tri-2-furylphosphine and triphenylarsine as palladium ligands: mechanistic and synthetic implications. *J. Am. Chem. Soc.* **1991**, *113*, 9585–9595 and references cited therein. (b) Liebeskind, L. S.; Fengi, R. W. 3-Stannylcyclobutenediones as nucleophilic cyclobutenedione equivalents. Synthesis of substituted cyclobutenediones and cyclobutenedione monoacetals and the beneficial effect of catalytic copper iodide on the Stille reaction. *J. Org. Chem.* **1990**, *55*, 5359–5364. (c) Casado, A. L.; Espinet, P. Quantitative Evaluation of the Factors Contributing to the “Copper Effect” in the Stille Reaction. *Organometallics* **2003**, *22*, 1305–1309.
- (18) For selected studies concerning the enhanced efficiency of Pd-catalyzed Stille couplings employing copper cocatalysts see: refs. 17b,c and (a) Farina, V.; Kapadia, S.; Krishnan, B.; Wang, C.; Liebeskind, L. S. On the Nature of the “Copper Effect” in the Stille Cross-Coupling. *J. Org. Chem.* **1994**, *59*, 5905–5911. (b) Han, X.; Stolz, B. M.; Corey, E. J. Cuprous Chloride Accelerated Stille Reactions. A General and Effective Coupling System for Sterically Congested Substrates and for Enantioselective Synthesis. *J. Am. Chem. Soc.* **1999**, *121*, 7600–7605. (c) Mee, S. P. H.; Lee, V.; Baldwin, J. E. Stille Coupling Made Easier—The Synergic Effect of Copper(I) Salts and the Fluoride Ion. *Angew. Chem. Int. Ed.* **2004**, *43*, 1132–1136. (d) Mee, S. P. H.; Lee, V.; Baldwin, J. E. Significant Enhancement of the Stille Reaction with a New Combination of Reagents—

- Copper(I) Iodide with Cesium Fluoride. *Chem. Eur. J.* **2005**, *11*, 3294–3308.
- (19) For selected studies concerning Pd-catalyzed Suzuki–Miyaura couplings employing copper cocatalysts see: (a) Boland, G. M.; Donnelly, D. M. X.; Finet, P.-J.; Rea, M. D. Synthesis of neoflavones by Suzuki arylation of 4-substituted coumarins. *J. Chem. Soc. Perkin Trans. 1* **1996**, 2591–2597. (b) Savarin, C.; Liebeskind, L. S. Nonbasic, Room Temperature, Palladium-Catalyzed Coupling of Aryl and Alkenyl Iodides with Boronic Acids Mediated by Copper(I) Thiophene-2-carboxylate (CuTC). *Org. Lett.* **2001**, *3*, 2149–2152. (c) Hodgson, P. B.; Salingue, F. H. The preparation of a stable 2-pyridylboronate and its reactivity in the Suzuki–Miyaura cross-coupling reaction. *Tetrahedron Lett.* **2004**, *45*, 685–687; (d) Yamamoto, Y.; Takizawa, M.; Yu, X.; Miyaura, N. Cyclic Triolborates: Air- and Water-Stable Ate Complexes of Organoboronic Acids. *Angew. Chem. Int. Ed.* **2008**, *47*, 928–931. (e) Deng, J. Z.; Paone, D. V.; Ginnetti, A. T.; Kurihara, H.; Dreher, S. D.; Weissman, S. A.; Stauffer, S. R.; Burgey, C. S. Copper-Facilitated Suzuki Reactions: Application to 2-Heterocyclic Boronates. *Org. Lett.* **2009**, *11*, 345–347. (f) Grob, J. E.; Nunez, J.; Dechantsreiter, M. A.; Hamann, L. G. Regioselective Synthesis and Slow-Release Suzuki–Miyaura Cross-Coupling of MIDA Boronate-Functionalized Isoxazoles and Triazoles. *J. Org. Chem.* **2011**, *76*, 10241–10248. (g) Knapp, D. M.; Gillis, E. P.; Burke, M. D. A General Solution for Unstable Boronic Acids: Slow-Release Cross-Coupling from Air-Stable MIDA Boronates. *J. Am. Chem. Soc.* **2009**, *131*, 6961–6963.
- (20) Peng, Y.; Li, W.-D. Z. *cine* Substitution and the Cu Effect in Stille Cross-Coupling Reactions: Mechanistic Perspectives and Synthetic Utility. *Eur. J. Org. Chem.* **2010**, 6703–6718.
- (21) (a) Liebeskind, L. S.; Wand, J. Synthesis of substituted cyclobutenediones by the palladium catalyzed cross-coupling of halocyclobutenediones with organostannanes. *Tetrahedron Lett.* **1990**, *31*, 4293–4296. (b) Liebeskind, L. S.; Yu, M. S.; Yu, R. H.; Wang, J.; Hagen, K. S. 4,4'-Bi(cyclobutene-1,2-diones): bisquaryls. *J. Am. Chem. Soc.* **1993**, *115*, 9048–9055. (c) Liebeskind, L. S.; Riesinger, S. W. Substituted quinone synthesis by palladium-copper cocatalyzed cross-coupling of stannylquinones with aryl and heteroaryl iodides. *J. Org. Chem.* **1993**, *58*, 408–413. (d) Allred, G. D.; Liebeskind, L. S. Copper-Mediated Cross-Coupling of Organostannanes with Organic Iodides at or below Room Temperature. *J. Am. Chem. Soc.* **1996**, *118*, 2748–2749.
- (22) Farina, V. New perspectives in the cross-coupling reactions of organostannanes. *Pure Appl. Chem.* **1996**, *68*, 73–78.
- (23) Other additives have been reported to remove $P(t\text{-Bu})_3$ in Pd-catalyzed reactions. See for example: Tan, Y.; Hartwig, J. F. Assessment of the Intermediacy of Arylpalladium Carboxylate Complexes in the Direct Arylation of Benzene: Evidence for C–H Bond Cleavage by “Ligandless” Species. *J. Am. Chem. Soc.* **2011**, *133*, 3308–3311.
- (24) Hobbs Jr, F. W. Palladium-catalyzed synthesis of alkynylamino nucleosides. A universal linker for nucleic acids. *J. Org. Chem.* **1989**, *54*, 3420–3422.
- (25) CCDC 1875861 (**9**) contains the supplementary crystallographic data. These data are available free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (26) Essentially no cross-coupled product was formed in reactions featuring either Au complex **1** or Cu complex **8** in the absence of Pd complex **2**.
- (27) The use of Cu complex **8** as an additive was not as effective as Au complex **1**. Experiments were also performed using CuI as an additive, which was also not as effective as Au complex **1** (see Supporting Information).
- (28) The possibility that boron/ gold transmetalation mechanisms may also be operative under these conditions cannot be excluded.
- (29) Each data point represents the average of two experiments with yields determined via GC with the aid of a calibrated internal standard.
- (30) Pullen, R.; Olding, A.; Smith, J. A.; Bissember, A. C. Capstone Laboratory Experiment Investigating Key Features of Palladium-Catalyzed Suzuki–Miyaura Cross-Coupling Reactions. *J. Chem. Educ.* **2018**, *95*, 2081–2085.
- (31) The results of the study reported in ref. 16 were consistent with transmetalation representing the rate-limiting step of the reaction.
- (32) It is unclear whether $[\text{Au}(\text{PPh}_3)(\text{NTf}_2)]$ (**5**) has the capacity to scavenge PPh_3 directly from $[\text{Pd}(\text{PPh}_3)_2(p\text{-tol})(\text{I})]$. For example, a number of new signals were observed when Au complex **5** was added to a THF- d_8 solution of $\text{trans-}[\text{Pd}(\text{PPh}_3)_2(p\text{-tol})(\text{I})]$, as judged by ^{31}P NMR spectroscopy. Espinet and coworkers suggest that CuI binds free PPh_3 in the reaction mixture rather than scavenging PPh_3 directly from $[\text{Pd}(\text{PPh}_3)_2(\text{aryl})(\text{I})]$ (ref. 17c).
- (33) Olbrich, F.; Lagow, R. J. A Novel Mononuclear Gold(I) Complex: Synthesis and X-Ray Structure of $[\text{Au}\{\text{P}(\text{C}_6\text{H}_5)_3\}_3][\text{SiF}_5] \cdot 1.5 \text{CH}_2\text{Cl}_2$. *Z. Anorg. Allg. Chem.* **1995**, *621*, 1929–1932.
- (34) The addition of $[\text{Au}(\text{PPh}_3)(\text{NTf}_2)]$ (**5**) to a THF- d_8 solution containing $[\text{Pd}(\text{PPh}_3)_4]$ and Bu_4NI provided similar results, as judged by ^{31}P NMR spectroscopy.
- (35) For early examples of phosphine-scavenging in olefin polymerizations, see: (a) Klabunde, U.; Itten, S. D. Nickel catalysis for ethylene homo- and co-polymerization. *J. Mol. Catal.* **1987**, *41*, 123–134. (b) Wang, C.; Friedrich, S.; Younkin, T. R.; Li, R. T.; Grubbs, R. H.; Bansleben, D. A.; Day, M. W. Neutral Nickel(II)-Based Catalysts for Ethylene Polymerization. *Organometallics* **1998**, *17*, 3149–3151.