

1992

Semi Williamson type matrices and the $W(2n, n)$ conjecture

Jennifer Seberry

University of Wollongong, jennie@uow.edu.au

Xian-Mo Zhang

University of Wollongong, xianmo@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/infopapers>



Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Seberry, Jennifer and Zhang, Xian-Mo: Semi Williamson type matrices and the $W(2n, n)$ conjecture 1992.
<https://ro.uow.edu.au/infopapers/1062>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Semi Williamson type matrices and the $W(2n, n)$ conjecture

Abstract

Four $(1, -1, 0)$ -matrices of order m , $X = (X_{ij})$, $Y = (Y_{ij})$, $Z = (Z_{ij})$, $U = (U_{ij})$ satisfying

(i) $XX^T + YY^T + ZZ^T + UU^T = 2mI_m$,

(ii) $x_{ij}^2 + y_{ij}^2 + z_{ij}^2 + u_{ij}^2 = 2$, $i, j = 1, \dots, m$,

(iii) X, Y, Z, U mutually amicable,

will be called semi Williamson type matrices of order m . In this paper we prove that if there exist Williamson type matrices of order n_1, \dots, n_k . then there exist semi Williamson type matrices of order $N = \prod_{j=1}^k n_j^{r_j}$, where r_j are non-negative integers. As an application, we obtain a $W(4N, 2N)$.

Although the paper presents no new $W(4n, 2n)$ for n , odd, $n < 3000$, it is a step towards proving the conjecture that there exists a $W(4n, 2n)$ for any positive integer n . This conjecture is a sub-conjecture of the Seberry conjecture [4, page 92] that $W(4n, k)$ exist for all $k = 0, 1, \dots, 4n$. In addition we find infinitely many new $W(2n, n)$, n odd and the sum of two squares.

Disciplines

Physical Sciences and Mathematics

Publication Details

Jennifer Seberry and Xian-Mo Zhang, Semi Williamson type matrices and the $W(2n, n)$ conjecture, Journal of Combinatorial Mathematics and Combinatorial Computing, 11, (1992), 65-71.

Semi Williamson Type Matrices and the $W(2n, n)$ Conjecture

Jennifer Seberry
and
Xian-Mo Zhang

Department of Computer Science
University College
University of New South Wales
Australian Defence Force Academy
Canberra, ACT 2600, AUSTRALIA

Abstract

Four $(1, -1, 0)$ -matrices of order m , $X = (x_{ij})$, $Y = (y_{ij})$, $Z = (z_{ij})$, $U = (u_{ij})$ satisfying

- (i) $XX^T + YY^T + ZZ^T + UU^T = 2mI_m$,
- (ii) $x_{ij}^2 + y_{ij}^2 + z_{ij}^2 + u_{ij}^2 = 2$, $i, j = 1, \dots, m$,
- (iii) X, Y, Z, U mutually amicable,

will be called semi Williamson type matrices of order m . In this paper we prove that if there exist Williamson type matrices of order n_1, \dots, n_k then there exist semi Williamson type matrices of order $N = \prod_{j=1}^k n_j^{r_j}$, where r_j are non-negative integers. As an application, we obtain a $W(4N, 2N)$.

Although the paper presents no new $W(4n, 2n)$ for n , odd, $n < 3000$, it is a step towards proving the conjecture that there exists a $W(4n, 2n)$ for any positive integer n . This conjecture is a sub-conjecture of the Seberry conjecture [4, page 92] that $W(4n, k)$ exist for all $k = 0, 1, \dots, 4n$. In addition we find infinitely many new $W(2n, n)$, n odd and the sum of two squares.

1 Introduction and Basic Definitions

Definition 1 Let A, B, C, D be four $(1, -1)$ -matrices of order n . If $AA^T + BB^T + CC^T + DD^T = 4nI_n$ and $UV^T = VU^T$ (U and V are amicable), where $U, V \in \{A, B, C, D\}$. We call A, B, C, D *Williamson type matrices* of order n .

Definition 2 Let W be a $(1, -1, 0)$ -matrix of order of order n satisfying $WW^T = cI_n$. We call W a *weighing matrix* of order n with weight c , denoted by $W(n, c)$.

Definition 3 Four $(1, -1, 0)$ -matrices of order m , $X = (x_{ij})$, $Y = (y_{ij})$, $Z = (z_{ij})$, $U = (u_{ij})$ satisfying

- (i) $XX^T + YY^T + ZZ^T + UU^T = 2mI_m$,
- (ii) $x_{ij}^2 + y_{ij}^2 + z_{ij}^2 + u_{ij}^2 = 2$, $i, j = 1, \dots, m$,
- (iii) X, Y, Z, U mutually amicable,

will be called *semi Williamson type matrices* of order m . In particular, if X, Y, Z, U are circulant and symmetric we call X, Y, Z, U *semi Williamson matrices* of order m .

Let $M = (M_{ij})$ and $N = (N_{gh})$ be orthogonal matrices with t^2 block M-structure (see [6]) of order tm and tn respectively, where M_{ij} is of order m ($i, j = 1, \dots, t$) and N_{gh} is of order n ($g, h = 1, 2, \dots, t$). We now define the operation \circ as the following:

$$M \circ N = \begin{bmatrix} L_{11} & L_{12} & \cdots & L_{1t} \\ L_{21} & L_{22} & \cdots & L_{2t} \\ \cdots & \cdots & \cdots & \cdots \\ L_{t1} & L_{t2} & \cdots & L_{tt} \end{bmatrix}$$

where M_{ij} , N_{ij} and L_{ij} are of order of m, n and mn , respectively and

$$L_{ij} = M_{i1} \times N_{1j} + M_{i2} \times N_{2j} + \cdots + M_{it} \times N_{tj},$$

where \times is Kronecker product, $i, j = 1, 2, \dots, t$. We call this the *strong Kronecker multiplication* of two matrices (see [7]).

Lemma 1 Let $A = [A_1, A_2, A_3, A_4]$ be a $(1, -1, 0)$ -matrix of order $m \times 4m$, where A_j is of order m , satisfying $\sum_{j=1}^4 A_j A_j^T = pI_m$ and $B^T = [B_1^T, B_2^T, B_3^T, B_4^T]$, where B_i is of order $n \times 4n$, be a $W(4n, q)$. Set $C = \sum_{j=1}^4 A_j \times B_j$. Then $CC^T = pqI_{mn}$.

Proof. $CC^T = (\sum_{j=1}^4 A_j \times B_j)(\sum_{j=1}^4 A_j^T \times B_j^T) = \sum_{j=1}^4 A_j A_j^T \times B_j B_j^T = (\sum_{j=1}^4 A_j A_j^T) \times qI_n = pI_m \times qI_n = pqI_{mn}$.

Notation 1 Write $OD(A, B, C, D) = \begin{bmatrix} A & B & C & D \\ D & C & -B & -A \\ B & -A & D & -C \\ C & -D & -A & B \end{bmatrix}$.

□

2 Preliminaries

Lemma 2 Let $a, b, c, d \in \{1, -1, 0\}$, $a^2 + b^2 + c^2 + d^2 = 2$ and $k, m, l, q \in \{1, -1\}$. Set $[x, y, z, u] = \frac{1}{2}[a, b, c, d]OD(k, m, l, q)$. Then $x, y, z, u \in \{1, -1, 0\}$, $x^2 + y^2 + z^2 + u^2 = 2$.

Proof. By Lemma 1, $x^2 + y^2 + z^2 + u^2 = \frac{1}{4} \cdot 2 \cdot 4 = 2$. Each of x, y, z, u is half the sum of four numbers, two of which are zero, and the other two of which are units. It follows that $x, y, z, u \in \{1, -1, 0\}$.

We note that the operation of Lemma 2 is norm preserving.

Lemma 3 If there exist Williamson type matrices of order m then there exist semi Williamson type matrices of order m .

Proof. Let A, B, C, D be the Williamson type matrices of order m then $\frac{1}{2}(A + B)$, $\frac{1}{2}(A - B)$, $\frac{1}{2}(C + D)$, $\frac{1}{2}(C - D)$ are semi Williamson type matrices. \square

Lemma 4 If there exist semi Williamson type matrices of order m and Williamson type matrices of order n then there exist semi Williamson type matrices of order mn .

Proof. Let $X = (x_{ij})$, $Y = (y_{ij})$, $Z = (z_{ij})$, $U = (u_{ij})$ be the semi Williamson type matrices of order m and $K = (k_{st})$, $L = (l_{st})$, $M = (m_{st})$, $Q = (q_{st})$ be the Williamson type matrices of order n . We now construct four matrices, say $R = (r_{\mu\nu})$, $S = (s_{\mu\nu})$, $V = (v_{\mu\nu})$, $W = (w_{\mu\nu})$, $i, j = 1, \dots, mn$, of order mn , where

$$[r_{\mu\nu}, s_{\mu\nu}, v_{\mu\nu}, w_{\mu\nu}] = \frac{1}{2}[x_{ij}, y_{ij}, z_{ij}, u_{ij}]OD(k_{st}, m_{st}, q_{st}, l_{st}).$$

By Lemma 2, $r_{\mu\nu}, s_{\mu\nu}, v_{\mu\nu}, w_{\mu\nu} \in \{1, -1, 0\}$ and $r_{\mu\nu}^2 + s_{\mu\nu}^2 + v_{\mu\nu}^2 + w_{\mu\nu}^2 = 2$, $\mu, \nu = 1, \dots, mn$. By Lemma 1, $RR^T + SS^T + VV^T + WW^T = \frac{1}{4}8mnI_{mn} = 2mnI_{mn}$. Since X, Y, Z, U are mutually amicable and K, L, M, Q are mutually amicable, R, S, V, W are also mutually amicable. \square

3 Main Results

Throughout this section we write $N = \prod_{j=1}^k n_j^{r_j}$, where r_j are non-negative integers.

Theorem 1 If there exist Williamson type matrices of order n_1, \dots, n_k then there exist semi Williamson type matrices of order N .

Proof. By Lemma 3, there exist semi Williamson type matrices of order n_1 . By Lemma 4, there exist semi Williamson type matrices of order $n_1 n_2$. Using Lemma 4 repeatedly, we prove the Theorem. \square

Corollary 1 *If there exist Williamson type matrices of order n_1, \dots, n_k then there exists a $W(4N, 2N)$.*

Proof. By Theorem 1, there exist semi Williamson type matrices of order N , say E, F, G, H . Then $A = OD(E, F, G, F)$ is a $W(4N, 2N)$. \square

Corollary 2 *If there exist Williamson type matrices of order n_1, \dots, n_k and an Hadamard matrix of order $4h$ then there exists a $W(4Nh, 2Nh)$.*

Proof. By Theorem 1, there exist semi Williamson type matrices of order N , say P, Q, R, S . Write $H = (H_{ij})$, $i, j = 1, 2, 3, 4$ for the Hadamard matrix of order $4h$, where H_{ij} is of order h . Set

$$B = \frac{1}{2}OD(P, Q, R, S) \circ (H_{ij}).$$

From (ii) of Definition 3, B is a $(1, -1, 0)$ -matrix of order $4Nh$. By Theorem 1, [7],

$$BB^T = 2NhI_{4Nh}.$$

Hence B is the required $W(4Nh, 2Nh)$. \square

4 Numerical Results

To construct $W(4n, 2n)$ we can use the known result that if there exist Hadamard matrices of order $4h_1$ and $4h_2$ then there exist two amicable and disjoint $W(4h_1h_2, 2h_1h_2)$ (see [7], [3]). Thus we obtain many $W(4n, 2n)$ whenever $n = h_1h_2$, where $4h_1$ and $4h_2$ are the order of Hadamard matrices. In particular, let $h_2 = 1$, we give the simple result that $W(4h, 2h)$ exists whenever an Hadamard matrix of order $4h$ exists (see [7], [5]). However Corollary 1 is new result. To show the advantages of Corollary 1 and Corollary 2, we now give new $W(4n, 2n)$. Let $a = 71 \cdot 79 \cdot 97$, $b = 71 \cdot 79$, $c = 71 \cdot 97$, $d = 79 \cdot 97$. Note Hadamard matrices of order $4b, 4c, 4d$ and $4a$ are not yet known and hence the method in [7] and [3] cannot be used. Since there exist Williamson type matrices of order 79, 97 and an Hadamard matrix of order 71, by Corollary 2, there exists a $W(4a, 2a)$. Similarly, we obtain new $W(4n, 2n)$, which cannot be obtained by using the method given in [7] or [3], for $n = 73 \cdot 83 \cdot 89$ and $83 \cdot 89 \cdot 103$. Also Corollary 1 and Corollary 2 give infinitely new $W(4h, 2h)$ directly for example $h = 5^j$ or $3^i 5^j 7^k$, where i, j, k are non-negative integers.

Corollary 1 has many uses. First, this is a step towards proving the conjecture that there exists a $W(4n, 2n)$ for any positive integer n . This conjecture is a sub-conjecture of the Seberry conjecture [4, page 92] that $W(4n, k)$ exist for all $k = 0, 1, \dots, 4n$. In addition we find infinitely many new $W(2n, n)$, n odd and the sum of two squares. It is interesting that unlike the product of Hadamard matrices (see [1], [3]), where the number of 2-factors will increase when the number of Hadamard matrices used to form

the product increases, the factor 4 in the order $4N = 4 \prod_{j=1}^k n_j^{r_j}$ of $W(4N, 2N)$ will be invariant no matter how large k and r_j become.

Furthermore, let W_1 be the $W(4N, 2N)$ for $N = \prod_{j=1}^k n_j^{r_j}$, where r_j are non-negative integers, mentioned in Corollary 1. Suppose we have another $W(4N, 2N)$, say W_2 , disjoint with W_1 . Using Craigen's [2] orthogonal pairs, we would obtain a powerful result: there exists an Hadamard matrix of order hN whenever there exists an Hadamard matrix of order h . In particular there exists an Hadamard matrix of order $8N$, $N = \prod_{j=1}^k n_j^{r_j}$,

where r_j are non-negative integers. $H = W_1 \times \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + W_2 \times \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$ is the

required Hadamard matrix.

The state of the $W(4n, 2n)$ conjecture, for small n , can be summarized by noting that a $W(2^t q, 2^{t-1} q)$ exists for q , odd, $q < 3000$ for precisely those q and t for which an Hadamard matrix exists in the Appendix of Seberry and Yamada [5].

The conjecture that a $W(2n, n)$ for every odd n where n is the sum of two squares has previously been resolved in the affirmative for $n = 5, 9, 13$ and 17 (see [4]).

Lemma 5 Let A_1, A_2, A_3, A_4 be type 1 $(1, -1)$ -matrices of order n satisfying

$$\sum_{i=1}^4 A_i A_i^T = 4n I_n \quad (1)$$

and

$$A_1 A_2^T + A_2 A_1^T + A_3 A_4^T + A_4 A_3^T = 0. \quad (2)$$

Then there exists a $W(2n, n)$.

Proof. Set $W = \frac{1}{2} \begin{bmatrix} A_1 + A_2 & A_3 + A_4 \\ A_3^T + A_4^T & -A_1^T - A_2^T \end{bmatrix}$ is a $W(2n, n)$. Then W is a $W(2n, n)$.
□

We note that if n is odd in Lemma 5 then by Corollary 2.11 [4] n is the sum of two squares. We call four $(1, -1)$ type 1 matrices that satisfy (1) and (2) *tight Williamson-like matrices*.

Corollary 3 Let $M = \prod_{j=1}^k p_j^{4r_j}$, where $p_j \equiv 3 \pmod{4}$, a prime and r_j is a non-negative integer, $j = 1, \dots, k$. Then there exists a $W(2n, n)$, where $n = 5 \cdot 9^t M$, $13 \cdot 9^t M$, $25 \cdot 9^t M$.

Proof. By Theorem 2, [10], there exist four type 1 $(1, -1)$ -matrices of order $5 \cdot 9^t$, $13 \cdot 9^t$, $25 \cdot 9^t$, satisfying (1) and (2). From [8], There exist four symmetric, mutually commutative type 1 $(1, -1)$ -matrices of order M , say B_1, B_2, B_3, B_4 , satisfying $\sum_{i=1}^4 B_i B_i^T = 4n I_n$, $B_1 B_2^T + B_3 B_4^T = 0$, $B_1 B_3^T + B_2 B_4^T = 0$, $B_1 B_4^T + B_2 B_3^T = 0$. By Theorem 1, [10], there exist four type 1 matrices of order $5 \cdot 9^t M$, $13 \cdot 9^t M$, $25 \cdot 9^t M$, satisfying (1) and (2). By Lemma 5, we have a $W(2n, n)$, where $n = 5 \cdot 9^t M$, $13 \cdot 9^t M$, $25 \cdot 9^t M$.
□

We now give tight Williamson-like matrices of order 5, 13 and 25. By the method given by Xia [9], we construct cyclic (1, -1) tight Williamson-like matrices of order 5 and 13 with first rows

+ - + + -, + + - + +, - - + + -, + + + + - and

+ + - - - + - - + - + +, - - + + + - + + + + - +,
+ - - + - + + + - - - + -, + - + + + + + - - + + - respectively.

From [9] we also construct type 1 tight Williamson-like matrices of order 25. Any element in the abelian group $Z_5 \oplus Z_5$ can be expressed as (a, b) , where $a, b \in Z_5$, and the additive addition in $Z_5 \oplus Z_5$ can be defined as $(a, b) + (c, d) = (a + b, c + d)$. Set

$$\begin{aligned} S_1 &= \{(0,0), (0,1), (1,2), (3,3), (0,3), (4,4), (3,4), (2,0), (2,2), (1,0), (1,4), (0,2), (3,0)\}, \\ S_2 &= \{(0,1), (4,0), (3,1), (4,4), (0,4), (4,2), (1,0), (1,1), (3,2)\}, \\ S_3 &= \{(1,2), (3,3), (1,3), (4,1), (3,4), (2,0), (2,3), (4,3), (1,4), (0,2), (2,4), (2,1)\}, \\ S_4 &= \{(3,3), (4,1), (0,3), (2,0), (4,3), (2,2), (0,2), (2,1), (3,0)\}. \end{aligned}$$

Hence the type 1 (1, -1) incidence matrices of S_1, S_2, S_3, S_4 form the tight Williamson-like matrices of order 25.

Finally we note that if $N + I$ is a symmetric conference matrix of order $n \equiv 2 \pmod{4}$ then $N + I, N - I, N + I, -N + I$ are tight Williamson-like matrices of order n .

References

- [1] S. S. Agayan. *Hadamard Matrices and Their Applications*, volume 1168 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-Heidelberg- New York, 1985.
- [2] R. Craigen. Constructing Hadamard matrices with orthogonal pairs. to appear, 1990.
- [3] R. Craigen, Jennifer Seberry, and Xian-Mo Zhang. Product of four Hadamard matrices. to appear in *Journal of Combinatorial Theory*, Ser. A, 1990.
- [4] A. V. Geramita and J. Seberry. *Orthogonal Designs: Quadratic Forms and Hadamard Matrices*. Marcel Dekker, New York-Basel, 1979.
- [5] Jennifer Seberry and Mieko Yamada. Hadamard matrices, sequences and block designs. In J. Dinitz and D. Stinson, editors, *Surveys in Contemporary Design Theory*, Wiley-Interscience Series in Discrete Mathematics. John Wiley, New York.
- [6] Jennifer Seberry and Mieko Yamada. On the products of Hadamard matrices, Williamson matrices and other orthogonal matrices using M-structures. *JCMCC*, 7:97-137, 1990.
- [7] Jennifer Seberry and Xian-Mo Zhang. Some orthogonal designs and complex Hadamard matrices by using two Hadamard matrices. to appear in *Austral. J. Comb.*, 1990.

- [8] Ming Yuan Xia. Some infinite classes of special Williamson matrices and difference sets. to appear in *Journal of Combinatorial Theory*, Ser. A.
- [9] Ming Yuan Xia. Supplementary difference sets and Hadamard matrices. *Acta Math. Sci.*, 4 (1):81-92 (in Chinese), 1984.
- [10] Ming Yuan Xia. Hadamard matrices. *Combinatorial Designs and Applications, Lecture Notes in Pure and Applied Mathematics*, Dekker, New York, 126:179-181, 1990.