



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Engineering and Information Sciences -
Papers: Part A

Faculty of Engineering and Information Sciences

2013

KMS states on the C^* -algebras of finite graphs

Astrid An Huef

University of Otago, astrid@uow.edu.au

Marcelo Laca

University of Victoria

Iain F. Raeburn

University of Otago, raeburn@uow.edu.au

Aidan D. Sims

University of Wollongong, asims@uow.edu.au

Publication Details

An Huef, A., Laca, M., Raeburn, I. F. & Sims, A. D. (2013). KMS states on the C^* -algebras of finite graphs. *Journal of Mathematical Analysis and Applications*, 405 (2), 388-399.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

KMS states on the C^* -algebras of finite graphs

Abstract

We consider a finite directed graph E , and the gauge action on its Toeplitz-Cuntz-Krieger algebra, viewed as an action of \mathbb{R} . For inverse temperatures larger than a critical value β_c , we give an explicit construction of all the KMS_β states. If the graph is strongly connected, then there is a unique KMS_{β_c} state, and this state factors through the quotient map onto $C^*(E)$. Our approach is direct and relatively elementary.

Keywords

algebras, c, states, finite, kms, graphs

Disciplines

Engineering | Science and Technology Studies

Publication Details

An Huef, A., Laca, M., Raeburn, I. F. & Sims, A. D. (2013). KMS states on the C^* -algebras of finite graphs. *Journal of Mathematical Analysis and Applications*, 405 (2), 388-399.

KMS STATES ON THE C^* -ALGEBRAS OF FINITE GRAPHS

ASTRID AN HUEF, MARCELO LACA, IAIN RAEBURN, AND AIDAN SIMS

ABSTRACT. We consider a finite directed graph E , and the gauge action on its Toeplitz-Cuntz-Krieger algebra, viewed as an action of \mathbb{R} . For inverse temperatures larger than a critical value β_c , we give an explicit construction of all the KMS_β states. If the graph is strongly connected, then there is a unique KMS_{β_c} state, and this state factors through the quotient map onto $C^*(E)$. Our approach is direct and relatively elementary.

Fix an integer $n \geq 2$, and consider the action α of \mathbb{R} lifted from the gauge action of \mathbb{T} on the Cuntz algebra \mathcal{O}_n . Olesen and Pedersen [15] showed that (\mathcal{O}_n, α) has a unique KMS state, which occurs at inverse temperature $\ln n$. Enomoto, Fujii and Watatani [5] extended this to the Cuntz-Krieger algebras \mathcal{O}_A . For an irreducible matrix A , they found that the unique KMS state has inverse temperature $\ln \rho(A)$, where $\rho(A)$ is the spectral radius of A , or equivalently the Perron-Frobenius eigenvalue of A .

There are now many generalisations of the Cuntz-Krieger algebras, and much is known about their KMS states. In particular, Exel and Laca [6] have conducted an extensive analysis of the KMS states on the Cuntz-Krieger algebras of infinite $\{0, 1\}$ -matrices. Their analysis is very general: they consider actions arising from embeddings of \mathbb{R} in the infinite torus \mathbb{T}^∞ , and they study also the Toeplitz extension of the Cuntz-Krieger algebra, which has many more KMS states. So it was something of a surprise when Kajiwara and Watatani [8] pinpointed a gap in our knowledge of KMS states for the gauge action on the Cuntz-Krieger algebras of finite graphs: because Exel and Laca assumed that their matrices had no zero rows, their results do not apply to graphs with sources. Kajiwara and Watatani showed that the existence of sources makes a big difference (see Corollary 6.1 below).

Here we describe the KMS states for the gauge action on the Toeplitz algebras of finite graphs. For a graph with vertex matrix A and $\beta > \ln \rho(A)$, our main theorem gives an explicit isomorphism of the simplex of KMS_β states on the Toeplitz algebra onto a simplex of codimension 1 in \mathbb{C}^{E^0} . Our methods are relatively elementary, and follow the general pattern of [12, 13].

After a brief review of background material, we begin in §2 by characterising KMS states on graph algebras in terms of their behaviour on the usual spanning family. In §3 we prove our main theorem about the KMS states on the Toeplitz algebra of a graph. In §4 we consider a strongly connected graph E , and prove that there is a unique $\text{KMS}_{\ln \rho(A)}$ state on $\mathcal{TC}^*(E)$, which factors through the $\text{KMS}_{\ln \rho(A)}$ state of $C^*(E)$ from [5]. In §5, we describe the ground and KMS_∞ states. In §6, we relate our results to those in [6] and [8]. We also discuss how they relate to the powerful machines of Laca-Neshveyev [11]

Date: 10 May 2011.

2010 Mathematics Subject Classification. 46L30, 46L55.

This research has been supported by the University of Otago, the Marsden Fund of the Royal Society of New Zealand, the Natural Sciences and Engineering Research Council of Canada, and the Australian Research Council.

for studying KMS states on Cuntz-Pimsner algebras and of Renault-Neshveyev [19, 14] for groupoid algebras. We finish with an appendix on the possible values of the spectral radius for vertex matrices of finite graphs.

1. BACKGROUND

We use the conventions of [18] for directed graphs E and their C^* -algebras $C^*(E)$. We also borrow the convention from the higher-rank graph literature in which we write, for example, E^*v for $\{\mu \in E^* : s(\mu) = v\}$, and vE^1w for $\{e \in E^1 : r(e) = v, s(e) = w\}$.

Suppose that E is a directed graph. A *Toeplitz-Cuntz-Krieger E -family* (P, S) consists of mutually orthogonal projections $\{P_v : v \in E^0\}$ and partial isometries $\{S_e : e \in E^1\}$ such that $S_e^*S_e = P_{s(e)}$ and

$$(1.1) \quad P_v \geq \sum_{e \in F} S_e S_e^* \quad \text{for every finite subset } F \text{ of } vE^1.$$

The definition used in [7] and [18] requires that the partial isometries $\{S_e : e \in E^1\}$ have mutually orthogonal ranges, but it turns out that this orthogonality follows from the other relations. To see this we need a simple lemma.

Lemma 1.1. *Suppose that P and Q are projections on a Hilbert space H and $\|P+Q\| \leq 1$. Then P and Q have orthogonal ranges.*

Proof. Take $h \in PH$. Then

$$\|h\|^2 \geq \|Ph + Qh\|^2 = (h + Qh | h + Qh) = \|h\|^2 + 3\|Qh\|^2,$$

and $\|Qh\| = 0$. So $Qh = 0$ for all $h \in PH$, and QH is orthogonal to PH . \square

Corollary 1.2. *Suppose that $\{P_v : v \in E^0\}$ are mutually orthogonal projections and $\{S_e : e \in E^1\}$ are partial isometries satisfying (1.1). Then the projections $\{S_e S_e^* : e \in E^1\}$ are mutually orthogonal.*

Proof. Suppose first that $r(e) = r(f) = v$, say. Then (1.1) with $F = \{e, f\}$ shows that $P_v \geq S_e S_e^* + S_f S_f^*$. Since $T \geq S \geq 0$ implies $\|T\| \geq \|S\|$, we have $1 = \|P_v\| \geq \|S_e S_e^* + S_f S_f^*\|$, and Lemma 1.1 implies that $S_e S_e^*$ and $S_f S_f^*$ are mutually orthogonal. On the other hand, if $r(e) \neq r(f)$, then applying (1.1) to singleton sets gives $S_e S_e^* \leq P_{r(e)}$ and $S_f S_f^* \leq P_{r(f)}$, and since $P_{r(e)}$ and $P_{r(f)}$ are mutually orthogonal, so are $S_e S_e^*$ and $S_f S_f^*$. \square

The Toeplitz algebra $\mathcal{TC}^*(E)$ is generated by a universal Toeplitz-Cuntz-Krieger family (p, s) . The existence of $\mathcal{TC}^*(E)$ was established in [7, Theorem 4.1], which says that the Toeplitz algebra $\mathcal{T}(X)$ of the associated graph correspondence X has the required property. Corollary 1.2 implies that $s_e^* s_f = \delta_{e,f} p_{s(e)}$, and then the usual argument for graph algebras (as in [18, Corollary 1.15], for example) gives the product formula

$$(1.2) \quad (s_\mu s_\nu^*)(s_\alpha s_\beta^*) = \begin{cases} s_{\mu\alpha'} s_\beta^* & \text{if } \alpha = \nu\alpha' \\ s_\mu s_{\beta\nu'}^* & \text{if } \nu = \alpha\nu' \\ 0 & \text{otherwise.} \end{cases}$$

From this formula, further standard arguments give

$$\mathcal{TC}^*(E) = \overline{\text{span}}\{s_\mu s_\nu^* : \mu, \nu \in E^*, s(\mu) = s(\nu)\}.$$

The Toeplitz algebra $\mathcal{TC}^*(E)$ carries a gauge action γ of \mathbb{T} satisfying $\gamma_z(s_\mu s_\nu^*) = z^{|\mu|-|\nu|} s_\mu s_\nu^*$, and a dynamics $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ which is lifted from γ via the map $t \mapsto e^{it}$. Since the quotient map q of $\mathcal{TC}^*(E)$ onto $C^*(E)$ is gauge-invariant, we write α also for the corresponding action on the graph algebra $C^*(E)$.

For every $\mu, \nu \in E^*$, the function $t \mapsto \alpha_t(s_\mu s_\nu^*) = e^{it(|\mu|-|\nu|)} s_\mu s_\nu^*$ on \mathbb{R} extends to an analytic function on all of \mathbb{C} . Since these elements span a dense subspace of $\mathcal{TC}^*(E)$, it follows from [17, Proposition 8.12.3] that a state ϕ of $\mathcal{TC}^*(E)$ is a KMS_β state of $(\mathcal{TC}^*(E), \alpha)$ for some $\beta \in \mathbb{R} \setminus \{0\}$ if and only if

$$(1.3) \quad \phi((s_\mu s_\nu^*)(s_\sigma s_\tau^*)) = \phi((s_\sigma s_\tau^*)\alpha_{i\beta}(s_\mu s_\nu^*)) = e^{-\beta(|\mu|-|\nu|)} \phi((s_\sigma s_\tau^*)(s_\mu s_\nu^*))$$

for all $\mu, \nu, \sigma, \tau \in E^*$. The KMS_β states of $(C^*(E), \alpha)$ come from the KMS_β states of $(\mathcal{TC}^*(E), \alpha)$ which factor through q .

As in [17], our KMS_0 states are the *invariant* traces (as opposed to *all* the traces, as in [2]). As in [3], we distinguish between KMS_∞ states, which are weak* limits of sequences of KMS_{β_n} states as $\beta_n \rightarrow \infty$, and ground states, which are states ϕ such that the functions $\phi_{a,b} : z \mapsto \phi(a\gamma_z(b))$ are bounded on the upper-half plane for $a, b \in \{s_\mu s_\nu^* : \mu, \nu \in E^*\}$. In the older literature, such as [2] or [17], the KMS_∞ states are defined to be the ground states. For general dynamical systems, every KMS_∞ state is a ground state (by [2, Theorem 5.3.23]), but not every ground state need be a KMS_∞ state (as happens in [12] and [13], for example.)

2. CHARACTERISING KMS STATES

Proposition 2.1. *Let E be a finite directed graph, and let $A \in M_{E^0}(\mathbb{N})$ be the vertex matrix with entries $A(v, w) = |vE^1 w|$. Let $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ be the gauge action, and define $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ by $\alpha_t = \gamma_{e^{it}}$. Let $\beta \in \mathbb{R}$.*

(a) *A state ϕ of $\mathcal{TC}^*(E)$ is a KMS_β state of $(\mathcal{TC}^*(E), \alpha)$ if and only if*

$$(2.1) \quad \phi(s_\mu s_\nu^*) = \delta_{\mu, \nu} e^{-\beta|\mu|} \phi(p_{s(\mu)}) \quad \text{for all } \mu, \nu \in E^*.$$

(b) *A state ϕ of $\mathcal{TC}^*(E)$ is a ground state of $(\mathcal{TC}^*(E), \alpha)$ if and only if*

$$(2.2) \quad \phi(s_\mu s_\nu^*) = 0 \quad \text{whenever } |\mu| > 0 \text{ or } |\nu| > 0.$$

(c) *Suppose that ϕ is a KMS_β state of $(\mathcal{TC}^*(E), \alpha)$, and define $m^\phi = (m_v^\phi) \in [0, \infty)^{E^0}$ by $m_v^\phi = \phi(p_v)$. Then m^ϕ is a probability measure on E^0 satisfying the subinvariance relation $Am^\phi \leq e^\beta m^\phi$.*

(d) *A KMS_β state ϕ of $(\mathcal{TC}^*(E), \alpha)$ factors through $C^*(E)$ if and only if $(Am^\phi)_v = e^\beta m_v^\phi$ whenever v is not a source.*

For part (d), we need a lemma. Recall that the graph algebra $C^*(E)$ is the quotient of $\mathcal{TC}^*(E)$ by the ideal J generated by

$$(2.3) \quad P := \left\{ p_v - \sum_{f \in vE^1} s_f s_f^* : v \in E^0 \text{ and } vE^1 \neq \emptyset \right\}.$$

We need to know that a state ϕ factors through $C^*(E)$ if and only if it vanishes on the generators of J (which is not obvious because ϕ is not a homomorphism). We have adapted the following lemma from [12, Lemma 10.3], and have tried to phrase the new version so that it might be useful in other computations of KMS states. Notice that

the sets P defined in (2.3) and $\mathcal{F} := \{s_\mu s_\nu^*\}$ in $\mathcal{TC}^*(E)$ have the properties required in Lemma 2.2.

Lemma 2.2. *Suppose (A, \mathbb{R}, α) is a dynamical system, and J is an ideal in A generated by a set P of projections which are fixed by α . Suppose that there is a family \mathcal{F} of analytic elements such that $\text{span } \mathcal{F}$ is dense in A , and such that for each $a \in \mathcal{F}$, there is a scalar-valued analytic function f_a satisfying $\alpha_z(a) = f_a(z)a$. If ϕ is a KMS_β state of (A, α) and $\phi(p) = 0$ for all $p \in P$, then ϕ factors through a state of A/J .*

Proof. It suffices to prove that $\phi(apb) = 0$ for all $a, b \in A$ and $p \in P$. Let $p \in P$. We have

$$0 \leq \phi(paa^*p) \leq \phi(p\|a\|^2p) = \|a\|^2\phi(p) = 0,$$

and hence ϕ vanishes on pAp . Now fix $a, b \in \mathcal{F}$. Since $\alpha_t(ap) = \alpha_t(a)p$ for $t \in \mathbb{R}$, the element ap is analytic with $\alpha_z(ap) = \alpha_z(a)p = f_a(z)ap$. Thus the KMS_β condition gives

$$\phi(apb) = \phi((ap)(pb)) = \phi(pb\alpha_{i\beta}(ap)) = f_a(i\beta)\phi(pbab) = 0,$$

and this extends to arbitrary a and b in A by linearity and continuity of ϕ . Now we use the linearity and continuity of ϕ again to see that ϕ vanishes on J . \square

Proof of Proposition 2.1. (a) Suppose that ϕ is a KMS_β state. If $\beta \neq 0$, then [17, Proposition 8.12.4] implies that ϕ is invariant for α and γ . This is also true for $\beta = 0$ by our convention that the KMS_0 states are invariant traces. For $|\mu| \neq |\nu|$, invariance gives

$$\phi(s_\mu s_\nu^*) = \int_{\mathbb{T}} \phi(\gamma_z(s_\mu s_\nu^*)) dz = \left(\int_{\mathbb{T}} z^{|\mu| - |\nu|} dz \right) \phi(s_\mu s_\nu^*) = 0.$$

For $|\mu| = |\nu|$ the product formula (1.2) gives $s_\nu^* s_\mu = \delta_{\nu, \mu} p_{s(\mu)}$, and the KMS condition gives

$$\phi(s_\mu s_\nu^*) = \phi(s_\nu^* \alpha_{i\beta}(s_\mu)) = e^{-\beta|\mu|} \phi(s_\nu^* s_\mu) = \delta_{\mu, \nu} e^{-\beta|\mu|} \phi(p_{s(\mu)}),$$

so ϕ satisfies (2.1).

Next suppose that ϕ satisfies (2.1). To see that ϕ is a KMS_β state, it suffices to check the KMS condition (1.3). (For $\beta = 0$, we need also to observe that any state ϕ satisfying (2.1) is automatically invariant for the gauge action — indeed, $\phi(s_\mu s_\nu^*) \neq 0$ implies $\mu = \nu$, and then $\gamma_z(s_\mu s_\nu^*) = s_\mu s_\nu^*$.) So we consider a pair of spanning elements $s_\mu s_\nu^*$ and $s_\sigma s_\tau^*$ in $\mathcal{TC}^*(E)$. Computations using the product formula (1.2) give

$$\begin{aligned} \phi(s_\mu s_\nu^* s_\sigma s_\tau^*) &= \begin{cases} \phi(s_{\mu\sigma'} s_{\tau'}^*) & \text{if } \sigma = \nu\sigma' \\ \phi(s_\mu s_{\tau\nu'}^*) & \text{if } \nu = \sigma\nu' \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} e^{-\beta|\tau|} \phi(p_{s(\tau)}) & \text{if } \sigma = \nu\sigma' \text{ and } \tau = \mu\sigma' \\ e^{-\beta|\mu|} \phi(p_{s(\mu)}) & \text{if } \nu = \sigma\nu' \text{ and } \mu = \tau\nu' \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Similarly,

$$\phi(s_\sigma s_\tau^* s_\mu s_\nu^*) = \begin{cases} e^{-\beta|\nu|} \phi(p_{s(\nu)}) & \text{if } \mu = \tau\mu' \text{ and } \nu = \sigma\mu' \\ e^{-\beta|\sigma|} \phi(p_{s(\sigma)}) & \text{if } \tau = \mu\tau' \text{ and } \sigma = \nu\tau' \\ 0 & \text{otherwise,} \end{cases}$$

and so

$$\phi(s_\sigma s_\tau^* \alpha_{i\beta}(s_\mu s_\nu^*)) = \begin{cases} e^{-\beta(|\mu|-|\nu|)} e^{-\beta|\nu|} \phi(p_{s(\nu)}) & \text{if } \mu = \tau\mu' \text{ and } \nu = \sigma\mu' \\ e^{-\beta(|\mu|-|\nu|)} e^{-\beta|\sigma|} \phi(p_{s(\sigma)}) & \text{if } \tau = \mu\tau' \text{ and } \sigma = \nu\tau' \\ 0 & \text{otherwise.} \end{cases}$$

If $\mu = \tau\mu'$ and $\nu = \sigma\mu'$, then $s(\mu) = s(\nu)$ and

$$\phi(s_\mu s_\nu^* s_\sigma s_\tau^*) = e^{-\beta|\mu|} \phi(p_{s(\mu)}) = e^{-\beta|\mu|} \phi(p_{s(\nu)}) = \phi(s_\sigma s_\tau^* \alpha_{i\beta}(s_\mu s_\nu^*)).$$

If $\tau = \mu\tau'$ and $\sigma = \nu\tau'$, then $s(\tau) = s(\sigma)$ and $|\mu| - |\nu| = |\mu\tau'| - |\nu\tau'| = |\tau| - |\sigma|$, so

$$\phi(s_\mu s_\nu^* s_\sigma s_\tau^*) = e^{-\beta|\tau|} \phi(p_{s(\tau)}) = e^{-\beta(|\mu|-|\nu|+|\sigma|)} \phi(p_{s(\sigma)}) = e^{-\beta(|\mu|-|\nu|)} \phi(s_\sigma s_\tau^* s_\mu s_\nu^*).$$

Otherwise both $\phi(s_\mu s_\nu^* s_\sigma s_\tau^*)$ and $\phi(s_\sigma s_\tau^* \alpha_{i\beta}(s_\mu s_\nu^*))$ are 0. Thus ϕ satisfies (1.3), and is a KMS_β -state.

(b) For every state ϕ and all μ, ν we have

$$|\phi(s_\mu \alpha_{a+ib}(s_\nu^*))| = |e^{-i(a+ib)|\nu|} \phi(s_\mu s_\nu^*)| = e^{b|\nu|} |\phi(s_\mu s_\nu^*)|.$$

If ϕ is a ground state and $\phi(s_\mu s_\nu^*) \neq 0$, this is bounded for $b > 0$, and hence $|\nu| = 0$; since $\phi(s_\nu s_\mu^*) = \overline{\phi(s_\mu s_\nu^*)}$, symmetry implies that $|\mu| = 0$ also.

Now suppose that ϕ satisfies (2.2). Then for $\sigma, \tau, \mu, \nu \in E^*$ we have

$$(2.4) \quad |\phi(s_\sigma s_\tau^* \alpha_{a+ib}(s_\mu s_\nu^*))| = e^{-b(|\mu|-|\nu|)} |\phi(s_\sigma s_\tau^* s_\mu s_\nu^*)|.$$

The right-hand side of (2.4) vanishes (and hence is trivially bounded) unless $\mu = \tau\mu'$ or $\tau = \mu\tau'$. If $\mu = \tau\mu'$, then $\phi(s_\sigma s_\tau^* s_\mu s_\nu^*) = \phi(s_{\sigma\mu'} s_\nu^*) = 0$ unless $|\nu| = |\sigma\mu'| = 0$, which implies $\mu' = s(\tau)$, $\mu = \tau$ and

$$|\phi(s_\sigma s_\tau^* \alpha_{a+ib}(s_\mu s_\nu^*))| = e^{-b|\mu|} \phi(s_\mu^* s_\mu),$$

which is bounded for $b > 0$. A similar argument applies when $\tau = \mu\tau'$, and hence ϕ is a ground state.

(c) Each m_v^ϕ is non-negative because ϕ is a positive functional. To see that m^ϕ is a probability measure on E^0 , note that $\sum_{v \in E^0} p_v$ is the identity of $\mathcal{TC}^*(E)$, and hence

$$1 = \phi(1) = \sum_{v \in E^0} \phi(p_v) = \sum_{v \in E^0} m_v^\phi.$$

Suppose $v \in E^0$ is not a source. We have $\phi(p_v) \geq \sum_{f \in vE^1} \phi(s_f s_f^*)$, and

$$(2.5) \quad \begin{aligned} \sum_{f \in vE^1} \phi(s_f s_f^*) &= \sum_{f \in vE^1} e^{-\beta} \phi(p_{s(f)}) = \sum_{f \in vE^1} e^{-\beta} m_{s(f)}^\phi \\ &= e^{-\beta} \sum_{w \in E^0} A(v, w) m_w^\phi = e^{-\beta} (Am^\phi)_v. \end{aligned}$$

Hence $(Am^\phi)_v \leq e^\beta \phi(p_v) = e^\beta m_v^\phi$.

Now suppose $v \in E^0$ is a source. Then $A(v, w) = 0$ for all $w \in E^0$, and

$$(Am^\phi)_v = \sum_{w \in E^0} A(v, w) m_w^\phi = 0 \leq e^\beta m_v^\phi.$$

Thus $(Am^\phi)_v \leq e^\beta m_v^\phi$ for all v .

(d) Choose $v \in E^0$ and suppose that v is not a source. By Lemma 2.2 it suffices to check that $\phi(p_v - \sum_{f \in vE^1} s_f s_f^*) = 0$ if and only if $(Am^\phi)_v = e^\beta m_v^\phi$. For this we use (2.1) and (2.5) to see that

$$e^\beta \phi\left(p_v - \sum_{f \in vE^1} s_f s_f^*\right) = e^\beta \left(\phi(p_v) - \sum_{f \in vE^1} e^{-\beta} \phi(p_{s(f)})\right) = e^\beta m_v^\phi - (Am^\phi)_v. \quad \square$$

3. KMS STATES AT LARGE INVERSE TEMPERATURES

In this section we study the KMS_β states of $(\mathcal{TC}^*(E), \alpha)$ for $\beta > \ln \rho(A)$. The import of this condition is that the series $\sum_{n=0}^{\infty} e^{-\beta n} A^n$ converges in the operator norm with sum $(I - e^{-\beta} A)^{-1}$ (by, for example, [4, §VII.3.1]). We use this observation several times in the proof of Theorem 3.1.

Theorem 3.1. *Let E be a finite directed graph with vertex matrix $A \in M_{E^0}(\mathbb{N})$. Let $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ be the gauge action and define $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ by $\alpha_t = \gamma_{e^{it}}$. Assume that $\beta > \ln \rho(A)$.*

- (a) *For $v \in E^0$, the series $\sum_{\mu \in E^*v} e^{-\beta|\mu|}$ either converges or is finite, with sum $y_v \geq 1$. Set $y := (y_v) \in [1, \infty)^{E^0}$, and consider $\epsilon \in [0, \infty)^{E^0}$. Then $m := (I - e^{-\beta} A)^{-1} \epsilon$ is a probability measure on E^0 if and only if $\epsilon \cdot y = 1$.*
- (b) *Suppose $\epsilon \in [0, \infty)^{E^0}$ satisfies $\epsilon \cdot y = 1$, and set $m := (I - e^{-\beta} A)^{-1} \epsilon$. Then there is a KMS_β state ϕ_ϵ of $(\mathcal{TC}^*(E), \alpha)$ satisfying*

$$(3.1) \quad \phi_\epsilon(s_\mu s_\nu^*) = \delta_{\mu,\nu} e^{-\beta|\mu|} m_{s(\mu)}.$$

- (c) *The map $\epsilon \mapsto \phi_\epsilon$ is an affine isomorphism of*

$$\Sigma_\beta := \{\epsilon \in [0, \infty)^{E^0} : \epsilon \cdot y = 1\}$$

onto the simplex of KMS_β states of $(\mathcal{TC}^(E), \alpha)$. The inverse of this isomorphism takes a KMS_β state ϕ to $(I - e^{-\beta} A)m^\phi$.*

Remark 3.2. Because $y_v \geq 1$, the extreme points of Σ_β are the vectors $\epsilon^u = (\epsilon_v^u) = (\delta_{u,v} y_u^{-1})$. Thus Σ_β is a simplex of dimension $|E^0| - 1$, as predicted by [6] (see §6.1).

Proof of Theorem 3.1(a). Let $v \in E^0$. Since $A^n(w, v)$ is the number of paths of length n from v to w , we have

$$(3.2) \quad \sum_{\mu \in E^*v} e^{-\beta|\mu|} = \sum_{n=0}^{\infty} \sum_{\mu \in E^n v} e^{-\beta n} = \sum_{n=0}^{\infty} \sum_{w \in E^0} e^{-\beta n} |wE^n v| = \sum_{n=0}^{\infty} \sum_{w \in E^0} e^{-\beta n} A^n(w, v).$$

(The sums in (3.2) are finite if $E^n v$ is empty for large n .) Since $\beta > \ln \rho(A)$, the series $\sum_{n=0}^{\infty} e^{-\beta n} A^n$ converges in the operator norm. Thus for every fixed $w \in E^0$ the series $\sum_{n=0}^{\infty} e^{-\beta n} A^n(w, v)$ converges, and hence the last sum in (3.2) converges. The sum is at least 1 because all the terms are non-negative and $e^{-\beta 0} A^0(v, v) = 1$.

The expansion $m = \sum_{n=0}^{\infty} e^{-\beta n} A^n \epsilon$ shows that $m \geq 0$, and we use the same expansion to compute

$$(3.3) \quad \begin{aligned} m(E^0) &= \sum_{v \in E^0} m_v = \sum_{v \in E^0} ((I - e^{-\beta} A)^{-1} \epsilon)_v \\ &= \sum_{v \in E^0} \left(\left(\sum_{n=0}^{\infty} e^{-\beta n} A^n \right) \epsilon \right)_v = \sum_{v \in E^0} \sum_{n=0}^{\infty} \sum_{w \in E^0} e^{-\beta n} A^n(v, w) \epsilon_w \end{aligned}$$

$$\begin{aligned}
&= \sum_{w \in E^0} \epsilon_w \left(\sum_{v \in E^0} \sum_{n=0}^{\infty} e^{-\beta n} |v E^n w| \right) = \sum_{w \in E^0} \epsilon_w \left(\sum_{\mu \in E^* w} e^{-\beta |\mu|} \right) \\
&= \epsilon \cdot y. \quad \square
\end{aligned}$$

Proof of Theorem 3.1 (b). We build our KMS_β states by representing $\mathcal{TC}^*(E)$ on $\ell^2(E^*)$. We write h_μ for the point mass at $\mu \in E^*$, and let $\{Q_v : v \in E^0\}$ and $\{T_e : e \in E^1\}$ be the partial isometries such that

$$Q_v h_\mu = \begin{cases} h_\mu & \text{if } v = r(\mu) \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad T_e h_\mu = \begin{cases} h_{e\mu} & \text{if } s(e) = r(\mu) \\ 0 & \text{otherwise.} \end{cases}$$

Then (Q, T) is a Toeplitz-Cuntz-Krieger family in $B(\ell^2(E^*))$, and there is a representation $\pi_{Q,T} : \mathcal{TC}^*(E) \rightarrow B(\ell^2(E^*))$ such that $\pi_{Q,T}(p_v) = Q_v$ and $\pi_{Q,T}(s_e) = T_e$ (in fact, it follows from [7, Theorem 4.1] that $\pi_{Q,T}$ is faithful). For $\mu \in E^*$ we set

$$(3.4) \quad \Delta_\mu := e^{-\beta |\mu|} \epsilon_{s(\mu)},$$

and note that $\Delta_\mu \geq 0$. We aim to define ϕ_ϵ by

$$(3.5) \quad \phi_\epsilon(a) = \sum_{\mu \in E^*} \Delta_\mu (\pi_{Q,T}(a) h_\mu | h_\mu) \quad \text{for } a \in \mathcal{TC}^*(E).$$

To see that (3.5) defines a state, we need to show that $\sum_{\mu \in E^*} \Delta_\mu = 1$. For $v \in E^0$ we have

$$\begin{aligned}
(3.6) \quad \sum_{\mu \in v E^*} \Delta_\mu &= \sum_{n=0}^{\infty} \sum_{\mu \in v E^n} e^{-\beta n} \epsilon_{s(\mu)} = \sum_{n=0}^{\infty} e^{-\beta n} \left(\sum_{w \in E^0} \sum_{\mu \in v E^n w} \epsilon_w \right) \\
&= \sum_{n=0}^{\infty} e^{-\beta n} \left(\sum_{w \in E^0} A^n(v, w) \epsilon_w \right) = \sum_{n=0}^{\infty} (e^{-\beta n} A^n \epsilon)_v,
\end{aligned}$$

which converges with sum $m_v = ((1 - e^{-\beta} A)^{-1} \epsilon)_v$ because $\beta > \ln \rho(A)$. We saw in part (a) that m is a probability measure, so $\sum_{\mu \in E^*} \Delta_\mu = \sum_{v \in E^0} m_v = 1$. This implies, first, that the series in (3.5) converges for all a , and hence defines a positive functional on $\mathcal{TC}^*(E)$, and, second, that $\phi_\epsilon(1) = 1$, so that ϕ_ϵ is a state.

To prove (3.1), let $\lambda \in E^*$. Then

$$(\pi_{Q,T}(s_\mu s_\nu^*) h_\lambda | h_\lambda) = (T_\nu^* h_\lambda | T_\mu^* h_\lambda) = \begin{cases} 1 & \text{if } \lambda = \mu \lambda' = \nu \lambda' \\ 0 & \text{otherwise.} \end{cases}$$

Since $\mu \lambda' = \nu \lambda'$ forces $\mu = \nu$, we have $\phi_\epsilon(s_\mu s_\nu^*) = 0$ if $\mu \neq \nu$. So suppose $\mu = \nu$. Then since (3.6) gives $\sum_{\mu \in v E^*} \Delta_\mu = m_v$, we have

$$\begin{aligned}
\phi_\epsilon(s_\mu s_\mu^*) &= \sum_{\lambda \in E^*} \Delta_\lambda (T_\mu^* h_\lambda | T_\mu^* h_\lambda) = \sum_{\lambda = \mu \lambda'} e^{-\beta |\mu \lambda'|} \epsilon_{s(\lambda')} \\
&= e^{-\beta |\mu|} \sum_{\lambda' \in s(\mu) E^*} \Delta_{\lambda'} = e^{-\beta |\mu|} m_{s(\mu)}.
\end{aligned}$$

Thus ϕ_ϵ satisfies (3.1). Since $\phi(p_v) = m_v$, ϕ_ϵ satisfies (2.1), and Proposition 2.1 implies that ϕ_ϵ is a KMS_β state. \square

Proof of Theorem 3.1 (c). To see that every KMS_β state ϕ has the form ϕ_ϵ , apply Proposition 2.1 (c) to see that $m^\phi = (\phi(p_v))$ is a subinvariant probability measure, and take $\epsilon := (I - e^{-\beta}A)m^\phi$. Then $m := (I - e^{-\beta}A)^{-1}\epsilon = m^\phi$, and comparing (2.1) with (3.1) shows that $\phi = \phi_\epsilon$.

The formula (3.1) also shows that the map $F : \epsilon \mapsto \phi_\epsilon$ is injective, and that F is weak* continuous from $\Sigma_\beta \subset \mathbb{R}^{E^0}$ to the state space of $\mathcal{TC}^*(E)$. Thus F is a homeomorphism of the compact space Σ_β onto the simplex of KMS_β states. The formulas (3.4) and (3.5) show that F is affine, and the formula for the inverse follows from the proof of surjectivity. \square

Remark 3.3. In the proof of Theorem 3.1 (c), we observed that a probability measure m on E^0 is subinvariant exactly when $\epsilon := (I - e^{-\beta}A)m$ is nonnegative. Thus parts (a) and (b) of Theorem 3.1 give a converse to Theorem 2.1 (c): the map $\phi \mapsto m^\phi$ defined by $m_v^\phi = \phi(p_v)$ is an isomorphism of the simplex of KMS_β states onto the simplex of subinvariant probability measures on E^0 .

Remark 3.4. Whether the sum in (3.2) is finite or infinite depends on the existence of cycles in E . If the index set E^*v is infinite for some v , then there is a cycle in E , Lemma A.1 implies that $\rho(A) \geq 1$, and hence Theorem 3.1 applies only to $\beta > \ln \rho(A) \geq 0$. The index set E^*v is finite for all v if and only if there are no cycles in E , in which case Lemma A.1 implies that $\rho(A) = 0$, and Theorem 3.1 applies to any real β . Thus when E has no cycles, there is a $(|E^0| - 1)$ -dimensional simplex of KMS_β states for all $\beta \in \mathbb{R}$.

4. KMS STATES AT THE CRITICAL INVERSE TEMPERATURE

The critical inverse temperature is $\beta = \ln \rho(A)$. As in the previous section, we can prove the existence of $\text{KMS}_{\ln \rho(A)}$ states for arbitrary graphs.

Proposition 4.1. *Suppose that E is a finite directed graph with vertex matrix A . If m is a probability measure on E^0 such that $Am \leq \rho(A)m$, then there is a $\text{KMS}_{\ln \rho(A)}$ state ϕ on $(\mathcal{TC}^*(E), \alpha)$ such that*

$$(4.1) \quad \phi(s_\mu s_\nu^*) = \delta_{\mu,\nu} \rho(A)^{-|\mu|} m_{s(\mu)}.$$

The state ϕ factors through a state of $C^(E)$ if and only if $(Am)_v = \rho(A)m_v$ for every vertex v which is not a source.*

Proof. Choose a sequence $\{\beta_n\} \subset (\ln \rho(A), \infty)$ such that $\beta_n \rightarrow \ln \rho(A)$. For each n , the vector m is a probability measure satisfying $Am \leq \rho(A)m \leq e^{\beta_n}m$. Hence $\epsilon_n := (I - e^{-\beta_n}A)m \in [0, \infty)^{E^0}$ and so the vector y of Theorem 3.1 (a) with $\beta = \beta_n$ satisfies $\epsilon_n \cdot y = 1$. Applying Theorem 3.1 (b) gives a KMS_{β_n} state ϕ_n satisfying

$$(4.2) \quad \phi_n(s_\mu s_\nu^*) = \delta_{\mu,\nu} e^{-\beta_n|\mu|} m_{s(\mu)}.$$

Since the state space of $\mathcal{TC}^*(E)$ is weak* compact we may assume by passing to a subsequence that the sequence $\{\phi_n\}$ converges to a state ϕ . Letting $n \rightarrow \infty$ in (4.2) shows that ϕ satisfies (4.1). Proposition 2.1 (a) implies that ϕ is a $\text{KMS}_{\ln \rho(A)}$ state. (Or we could apply the general result about limits of KMS states in [2, Proposition 5.3.23].) The last assertion follows from Proposition 2.1 (d). \square

Corollary 4.2. *Suppose that E is a finite directed graph with vertex matrix A . Then $(\mathcal{TC}^*(E), \alpha)$ has a $\text{KMS}_{\ln \rho(A)}$ state.*

Proof. Choose a decreasing sequence $\{\beta_n\}$ with $\beta_n \rightarrow \ln \rho(A)$. Then Theorem 3.1 (a) implies that there are probability measures m^n on E^0 such that $Am^n \leq e^{\beta_n} m^n$. By passing to a subsequence, we may suppose that m^n converges pointwise to a probability measure m , and then $Am \leq \rho(A)m$. So Proposition 4.1 gives a $\text{KMS}_{\ln \rho(A)}$ state. \square

To get uniqueness at the critical inverse temperature $\beta = \ln \rho(A)$, we impose some restrictions on E . We say that E is *strongly connected* if vE^*w is nonempty for every $v, w \in E^0$; equivalently, if the vertex matrix A is irreducible in the sense of [20, Chapter 1]. Thus [20, Theorem 1.5] implies that $\rho(A)$ is an eigenvalue of A for which there is an eigenvector $x = (x_v)$ with $x_v > 0$ for all v . The eigenvector x such that $\sum_{v \in E^0} x_v = 1$ is the *unimodular Perron-Frobenius eigenvector* of A . The following is our most satisfying result on uniqueness, but later we will improve it by allowing sources (Corollary 6.1). Notice in particular that parts (a) and (c) complete the description of the KMS states on $\mathcal{TC}^*(E)$ when E is strongly connected.

Theorem 4.3. *Let E be a finite directed graph and suppose that E is strongly connected. Let $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ be the gauge action and define $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ by $\alpha_t = \gamma_{e^{it}}$. Let x be the unimodular Perron-Frobenius eigenvector of the vertex matrix A .*

(a) *The system $(\mathcal{TC}^*(E), \alpha)$ has a unique $\text{KMS}_{\ln \rho(A)}$ state ϕ . This state satisfies*

$$(4.3) \quad \phi(s_\mu s_\nu^*) = \delta_{\mu, \nu} \rho(A)^{-|\mu|} x_{s(\mu)},$$

and factors through a $\text{KMS}_{\ln \rho(A)}$ state $\bar{\phi}$ of $(C^(E), \alpha)$.*

(b) *The state $\bar{\phi}$ is the only KMS state of $(C^*(E), \alpha)$.*

(c) *If $\beta < \ln \rho(A)$, then $(\mathcal{TC}^*(E), \alpha)$ has no KMS_β states.*

Proof. (a) We proved existence of ϕ in Corollary 4.2, and ϕ factors through $C^*(E)$ because $Ax = \rho(A)x$. To establish uniqueness, suppose that ψ is a $\text{KMS}_{\ln \rho(A)}$ state. Then Proposition 2.1 (c) says that $m^\psi = (\psi(p_v))$ is a probability measure satisfying $Am^\psi \leq \rho(A)m^\psi$. Now the forward implication in the last assertion of [20, Theorem 1.6] implies that $m^\psi = x$, and together the formulas (2.1) and (4.3) imply that $\psi = \phi$.

(b) Suppose that ψ is a KMS state of $(C^*(E), \alpha)$, with inverse temperature β , say. Then Proposition 2.1 (d) implies that $Am^{\psi \circ q} = e^\beta m^{\psi \circ q}$; since A is irreducible, the backward implication in the last assertion of [20, Theorem 1.6] implies that $e^\beta = \rho(A)$. Now the uniqueness in part (a) implies that $\psi \circ q = \phi = \bar{\phi} \circ q$, and $\psi = \bar{\phi}$.

(c) Suppose that ϕ is a KMS_β state of $(\mathcal{TC}^*(E), \alpha)$. Then Proposition 2.1 (c) implies that $m^\phi := (\phi(p_v))$ satisfies $Am^\phi \leq e^\beta m^\phi$. In other words, m^ϕ is subinvariant. Since $m^\phi \geq 0$ pointwise, [20, Theorem 1.6] implies that $e^\beta \geq \rho(A)$, or equivalently $\beta \geq \ln \rho(A)$. \square

Remark 4.4. The irreducibility of A was crucial in the proof of uniqueness. A similar phenomenon occurs in [13, Theorem 5.3], where an extra hypothesis (there, that a given integer matrix is a dilation matrix) is needed to ensure uniqueness at the critical inverse temperature.

Remark 4.5. A strongly connected graph contains at least one cycle. If it contains more than one, then Lemma A.1 implies that $\rho(A) > 1$, and $C^*(E)$ has no KMS_0 states. However, if E consists of a single cycle, then $\rho(A) = 1$, and Theorem 4.3 implies that $C^*(E)$ has a unique KMS_0 state. We check that this is consistent with what we know about cycles. Suppose that E is a cycle with n edges. Then $C^*(E)$ is isomorphic to $C(\mathbb{T}, M_n(\mathbb{C}))$, and the gauge action acts transitively on the spectrum \mathbb{T} (see [18, Example 2.14], for

example). Thus there is a unique invariant measure, and integrating the usual normalised trace against this measure gives a unique invariant trace on $C(\mathbb{T}, M_n(\mathbb{C}))$.

5. GROUND STATES AND KMS_∞ STATES

Proposition 5.1. *Let E be a finite directed graph, let $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ be the gauge action and define $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{TC}^*(E)$ by $\alpha_t = \gamma_{e^{it}}$. Suppose that ϵ is a probability measure on E^0 . Then there is a KMS_∞ state ϕ_ϵ satisfying*

$$(5.1) \quad \phi_\epsilon(s_\mu s_\nu^*) = \begin{cases} 0 & \text{unless } |\mu| = |\nu| = 0 \text{ and } \mu = \nu \\ \epsilon_v & \text{if } \mu = \nu = v \in E^0. \end{cases}$$

Every ground state of $(\mathcal{TC}^*(E), \alpha)$ is a KMS_∞ state, and the map $\epsilon \mapsto \phi_\epsilon$ is an affine isomorphism of the simplex of probability measures on E^0 onto the set of ground states of $(\mathcal{TC}^*(E), \alpha)$.

Proof. Choose a sequence $\beta_j \rightarrow \infty$ as $j \rightarrow \infty$ with each $\beta_j > \ln \rho(A)$. For each j , define (y_v^j) as in Theorem 3.1 (a) by $y_v^j = \sum_{\alpha \in E^*v} e^{-\beta_j |\alpha|}$. Set $\epsilon_v^j := \epsilon_v (y_v^j)^{-1}$, and let ϕ_j be the KMS_{β_j} state ϕ_{ϵ^j} of $(\mathcal{TC}^*(E), \alpha)$ described in Theorem 3.1 (b). Since the state space is weak* compact we may assume that ϕ_j converges in the weak* topology to a state ϕ_ϵ .

Set $m^j := (I - e^{-\beta_j} A)^{-1} \epsilon^j$, and take μ, ν in E^* . Then

$$\phi_j(s_\mu s_\nu^*) = \delta_{\mu, \nu} e^{-\beta_j |\mu|} m_{s(\mu)}^j.$$

This is always 0 if $\mu \neq \nu$, so suppose that $\mu = \nu$. If $|\mu| > 0$, then $e^{-\beta_j |\mu|} \rightarrow 0$, and hence $\phi_j(s_\mu s_\mu^*) \rightarrow 0$. So suppose that $\mu = \nu = v$ is a vertex. An application of the dominated convergence theorem shows that $y_v^j \rightarrow 1$ as $j \rightarrow \infty$. Hence $\epsilon_v^j = \epsilon_v (y_v^j)^{-1} \rightarrow \epsilon_v$. Since $(I - e^{-\beta_j} A)^{-1} \rightarrow I$ in the operator norm, we have $m_v^j \rightarrow \epsilon_v$, and hence $\phi_j(p_v) \rightarrow \epsilon_v$. Since $\phi_j(p_v) \rightarrow \phi_\epsilon(p_v)$, we deduce that $\phi_\epsilon(p_v) = \epsilon_v$, and ϕ_ϵ satisfies (5.1).

Since $\phi_\epsilon(s_\mu s_\nu^*) = 0$ whenever $|\mu| > 0$ or $|\nu| > 0$, Proposition 2.1 (b) implies that ϕ_ϵ is a ground state. Now let ϕ be a ground state, and $\epsilon_v := \phi(p_v)$. Then ϵ is a probability measure on E^0 , and $\phi = \phi_\epsilon$ because ϕ is determined by its values on the vertex projections (by Proposition 2.1 (b) again). Thus $\epsilon \mapsto \phi_\epsilon$ maps the simplex of probability measures onto the ground states, and it is clearly affine and injective. Since each ϕ_ϵ is by construction a KMS_∞ state, it follows that every ground state is a KMS_∞ state. \square

6. CONNECTIONS WITH THE LITERATURE

6.1. Cuntz-Krieger algebras. When E has no sources, we could in principle deduce Theorem 3.1 from the work of Exel and Laca on the Cuntz-Krieger algebras of $\{0, 1\}$ -matrices. To do this, we apply [6, Theorem 18.4] to the edge matrix $B \in M_{E^1}(\mathbb{N})$ of E defined by

$$B(e, f) := \begin{cases} 1 & \text{if } s(e) = r(f) \\ 0 & \text{otherwise.} \end{cases}$$

We need to assume that E has no sources, because an edge e such that $s(e)$ is a source would give a row of zeroes in B , which is not allowed in [6]. Edges with the same range give equal columns of B , so provided there are no sinks, the distinct columns of B are in one-to-one correspondence with the vertex set E^0 , and the set Ω_e in [6, Theorem 18.4] can be identified with E^0 . The condition $\epsilon \cdot y = 1$ is phrased in [6] as $Z(\beta, \epsilon) = 1$, but we think it would be hard to dig our formula for ϕ_ϵ out of [6].

6.2. Graphs with sources. Kajiwara and Watatani [8] have recently considered the KMS states on the graph algebras $C^*(E)$ of arbitrary finite graphs, and have shown in particular that sources give rise to KMS_β states [8, Theorem 4.4].

Recall that a subset H of E^0 is *saturated* if $s(vE^1) \subset H \implies v \in H$. The *saturation* of a hereditary set $S \subset E^0$ is the smallest saturated set H containing S . The saturation is itself hereditary. For a saturated hereditary set H , the C^* -algebra $C^*(E \setminus H)$ of the graph in (b) below is a quotient of $C^*(E)$ (see [18, Theorem 4.9]).

Corollary 6.1. *Let E be a finite directed graph with vertex matrix A . Let $\gamma : \mathbb{T} \rightarrow \text{Aut } C^*(E)$ be the gauge action and define $\alpha : \mathbb{R} \rightarrow \text{Aut } C^*(E)$ by $\alpha_t = \gamma_{e^{it}}$.*

- (a) *Assume that $\beta > \ln \rho(A)$, and that ϵ belongs to the simplex Σ_β of Theorem 3.1. Then ϕ_ϵ factors through a state of $C^*(E)$ if and only if $\epsilon_v = 0$ whenever v is not a source.*
- (b) *Let H be the saturation of the set of sources in E . Suppose that E has no sinks and that the graph $E \setminus H := (E^0 \setminus H, E^1 \setminus s^{-1}(H), r, s)$ is strongly connected. Let x be the unimodular Perron-Frobenius eigenvector for the vertex matrix $A_{E \setminus H}$ of $E \setminus H$. Then there is a unique $\text{KMS}_{\ln \rho(A)}$ state ϕ of $(C^*(E), \alpha)$, and*

$$(6.1) \quad \phi(s_\mu s_\nu^*) = \begin{cases} \delta_{\mu,\nu} \rho(A)^{-|\mu|} x_{s(\mu)} & \text{if } s(\mu) \in E^0 \setminus H \\ 0 & \text{if } s(\mu) \in H. \end{cases}$$

Proof. (a) We set $m := (I - e^{-\beta} A)^{-1} \epsilon$. Proposition 2.1 (d) implies that ϕ_ϵ factors through a state of $C^*(E)$ if and only if $m_v = (e^{-\beta} A m)_v$ whenever v is not a source. Since $\epsilon = (I - e^{-\beta} A)m$, this is equivalent to $\epsilon_v = 0$ whenever v is not a source.

(b) Corollary 4.2 implies that there is a $\text{KMS}_{\ln \rho(A)}$ state ψ on $\mathcal{T}C^*(E)$. Then Proposition 2.1 (c) says that $(m_v^\psi) := (\psi(p_v))$ gives a probability measure m^ψ satisfying the subinvariance relation $A m^\psi \leq \rho(A) m^\psi$. We will analyse this subinvariance relation by writing A in a particular block form.

To do this, let S be the set of sources in E . Recall, from [1, Remark 3.1] for example, that we can construct the saturation H of the hereditary set S inductively. We set $S_0 := S$, and for $k \geq 0$ set

$$S_{k+1} = S_k \cup \{v \in E^0 : s(vE^1) \subset S_k\}.$$

Since E is finite, we eventually have $S_{n+1} = S_n$, and then $H = S_n$. We now order E^0 by listing first the vertices in $E^0 \setminus H$, then those in $H \setminus S_{n-1}$, then those in $S_{n-1} \setminus S_{n-2}$ and so on, finishing with the sources S . This ordering gives a block decomposition

$$A = \begin{pmatrix} A_{E \setminus H} & B \\ 0 & A_H \end{pmatrix}.$$

Since $w \in S_{k+1}$ and $A(w, v) \neq 0$ imply $v \in S_k$, the matrix A_H is strictly upper triangular, in the sense that all entries on or below the diagonal are 0.

We write $m^\psi = (m^{E \setminus H}, m^H)$ in block form. Subinvariance says that

$$(6.2) \quad A_{E \setminus H} m^{E \setminus H} + B m^H \leq \rho(A) m^{E \setminus H}, \text{ and}$$

$$(6.3) \quad A_H m^H \leq \rho(A) m^H.$$

Since A_H is strictly upper triangular, $\rho(A_{E \setminus H}) = \rho(A)$. Thus, since B and m^H are non-negative, (6.2) implies that $A_{E \setminus H} m^{E \setminus H} \leq \rho(A_{E \setminus H}) m^{E \setminus H}$. Now the last assertion in [20,

Theorem 1.6] implies that $A_{E \setminus H} m^{E \setminus H} = \rho(A_{E \setminus H}) m^{E \setminus H}$, and $m^{E \setminus H}$ is a Perron-Frobenius eigenvector for $A_{E \setminus H}$.

Since $A_{E \setminus H} m^{E \setminus H} = \rho(A_{E \setminus H}) m^{E \setminus H}$, (6.2) implies that $Bm^H = 0$. We next claim that $m^H = 0$, or equivalently that $m_v^\psi = 0$ for every vertex $v \in H$. First we consider $v \in S_n \setminus S_{n-1}$. Since E has no sinks, there is an edge e with $s(e) = v$. The range $r(e)$ cannot be in H , because otherwise $r(e) \in S_n$ but $s(e)$ is not in S_{n-1} . Thus $B(r(e), v) > 0$, and $(Bm^H)_{r(e)} = 0$ implies $m_{r(e)}^H = 0$. Thus $m_v^\psi = 0$ for all $v \in S_n \setminus S_{n-1}$. Now we repeat the argument with $w \in S_{n-1} \setminus S_{n-2}$, and find that $s(e) = w$ implies that $r(e)$ is in either $E \setminus H$ or $S_n \setminus S_{n-1}$; thus one of $B(r(e), w)m_w = 0$ or $0 \leq A_H(r(e), w)m_w^\psi \leq \rho(A)m_{r(e)}^\psi$ forces $m_w^\psi = 0$. A finite induction argument gives $m_v^\psi = 0$ for all $v \in H$, as claimed.

Since m^ψ is a probability measure on E^0 , we deduce that $m^{E \setminus H}$ is a probability measure too. Thus $m^{E \setminus H}$ is the unimodular Perron-Frobenius eigenvector x for $A_{E \setminus H}$. Now the vector m^ψ has block form $(x, 0)$, and we have $Am^\psi = \rho(A)m^\psi$. Thus it follows from Proposition 2.1 (d) that ψ factors through a state ϕ of $C^*(E)$, and this is a $\text{KMS}_{\ln \rho(A)}$ state of $(C^*(E), \alpha)$. The formula in Proposition 2.1 (a) implies that ϕ satisfies (6.1).

To see uniqueness, suppose ϕ' is a $\text{KMS}_{\ln \rho(A)}$ state of $(C^*(E), \alpha)$. Then we can run the argument of the preceding five paragraphs with $\psi = \phi' \circ q$, and deduce that ϕ' also satisfies (6.1). Thus ϕ' and ϕ agree on the spanning elements $s_\mu s_\nu^*$, and hence are equal. \square

Corollary 6.1 (a) implies that for $\beta > \ln \rho(A)$ the simplex of KMS_β states of $(C^*(E), \alpha)$ has dimension $|S|$ where S is the set of sources. Corollary 6.1 (a) contains [8, Theorem 4.4] because in both situations considered there, the range of inverse temperatures is the same as ours. Our result applies to arbitrary finite graphs, and not just ones whose vertex matrix has entries in $\{0, 1\}$, as in [8]. The uniqueness in part (b) appears to be new.

6.3. Cuntz-Pimsner algebras. When we were proving Theorem 3.1 we were guided by the machinery developed in [11], and in particular by the construction in the proof of [11, Theorem 2.1]. We believe that our direct approach will be more accessible to more readers, but putting our calculations in the context of [11] might provide an illuminating example for those interested in the general machine.

To apply the ideas of [11], we view $\mathcal{T}C^*(E)$ as the Toeplitz algebra $\mathcal{T}(X)$ of the graph correspondence X as in [7, Theorem 4.1] and [18, §8]. We consider only the Toeplitz algebra because there are competing definitions of the Cuntz-Pimsner algebra \mathcal{O}_X in the literature, and when E has sources the one used in [11] is not the one which gives $C^*(E)$ ¹.

In the conventions of [18], X has underlying set $C(E^1)$, with module actions given by $(a \cdot x \cdot b)(e) = a(r(e))x(e)b(s(e))$ and inner product by $\langle x, y \rangle(v) = \sum_{s(e)=v} \overline{x(e)}y(e)$. We can similarly realise the tensor powers $X^{\otimes n}$ as bimodules with underlying set $C(E^n)$. The dynamics on $\mathcal{T}(X)$ is implemented by the unitaries U_t^n on $X^{\otimes n} = C(E^n)$ such that $U_t^n \delta_\mu = e^{itn} \delta_\mu$, and hence the operator $\Gamma(e^{-\beta D})$ on the Fock module $\mathcal{F}(X) = \bigoplus_{n \geq 0} X^{\otimes n}$ considered in [11] is multiplication by $e^{-\beta n}$ on $X^{\otimes n}$.

A function $\epsilon \in \Sigma_\beta$ gives a trace $\tau_\epsilon : f \mapsto \sum_{v \in E^0} f(v)\epsilon_v$ on $C(E^0)$. For each n , the identity operator on $C(E^n) = X^{\otimes n}$ satisfies $1_n = \sum_{\lambda \in E^n} \Theta_{\delta_\lambda, \delta_\lambda}$. Thus the induced trace

¹The ideal I_X such that $\mathcal{O}_X = \mathcal{T}(X)/I_X$ in [11] is $\phi^{-1}(\mathcal{K}(X))$ rather than $J_X := \phi^{-1}(\mathcal{K}(X)) \cap (\ker \phi)^\perp$ (see the discussion in [18, Chapter 8] or [8, §4]). Kajiwara and Watatani deduce their results about KMS states on $C^*(E)$ for E with sources from a version of [11, Theorem 2.5] for algebras of the form $\mathcal{T}(X)/J_X$ [8, Theorem 3.17].

$F^n \tau_\epsilon$ on $A = C(E^0) \subset \mathcal{L}(\mathcal{F}(X))$ constructed in [11, Theorem 1.1] is given by

$$(F^n \tau_\epsilon)(\delta_w) = \sum_{\lambda \in E^n} \tau_\epsilon(\langle \delta_\lambda, (\delta_w \Gamma(e^{-\beta D}))(\delta_\lambda) \rangle) = \sum_{\lambda \in E^n} \tau_\epsilon(\langle \delta_\lambda, e^{-\beta n} \delta_w \cdot \delta_\lambda \rangle),$$

where the dot in $\delta_w \cdot \delta_\lambda$ is the left action of $\delta_w \in C(E^0)$ on $\delta_\lambda \in C(E^n)$. We can compute

$$\langle \delta_\lambda, e^{-\beta n} \delta_w \cdot \delta_\lambda \rangle(v) = \sum_{\nu \in E^n v} \overline{\delta_\lambda(\nu)} e^{-\beta n} (\delta_w \cdot \delta_\lambda)(\nu) = \begin{cases} e^{-\beta n} \delta_w(r(\lambda)) & \text{if } s(\lambda) = v \\ 0 & \text{otherwise,} \end{cases}$$

and therefore

$$\begin{aligned} (F^n \tau_\epsilon)(\delta_w) &= \sum_{\lambda \in E^n} \sum_{v \in E^0} \langle \delta_\lambda, e^{-\beta n} \delta_w \cdot \delta_\lambda \rangle(v) \epsilon_v \\ &= \sum_{v \in E^0} \sum_{\lambda \in E^n v} e^{-\beta n} \delta_w(r(\lambda)) \epsilon_v = \sum_{v \in E^0} e^{-\beta n} A^n(w, v) \epsilon_v. \end{aligned}$$

Thus the subinvariant vector $m = \sum_{n \geq 0} e^{-\beta n} A^n \epsilon = (I - e^{-\beta} A)^{-1} \epsilon$ in Theorem 3.1 is the trace $\sum_{n \geq 0} F^n \tau_\epsilon$ induced by the Fock module in the proof of [11, Theorem 2.1].

Laca and Neshveyev construct their KMS_β state as follows. The trace τ_ϵ on $C(E^0)$ gives a finite-dimensional Hilbert space H_ϵ which carries a representation M of $C(E^0)$ by multiplication operators. Rieffel induction then gives a representation $\mathcal{F}(X)$ -Ind M of $\mathcal{L}(\mathcal{F}(X))$ on $\mathcal{F}(X) \otimes_{C(E^0)} H_\epsilon$. As above, the identity on $X^{\otimes n} = C(E^n)$ is given by $1_n = \sum_{\lambda \in E^n} \Theta_{\delta_\lambda, \delta_\lambda}$. Thus [11, Theorem 1.1(ii)] implies that the trace $\text{Tr}_{\tau_\epsilon}$ satisfies

$$\text{Tr}_{\tau_\epsilon}(T) = \sum_{\lambda \in E^n} \tau_\epsilon(\langle \delta_\lambda, T \delta_\lambda \rangle) = \sum_{\lambda \in E^n} \langle \delta_\lambda, T \delta_\lambda \rangle \epsilon_{s(\lambda)}.$$

Thus the KMS_β state ϕ in the proof of [11, Theorem 2.1] is given by

$$\phi(a) = \sum_{\lambda \in E^n} \langle \delta_\lambda, a \Gamma(e^{-\beta D}) \delta_\lambda \rangle \epsilon_{s(\lambda)} = \sum_{\lambda \in E^n} e^{-\beta n} \langle \delta_\lambda, a \delta_\lambda \rangle \epsilon_{s(\lambda)}.$$

For $a = s_\mu s_\nu^*$, the inner product $\langle \delta_\lambda, a \delta_\lambda \rangle$ vanishes unless $\mu = \nu$ and $\lambda = \mu \lambda'$, and then $\langle \delta_\lambda, a \delta_\lambda \rangle = 1$. Thus

$$\begin{aligned} \phi(s_\mu s_\mu^*) &= e^{-\beta |\mu|} \sum_{\lambda' \in s(\mu) E^*} e^{-\beta |\lambda'|} \epsilon_{s(\lambda')} = e^{-\beta |\mu|} \sum_{n \geq 0} (e^{-\beta n} A^n \epsilon)_{s(\mu)} \\ &= e^{-\beta |\mu|} ((1 - e^{-\beta n} A^n)^{-1} \epsilon)_{s(\mu)} = e^{-\beta |\mu|} m_{s(\mu)}. \end{aligned}$$

Thus the state constructed in [11, Theorem 2.1] is exactly the same as ours. In fact, the representation $\mathcal{F}(X)$ -Ind M is unitarily equivalent to the representation $\pi_{Q,T}$ on $\ell^2(E^*)$ in the proof of Theorem 3.1 via the unitary which sends $\delta_\mu \otimes \delta_{s(\mu)}$ to $\epsilon_{s(\mu)}^{1/2} h_\mu$.

6.4. Groupoid algebras. In [19, Proposition II.5.4] Renault describes the KMS states for the C^* -algebras of principal groupoids, which are the groupoids with no isotropy (see also [9]). Neshveyev has recently extended Renault's result to groupoids with isotropy [14, Theorem 1.3]. Since graph algebras have a groupoid model, Neshveyev's result applies in our situation. We will see that this approach requires calculations similar to those in the proof of Theorem 3.1, but they may provide an instructive example for those interested in Neshveyev's approach. To avoid having to adjust the standard groupoid model of [10] or [16], we assume that E has no sources.

The groupoid model G for the Toeplitz algebra has unit space $G^{(0)} := E^* \cup E^\infty$, and the sets $Z(\alpha) := \{\alpha\beta : \beta \in G^{(0)}\}$ for $\alpha \in E^*$ form a basis of open compact sets for the topology on $G^{(0)}$ (see [16, Proposition 3.3]). We have

$$G = \{(\alpha y, |\gamma| - |\alpha|, \gamma y) : y \in G^{(0)}, \alpha, \gamma \in E^*, s(\alpha) = s(\gamma) = r(y)\},$$

$r(y, k, z) = y$ and $s(y, k, z) = z$. The sets

$$Z(\alpha, \gamma) := \{(\alpha y, |\gamma| - |\alpha|, \gamma y) : y \in G^{(0)}, s(\alpha) = s(\gamma) = r(y)\}$$

form a basis of compact-open sets for the topology on G . When we view $\mathcal{T}C^*(E)$ as $C^*(G)$, our dynamics is the one studied in [14] for the cocycle $c : G \rightarrow \mathbb{R}$ defined by $c(y, k, z) = k$. To construct KMS states on $C^*(G)$ using Neshveyev's theorem, we need to find quasi-invariant measures on $G^{(0)}$ and check some conditions involving the isotropy.

Fix $\beta > \ln \rho(A)$ and ϵ as in Theorem 3.1 (b). Consider the numbers $\{\Delta_\alpha : \alpha \in E^*\}$ constructed in the proof of that theorem. Since $\sum_{\alpha \in E^*} \Delta_\alpha = 1$, the discrete measure $\sum_{\alpha \in E^*} \Delta_\alpha \delta_\alpha$ is a probability measure on E^* , and hence gives a probability measure μ on $G^{(0)}$ such that $\mu(Z(\alpha)) = \sum_{r(\lambda)=s(\alpha)} \Delta_{\alpha\lambda}$ and $\mu(E^\infty) = 0$. To see that μ is quasi-invariant, consider the basic open set $U = Z(\alpha, s(\alpha)) = \{(\alpha y, -|\alpha|, y)\}$. Then r and s are homeomorphisms on U , and $T := s|_U \circ (r|_U)^{-1}$ maps αy to y . For $\lambda \in E^*$, we have

$$T_*\mu(Z(\lambda)) = \mu(T^{-1}(Z(\lambda))) = \begin{cases} 0 & \text{if } r(\lambda) \neq s(\alpha) \\ \mu(Z(\alpha\lambda)) & \text{if } r(\lambda) = s(\alpha). \end{cases}$$

Now the calculation

$$\begin{aligned} \mu(Z(\alpha\lambda)) &= \sum_{r(\gamma)=s(\lambda)} \Delta_{\alpha\lambda\gamma} = \sum_{r(\gamma)=s(\lambda)} e^{-\beta|\alpha\lambda\gamma|} \epsilon_{s(\gamma)} \\ &= e^{-\beta|\alpha|} \sum_{r(\gamma)=s(\lambda)} e^{-\beta|\lambda\gamma|} \epsilon_{s(\gamma)} = e^{-\beta|\alpha|} \mu(Z(\lambda)) \end{aligned}$$

implies that $\frac{dT_*\mu}{d\mu} = e^{-\beta|\alpha|}$, and hence that μ is quasi-invariant with the correct cocycle (see the discussion preceding [14, Theorem 1.3]).

In the groupoid G , a unit y has nonzero isotropy group $G_y^y = \{k : (y, k, y) \in G\}$ only if y is an infinite path. Thus $\mu(\{y : G_y^y \neq \{0\}\}) = 0$, and the measurable fields of states described in conditions (ii) and (iii) of [14, Theorem 1.3] give no extra KMS states (see the comments following that theorem in [14]). So Neshveyev's theorem gives a KMS_β state ψ_ϵ on $\mathcal{T}C^*(E)$ which is the composition of the state μ_* on $C(G^{(0)})$ with the expectation E of $C^*(G)$ onto $C(G^{(0)})$. We can check on elements of the form $\chi_{Z(\alpha)} = E(s_\alpha s_\alpha^*)$ that ψ_ϵ is the state ϕ_ϵ in Theorem 3.1.

We now suppose that E is strongly connected, and consider states on $C^*(E)$. The groupoid model for $C^*(E)$ is the reduction of G to E^∞ , and we can construct measures on E^∞ by constructing functionals on $C(E^\infty) = \varinjlim C(E^n)$. Suppose that $\rho(A) > 1$, and x is the unimodular Perron-Frobenius eigenvector A . Then the relation $\mu_*(Z(\alpha)) = \rho(A)^{-|\alpha|} x_{s(\alpha)}$ completely characterises a quasi-invariant measure μ on E^∞ . The paths y with nonzero isotropy have the form $\alpha\gamma\gamma\gamma\cdots$ for some $\alpha, \gamma \in E^*$ with $s(\gamma) = r(\gamma) = s(\alpha)$; since E is finite, there are countably many such paths, and since $\{\alpha\gamma\gamma\gamma\cdots\} = \bigcap_{n=1}^\infty Z(\alpha\gamma^n)$ we have $\mu(\{\alpha\gamma\gamma\gamma\cdots\}) \leq \rho(A)^{-|\alpha|-n}$ for all n , and $\mu(\{\alpha\gamma\gamma\gamma\cdots\}) = 0$. Thus the units with nonzero isotropy have measure zero, and again the measurable fields of states in [14, Theorem 1.3, (ii) and (iii)] give no additional KMS states.

Now suppose that $\rho(A) = 1$, in which case Lemma A.1 implies that E consists of a single cycle, say with n edges. Then the path space E^∞ has n points, the Perron-Frobenius eigenvector x is constant, and μ is the uniformly distributed probability measure. The isotropy group at $y \in E^\infty$ is $G_y^y = \{(y, nk, y) : k \in \mathbb{Z}\} \cong n\mathbb{Z}$, so Neshveyev's condition (iii) kicks in. His cocycle $c : (y, k, z) \mapsto k$ is injective on G_y^y , and hence the only state ϕ of $C^*(G_y^y) = C^*(n\mathbb{Z})$ such that $\phi(u_{nk}) = 0$ for all $k \in c^{-1}(0)$ is evaluation at 0 on $C_c(n\mathbb{Z}) \subset C^*(n\mathbb{Z})$. So there is just one measurable field satisfying (iii), and Neshveyev's theorem gives just one KMS_0 state on $C^*(E)$. This is consistent, because he too requires his KMS_0 states to be invariant (just before [14, Theorem 1.3]), and he makes it clear in the proof of the theorem that the purpose of (iii) is to ensure invariance.

APPENDIX A. THE SPECTRAL RADIUS OF A VERTEX MATRIX

We now discuss some properties of the vertex matrices of finite graphs which we have used at various points. These are non-negative matrices in the sense of [20, Chapter 1], and have integer entries, so there must be a good chance that these results are known. But it is easy enough to give quick proofs.

Lemma A.1. *Suppose that E is a finite graph, and let $A \in M_{E^0}(\mathbb{N})$ be the vertex matrix with entries $A(v, w) = |vE^1w|$.*

- (a) *If E contains at least one cycle, then the spectral radius of A satisfies $\rho(A) \geq 1$.*
- (b) *If E contains no cycles, then $\rho(A) = 0$.*
- (c) *If E consists of disjoint cycles, then $\rho(A) = 1$.*
- (d) *If E is strongly connected and not a cycle, then $\rho(A) > 1$.*

Proof. (a) For each $v \in E^0$ which lies on a cycle, we consider the set C_v of vertices w which lie on a return path based at v . Two such C_v and C_u are either equal or disjoint; the distinct subsets C_1, \dots, C_n are the classes (essential and inessential) which Seneta discusses in [20, Page 12]. By Seneta's structure theory, we can order the vertices so that A is a block upper triangular matrix whose square diagonal blocks are 1×1 blocks of the form (0) and the irreducible submatrices A_i of A associated to the classes C_i . (Our matrix is upper rather than lower triangular because of the way we defined the vertex matrix.) Since the determinant of a block-triangular matrix is the product of the determinants of the diagonal blocks, the spectrum of A is either $\bigcup_{i=1}^n \sigma(A_i)$ or $(\bigcup_{i=1}^n \sigma(A_i)) \cup \{0\}$. So it suffices for us to prove that, if A is irreducible, as the A_i are, then $\rho(A) \geq 1$.

By irreducibility, for each $v \in E^0$ there exists $m_v > 0$ such that $A^{m_v}(v, v) > 0$, and since A is an integer matrix, $A^{m_v}(v, v) \geq 1$. Now with $m := \prod_{v \in E^0} m_v$ we have $A^m(w, w) \geq 1$ for all $w \in E^0$. Thus the trace of A^m satisfies $\text{Tr } A^m \geq |E^0|$. Since the trace is the sum of the complex eigenvalues counted according to multiplicity, at least one of these eigenvalues has absolute value at least 1, giving $\rho(A^m) \geq 1$. The spectral radius formula implies that $\rho(A^m) = \rho(A)^m$, so we must have $\rho(A) \geq 1$ too.

(b) In this case the structure theory of [20, §1.2] says that we can order the vertices so that A is upper triangular with 0s down the diagonal, and hence $\sigma(A) = \{0\}$.

(c) In this case A is a permutation matrix, and every eigenvalue is a root of unity.

(d) Since E is strongly connected, every vertex receives an edge. Since E is not a cycle, at least one vertex v receives 2 edges, e and f , say. There are paths from v to $s(e)$ and $s(f)$, and hence there are cycles μ and ν based at v such that $\mu_1 = e$ and $\nu_1 = f$. Now $\mu^{|\nu|}$ and $\nu^{|\mu|}$ are distinct elements of $vE^{|\mu||\nu|}v$, and we have $A^{|\mu||\nu|}(v, v) \geq 2$. With m as

in the proof of (a), $n := |\mu| |\nu| m$ satisfies $A^n(w, w) \geq 1$ for all w and $A^n(v, v) > 1$, and the arguments at the end of the proof of (a) show that $\rho(A) > 1$. \square

REFERENCES

- [1] T. Bates, J.H. Hong, I. Raeburn and W. Szymański, The ideal structure of the C^* -algebras of infinite graphs, *Illinois J. Math.* **46** (2002), 1159–1176.
- [2] O. Bratteli and D.W. Robinson, *Operator Algebras and Quantum Statistical Mechanics II*, second ed., Springer-Verlag, Berlin, 1997.
- [3] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields, and Motives*, Colloquium Publications, vol. 55, Amer. Math. Soc., Providence, 2008.
- [4] N. Dunford and J.T. Schwartz, *Linear Operators, Part I*, Interscience, New York, 1958.
- [5] M. Enomoto, M. Fujii and Y. Watatani, KMS states for gauge action on \mathcal{O}_A , *Math. Japon.* **29** (1984), 607–619.
- [6] R. Exel and M. Laca, Partial dynamical systems and the KMS condition, *Comm. Math. Phys.* **232** (2003), 223–277.
- [7] N.J. Fowler and I. Raeburn, The Toeplitz algebra of a Hilbert bimodule, *Indiana Univ. Math. J.* **48** (1999), 155–181.
- [8] T. Kajiwara and Y. Watatani, KMS states on finite-graph C^* -algebras, arXiv:1007.4248.
- [9] A. Kumjian and J. Renault, KMS states on C^* -algebras associated to expansive maps, *Proc. Amer. Math. Soc.* **134** (2006), 2067–2078.
- [10] A. Kumjian, D. Pask, I. Raeburn and J. Renault, Graphs, groupoids and Cuntz-Krieger algebras, *J. Funct. Anal.* **144** (1997), 505–541.
- [11] M. Laca and S. Neshveyev, KMS states of quasi-free dynamics on Pimsner algebras, *J. Funct. Anal.* **211** (2004), 457–482.
- [12] M. Laca and I. Raeburn, Phase transition on the Toeplitz algebra of the affine semigroup over the natural numbers, *Adv. Math.* **225** (2010), 643–688.
- [13] M. Laca, I. Raeburn and J. Ramagge, Phase transition on Exel crossed products associated to dilation matrices, *J. Funct. Anal.* **261** (2011), 3633–3664.
- [14] S. Neshveyev, KMS states on the C^* -algebras of non-principal groupoids, *J. Operator Theory*, to appear; arXiv:1106.5912.
- [15] D. Olesen and G.K. Pedersen, Some C^* -dynamical systems with a single KMS state, *Math. Scand.* **42** (1978), 111–118.
- [16] A.L.T. Paterson, Graph inverse semigroups, groupoids and their C^* -algebras, *J. Operator Theory* **48** (2002), 645–662.
- [17] G.K. Pedersen, *C^* -Algebras and their Automorphism Groups*, London Math. Soc. Monographs, vol. 14, Academic Press, London, 1979.
- [18] I. Raeburn, *Graph Algebras*, CBMS Regional Conference Series in Mathematics, vol. 103, Amer. Math. Soc., Providence, 2005.
- [19] J. Renault, *A Groupoid Approach to C^* -Algebras*, Lecture Notes in Math., vol. 793, Springer-Verlag, Berlin, 1980.
- [20] E. Seneta, *Non-Negative Matrices and Markov Chains*, second ed., Springer-Verlag, New York, 1981.

ASTRID AN HUEF AND IAIN RAEBURN, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTAGO, PO BOX 56, DUNEDIN 9054, NEW ZEALAND

E-mail address: astrid@maths.otago.ac.nz, iraeburn@maths.otago.ac.nz

MARCELO LACA, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA, VICTORIA, BC V8W 3P4, CANADA

E-mail address: laca@math.uvic.ca

AIDAN SIMS, SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF WOLLONGONG, NSW 2522, AUSTRALIA

E-mail address: asims@uow.edu.au