

2009

Stemofoline ethyl acetate solvate

Pitchaya Mungkornasawakul
University of Wollongong, pitchaya@uow.edu.au

Stephen G. Pyne
University of Wollongong, spyne@uow.edu.au

Alison T. Ung
alison_ung@uow.edu.au

Araya Jatisatienr
Chiang Mai University, Thailand

Anthony C. Willis
Australian National University

Follow this and additional works at: <https://ro.uow.edu.au/scipapers>



Part of the [Life Sciences Commons](#), [Physical Sciences and Mathematics Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Mungkornasawakul, Pitchaya; Pyne, Stephen G.; Ung, Alison T.; Jatisatienr, Araya; and Willis, Anthony C.: Stemofoline ethyl acetate solvate 2009.
<https://ro.uow.edu.au/scipapers/1138>

Stemofoline ethyl acetate solvate

Abstract

Crystals of the title compound, C₂₂H₂₉N₂O₅, ([systematic name: (2R,3R,5R,5aS,6R,8aR,9S)-(5Z)-5-[3-butyltetrahydro-6-methyl-2,5-methano-4,3,8a-[1]propanyl[3]ylidene-furo[3,2-j][1,4]oxazepin-7(5H)-ylidene]-4-methoxy-3-methylfuran-2(5H)-one ethyl acetate solvate) were isolated from the root extracts of *Stemona aphylla* (Stemonaceae). The structure closely resembles those of stemofoline derivatives which have previously been reported. Intermolecular contacts are observed between some C-bonded H atoms and nearby O atoms, perhaps indicating weak interactions which could influence the packing of species within the unit cell.

Keywords

CMMB

Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

Mungkornasawakul, P., Pyne, S. G., Ung, A. T., Jatisatienr, A. & Willis, A. C. (2009). Stemofoline ethyl acetate solvate. *Acta Crystallographica Section E: Structure Reports Online*, 65 (8), o1878-o1879.

Stemofoline ethyl acetate solvate

Pitchaya Mungkornasawakul,^a Stephen G. Pyne,^{a*}
Alison T. Ung,^b Araya Jatisatienr^c and Anthony C. Willis^d^aSchool of Chemistry, University of Wollongong, Wollongong, NSW 2522, Australia,^bDepartment of Chemistry and Forensic Science, University of Technology Sydney, PO Box 123, Broadway, Sydney, NSW 2007, Australia, ^cDepartment of Biology, Chang Mai University, Chang Mai 50202, Thailand, and ^dResearch School of Chemistry, The Australian National University, Canberra, ACT 0200, Australia

Correspondence e-mail: spyne@uow.edu.au, willis@rsc.anu.edu.au


Received 1 July 2009; accepted 9 July 2009

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.032; wR factor = 0.119; data-to-parameter ratio = 10.4.

Crystals of the title compound, $\text{C}_{22}\text{H}_{29}\text{NO}_5 \cdot \text{C}_4\text{H}_8\text{O}_2$, [systematic name: (2*R*,3*R*,5*R*,5*aS*,6*R*,8*aR*,9*S*)-(5*Z*)-5-[3-butyl-tetrahydro-6-methyl-2,5-methano-4,3,8a-[1]propanyl[3]ylidene-furo[3,2-*f*][1,4]oxazepin-7(5*H*)-ylidene]-4-methoxy-3-methylfuran-2(5*H*)-one ethyl acetate solvate] were isolated from the root extracts of *Stemona aphylla* (Stemonaceae). The structure closely resembles those of stemofoline derivatives which have previously been reported. Intermolecular contacts are observed between some C-bonded H atoms and nearby O atoms, perhaps indicating weak interactions which could influence the packing of species within the unit cell.

Related literature

For the single-crystal X-ray structure and absolute configuration of stemofoline as the hydrobromide monohydrate, see: Irie *et al.* (1970). For two stemofoline alkaloids with structural modifications in the butyl side chain, see: Seger *et al.* (2004). For the isolation of stemofoline from the root extracts of *Stemona aphylla* (Stemonaceae), see: Mungkornasawakul *et al.* (2009). For details of the weighting scheme used, see: Watkin (1994); Prince (1982).



Experimental

Crystal data

 $\text{C}_{22}\text{H}_{29}\text{NO}_5 \cdot \text{C}_4\text{H}_8\text{O}_2$ $M_r = 475.58$ Orthorhombic, $P2_12_12_1$ $a = 10.3908$ (1) Å $b = 10.6549$ (2) Å $c = 22.4143$ (4) Å $V = 2481.55$ (7) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.09$ mm⁻¹ $T = 200$ K $0.45 \times 0.08 \times 0.06$ mm

Data collection

Nonius KappaCCD diffractometer

Absorption correction: integration

via Gaussian method (Coppens,

1970) implemented in *maXus*(Mackay *et al.*, 2000) $T_{\min} = 0.976$, $T_{\max} = 0.996$

41745 measured reflections

3209 independent reflections

2149 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.060$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.032$ $wR(F^2) = 0.119$ $S = 0.91$

3208 reflections

307 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.30$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.25$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C17—H171 \cdots O4	0.97	2.33	3.046 (4)	130
C22—H222 \cdots O5 ⁱ	0.97	2.43	3.374 (4)	165
C1—H11 \cdots O6	0.98	2.66	3.485 (4)	142
C10—H101 \cdots O6	0.97	2.70	3.477 (3)	138

Symmetry code: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$

Data collection: *COLLECT* (Nonius, 2001); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *ORTEPII* (Johnson, 1976) in *TEXSAN* (Molecular Structure Corporation, 1997); software used to prepare material for publication: *CRYSTALS*.

We acknowledge the Australian Research Council and the Endeavour Fellowship Scheme for supporting this research.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2635).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
- Coppens, P. (1970). *The Evaluation of Absorption and Extinction in Single-Crystal Structure Analysis*, in *Crystallographic Computing*, edited by F. R. Ahmed, S. R. Hall & C. P. Huber, pp. 255–270. Copenhagen: Munksgaard.
- Irie, H., Masaki, N., Ohno, K., Osaki, K., Taga, T. & Uyco, S. (1970). *J. Chem. Soc. Chem. Commun.* p. 1066.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Mackay, S., Gilmore, C. J., Edwards, C., Stewart, N. & Shankland, K. (2000). *maXus*. Nonius BV, The Netherlands, MacScience, Japan, and The University of Glasgow, Scotland.
- Molecular Structure Corporation (1997). *TEXSAN*. MSC, The Woodlands, Texas, USA.
- Mungkornasawakul, P., Chaibong, S., Sastraruji, T., Jatisatienr, A., Jatisatienr, C., Pyne, S. G., Ung, A. T., Korth, J. & Lic, W. (2009). *J. Nat. Prod.* **72**, 848–851.

Nonius (2001). *COLLECT*. Nonius BV, Delft, The Netherlands.

Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.

Prince, E. (1982). In *Mathematical Techniques in Crystallography and Materials Science*. New York: Springer-Verlag.

Segcr, C., Merciter, K., Kalteneqger, E., Pacher, T., Greger, H. & Hofer, O. (2004). *Chem. Biodivers.* **1**, 265–279.

Watkin, D. (1994). *Acta Cryst.* **A50**, 411–437.

supplementary materials

Acta Cryst. (2009). E65, o1878-o1879 [doi:10.1107/S1600536809026889]

Stemofoline ethyl acetate solvate

P. Mungkornasawakul, S. G. Pyne, A. T. Ung, A. Jatisatienr and A. C. Willis

Comment

The crystallographic asymmetric unit consists of one $C_{22}H_{29}NO_5$ molecule and one ethyl acetate molecule of crystallization. Examination of distances and angles around the N atom shows N4—C5 to be slightly shorter than the other two distances and for C5—N4—C9a to be significantly larger than the other two angles. These features were also observed in the closely related compounds reported by Seger *et al.* (2004). Unfortunately, coordinates are not available for the structure of stemofoline hydrobromide hydrate (Irie *et al.*, 1970).

Experimental

Stemofoline was isolated from the root extracts of *Stemona aphylla* (Stemonaceae) as reported earlier (Mungkornasawakul *et al.*, 2009) and was crystallized from ethyl acetate/40–60°C petroleum ether. The sample had m.p. 71.5–73.2 °C (lit., Irie *et al.* (1970), 87–89 °C) and $[\alpha]_D^{23}$ 269.1 (c 0.18, $CHCl_3$) (lit., Irie *et al.* (1970), $[\alpha]_D$ 273 (MeOH)).

Refinement

The compound is enantiometrically pure but the anomalous dispersion terms are very low for all elements in the structure and so the absolute configuration can not be determined in this experiment. Consequently Friedel-pair reflections have been averaged and the Flack parameter has not been refined. The molecule is presented with the same absolute configuration as that determined by X-ray crystallography for a closely related compound (Seger *et al.*, 2004). It is also consistent with that reported for stemofoline.HBr.H₂O (Irie *et al.*, 1970), though the absolute structure determination performed there could possibly be regarded as not totally reliable.

All hydrogen atoms were observed in difference electron density maps prior to their inclusion in the structure. They were included at calculated positions and then were initially refined with soft restraints on the bond lengths and angles to regularize their geometry (C—H in the range 0.93–0.98 Å) and with $U_{iso}(H)$ in the range 1.2–1.5 times U_{eq} of the parent atom, after which the positions were refined with riding constraints.

One strong reflection with poor agreement between F_o and F_c was removed from the refinement as being unreliably measured.

The largest features in the final difference electron density map are located along C—C bonds.

supplementary materials

Figures




Fig. 1. The molecular structure of stemofoline ethyl acetate solvate, with the atom labelling scheme. Displacement ellipsoids for non-H atoms are drawn at the 30% probability level.



Fig. 2. Unit-cell packing diagram of stemofoline ethyl acetate solvate.

(2*R*,3*R*,5*R*,5*aS*, 6*R*,8*aR*,9*S*)-(5*Z*)-5-[3-butyltetrahydro-6-methyl- 2,5-methano-4,3,8*a*-[1]propanyl[3]ylidenefuro[3,2-*f*][1,4]oxazepin-7(5*H*)-ylidene]-4-methoxy-3-methylfuran-2(5*H*)-one ethyl acetate solvate

Crystal data

$C_{22}H_{29}NO_5 \cdot C_4H_8O_2$

$M_r = 475.58$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 10.3908$ (1) Å

$b = 10.6549$ (2) Å

$c = 22.4143$ (4) Å

$V = 2481.55$ (7) Å³

$Z = 4$

$F_{000} = 1024$

$D_x = 1.273$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 36203 reflections

$\theta = 2.6$ – 27.5°

$\mu = 0.09$ mm⁻¹

$T = 200$ K

Needle, colourless

$0.45 \times 0.08 \times 0.06$ mm

Data collection

Nonius KappaCCD diffractometer

Monochromator: graphite

$T = 200$ K

φ and ω scans

Absorption correction: integration via Gaussian method (Coppens, 1970) implemented in maXus (Mackay *et al.*, 2000)

$T_{\min} = 0.976$, $T_{\max} = 0.996$

41745 measured reflections

3209 independent reflections

2149 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.060$

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 2.6^\circ$

$h = -13 \rightarrow 13$

$l = -29 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

Method, part 1, Chebychev polynomial, (Watkin, 1994, Prince, 1982) [weight] = 1.0/[A₀*T₀(x) + A₁*T₁(x) ... + A_{n-1}]*T_{n-1}(x)]
 where A_i are the Chebychev coefficients listed below and x = F/Fmax Method = Robust Weighting (Prince, 1982) W = [weight] * [1-(deltaF/6*sigma*maF)²]² A_i are: 38.7 60.5 31.2 8.38
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.119$
 $S = 0.91$
 3208 reflections
 307 parameters
 Primary atom site location: structure-invariant direct methods
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$
 Extinction correction: None

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.43611 (16)	0.73200 (17)	0.47018 (7)	0.0301
O2	0.27062 (18)	0.79684 (16)	0.52966 (7)	0.0295
O3	0.30052 (17)	0.82685 (17)	0.64855 (8)	0.0315
O4	0.3242 (2)	0.49666 (18)	0.65754 (9)	0.0430
O5	0.3409 (2)	0.9075 (2)	0.73957 (9)	0.0469
O6	0.4158 (3)	0.2904 (2)	0.50427 (12)	0.0612
O7	0.4878 (2)	0.1608 (2)	0.57490 (9)	0.0427
N4	0.25605 (19)	0.6497 (2)	0.35879 (8)	0.0278
C1	0.4196 (3)	0.5479 (3)	0.40840 (11)	0.0331
C2	0.4559 (2)	0.6861 (2)	0.41002 (10)	0.0302
C3	0.3504 (2)	0.7525 (2)	0.37202 (10)	0.0287
C5	0.1296 (2)	0.7117 (3)	0.35629 (11)	0.0344
C6	0.1362 (3)	0.8327 (3)	0.39472 (11)	0.0355
C7	0.2744 (2)	0.8344 (2)	0.41800 (10)	0.0286
C8	0.2994 (2)	0.7520 (2)	0.47146 (10)	0.0265
C9	0.2300 (2)	0.6269 (2)	0.46769 (10)	0.0257
C9a	0.2729 (2)	0.5612 (2)	0.40984 (10)	0.0290
C10	0.2550 (2)	0.5713 (2)	0.53031 (10)	0.0268
C11	0.2737 (2)	0.6909 (2)	0.56580 (10)	0.0267
C12	0.2931 (2)	0.7068 (2)	0.62431 (11)	0.0280
C13	0.3204 (2)	0.6191 (3)	0.67198 (11)	0.0326
C14	0.3440 (3)	0.6821 (3)	0.72309 (11)	0.0347
C15	0.3298 (2)	0.8144 (3)	0.70898 (11)	0.0346
C16	0.3825 (3)	0.6425 (3)	0.78503 (12)	0.0495
C17	0.1423 (3)	0.4895 (3)	0.55107 (12)	0.0376
C18	0.3966 (3)	0.8207 (3)	0.31649 (11)	0.0357
C19	0.4653 (3)	0.7364 (3)	0.27164 (11)	0.0377
C20	0.5063 (3)	0.8045 (3)	0.21512 (12)	0.0452
C21	0.5627 (4)	0.7165 (4)	0.16876 (13)	0.0574
C22	0.3592 (4)	0.4074 (3)	0.70266 (16)	0.0556
C23	0.6093 (4)	0.3409 (4)	0.55692 (18)	0.0651
C24	0.4940 (3)	0.2645 (3)	0.54151 (14)	0.0416

supplementary materials

C25	0.3810 (3)	0.0762 (3)	0.56316 (15)	0.0440
C26	0.2641 (3)	0.1114 (3)	0.59907 (17)	0.0558
H11	0.4519	0.5013	0.4432	0.0396*
H12	0.4479	0.5081	0.3715	0.0395*
H21	0.5450	0.7060	0.3962	0.0365*
H911	0.2270	0.4811	0.4032	0.0338*
H51	0.0633	0.6534	0.3702	0.0410*
H52	0.1109	0.7355	0.3144	0.0413*
H61	0.0745	0.8283	0.4278	0.0416*
H62	0.1195	0.9067	0.3706	0.0420*
H71	0.3101	0.9183	0.4223	0.0337*
H91	0.1369	0.6445	0.4656	0.0306*
H101	0.3345	0.5243	0.5313	0.0325*
H161	0.3834	0.7165	0.8100	0.0738*
H162	0.3207	0.5819	0.8013	0.0742*
H163	0.4664	0.6037	0.7857	0.0738*
H171	0.1554	0.4602	0.5914	0.0551*
H172	0.0625	0.5382	0.5495	0.0547*
H173	0.1341	0.4163	0.5249	0.0551*
H181	0.4556	0.8879	0.3283	0.0420*
H182	0.3199	0.8577	0.2961	0.0419*
H191	0.5420	0.7010	0.2907	0.0449*
H192	0.4080	0.6653	0.2615	0.0455*
H201	0.5699	0.8696	0.2253	0.0540*
H202	0.4287	0.8439	0.1984	0.0544*
H211	0.5834	0.7639	0.1322	0.0857*
H212	0.6415	0.6757	0.1828	0.0859*
H213	0.4996	0.6521	0.1586	0.0860*
H221	0.3518	0.3231	0.6844	0.0821*
H222	0.4473	0.4219	0.7153	0.0819*
H223	0.3003	0.4131	0.7367	0.0818*
H231	0.6081	0.4193	0.5351	0.0959*
H232	0.6114	0.3581	0.5997	0.0961*
H233	0.6878	0.2943	0.5464	0.0963*
H251	0.4091	-0.0102	0.5747	0.0533*
H252	0.3601	0.0796	0.5197	0.0528*
H261	0.1959	0.0507	0.5919	0.0841*
H262	0.2843	0.1122	0.6421	0.0837*
H263	0.2342	0.1963	0.5874	0.0839*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0236 (7)	0.0391 (9)	0.0275 (8)	-0.0033 (7)	-0.0031 (7)	0.0015 (7)
O2	0.0370 (8)	0.0293 (8)	0.0223 (7)	-0.0010 (7)	-0.0001 (7)	0.0000 (7)
O3	0.0339 (9)	0.0344 (9)	0.0263 (8)	-0.0039 (7)	-0.0009 (7)	-0.0017 (7)
O4	0.0571 (12)	0.0361 (10)	0.0359 (9)	0.0023 (9)	-0.0096 (9)	0.0058 (8)
O5	0.0530 (12)	0.0514 (12)	0.0363 (10)	-0.0041 (10)	-0.0027 (9)	-0.0099 (9)

supplementary materials

O6	0.0657 (15)	0.0521 (14)	0.0658 (15)	0.0054 (12)	-0.0062 (13)	0.0190 (12)
O7	0.0420 (10)	0.0406 (10)	0.0456 (10)	-0.0051 (9)	-0.0037 (8)	0.0086 (9)
N4	0.0241 (9)	0.0348 (10)	0.0246 (9)	-0.0017 (8)	-0.0019 (7)	0.0005 (8)
C1	0.0330 (12)	0.0360 (13)	0.0304 (11)	0.0083 (10)	0.0025 (10)	0.0004 (10)
C2	0.0248 (11)	0.0404 (14)	0.0255 (10)	-0.0014 (10)	0.0004 (9)	0.0027 (10)
C3	0.0278 (11)	0.0322 (11)	0.0262 (10)	0.0006 (9)	-0.0008 (9)	0.0007 (10)
C5	0.0268 (11)	0.0450 (13)	0.0314 (11)	0.0020 (10)	-0.0069 (9)	-0.0007 (11)
C6	0.0355 (12)	0.0406 (13)	0.0303 (11)	0.0113 (11)	-0.0043 (10)	0.0025 (11)
C7	0.0318 (12)	0.0289 (11)	0.0253 (10)	-0.0009 (10)	-0.0016 (9)	0.0019 (10)
C8	0.0257 (10)	0.0286 (11)	0.0252 (10)	0.0002 (9)	-0.0017 (9)	-0.0030 (10)
C9	0.0238 (10)	0.0282 (11)	0.0250 (10)	-0.0008 (9)	-0.0009 (8)	0.0005 (9)
C9a	0.0304 (12)	0.0301 (11)	0.0265 (10)	-0.0016 (9)	0.0023 (9)	-0.0003 (9)
C10	0.0262 (10)	0.0275 (11)	0.0267 (10)	-0.0010 (9)	-0.0006 (9)	-0.0006 (9)
C11	0.0217 (10)	0.0307 (11)	0.0275 (10)	-0.0019 (9)	0.0012 (8)	0.0023 (9)
C12	0.0245 (10)	0.0326 (11)	0.0269 (11)	-0.0017 (9)	0.0009 (9)	-0.0013 (10)
C13	0.0274 (11)	0.0401 (13)	0.0304 (11)	-0.0012 (10)	-0.0006 (9)	0.0026 (10)
C14	0.0305 (12)	0.0492 (15)	0.0245 (10)	-0.0014 (11)	-0.0005 (9)	0.0041 (10)
C15	0.0278 (12)	0.0490 (15)	0.0271 (11)	-0.0034 (11)	-0.0020 (9)	-0.0010 (11)
C16	0.0594 (19)	0.0608 (19)	0.0281 (12)	-0.0049 (16)	-0.0091 (12)	0.0073 (13)
C17	0.0400 (14)	0.0379 (13)	0.0350 (12)	-0.0111 (11)	-0.0010 (11)	0.0053 (11)
C18	0.0366 (13)	0.0434 (14)	0.0270 (11)	-0.0039 (12)	0.0030 (10)	0.0071 (11)
C19	0.0356 (13)	0.0450 (15)	0.0325 (12)	-0.0036 (12)	0.0009 (10)	0.0062 (11)
C20	0.0440 (15)	0.0642 (18)	0.0275 (11)	-0.0026 (14)	0.0045 (11)	0.0076 (13)
C21	0.0618 (19)	0.076 (2)	0.0344 (14)	-0.0067 (18)	0.0077 (14)	-0.0012 (15)
C22	0.071 (2)	0.0446 (16)	0.0517 (17)	0.0004 (16)	-0.0158 (17)	0.0136 (14)
C23	0.073 (2)	0.0541 (19)	0.068 (2)	-0.0221 (19)	0.0059 (19)	-0.0021 (18)
C24	0.0468 (16)	0.0331 (13)	0.0451 (14)	0.0039 (12)	0.0067 (12)	-0.0018 (12)
C25	0.0415 (15)	0.0349 (13)	0.0558 (17)	-0.0041 (12)	-0.0001 (13)	0.0016 (13)
C26	0.0461 (17)	0.0518 (18)	0.070 (2)	-0.0022 (15)	0.0117 (16)	0.0055 (16)

Geometric parameters (Å, °)

O1—C2	1.449 (3)	C10—H101	0.966
O1—C8	1.436 (3)	C11—C12	1.338 (3)
O2—C8	1.421 (3)	C12—C13	1.447 (3)
O2—C11	1.389 (3)	C13—C14	1.350 (4)
O3—C12	1.392 (3)	C14—C15	1.452 (4)
O3—C15	1.394 (3)	C14—C16	1.505 (3)
O4—C13	1.345 (3)	C16—H161	0.967
O4—C22	1.435 (3)	C16—H162	0.980
O5—C15	1.212 (3)	C16—H163	0.965
O6—C24	1.197 (4)	C17—H171	0.965
O7—C24	1.336 (4)	C17—H172	0.978
O7—C25	1.453 (3)	C17—H173	0.980
N4—C3	1.499 (3)	C18—C19	1.525 (4)
N4—C5	1.471 (3)	C18—H181	0.980
N4—C9a	1.493 (3)	C18—H182	1.000
C1—C2	1.520 (4)	C19—C20	1.521 (4)
C1—C9a	1.531 (3)	C19—H191	0.980

supplementary materials

C1—H11	0.984	C19—H192	0.990
C1—H12	0.976	C20—C21	1.518 (5)
C2—C3	1.559 (3)	C20—H201	0.985
C2—H21	0.998	C20—H202	0.983
C3—C7	1.565 (3)	C21—H211	0.987
C3—C18	1.520 (3)	C21—H212	0.979
C5—C6	1.552 (4)	C21—H213	0.975
C5—H51	0.978	C22—H221	0.991
C5—H52	0.993	C22—H222	0.970
C6—C7	1.528 (3)	C22—H223	0.981
C6—H61	0.981	C23—C24	1.489 (5)
C6—H62	0.971	C23—H231	0.968
C7—C8	1.508 (3)	C23—H232	0.977
C7—H71	0.972	C23—H233	0.984
C8—C9	1.519 (3)	C25—C26	1.504 (4)
C9—C9a	1.539 (3)	C25—H251	1.000
C9—C10	1.545 (3)	C25—H252	0.999
C9—H91	0.986	C26—H261	0.973
C9a—H911	0.989	C26—H262	0.987
C10—C11	1.515 (3)	C26—H263	0.991
C10—C17	1.532 (3)		
O1...O2 ⁱ	3.489 (2)	O5...C22 ^{iv}	3.375 (5)
O2...C25 ⁱⁱ	3.277 (4)	O6...C10	3.477 (3)
O2...C2 ⁱⁱⁱ	3.543 (3)	O6...C1	3.485 (4)
O3...C26 ⁱⁱ	3.250 (4)	O7...C16 ^v	3.422 (3)
O3...C25 ⁱⁱ	3.379 (4)	C2...C11 ⁱ	3.594 (3)
O3...C5 ⁱ	3.446 (3)	C5...C15 ⁱⁱⁱ	3.453 (3)
C2—O1—C8	102.09 (17)	C11—C12—C13	132.2 (2)
C8—O2—C11	104.91 (18)	C12—C13—O4	117.0 (2)
C12—O3—C15	107.7 (2)	C12—C13—C14	109.9 (2)
C13—O4—C22	118.7 (2)	O4—C13—C14	133.0 (2)
C24—O7—C25	116.6 (2)	C13—C14—C15	106.2 (2)
C3—N4—C5	105.30 (19)	C13—C14—C16	133.7 (3)
C3—N4—C9a	103.48 (17)	C15—C14—C16	120.0 (2)
C5—N4—C9a	114.66 (18)	C14—C15—O3	109.1 (2)
C2—C1—C9a	99.0 (2)	C14—C15—O5	131.5 (2)
C2—C1—H11	112.7	O3—C15—O5	119.5 (3)
C9a—C1—H11	111.7	C14—C16—H161	107.8
C2—C1—H12	111.5	C14—C16—H162	110.6
C9a—C1—H12	111.0	H161—C16—H162	109.2
H11—C1—H12	110.5	C14—C16—H163	112.0
C1—C2—O1	108.29 (19)	H161—C16—H163	109.4
C1—C2—C3	104.6 (2)	H162—C16—H163	107.8
O1—C2—C3	104.78 (18)	C10—C17—H171	111.1
C1—C2—H21	115.4	C10—C17—H172	109.6
O1—C2—H21	110.4	H171—C17—H172	109.0
C3—C2—H21	112.8	C10—C17—H173	109.7

supplementary materials

C2—C3—N4	103.71 (19)	H171—C17—H173	108.3
C2—C3—C7	104.38 (18)	H172—C17—H173	109.1
N4—C3—C7	102.02 (18)	C3—C18—C19	114.0 (2)
C2—C3—C18	116.2 (2)	C3—C18—H181	109.0
N4—C3—C18	113.22 (19)	C19—C18—H181	108.5
C7—C3—C18	115.6 (2)	C3—C18—H182	108.1
N4—C5—C6	108.20 (19)	C19—C18—H182	107.7
N4—C5—H51	109.4	H181—C18—H182	109.5
C6—C5—H51	112.4	C18—C19—C20	113.5 (2)
N4—C5—H52	109.0	C18—C19—H191	108.6
C6—C5—H52	108.7	C20—C19—H191	108.6
H51—C5—H52	109.1	C18—C19—H192	108.7
C5—C6—C7	103.9 (2)	C20—C19—H192	110.0
C5—C6—H61	110.5	H191—C19—H192	107.1
C7—C6—H61	110.9	C19—C20—C21	112.6 (3)
C5—C6—H62	111.0	C19—C20—H201	109.3
C7—C6—H62	110.3	C21—C20—H201	109.6
H61—C6—H62	110.0	C19—C20—H202	106.9
C6—C7—C3	104.02 (19)	C21—C20—H202	108.7
C6—C7—C8	115.2 (2)	H201—C20—H202	109.7
C3—C7—C8	96.39 (18)	C20—C21—H211	109.6
C6—C7—H71	113.8	C20—C21—H212	112.1
C3—C7—H71	112.7	H211—C21—H212	108.2
C8—C7—H71	113.0	C20—C21—H213	109.6
C7—C8—O1	103.93 (19)	H211—C21—H213	108.2
C7—C8—O2	119.8 (2)	H212—C21—H213	108.9
O1—C8—O2	106.06 (17)	O4—C22—H221	106.9
C7—C8—C9	112.66 (18)	O4—C22—H222	109.8
O1—C8—C9	109.78 (19)	H221—C22—H222	109.7
O2—C8—C9	104.25 (18)	O4—C22—H223	110.5
C8—C9—C9a	107.95 (18)	H221—C22—H223	109.2
C8—C9—C10	101.88 (18)	H222—C22—H223	110.6
C9a—C9—C10	122.82 (19)	C24—C23—H231	110.1
C8—C9—H91	107.6	C24—C23—H232	110.4
C9a—C9—H91	109.3	H231—C23—H232	109.6
C10—C9—H91	106.4	C24—C23—H233	109.6
C9—C9a—C1	110.4 (2)	H231—C23—H233	108.9
C9—C9a—N4	108.94 (19)	H232—C23—H233	108.1
C1—C9a—N4	99.16 (19)	C23—C24—O7	111.2 (3)
C9—C9a—H911	112.3	C23—C24—O6	125.6 (3)
C1—C9a—H911	113.4	O7—C24—O6	123.3 (3)
N4—C9a—H911	111.9	O7—C25—C26	111.4 (3)
C9—C10—C11	100.13 (17)	O7—C25—H251	107.5
C9—C10—C17	111.4 (2)	C26—C25—H251	109.1
C11—C10—C17	114.6 (2)	O7—C25—H252	108.7
C9—C10—H101	111.4	C26—C25—H252	109.7
C11—C10—H101	108.3	H251—C25—H252	110.3
C17—C10—H101	110.6	C25—C26—H261	109.6
C10—C11—O2	111.99 (19)	C25—C26—H262	110.7

supplementary materials

C10—C11—C12	129.8 (2)	H261—C26—H262	108.8
O2—C11—C12	118.2 (2)	C25—C26—H263	109.8
O3—C12—C11	120.5 (2)	H261—C26—H263	109.5
O3—C12—C13	107.1 (2)	H262—C26—H263	108.4
O1—C2—C1—C9a	75.1 (2)	C2—C3—N4—C5	147.5 (2)
O1—C2—C3—N4	-107.2 (2)	C2—C3—N4—C9a	26.8 (2)
O1—C2—C3—C7	-0.7 (2)	C2—C3—C7—C6	-146.3 (2)
O1—C2—C3—C18	127.9 (2)	C2—C3—C7—C8	-28.3 (2)
O1—C8—O2—C11	-77.4 (2)	C2—C3—C18—C19	59.5 (3)
O1—C8—C7—C3	49.8 (2)	C3—N4—C5—C6	-25.4 (2)
O1—C8—C7—C6	158.7 (2)	C3—N4—C9a—C9	65.3 (2)
O1—C8—C9—C10	72.6 (2)	C3—C2—O1—C8	31.6 (2)
O1—C8—C9—C9a	-57.9 (2)	C3—C2—C1—C9a	-36.3 (2)
O2—C8—O1—C2	179.6 (2)	C3—C7—C6—C5	23.2 (2)
O2—C8—C7—C3	167.9 (2)	C3—C7—C8—C9	-68.9 (2)
O2—C8—C7—C6	-83.2 (3)	C3—C18—C19—C20	177.6 (2)
O2—C8—C9—C10	-40.6 (2)	C5—N4—C3—C7	39.3 (2)
O2—C8—C9—C9a	-171.2 (2)	C5—N4—C3—C18	-85.6 (2)
O2—C11—C10—C9	-4.1 (2)	C5—N4—C9a—C9	-48.8 (2)
O2—C11—C10—C17	-123.5 (2)	C5—C6—C7—C8	-81.0 (2)
O2—C11—C12—O3	4.6 (3)	C6—C5—N4—C9a	87.6 (2)
O2—C11—C12—C13	-169.4 (2)	C6—C7—C3—C18	84.6 (2)
O3—C12—C11—C10	-176.2 (2)	C6—C7—C8—C9	39.9 (3)
O3—C12—C13—O4	-177.0 (2)	C7—C3—N4—C9a	-81.4 (2)
O3—C12—C13—C14	-0.2 (3)	C7—C3—C18—C19	-177.6 (2)
O3—C15—C14—C13	-1.4 (3)	C7—C8—O2—C11	165.6 (2)
O3—C15—C14—C16	176.9 (2)	C7—C8—C9—C10	-172.1 (2)
O4—C13—C12—C11	-2.4 (4)	C7—C8—C9—C9a	57.4 (2)
O4—C13—C14—C15	177.0 (2)	C8—O2—C11—C10	-21.4 (2)
O4—C13—C14—C16	-0.9 (5)	C8—O2—C11—C12	157.9 (2)
O5—C15—O3—C12	-179.8 (2)	C8—C7—C3—C18	-157.3 (2)
O5—C15—C14—C13	179.9 (3)	C8—C9—C10—C11	26.1 (2)
O5—C15—C14—C16	-1.9 (4)	C8—C9—C10—C17	147.8 (2)
O6—C24—O7—C25	1.4 (4)	C9—C8—O2—C11	38.5 (2)
N4—C3—C2—C1	6.7 (2)	C9—C10—C11—C12	176.6 (2)
N4—C3—C7—C6	-38.7 (2)	C10—C11—C12—C13	9.8 (4)
N4—C3—C7—C8	79.4 (2)	C11—C10—C9—C9a	146.8 (2)
N4—C3—C18—C19	-60.4 (3)	C11—C12—O3—C15	-176.0 (2)
N4—C5—C6—C7	0.6 (3)	C11—C12—C13—C14	174.4 (3)
N4—C9a—C1—C2	52.8 (2)	C12—O3—C15—C14	1.2 (2)
N4—C9a—C9—C8	-51.8 (2)	C12—C11—C10—C17	57.2 (3)
N4—C9a—C9—C10	-169.7 (2)	C12—C13—O4—C22	176.0 (2)
C1—C2—O1—C8	-79.7 (2)	C12—C13—C14—C15	0.9 (3)
C1—C2—C3—C7	113.1 (2)	C12—C13—C14—C16	-177.0 (3)
C1—C2—C3—C18	-118.2 (2)	C13—C12—O3—C15	-0.6 (2)
C1—C9a—N4—C3	-50.1 (2)	C14—C13—O4—C22	0.1 (4)
C1—C9a—N4—C5	-164.2 (2)	C17—C10—C9—C9a	-91.5 (2)
C1—C9a—C9—C8	56.1 (2)	C18—C3—N4—C9a	153.7 (2)
C1—C9a—C9—C10	-61.8 (3)	C18—C19—C20—C21	-174.5 (3)

supplementary materials

C2—O1—C8—C7	-53.2 (2)	C23—C24—O7—C25	-178.5 (3)
C2—O1—C8—C9	67.5 (2)	C24—O7—C25—C26	-87.9 (3)
C2—C1—C9a—C9	-61.4 (2)		

Symmetry codes: (i) $x+1/2, -y+3/2, -z+1$; (ii) $x, y+1, z$; (iii) $x-1/2, -y+3/2, -z+1$; (iv) $-x+1, y+1/2, -z+3/2$; (v) $-x+1, y-1/2, -z+3/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C17—H171 \cdots O4	0.97	2.33	3.046 (4)	130
C22—H222 \cdots O5 ^y	0.97	2.43	3.374 (4)	165
C1—H11 \cdots O6	0.98	2.66	3.485 (4)	142
C10—H101 \cdots O6	0.97	2.70	3.477 (3)	138

Symmetry codes: (v) $-x+1, y-1/2, -z+3/2$.

Fig. 1




Fig. 2

