



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Illawarra Health and Medical Research Institute

Faculty of Science, Medicine and Health

2018

Predictors for misreporting sodium and potassium intakes by overweight and obese participants in a food-based clinical trial: implications for practice

Vivienne Guan

University of Wollongong, xg885@uowmail.edu.au

Yasmine Probst

University of Wollongong, yasmine@uow.edu.au

Elizabeth Neale

University of Wollongong, elizan@uow.edu.au

Linda C. Tapsell

University of Wollongong, ltapsell@uow.edu.au

Publication Details

Guan, V. X., Probst, Y. C., Neale, E. P. & Tapsell, L. C. (2018). Predictors for misreporting sodium and potassium intakes by overweight and obese participants in a food-based clinical trial: implications for practice. *European Journal of Clinical Nutrition*, Online First 1-5.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Predictors for misreporting sodium and potassium intakes by overweight and obese participants in a food-based clinical trial: implications for practice

Abstract

This study compared self-reported sodium and potassium intakes with urinary biomarkers and identified predictive factors. Secondary analysis of the 3-month intensive phase of the HealthTrack study with control (C), interdisciplinary intervention (I), intervention plus 30 g walnuts/day (IW) arms (n = 149). Dietary data was derived from diet history (DH) interviews and biomarker measures from urine. Urine-derived sodium (all, p = 0.000) and potassium (C: p = 0.011; I: p = 0.000; IW: p = 0.004) measures were significantly greater than self-reported intakes over the three months. Multiple linear regression showed body weight at baseline, body mass index (BMI) at baseline, and combined BMI at baseline and DH interviewer significantly negatively predicted the differences in sodium intake and excretion for C ($\beta = -21.226$, p = 0.016), I ($\beta = -106.140$, p = 0.002) and IW (F (9.530, 2df), p = 0.000), respectively. Where intakes of sodium and potassium are of interest in a trial, both reported intake and urinary biomarker measures are recommended.

Disciplines

Medicine and Health Sciences

Publication Details

Guan, V. X., Probst, Y. C., Neale, E. P. & Tapsell, L. C. (2018). Predictors for misreporting sodium and potassium intakes by overweight and obese participants in a food-based clinical trial: implications for practice. *European Journal of Clinical Nutrition*, Online First 1-5.

1 **Predictors for misreporting sodium and potassium intakes by overweight and obese**
2 **participants in a food-based clinical trial: Implications for practice**

3 **Running title:** Predictors for misreporting sodium and potassium intakes

4 Vivienne X Guan^{1,2}, Yasmine C Probst^{1,2}, Elizabeth P Neale^{1,2}, Linda C Tapsell^{1,2}

5 1. School of Medicine, Faculty of Science, Medicine and Health, University of Wollongong,
6 Wollongong, New South Wales 2522, Australia

7 2. Illawarra Health and Medical Research Institute, University of Wollongong, Wollongong,
8 New South Wales 2522, Australia

9 **Corresponding author**

10 Vivienne X Guan

11 Address: C/O University of Wollongong, Wollongong, New South Wales 2522, Australia

12 E-mail address: xg885@uowmail.edu.au

13 Tel: +61 2 4221 8169

14 Fax: +61 2 4221 4844

15 **Abstract**

16 This study compared self-reported sodium and potassium intakes with urinary biomarkers
17 and identified predictive factors. Secondary analysis of the 3-month intensive phase of the
18 HealthTrack study with control (*C*), interdisciplinary intervention (*I*), intervention plus 30g
19 walnuts/day (*IW*) arms (n=149). Dietary data was derived from diet history (DH) interviews
20 and biomarker measures from urine. Urine-derived sodium (all, p=0.000) and potassium (*C*:
21 p=0.011; *I*: p=0.000; *IW*: p=0.004) measures were significantly greater than self-reported
22 intakes over the three months. Multiple linear regression showed body weight at baseline,
23 body mass index (BMI) at baseline, and combined BMI and DH interviewer at baseline
24 significantly negatively predicted the difference in sodium intake and excretion for *C* ($\beta =$ -
25 21.226, p = 0.016), *I* ($\beta =$ -106.140, p = 0.002) and *IW* (F (9.530, 2df), p=0.000) ,
26 respectively. Where intakes of sodium and potassium are of interest in a trial, both reported
27 and urinary biomarker measures are recommended.

28

29 **Introduction**

30 Well-designed randomised controlled trials (RCTs) provide the highest level of evidence for
31 developing dietary recommendations in nutrition(1). Where food-based trials are undertaken,
32 the relationship between foods and nutrients needs to be considered to reflect the accuracy of
33 reported food intakes. Moreover, the measurement of self-reported dietary intake data
34 appears to be influenced by the dietary intervention itself. This intervention related bias may
35 impact on the interpretation of the findings(2, 3). The aim of this study was to compare
36 estimates of intake for dietary sodium and potassium with biomarker measures for urinary
37 excretion of sodium and potassium. A secondary aim was to determine factors associated
38 with differences between the measures in the context of a RCT for weight loss.

39 **Methods**

40 The present study is a secondary analysis of the 3-month intensive phase of the HealthTrack
41 study (ANZCTR #12614000581662) for weight loss, involving three arms: a control (*C*), a
42 novel interdisciplinary lifestyle intervention (*I*) and intervention plus 30g walnuts/day (*IW*).
43 All participants provided written informed consent. Detailed study protocols (4) and primary
44 results (5) are described elsewhere. Detailed methods of data handling are provided in the
45 **Supplementary Information**. In brief, dietary intakes reflecting usual weekly food
46 consumption over the previous three-month period was collected using a diet history (DH)
47 interview at baseline and 3 months. The plausibility of self-reported dietary intake has been
48 reported elsewhere(6). Overweight and obese participants (Body Mass Index (BMI) 25-
49 40kg/m²) collected 24-hour urine samples at baseline and 3 months. The incomplete urine
50 samples (<500mL or creatinine ratio outside 0.6-1.4) and participants taking diuretics were
51 excluded. Urine-derived intakes were calculated. The usual urine-derived intake of sodium
52 and potassium in the three month intensive phase of the HealthTrack was calculated using the

53 Multiple Source Method(7). Bland-Altman plots were developed for self-reported vs. urinary
54 measures and multiple linear regression used to determine predictive factors.

55 **Results**

56 A total of 149 participants were included in this analysis (**Supplementary Figure 1**). The
57 values of age and BMI were 44.2 ± 7.6 years and 32.0 ± 4.2 kg/m², respectively; and 69.1% of
58 participants were female. Participant data by study arms is presented in **Table 1**.

59 Median urine-derived sodium and potassium intakes were significantly greater than median
60 self-reported intakes in all arms at three months (sodium: *C*: $p=0.000$; *I*: $p=0.000$; *IW*:
61 $p=0.000$; potassium: *C*: $p=0.011$; *I*: $p=0.000$; *IW*: $p=0.004$). Correlations between self-
62 reported and urine-derived sodium measures were $r = 0.213$ ($p=0.140$) for *C*, $r=0.146$
63 ($p=0.340$) for *I* and $r=0.298$ ($p=0.026$) for *IW*. Correlations between potassium measures
64 were $r=0.188$ ($p=0.200$) for *C*, $r=0.597$ ($p=0.000$) for *I* and $r=0.350$ ($p=0.008$) for *IW*.

65 Bland-Altman plots demonstrated participants in all the arms generally underreported their
66 sodium and potassium intakes compared with urinary measures over the three months. The
67 range between the upper and lower limits of agreements for sodium and potassium intakes
68 was wide for all study arms. For sodium, in *I* the underestimation was greater at higher
69 intakes (**Figure 1A**), whereas for *IW* underestimation was less at higher intakes (**Figure 1B**).
70 The differences in estimated potassium intakes during the three months between self-reported
71 and urine-derived intakes were fairly consistent for all study arms.

72 The regression model established that body weight at baseline significantly predicted the
73 difference in sodium intakes during three months for *C* ($\beta=-21.226$, $t=-2.511$, $p=0.016$), but
74 BMI at baseline ($\beta=-106.140$, $t= -3.258$, $P = 0.002$) for *I*. BMI at baseline ($\beta=-0.017$, $t=-$
75 3.654 , $p=0.001$) and the DH interviewer ($\beta=-0.019$, $t=-2.382$, $p=0.021$) significantly predicted

76 the difference in sodium intakes between self-reported and urine-derived data at three months
77 for *IW*. The model was significant, ($F(9.530, 2df)$, $p=0.000$) explaining 26% variability of
78 the difference between sodium measurements for *IW*. There were no identified predictors for
79 the difference in potassium intakes between self-reported and urine-derived data for all arms.

80 **Discussion**

81 The present analysis was aimed to determine the specific observed bias related to intakes of
82 sodium and potassium. As both sodium and potassium are fairly ubiquitous in the food
83 supply, it was expected that participants in all study arms underreported intakes. The finding
84 for sodium intakes are consistent with other research (3), but in this study, the under-
85 reporting of potassium intakes compared to biomarker values was not anticipated (3). This
86 discrepancy may be attributed to differences in dietary advice in the intervention arms,
87 focused on reduced sodium intakes (3), versus energy reductions in the HealthTrack study
88 overall. Additionally, while potassium is mainly sourced from consumed foods, sodium is
89 also added to foods, particularly for processed and convenience foods. The results also
90 suggested that the intra-individual variances of urine-derived sodium intake were greater than
91 those for potassium intake. Thus a greater underreporting of sodium from DH interviews
92 compared to potassium is likely. This aligns with literature previously demonstrating the
93 challenges in estimation of sodium intake (8). It confirms that biomarkers and/or additional
94 dietary assessment tools may be required to strengthen the quality of data in food-based
95 RCTs if there is a concurrent interest in nutrients such as sodium and potassium.

96 The findings of this study also confirm that dietary interventions themselves may influence
97 accurate assessment of dietary intake data (2, 3), particularly using DH interviews. One of the
98 main activities of individualised dietary advice provided by dietitians is goal setting(9), and
99 in this case losing weight was the primary goal. Anecdotal evidence suggests that those who

100 did not achieve their weight loss goals required dietitians to ask more probing questions
101 during the DH interviews. Additionally, the proportions of female and overweight and obese
102 participants were high in the present sample, who tended to misreport dietary intake(10).
103 Thus, participants may be more conscious of their body weights measurements, which in turn
104 may influence their dietary intake reporting.

105 In conclusion, dietary advice needs to be provided in terms of foods but reporting of food
106 intake is inaccurate, especially with respect to sodium which can be added to food. Where
107 intakes of sodium and potassium are of interest in a trial, both reported measures and urinary
108 biomarkers are required to identify reported food sources, enhance advice provision and to
109 ensure the accuracy of nutrient intake estimates.

110 *Acknowledgements*

111 The authors would like to thank all the participants and the HealthTrack study team.

112 *Conflict of Interest Disclosure*

113 No potential conflict of interest was reported by the authors.

114 *Funding*

115 The HealthTrack study was funded by the Illawarra Health and Medical Research Institute
116 and California Walnut Commission. The funding bodies had no involvement in the data
117 analysis, data interpretation, or writing of the manuscript.

118 **References**

- 119 1. Weaver CM, Miller JW. Challenges in conducting clinical nutrition research.
120 Nutrition Reviews. 2017;75(7):491-9.
- 121 2. Natarajan L, Pu M, Fan J, Levine RA, Patterson RE, Thomson CA, et al.
122 Measurement Error of Dietary Self-Report in Intervention Trials. American Journal of
123 Epidemiology. 2010;172(7):819-27.

- 124 3. Espeland MA, Kumanyika S, Wilson AC, Wilcox S, Chao D, Bahnson J, et al.
125 Lifestyle Interventions Influence Relative Errors in Self-Reported Diet Intake of Sodium and
126 Potassium. *Annals of Epidemiology*. 2001;11(2):85-93.
- 127 4. Tapsell LC, Lonergan M, Martin A, Batterham MJ, Neale EP. Interdisciplinary
128 lifestyle intervention for weight management in a community population (HealthTrack
129 study): Study design and baseline sample characteristics. *Contemporary Clinical Trials*.
130 2015;45, Part B:394-403.
- 131 5. Tapsell LC, Lonergan M, Batterham MJ, Neale EP, Martin A, Thorne R, et al. Effect
132 of interdisciplinary care on weight loss: a randomised controlled trial. *BMJ Open*. 2017;7(7).
- 133 6. Hocking R, Neale E, Tapsell L, Probst Y. The impact of dietary assessment methods
134 on measurement of macronutrient intake in overweight adults: a secondary analysis of the
135 HealthTrack study (in preparation).
- 136 7. Harttig U, Haubrock J, Knuppel S, Boeing H. The MSM program: web-based
137 statistics package for estimating usual dietary intake using the Multiple Source Method.
138 *European Journal of Clinical Nutrition*. 2011;65(S1):S87-S91.
- 139 8. Cogswell ME, Maalouf J, Elliott P, Loria CM, Patel S, Bowman BA. Use of Urine
140 Biomarkers to Assess Sodium Intake: Challenges and Opportunities. *Annu Rev Nutr*.
141 2015;35(1):349-87.
- 142 9. Holli BB, Beto JA. *Nutrition Counseling and Education Skills for Dietetics*
143 *Professionals*, 6th edition. US: Lippincott Williams and Wilkins 2012.
- 144 10. Scagliusi FB, Ferriolli E, Pfrimer K, Laureano C, Cunha CSF, Gualano B, et al.
145 Characteristics of women who frequently under report their energy intake: a doubly labeled
146 water study. *European Journal of Clinical Nutrition*. 2009;63(10):1192-9.

147 ***Figure legends***

148 Figure 1A: Bland-Altman plot for assessing bias between self-reported dietary intake and
149 urine derived intake for sodium intake for intervention arm during 3 months of intensive
150 phase of the HealthTrack study (secondary analysis cohort)

151 Figure 1B: Bland-Altman plot for assessing bias between self-reported dietary intake and
152 urine derived intake for Log10 value of sodium intake for intervention + walnut arm during 3
153 months of intensive phase of the HealthTrack study (secondary analysis cohort)

154

155 **Table 1: Demographic characteristics, dietary nutrient intakes and biomarkers by arms during 3 months of intensive phase of the**
 156 **HealthTrack study (secondary analysis cohort)**

	Control (n=48)	Intervention (n=45)	Intervention + Walnuts (n=56)	p value
Age (y) ¹	44.5±7.3	45.8±6.5	42.8±8.5	0.128 ²
Gender n (%)				0.893 ³
Male	16 (33.3)	13 (28.9)	17 (30.4)	
Female	32 (66.7)	32 (71.1)	39 (69.6)	
Levels of education n (%)				0.118 ³
No school certificate or other qualifications	3 (6.3)	2 (4.4)	0 (0)	
School or intermediate certificate (or equivalent)	0 (0)	5 (11.1)	1 (1.8)	
Higher school or leaving certificate (or equivalent)	3 (6.3)	2 (4.4)	4 (7.1)	

Trade/apprenticeship (e.g. hairdresser, chef)	4 (8.3)	5 (11.1)	3 (5.4)	
Certificate/diploma (e.g. child care, technician)	13 (27.1)	6 (13.3)	18 (32.1)	
University degree	10 (20.8)	11 (24.4)	20 (35.7)	
Post graduate degree	15 (31.3)	14 (31.3)	10 (17.9)	
Body weight (kg)¹				
At baseline	90.4±16.8	91.0±15.9	90.2±15.7	0.963 ²
At 3 months	89.1±16.7	87.8±15.8	87.4±15.0	0.848 ²
BMI (kg/m²)¹				
At baseline	32.3±4.2	32.1±4.1	31.8±4.3	0.888 ²
At 3 months	31.8±4.2	30.9±4.3	30.9±4.3	0.515 ²
Waist circumference (cm)¹				
At baseline	104.0±13.6	102.8±11.1	102.3±12.2	0.776 ²
At 3 months	101.3±14.0	99.2±12.5	98.2±11.9	0.466 ²
Energy intake (kJ/d)⁴				

At baseline	8612.4 (7152.4 – 11021.7)	9165.0 (7820.4 – 11636.6)	9208.7 (7337.7 – 10720.4)	0.510 ⁵
During 3 months	7337.4 (6660.9 – 8972.0)	6635.6 (5933.0 – 7614.3)	7252.5 (6310.4 – 8821.1)	0.014 ⁵
Estimated energy requirement(kJ/d)⁴	8229.7 (7243.3 – 9600.4)	8285.4 (7479.7 – 9446.5)	8276.8 (7544.0 – 9818.6)	0.946 ⁵
Dietary sodium intake (mg/d)⁴				
At baseline	2008.6 (1557.7 – 2754.1)	2587.6 (1990.2 – 3014.4)	2634.9 (1964.5 – 3200.3)	0.120 ⁵
During 3 months	2060.4 (1566.5 – 2464.2)	1717.7 (1393.6 – 2064.9)	1703.1 (1418.8 – 2182.6)	0.054 ⁵
Dietary potassium intake (mg/d)⁴				
At baseline	3792.4 (3125.3 – 4666.6)	3821.0 (3304.7 – 4569.9)	3684.1 (3077.7 – 4317.0)	0.464 ⁵
During 3 months	3608.5 (2979.9 – 4224.0)	3320.2 (2990.2 – 3741.6)	3578.2 (3019.4 – 4153.1)	0.336 ⁵
Urinary sodium excretion (mg/d)⁴				
At baseline	3757.6 (2701.2 – 4981.1)	3450.0 (2527.3 – 4573.3)	4011.6 (3162.5 – 5141.6)	0.223 ⁵
At 3 months	3182.6 (2300.0 – 4366.0)	3102.3 (2326.7 – 4787.2)	3035.5 (2139.5 – 3744.2)	0.431 ⁵
Urine-derived sodium intake during 3 months (mg/d)^{4,6}	3759.0 (3142.2 – 4176.6)	3551.4 (2999.8 – 4186.0)	3653.2 (3211.2 – 4151.6)	0.819 ⁵
Intra-individual variance(mg/d) ⁶	3.60	1.37	0.34	-
Urinary potassium excretion (mg/d)⁴				

At baseline	4153.2 (3013.6 – 5356.2)	3748.1 (3393.5 – 4533.1)	4001.3 (2773.1 – 5267.5)	0.588 ⁵
At 3 month	3570.8 (3152.9 – 4457.1)	3798.7 (3064.3 – 4406.5)	4077.3 (2139.5 – 3744.2)	0.779 ⁵
Urine-derived potassium intake during 3 months (mg/d)^{4,6}	4015.3 (3415.2 – 4642.0)	3962.3 (3501.4 – 4388.4)	3803.6 (3322.8 – 4658.5)	0.858 ⁵
Intra-individual variance(mg/d) ⁶	0.12	0.01	0.07	-

157 ¹ Data presented as mean ± SD

158 ² Derived by one-way analysis of variance between arms

159 ³ Derived by Pearson's chi-square test for differences in portions between arms

160 ⁴ Data presented as median (interquartile range)

161 ⁵ Derived by Kruskal-Wallis H test between arms

162 ⁶ Derived from 24-h urinary data calculated by the Multiple Source Method(7), by computing 24-h urinary data at baseline and at 3 months

163

164




Figure 1A: Bland-Altman plot for assessing bias between self-reported dietary intake and urine derived intake for sodium intake for intervention arm during 3 months of intensive phase of the HealthTrack study (secondary analysis cohort) ¹.

¹ Solid line represented the mean difference; dotted lines represented upper and lower limits of agreement (mean \pm 1.96 SDs)




Figure 1B: Bland-Altman plot for assessing bias between self-reported dietary intake and urine derived intake for Log10 value of sodium intake for intervention + walnut arm during 3 months of intensive phase of the HealthTrack study (secondary analysis cohort)¹.

¹ Solid line represented the mean difference; dotted lines represented upper and lower limits of agreement (mean \pm 1.96 SDs)

Supplementary information - Detailed methods

The HealthTrack study

The present study is a secondary analysis of the 3-month intensive phase of the HealthTrack study. Detailed study protocols (4) and primary results (5) are described elsewhere. In brief, the HealthTrack study was a 12-month randomised controlled trial (Australian and New Zealand Clinical Trial Registry, ANZCTR N 12614000581662). Ethics approval for the HealthTrack study was obtained from the University of Wollongong/Illawarra Shoalhaven Local Health District Human Research Ethics Committee (HE13/189). The aim of the trial was to test the effects of a novel interdisciplinary lifestyle intervention delivered by a dietitian compared with usual care delivered by a nurse. The primary outcomes was weight loss in overweight and obese adults. There were three arms involved in the study, a control arm providing usual care based on the Australian Guide to Health Eating (C) (11), an intervention arm receiving interdisciplinary intervention delivered by individualised dietary advice (I) and a third arm receiving the intervention plus 30g walnuts per day (IW). A computer-generated randomisation sequence was applied for randomization by an independent investigator. Participants and DH interviewers were blinded to the randomised groups. Individualised dietary advice in the intervention groups was achieved by personalised target requirements and participants' usual food habits. Estimated energy requirements were calculated using the Mifflin-St Jeor equation (12). Recruited participants were aged 25-54 years, with a Body Mass Index (BMI) of 25-40kg/m², and residents of the Illawarra region. Participants who were unable to communicate in the English language, had an impaired ability to participate in the study, other medical conditions thought to limit survival to 1 year, suffered from immunodeficiency, reported illegal drug use or regular alcohol intake associated with alcoholism (>50 g/day), or difficulties or major impediments to participating in components of the study were excluded. Participants' level of education was self-reported

during a screening survey. Body weight, height, waist circumference were measured by using standard procedures (4). BMI was also calculated.

Urine derived dietary intake

Participants were asked to collect 24-hour urine samples to assess sodium and potassium excretion at baseline and 3 months. Detailed instructions were provided by Accredited Practising Dietitians (APDs) and standard plastic containers were distributed to all participants. Participants were instructed to discard the first urine of the day and collect the rest over a 24 hour period. The collected samples were stored at 2°C to 8°C by Southern IML Pathology. The total volume of urine was measured, recorded and indirect ion-specific electrodes were used to determine sodium and potassium concentrations. The Jaffe reaction colorimetric method was applied to determine creatinine concentration (13). Participants who were taking diuretics and those with incomplete urine samples were excluded from analyses. Total urine volume of a sample less than 500 mL (14) and/or a creatinine ratio out of range of 0.6-1.4(15) was used to identify incomplete urine samples. The creatinine ratio was calculated as observed creatinine excretion divided by expected creatinine excretion, with expected creatinine excretion estimated using the Joossens & Geboers algorithm (expected creatinine excretion = body weight (kg) * 24 for males or 21 for females) (15). In the case that body weight was not collected, the sample was discarded if the creatinine of the sample was ≤ 6.0 mmol/day and the urine volume was < 1000 mL/day(16). The overall recovery rates of urinary sodium and potassium are reported to be 86% and 77%, respectively (17, 18). Thus, the 24-hour urine-derived intakes of sodium and potassium were the result of 24-hour urinary sodium and potassium divided by 0.86 and 0.77, respectively.

Self-reported dietary intakes

Dietary intake data reflecting usual weekly food consumption at baseline and at three months was collected using a DH interview performed by APDs, conducted between May 2014 and July 2015. The dietary intake data collection protocol was validated previously (19).

Participants were asked to recall their usual intake over the past three-month period at the three month time point. A food checklist was used to assess commonly omitted food items during the interviews. Collected food intake data was coded to FoodWorks Professional nutrient analysis software (Xyris, Springhill QLD, Australia, Version 7, 2007) supported by the AUSNUT 2007 food composition database (20). The plausibility of self-reported dietary intake has been examined and reported elsewhere(6).

Statistical methods

The usual urine-derived intake of sodium and potassium in the three month intensive phase of the HealthTrack were calculated by using the Multiple Source Method (7). To account for within-person variability of intakes, the 24-h urine-derived intakes of sodium and potassium at baseline and three months were computed. Statistical analyses were performed using IBM SPSS Statistics (version 21.0, IBM Corp, Chicago IL, 2012). Normally distributed data was presented as mean and standard deviation and as median and interquartile range for skewed data. To assess difference in demographic characteristics, dietary nutrient intakes and biomarkers between study arms, one-way analysis of variance was used for normally distributed data and Kruskal-Wallis H test for nonparametric data. The post-hoc test was used to identify where the differences were within arms. A Pearson's chi-square test was applied to compare differences in the proportion of gender, levels of education and DH interviewers between arms. The Wilcoxon signed rank test was conducted to assess the difference in self-reported dietary intake and urine-derived intake of sodium and potassium between baseline and three months in each arm, and Spearman's correlation was used to examine the relationship between self-reported dietary intake and urine-derived intake of sodium and


potassium. Bland-Altman plots (21) were used to investigate relative agreement between self-reported dietary intakes and urine-derived intakes by plotting the difference of intake of each participant against the mean of both measurements. The 95% limits of agreement in Bland-Altman plots were calculated by the mean difference $\pm 1.96 * SD$. Stepwise multiple linear regression was used to identify the predictors of the difference between dietary and urinary sodium and potassium measures. Covariates included age (y), gender, levels of education, body weight at baseline (kg), BMI at baseline (kg/m^2), waist circumference at baseline (cm), body weight change at 3 months (kg) and DH interviewers who conducted DH interview at 3 months. Normality, linearity, multicollinearity and homoscedasticity of the identified models were assessed to ensure there were no violations of assumptions. The distribution of differences between self-reported and urine-derived intakes of sodium and potassium were estimated prior to performing Bland-Altman plots and stepwise multiple linear regression analysis using the Shapiro-Wilk test. A logarithmic transformation of original data was conducted for differences that were not normally distributed for the Bland-Altman plots and stepwise multiple linear regression analysis. Statistical significance was determined as a two-sided P values less than 0.05.

References

1. Weaver CM, Miller JW. Challenges in conducting clinical nutrition research. *Nutrition Reviews*. 2017;75(7):491-9.
2. Natarajan L, Pu M, Fan J, Levine RA, Patterson RE, Thomson CA, et al. Measurement Error of Dietary Self-Report in Intervention Trials. *American Journal of Epidemiology*. 2010;172(7):819-27.
3. Espeland MA, Kumanyika S, Wilson AC, Wilcox S, Chao D, Bahnson J, et al. Lifestyle Interventions Influence Relative Errors in Self-Reported Diet Intake of Sodium and Potassium. *Annals of Epidemiology*. 2001;11(2):85-93.

4. Tapsell LC, Lonergan M, Martin A, Batterham MJ, Neale EP. Interdisciplinary lifestyle intervention for weight management in a community population (HealthTrack study): Study design and baseline sample characteristics. *Contemporary Clinical Trials*. 2015;45, Part B:394-403.
5. Tapsell LC, Lonergan M, Batterham MJ, Neale EP, Martin A, Thorne R, et al. Effect of interdisciplinary care on weight loss: a randomised controlled trial. *BMJ Open*. 2017;7(7).
6. Hocking R, Neale E, Tapsell L, Probst Y. The impact of dietary assessment methods on measurement of macronutrient intake in overweight adults: a secondary analysis of the HealthTrack study (in preparation).
7. Harttig U, Haubrock J, Knuppel S, Boeing H. The MSM program: web-based statistics package for estimating usual dietary intake using the Multiple Source Method. *European Journal of Clinical Nutrition*. 2011;65(S1):S87-S91.
8. Cogswell ME, Maalouf J, Elliott P, Loria CM, Patel S, Bowman BA. Use of Urine Biomarkers to Assess Sodium Intake: Challenges and Opportunities. *Annu Rev Nutr*. 2015;35(1):349-87.
9. Holli BB, Beto JA. *Nutrition Counseling and Education Skills for Dietetics Professionals*, 6th edition. US: Lippincott Williams and Wilkins 2012.
10. Scagliusi FB, Ferriolli E, Pfrimer K, Laureano C, Cunha CSF, Gualano B, et al. Characteristics of women who frequently under report their energy intake: a doubly labeled water study. *European Journal of Clinical Nutrition*. 2009;63(10):1192-9.
11. National Health and Medical Research Council. *Australian Dietary Guidelines*. Canberra: NHMRC; 2013.
12. Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive equation for resting energy expenditure in healthy individuals. *The American Journal of Clinical Nutrition*. 1990;51(2):241-7.

13. Bonsnes RW, Taussky HH. On the colorimetric determination of creatinine by the jaffe reaction. *J Biol Chem.* 1945;158(3):581-91.
14. Wang C-Y, Cogswell ME, Loria CM, Chen T-C, Pfeiffer CM, Swanson CA, et al. Urinary Excretion of Sodium, Potassium, and Chloride, but Not Iodine, Varies by Timing of Collection in a 24-Hour Calibration Study. *The Journal of Nutrition.* 2013;143(8):1276-82.
15. Joossens JV, Geboers J. Monitoring salt intake of the population: methodological considerations. In: GG De Backer, HT Pedoe, P Ducimetiere, editors. *Surveillance of the Dietary Habits of the Population with Regard to Cardiovascular Diseases, EURONUT Report 2.* Wageningen: Department of Human Nutrition, Agricultural University; 1984. p. 61–73.
16. Reinivuo H, Valsta LM, Laatikainen T, Tuomilehto J, Pietinen P. Sodium in the Finnish diet: II Trends in dietary sodium intake and comparison between intake and 24-h excretion of sodium. *European Journal of Clinical Nutrition.* 2006;60(10):1160-7.
17. Holbrook JT, Patterson KY, Bodner JE, Douglas LW, Veillon C, Kelsay JL, et al. Sodium and potassium intake and balance in adults consuming self-selected diets. *The American Journal of Clinical Nutrition.* 1984;40(4):786-93.
18. Tasevska N, Runswick SA, Bingham SA. Urinary Potassium Is as Reliable as Urinary Nitrogen for Use as a Recovery Biomarker in Dietary Studies of Free Living Individuals. *The Journal of Nutrition.* 2006;136(5):1334-40.
19. Martin GS, Tapsell LC, Denmeade S, Batterham MJ. Relative validity of a diet history interview in an intervention trial manipulating dietary fat in the management of Type II diabetes mellitus. *Preventive Medicine.* 2003;36(4):420-8.
20. Food Standards Australia New Zealand. *AUSNUT 2007* Canberra: FSANZ; 2010.
21. Martin Bland J, Altman D. Statistical methods for assessing agreement between two methods of clinical measurement. *The Lancet.* 1986;327(8476):307-10.



Supplementary Figure1: Participant flow of urinary collection and analyses in the HealthTrack study