

2019

What Do Individuals With Borderline Personality Disorder Want From Treatment? A Study of Self-generated Treatment and Recovery Goals

Fiona Y. Ng

University of Wollongong, fionan@uow.edu.au

Phoebe Carter

University of Wollongong, pcarter@uow.edu.au

Marianne E. Bourke

University of Wollongong, mbourke@uow.edu.au

Brin F. S. Grenyer

University of Wollongong, grenyer@uow.edu.au

Publication Details

Ng, F. Y. Y., Carter, P. E., Bourke, M. E. & Grenyer, B. F. S. (2019). What Do Individuals With Borderline Personality Disorder Want From Treatment? A Study of Self-generated Treatment and Recovery Goals. *Journal of Psychiatric Practice*, 25 (2), 148-155.

What Do Individuals With Borderline Personality Disorder Want From Treatment? A Study of Self-generated Treatment and Recovery Goals

Abstract

Outcome measurement has progressed in the field of personality disorders. While the majority of trials have evaluated outcomes on the basis of symptom and diagnostic indices, what is considered a meaningful and valued outcome to individuals has seldom been investigated. Self-generated treatment goals were collected from 102 individuals seeking treatment for borderline personality disorder and independently coded by 2 raters. Responses were content-analyzed to determine the categories of goals people want for treatment. A total of 464 individual goal units across 4 main goal types emerged in the content analysis: reducing symptoms, improved well-being, better interpersonal relationships, and having a greater sense of self. Although the reduction of symptoms was the most commonly reported goal, 88.2% reported wanting better psychosocial functioning, including improvements in relationships, vocation, and self-understanding. The existence of the wide range of goals suggests that there is a need for clinicians to establish a collaborative formulation of treatment goals with individuals to ensure that treatment is personalized and meaningful.

Disciplines

Medicine and Health Sciences

Publication Details

Ng, F. Y. Y., Carter, P. E., Bourke, M. E. & Grenyer, B. F. S. (2019). What Do Individuals With Borderline Personality Disorder Want From Treatment? A Study of Self-generated Treatment and Recovery Goals. *Journal of Psychiatric Practice*, 25 (2), 148-155.

1 **VERSION ACCEPTED FOR PUBLICATION**

2 **CITATION:**

3 Ng, F.Y.Y., Carter, P.E., Bourke, M.E., & Grenyer, B.F.S. (2019). What Do Individuals With
4 Borderline Personality Disorder Want From Treatment? A Study of Self-generated Treatment
5 and Recovery Goals. *Journal of Psychiatric Practice*, 25(2), 148-155. doi:
6 10.1097/PRA.0000000000000369
7 [https://journals.lww.com/practicalpsychiatry/Abstract/2019/03000/What_Do_Individuals_With_](https://journals.lww.com/practicalpsychiatry/Abstract/2019/03000/What_Do_Individuals_With_Borderline_Personality.12.aspx)
8 [Borderline_Personality.12.aspx](https://journals.lww.com/practicalpsychiatry/Abstract/2019/03000/What_Do_Individuals_With_Borderline_Personality.12.aspx)

9
10 **Practitioner's Corner**

11 **What Do Individuals With Borderline Personality Disorder Want From Treatment? A**
12 **Study of Self-generated Treatment and Recovery Goals**

13 Fiona Y.Y. Ng, BHltSci(Hons), Phoebe Carter, DPsyc, Marianne E. Bourke, PhD, Brin FS
14 Grenyer, PhD

15 NG, CARTER, BOURKE, and GRENYER: School of Psychology and Illawarra Health and
16 Medical Research Institute, University of Wollongong, Australia

17 Please send correspondence to: Professor Brin FS Grenyer, Building 41, Room 137, University
18 of Wollongong, NSW 2522 Wollongong, Australia (email: grenyer@uow.edu.au).

19 This research was conducted with the support of the Australian Government Research Training
20 Program Scholarship awarded to FYYN, and NSW Ministry of Health support to the Project Air
21 Strategy for Personality Disorders.

22
23 The authors declare no conflicts of interest.

Abstract

Outcome measurement has progressed in the field of personality disorders. While the majority of trials have evaluated outcomes on the basis of symptom and diagnostic indices, what is considered a meaningful and valued outcome to individuals has been seldom investigated. Self-generated treatment goals were collected from 102 individuals seeking treatment for borderline personality disorder (BPD) and independently coded by two raters. Responses were content-analysed to determine the categories of goals people want for treatment. A total of 464 individual goal units across 4 main goal types emerged in the content analysis: reducing symptoms, improved wellbeing, better interpersonal relationships, and having a greater sense of self. Although the reduction of symptoms was the most commonly reported goal, 88.2% reported wanting better psychosocial functioning, including improvements in relationships, vocation, and self-understanding. The existence of the wide range of goals suggests that there is a need for clinicians to establish a collaborative formulation of treatment goals with individuals to ensure treatment is personalized and meaningful.

KEY WORDS: borderline personality disorder, treatment goals, recovery, qualitative, personalized treatment

1 The examination of outcomes in the field of personality disorders has progressed since the first
2 randomized control trial (RCT) published in 1991.¹ In a recent systematic review, 33 RCTs were
3 identified that were designed to evaluate the efficacy of interventions for people with borderline
4 personality disorder (BPD).² Specialized interventions for BPD have treatment goals that target changes
5 in behavior, such as in dialectical behavior therapy¹ or the specific modification of representations and
6 understanding of self and other, for example, in schema therapy, transference focused psychotherapy, and
7 mentalization based treatment.³⁻⁵ However, measures used in intervention trials usually measure only the
8 key symptoms and service use. Consumer reports suggest that we need to go beyond symptom change⁶
9 and measure a broader set of recovery goals. This has been supported by the literature, which has reported
10 a disconnect between service targets and personal goals of individuals with BPD,⁶ and the recognition that
11 recovery extends beyond symptom remission.⁷

12 Given the international shift toward recovery-oriented mental health servicing and the provision
13 of person-centred care,⁸ questions remain concerning what individuals perceive to be important to them at
14 the start of treatment. Various attempts to personalize treatment and focus on service user generated goals
15 in other diagnoses have been made. The Camberwell Assessment of Need (CAN)⁹ is one example and
16 measures the met and unmet needs of individuals across 22 health and social domains. The aims of the
17 CAN diverge from conventional clinical assessment, as it differentiates between the met needs (ie, met
18 through the provision of services) and unmet needs, which are areas identified as requiring further
19 intervention or support. Limited research has been conducted into what individuals with BPD value. In a
20 study examining the met and unmet needs of people with personality disorders, 8 key areas of unmet need
21 were identified: “self-care, psychotic symptoms, psychological distress, risk to self, risk to others, alcohol
22 use, sexual expression and budgeting.”^{10, p541} While the majority of these unmet needs reflect the
23 symptomatic difficulties known to be experienced by individuals with personality disorder, this finding
24 also provides an indication of the domains that require greater investigation.

25 One approach to personalizing treatment and focusing on the goals generated by service users has
26 been through understanding the target complaints of individuals at the start of therapy. Measures such as
27 Battle’s Target Complaints Measure¹¹ provide an opportunity for individuals to spontaneously formulate
28 and identify their own goals to guide the direction of therapy. The use of personalized treatment goals has
29 been identified as producing larger effect sizes than symptom checklists when evaluating the
30 effectiveness of psychotherapy in clinical trials.¹² The identification of specific categories of goals that
31 individuals value may be important in understanding treatment needs and developing new ways of
32 personalizing treatment. The goal of this study was to examine the personally meaningful treatment goals
33 of individuals seeking treatment for BPD.

METHOD

Study Design and Participants

This qualitative study utilized data collected from individuals who were seeking treatment for BPD at a community-based psychotherapy program. Individuals were assessed for suitability for the program and were only admitted if they were over 18 years of age and had a primary diagnosis of DSM-IV BPD, diagnosed using the Structured Clinical Interview for DSM (SCID-I and SCID-II)^{13,14} by 2 trained doctoral level clinical psychologists. Individuals were excluded from the program if there was indication of substance abuse, or they met criteria for a primary diagnosis of schizophrenia, schizoaffective disorder, bipolar disorder, major depressive disorder with psychotic features, or a history of neurological disorder. All participants were fluent in English and gave explicit informed written consent (including consent for the audio recording of clinical assessments) following approval from the University of Wollongong Social Sciences Human Ethics Committee.

Procedure

Participants were entering a year-long program of treatment. Individual goals for treatment were self-generated by participants at the first assessment session, guided by using the Target Complaints Measure.¹¹ Goals could be both specific and more general and long-term in focus and were not delimited by clinicians in any way. The Target Complaints Measure is a semi-structured clinician guided interview, which was used as part of the intake assessment session to ascertain each participant's treatment goals or chief complaints.¹¹ Participants were told "I want you to tell me in your own words the most important problems that you have that you want help with to change by coming here. These are the kind of goals you might have for your treatment" as specified by the Target Complaints Measure.¹¹ Participants were prompted to provide up to 3 goals and to self-rate how severe these were as an issue or problem for them on a scale of 0 (not a problem/least severe issue for me) to 10 (the worst/most severe issue for me).

Data Analysis

An inductive conventional content analytic approach to understanding the goals of individuals and the development of goal categories was used. This followed a 3-step approach as described by Hsieh and Shannon.¹⁵

Participants' descriptions of goals were audio-recorded and transcribed verbatim. Researchers immersed themselves in the data by reading and reflecting on participant responses to gain an overall understanding.

1 First, participant responses were tagged with codes, referred to as goal units, to accurately describe the
2 data. Due to the recognition that multiple goals could be present within an individual goal, some goals
3 could be represented by more than one code. Thus, although the Target Complaints Measure¹¹ specified
4 up to 3 goals, some participants provided more than 3 goals within their descriptions. Second, similar or
5 related codes were condensed into goal categories that allowed for both homogeneity within the group
6 and heterogeneity between groups. Lastly, goal categories were grouped into meaningful themes to
7 represent participant responses. The coding process was supported by the use of the NVivo 10 software
8 for qualitative data analysis. The data was initially independently coded and categorized by 2 researchers,
9 this was followed by the discussion and review of codes by a third researcher who is an expert in
10 personality disorders. The trustworthiness of the data was ensured by having consistent discussion about
11 codes and findings emerging from the data with the wider research team to ensure that concepts were not
12 overlooked within the data. Through multiple discussions and reviews, the coding and categorization of
13 codes were refined. Discrepancies among the coding and the subsequent categories between researchers
14 were discussed and resolved through consensus. Illustrative quotes of the goals were provided to support
15 and provide evidence for the interpretations of the researchers. The researchers analyzing the data were
16 independent from the clinicians providing psychological care to the participants.

18 **RESULTS**

19 **Characteristics of Participants**

20 A total of 102 consecutively recruited participants seeking treatment who met criteria for a primary DSM-
21 IV-TR diagnosis of BPD were invited to participate. All participants gave written informed consent to the
22 study. Table 1 outlines the demographic characteristics of the participants.

23 **Treatment Goals Identified by Participants**

24 Overall, participants identified a total of 268 goals, with an average of 2.8 goals per participant. All
25 participants (100%) were able to report 1 goal, 100 participants (98%) reported 2 goals, and 86
26 participants (84%) were able to report 3 goals. The majority of goals identified had multiple components.
27 The goals reported were then analysed into constituent units, for a total of 464 individual goal units (See
28 Table 2) or 4.5 goal units per participant. Reported goals could include the same individual goal unit on
29 multiple occasions in their descriptions, however, this was only counted once. Therefore, goal units
30 identified in Table 2 are indicative of the number of participants endorsing a specific goal unit.

Findings from the content analysis reveal 4 key themes associated with treatment and personally meaningful goals for recovery. The reduction of symptoms was the most commonly reported goal by participants (n=88, 86.3%), followed by the desire to improve wellbeing (n=64, 62.7%), having better interpersonal relationships (n=54, 52.9%), and having a greater sense of self (n=40, 39.2%). Although reducing symptoms was the most commonly reported theme, 90 participants (88.2%) also reported at least one goal pertaining to a psychosocial goal category. Goals reported by participants were identified as not being mutually exclusive, so that achievement of goals in one area could contribute to improvements in other areas.

Goal theme 1: Reducing symptoms

The goal of reducing symptoms was the most commonly cited theme in the study, where reducing suicidality and impulsivity and depressive and anxiety symptoms were some of the most highly reported goal categories. Participants discussed the impact of symptoms on daily functioning and self-perceptions. *"I'd certainly like to manage my depression better, so that I don't end up back in hospital again. I'd like to be able to explore things that may be affecting me as an adult so that I can understand why I feel the way about things that don't make sense. I just want to get on with my life, be a whole person rather than be in fragments."* (Individual 5091)

The experiences of symptoms were sometimes interrelated so that the experience of depressive or anxiety symptoms corresponded with a desire to engage in self-harming behaviors or increased suicidality. The reduction of symptoms had a compounding effect on a person's ability to engage in and achieve other psychosocial goals. *"I want to be able to deal with the depression and cope with distress... I'd like to get to a point where I can go back to do some study or do some work."* (Individual 3054)

Goal theme 2: Improving wellbeing

Goals pertaining to improving wellbeing were global in nature, and individual differences contributed to the heterogeneity of the goals. The desire to improve coping style was one of the most highly reported goal categories by participants and widely reflected the overall desire to improve symptomatically. The ability to effectively manage emotions and thoughts was believed to contribute to improved quality of life and emotional experience. *"To learn how to control the fuzziness that leads to those instances and slowly reducing the want, need and the action of self-harm"* (Individual 5086). Goals associated with improving current life situations and financial situation were also at times interconnected with symptoms and interpersonal relationships. Life situations mentioned were broad and included court cases, assault, divorce, and the loss of a child. *"At the moment, the involvement of court case for sexual assault is very stressful... It has restricted parts of my life, through avoiding people and avoiding relationships. Want to*

1 *get on with life and put it behind me”* (Individual 3051). Despite goals to improve life situations, one
2 participant articulated that these may be considered “*general life problems*” (Individual 5076),
3 highlighting the common experiences of individuals.

4 Vocational pursuits such as engagement in paid work and education were valued goals. Despite
5 the desire to be connected with society, the emotional intensity experienced by participants was identified
6 as a barrier. “*Being employable, but when you look at my CV, it’s like what have you been doing all these*
7 *years? Getting a part time job is really important. If I took on a full time position, I would let down my*
8 *employer and myself because it has been a few years since I have been in work*” (Individual 5100). Yet, it
9 was recognized that participation in a structured vocation may not be suitable for all individuals, so that
10 assisting individuals to take part in personally meaningful activities would be a valued target of treatment.
11 “*I really want to do dancing. Dancing used to really help me... I think it is teaching myself to go there*
12 *and not matter whether I will be put down for it.*” (Individual 5151)
13

14 ***Goal theme 3: Better interpersonal relationships***

15 Better interpersonal relationships were another key theme associated with developing a sense of
16 connectedness with others, improving current relationships, and developing interpersonal skills.
17 Connectedness was described by participants as being on a continuum including developing relationships,
18 connecting with others, and relating to others. “*Just being able to feel like I fit somewhere, I feel like I*
19 *don’t fit. I’m too scared to go out and meet new people... It is a big problem cause I don’t do anything.*’
20 (Individual 5106)

21 Difficulty relating to others was acknowledged and along with the goal that it could be improved
22 through a process of developing greater communication, engagement, and understanding of others. The
23 reported goal of developing and improving interpersonal skills reflected the need to overcome grief and
24 loneliness, to be more assertive, and establish trust with others to more effectively initiate or engage in
25 relationships. “*Be able to go with my own judgement or my own decision, instead of running to my father*
26 *all the time and his opinion—like decision making and assertiveness. I don’t trust my own judgement and*
27 *I am not very assertive either*” (Individual 5078). Relationships were mainly discussed by participants in
28 the context of their significant others, friends, family, and mothers. Participants also identified specific
29 goals related to improving parenting capacity.
30

31 ***Goal theme 4: Greater sense of self***

32 The theme of developing a greater sense of self was associated with improving attitudes toward self and
33 increasing personal awareness. The goal of increasing personal awareness was expressed as achievable
34 through developing greater self-understanding and being able to conceive of what might be a meaningful

1 life direction or goal, and have motivation to move in that direction. *“Be more of a whole*
2 *person...learning some tools that will help me be motivated to get out and do things and enjoy life instead*
3 *of dragging myself through it, all the time.”* (Individual 5091)

4 Some participants broadly discussed goals to “get to know who I am,” while others discussed a
5 desire to shift away from a “victim persona” developed from experiences of trauma and to no longer be
6 viewed only through the lens of their BPD diagnosis. *“I have childhood issues and I’m hanging onto*
7 *them. I’m dealing with them really well... but still need help to deal with some of those issues, how to not*
8 *be a victim”* (Individual 3054). Goals pertaining to the development of self-esteem and self-worth were
9 discussed, as were the negative consequences of poor self-esteem and self-worth *“If I consciously self-*
10 *harm, it is because of my self-esteem. I just hate myself”* (Individual 5090).

11 Some participants recognized that improving self-esteem may be an ongoing journey, and that a
12 person’s attitude toward him- or herself is inextricably linked with increasing personal awareness. The
13 ability to separate oneself from others in order to develop a sense of who one is and a sense of
14 genuineness was also identified by some participants. *“Getting to know me... I want to be more*
15 *consistent. I’ve gotten to the point where I push people away because I can’t be me and I am sort of*
16 *resenting them for it, even if they are not doing it”* (Individual 5113).

18 **DISCUSSION**

19 This study explored the views of individuals seeking treatment for BPD on their personally meaningful
20 goals for treatment. Participants identified goals in 4 main categories: reducing symptoms, improving
21 wellbeing, better interpersonal relationships, and having a greater sense of self. Personally meaningful
22 treatment goals identified in this study extend beyond the reduction of symptoms to include those of a
23 psychosocial nature. This supports calls to expand outcome measures to monitor progress and include
24 aspects that are global in nature, such as subjective wellbeing and the views of individuals seeking
25 treatment.^{16,17} The goal themes identified in this study were consistent with research examining the lived
26 experience of individuals with personality disorders^{6,18,19} and reflected some of the domains present in the
27 Camberwell Assessment of Need.⁹ Domains of psychopathology in BPD were also reflected in the
28 identified goals,²⁰ including difficulties in relational functioning, emotion dysregulation, and
29 understanding self and others. However, the identified goal themes and categories expand on the work to
30 date in the literature by providing greater insights into the specific aspects that may be important to
31 individuals that could be potentially targeted during treatment.

1 The identification of symptom reduction as the most cited theme was not surprising given the
2 severe nature of BPD and that individuals were at the start of treatment. Interestingly, studies of the lived
3 experience of personality disorders have conceptualized recovery as the reconciliation of self and other
4 representations through the development of a sense of self that could be achieved through the engagement
5 of interpersonal relationships and the community.^{6,21} Although these themes are reflected in the findings
6 from our study, fewer than half the participants reported goals associated with developing a greater sense
7 of self. This may be associated with the sample being at the start of treatment, in contrast to other studies
8 where participants were engaging in a specialist intervention and therefore were more aware of their
9 underlying difficulties. This finding may also reflect the shifting nature of treatment goals and suggest
10 that routine monitoring of goals may be required.

11 The treatment goals that were reported were not mutually exclusive, so that participants believed
12 that improvements in one goal would contribute to the achievement of other goals. This suggests that
13 there may be multiple processes and challenges involved in achieving desired recovery outcomes in a
14 personally meaningful manner. Although the identification of these processes and challenges were
15 beyond the scope of this study, understanding these will have implications for clinical practice and can
16 provide guidance for the development of recovery-oriented mental health services for BPD.

17 **Implications for Clinical Practice**

18 The narrow treatment targets of interventions for BPD have been reported as a limitation to care by
19 individuals with BPD.⁶ Although this study identified similarities between individual treatment goals and
20 the typical targets of interventions, some identified goal categories and units did reflect that a wider focus
21 may not be captured in psychotherapeutic interventions and treatment manuals for BPD. Given the
22 findings reported here, there is room for treatment manuals to focus more broadly on goals identified by
23 individuals. Having broader treatment targets may also have the effect of generating greater motivation
24 for behavioral change and improving treatment engagement. In addition, the therapeutic alliance between
25 clinicians and individuals could also profit from greater awareness of individual goals.

26 The development of new methods of integrating existing psychotherapeutic evidence-based
27 approaches with psychosocial interventions may be important in assisting individuals with BPD achieve
28 their desired outcomes.²² The findings from this study provide a basis for understanding areas of
29 importance to individuals with BPD. Evidence-based social interventions and psychosocial rehabilitation
30 interventions such as illness management and recovery,²³ assertive community treatment,²⁴ or individual
31 placement and support²⁵ may help support individuals with BPD in achieving goals that extend beyond
32 the scope of the current manualized interventions with an evidence base. In addition, developing the
33

1 capacity of individuals with lived experience to become peer support workers may also present a unique
2 opportunity for individuals with similar experiences to learn from each other.²⁶

3 The development of enhanced therapeutic interventions that target specific goals of interest to
4 individuals with BPD may also be relevant. One recent example of such an intervention pertains to
5 improving the parenting capacity of individuals with BPD who are parents.²⁷ Continual evaluation of the
6 integration of these interventions with evidence-based interventions should be completed using multiple
7 measures and methodologies.

9 **Limitations and Future Research**

10 Treatment goals reported by participants in our study were framed in a clinically oriented manner, so that
11 goals predominantly focused on the symptoms and problems participants wanted to overcome. Although
12 this can be attributed to the context in which goals were formulated, the identified goals may also be
13 reflective of individuals who are at the start of their recovery journey. The goals, however, provide a good
14 indicator of the valued outcomes through the perspectives of individuals seeking treatment. The wide
15 range of individual goal units (N=464) identified is also indicative of the personal nature of treatment
16 goals and the need for mental health clinicians to ask individuals what their goals are for treatment,
17 particularly given the links between goal consensus, collaboration, and attainment.^{28,29} Goals for treatment
18 and recovery are not static. This is reflected in the non-linear trajectory of recovery.³⁰ More research
19 examining changes in goal content longitudinally may provide a more nuanced understanding of the
20 differences between individuals who may be at different stages of their recovery and whether treatments
21 received are assisting individuals to attain their goals.

1 REFERENCES

- 2 1. Linehan MM, Armstrong HE, Suarez A, et al. Cognitive-behavioral treatment of chronically
3 parasuicidal borderline patients. *Arch Gen Psychiatry*. 1991;48:1060–1064.
- 4 2. Cristea IA, Gentili C, Cotet CD, et al. Efficacy of psychotherapies for borderline personality
5 disorder. *JAMA Psychiatry*. 2017;74:319–328.
- 6 3. Bateman A, Fonagy P. Mentalization based treatment for borderline personality disorder. *World*
7 *Psychiatry*. 2010;9:11–5.
- 8 4. Kellogg SH, Young JE. Schema therapy for borderline personality disorder. *J Clin Psychol*.
9 2006;62:445–458.
- 10 5. Kernberg OF, Yeomans FE, Clarkin JF, et al. Transference focused psychotherapy: overview and
11 update. *Int J Psychoanal*. 2008;89:601–620.
- 12 6. Katsakou C, Marougka S, Barnicot K, et al. Recovery in borderline personality disorder (BPD): A
13 qualitative study of service users' perspectives. *Plos One*. 2012;7:e36517.
- 14 7. Ng FYY, Bourke ME, Grenyer BFS. Recovery from borderline personality disorder: a systematic
15 review of the perspectives of consumers, clinicians, family and carers. *Plos One*. 2016;11:e0160515.
- 16 8. Le Boutillier C, Leamy M, Bird V, et al. What does recovery mean in practice? A qualitative
17 analysis of international recovery-oriented practice guidance. *Psychiatr Serv*. 2011;62:1470–1476.
- 18 9. Phelan M, Slade M, Thornicroft G, et al. The Camberwell Assessment of Need: the validity and
19 reliability of an instrument to assess the needs of people with severe mental illness. *Br J Psychiatry*.
20 1995;167:589–595.
- 21 10. Hayward M, Slade M, Moran PA. Personality disorders and unmet needs among psychiatric
22 inpatients. *Psychiatr Serv*. 2006;57:538–543.
- 23 11. Battle CC, Imber SD, Hoehn-Saric R, et al. Target complaints as criteria of improvement. *Am J*
24 *Psychother*. 1966;20:184–192.
- 25 12. Lindhiem O, Bennett CB, Orimoto TE, et al. A meta-analysis of personalized treatment goals in
26 psychotherapy: a preliminary report and call for more studies. *Clin Psychol (New York)*. 2016;23:165–
27 176.
- 28 13. First MB, Spitzer RL, Gibbon M, et al. User's guide for the Structured Clinical Interview for
29 DSM-IV Axis I disorders SCID-I: Clinician Version. Washington: American Psychiatric Press; 1997.
- 30 14. First MB, Gibbon M, Spitzer RL, et al. User's Guide for the Structured Clinical Interview for
31 DSM-IV Axis II Personality Disorders. Washington: American Psychiatric Press; 1997.
- 32 15. Hsieh HF, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res*.
33 2005;15:1277–1288.
- 34 16. Howard KI, Lueger RJ, Maling MS, et al. A phase model of psychotherapy outcome: causal
35 mediation of change. *J Consult Clin Psychol*. 1993;61:678–685.
- 36 17. Thornicroft G, Tansella M. Growing recognition of the importance of service user involvement in mental
37 health service planning and evaluation. *Epidemiol Psichiatr Soc*. 2005;14:1–3.
- 38 18. Gillard S, Turner, K, Neffgen M. Understanding recovery in the context of lived experience of
39 personality disorders: a collaborative, qualitative research study. *BMC Psychiatry*. 2015;15:1–13.
- 40 19. Shepherd A, Sanders C, Shaw J. Seeking to understand lived experiences of personal recovery in
41 personality disorder in community and forensic settings: a qualitative methods investigation. *BMC*
42 *Psychiatry*. 2017;17:1–10.
- 43 20. Sanislow CA, Grilo CM, Morey LC, et al. Confirmatory factor analysis of DSM-IV criteria for
44 borderline personality disorder: findings from collaborative longitudinal personality disorders study. *Am*
45 *J Psychiatry*. 2002;159:284–290.
- 46 21. Turner K, Lovell K, Brooker A. '... and they all lived happily ever after': 'Recovery' or discovery
47 of the self in personality disorder? *Psychodyn Pract*. 2011;17:341–346.
- 48 22. Frese FJ, Stanley J, Kress K, et al. Integrating evidence-based practices and the recovery model.
49 *Psychiatr Serv*. 2001;52:1462–1468.

23. Mueser KT, Corrigan PW, Hilton DW, et al. Illness management and recovery: a review of the research. *Psychiatr Serv*. 2002;53:1272–1284.
24. Stein LI, Test MA. Alternative to mental hospital treatment: I. Conceptual model, treatment program, and clinical evaluation. *Arch Gen Psychiatry*. 1980;37:392–397.
25. Burns T, Catty J, Becker T, et al. The effectiveness of supported employment for people with severe mental illness: a randomised controlled trial. *The Lancet*. 2007;370:1146–1152.
26. Repper J, Carter T. A review of the literature on peer support in mental health services. *J Ment Health*. 2011;20:392–411.
27. McCarthy KL, Lewis KL, Bourke ME, et al. A new intervention for people with borderline personality disorder who are also parents: a pilot study of clinician acceptability. *Borderline Personal Disord Emot Dysregul*. 2016;3:1–9.
28. Tryon GS, Winograd G. Goal consensus and collaboration. In: Norcross, JC, ed. *Psychotherapy Relationships That Work: Evidence-based Responsiveness*, 2nd ed. New York: Oxford University Press; 2011: pp. 153–167.
29. Clarke SP, Oades LG, Crowe TP, et al. The role of symptom distress and goal attainment in promoting aspects of psychological recovery for consumers with enduring mental illness. *J Ment Health*. 2009;18:389–397.
30. Slade M. *Personal Recovery and Mental Illness: A Guide for Mental Health Professionals*. New York: Cambridge University Press; 2009.

1 **Table 1: Characteristics of Participants (N = 102)**

Characteristic	Range	n (%) or Mean (SD)
Female		89 (87.3%)
Age	18–56 years	29.7 years (8.84)
Relationship status:		
Single		57 (55.9%)
Married		20 (19.6%)
De-facto		11 (10.8%)
Divorced		7 (6.9%)
Separated		7 (6.9%)
Years of education	7.5–19 years	12.1 years (2.58)

2

3

4

1 **Table 2: Formulated themes and frequencies of participant-generated goals verbalized at the**
2 **commencement of treatment for borderline personality disorder (N=102; 464 individual goal units)**

Formulated theme and goal categories	Participant-generated goal statements	Number of participants endorsing the theme (% of all participants)
Goal Theme: Reducing Symptoms (5 goal categories, 17 goal units)		88 (86.3%)
<i>Suicidality and Impulsivity</i> (51 participants, 50.0%)	Self-harm behaviors/thoughts	22 (21.6%)
	Suicidality	17 (16.7%)
	Anger	15 (14.7%)
	Drug and alcohol misuse	7 (6.9%)
	Gambling urges	2 (2.0%)
	Shoplifting	2 (2.0%)
<i>Depressive Symptoms</i> (42 participants, 41.2%)	Negative mood/thoughts	36 (35.3%)
	Mood swings	6 (5.9%)
<i>Anxiety Symptoms</i> (40 participants, 39.2%)	General anxiety	17 (16.7%)
	Posttraumatic stress/trauma	17 (16.7%)
	Panic attacks	7 (6.9%)
	Social anxiety	5 (4.9%)
	Specific phobia	5 (4.9%)
<i>Eating Related Issues</i> (11 participants, 10.8%)	Disordered eating	7 (6.9%)
	Weight loss	4 (3.9%)

<i>Transient Symptoms</i> (8 participants, 7.8%)	Dissociation Hallucinations	6 (5.9%) 2 (2.0%)
Goal Theme: Improving Wellbeing (4 goal categories, 14 goal units)		64 participants (62.7%)
<i>Coping Style</i> (46 participants, 45.1%)	Having control over emotions Improve coping style Coping with distress/stress Having control over thoughts Improve functioning and use of skills General sense of control	25 (24.5%) 17 (16.7%) 15 (14.7%) 7 (6.9%) 8 (7.8%) 2 (2.0%)
<i>Vocation</i> (21 participants, 20.6%)	Engaging in paid work Engaging in activities Education	9 (8.8%) 8 (7.8%) 5 (4.9%)
<i>Current Life Situations</i> (18 participants, 17.6%)	Solve specific life situations Financial situation	14 (13.7%) 4 (3.9%)
<i>Physical Health</i> (9 participants, 8.8%)	Improve physical health Stay out of hospital Come off medication	5 (4.9%) 3 (2.9%) 1 (1%)

Goal Theme: Better Interpersonal Relationships (4 goal categories, 14 goal units)		54 participants (52.9%)
<i>Interpersonal skills</i> (23 participants, 22.5%)	Overcome my grief and loneliness Being assertive with others Trusting others Become independent Reduce abandonment fears	10 (9.8%) 5 (4.9%) 4 (3.9%) 3 (2.9%) 2 (2.0%)
<i>Improving Current Relationships</i> (22 participants, 21.6%)	Relationship with my significant other Relationship with my friends Relationship with my family Relationship with my mother	6 (5.9%) 6 (5.9%) 6 (5.9%) 5 (4.9%)
<i>Connectedness</i> (21 participants, 20.6%)	Connecting with others Developing relationships Relating to others	12 (11.8%) 7 (6.9%) 5 (4.9%)
<i>Parenting</i> (8 participants, 7.8%)	Develop my parenting skills Have contact and a better relationship with my children	6 (5.9%) 4 (3.9%)
Goal Theme: Greater sense of self (2 goal categories, 11 goal units)		40 participants (39.2%)
<i>Attitudes Toward Self</i>	Having self-esteem and self-worth	12 (11.8%)

(27 participants, 26.5%)	Sense of self	9 (8.8%)
	Having self-confidence	4 (3.9%)
	Become self-accepting	3 (2.9%)
	Improve self-image and reduce perfectionism	3 (2.9%)
<i>Personal Awareness</i>	Understanding myself	10 (9.8%)
(23 participants, 22.5%)	Develop my goals and motivation	8 (7.8%)
	Identifying my vulnerabilities	5 (4.9%)
	Reducing feelings of emptiness	4 (3.9%)
	Having a sense of purpose	2 (2.0%)
	Having my own opinions	2 (2.0%)

1

2

3

4