



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Engineering and Information Sciences -
Papers: Part B

Faculty of Engineering and Information Sciences

2018

On the Global Limiting Absorption Principle for Massless Dirac Operators

Alan L. Carey

University of Wollongong, acarey@uow.edu.au

Fritz Gesztesy

Baylor University

Jens Kaad

University of Southern Denmark

Galina Levitina

University of New South Wales

Roger Nichols

University of Tennessee at Chattanooga

See next page for additional authors

Publication Details

Carey, A., Gesztesy, F., Kaad, J., Levitina, G., Nichols, R., Potapov, D. & Sukochev, F. (2018). On the Global Limiting Absorption Principle for Massless Dirac Operators. *Annales Henri Poincaré*, 19 (7), 1993-2019.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

On the Global Limiting Absorption Principle for Massless Dirac Operators

Abstract

We prove a global limiting absorption principle on the entire real line for free, massless Dirac operators (Formula presented.) for all space dimensions (Formula presented.), (Formula presented.). This is a new result for all dimensions other than three, in particular, it applies to the two-dimensional case which is known to be of some relevance in applications to graphene. We also prove an essential self-adjointness result for first-order matrix-valued differential operators with Lipschitz coefficients.

Disciplines

Engineering | Science and Technology Studies

Publication Details

Carey, A., Gesztesy, F., Kaad, J., Levitina, G., Nichols, R., Potapov, D. & Sukochev, F. (2018). On the Global Limiting Absorption Principle for Massless Dirac Operators. *Annales Henri Poincare*, 19 (7), 1993-2019.

Authors

Alan L. Carey, Fritz Gesztesy, Jens Kaad, Galina Levitina, Roger Nichols, Denis Potapov, and F Sukochev

ON THE GLOBAL LIMITING ABSORPTION PRINCIPLE FOR MASSLESS DIRAC OPERATORS

ALAN CAREY, FRITZ GESZTESY, JENS KAAD, GALINA LEVITINA, ROGER NICHOLS,
DENIS POTAPOV, AND FEDOR SUKOCHEV

ABSTRACT. We prove a global limiting absorption principle on the entire real line for free, massless Dirac operators $H_0 = \alpha \cdot (-i\nabla)$ for all space dimensions $n \in \mathbb{N}$, $n \geq 2$. This is a new result for all dimensions other than three, in particular, it applies to the two-dimensional case which is known to be of some relevance in applications to graphene.

We also prove an essential self-adjointness result for first-order matrix-valued differential operators with Lipschitz coefficients.

1. INTRODUCTION

1.1. Motivation and background. This paper was motivated by recent investigations of the Witten index (a possible substitute for the Fredholm index) for classes of non-Fredholm operators, a prime example of which being the massless Dirac operator, see, for instance, [14]–[17]. Without going into the somewhat lengthy details here, we mention that the Witten index can be expressed in terms of the spectral shift function associated with the pair (H, H_0) , where $H_0 = \alpha \cdot (-i\nabla)$ represents the free massless Dirac operator and $H = H_0 + V$ the massless interacting Dirac operator. For a detailed introduction to this circle of ideas we refer to the survey [16].

To briefly set the stage, suppose that \mathcal{H}_+ is a Banach space that embeds continuously and densely into the complex, separable Hilbert space \mathcal{H} so that \mathcal{H} embeds continuously and densely into $\mathcal{H}_- := \mathcal{H}_+^*$. If $S : \text{dom}(H) \subset \mathcal{H} \rightarrow \mathcal{H}$ is a self-adjoint operator, then (as we discuss in Section 3)

$$\|(S - (\lambda \pm i\mu)I_{\mathcal{H}})^{-1}\|_{\mathcal{B}(\mathcal{H}_+, \mathcal{H}_-)} \leq C_1, \quad \lambda \in \mathbb{R}, \mu \in (0, \infty), \quad (1.1)$$

holds for some constant $C_1 \in (0, \infty)$ (independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$) if and only if

$$|(\psi, (S - (\lambda \pm i\mu)I_{\mathcal{H}})^{-1}\psi)_{\mathcal{H}}| \leq C_2 \|\psi\|_{\mathcal{H}_+}^2, \quad \psi \in \mathcal{H}_+, \lambda \in \mathbb{R}, \mu \in (0, \infty), \quad (1.2)$$

holds for some constant $C_2 \in (0, \infty)$ (independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$).

Such estimates yield the absence of any singular spectrum of S and permit perturbative arguments that yield unitary equivalence of S and additive perturbations

Date: February 25, 2018.

2010 Mathematics Subject Classification. Primary: 35P25, 35Q40, 81Q10; Secondary: 47A10, 47A40.

Key words and phrases. Dirac operators, limiting absorption principle, essential self-adjointness.

A.C., G.L., and F.S. gratefully acknowledge financial support from the Australian Research Council. J.K. is supported by the DFF-research project 2 “Automorphisms and invariants of operator algebras,” no. 7014-00145B and by the Villum foundation (Grant 7423).

$S + W$, assuming that W is “sufficiently small” (in a sense to be discussed in Theorem 3.15). They also permit a discussion of Birman–Schwinger-type operators (including their boundary values on the real axis) which are of fundamental importance in expressing the Witten index in terms of the spectral shift function for the pair $(S, S + W)$ (cf. Theorem 3.15 in the concrete case of the pair (H, H_0)).

In particular, we will employ the fact (1.1), (1.2) to prove the principal goal of this paper, a uniform (or global) limiting absorption principle for $H_0 = \alpha \cdot (-i\nabla)$ for all space dimensions $n \in \mathbb{N}$, $n \geq 2$. More precisely, given $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$, we will prove in Theorem 3.13 that there exists a $C \in (0, \infty)$ (independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$), such that

$$\begin{aligned} \|(H_0 - (\lambda \pm i\mu)I_{[L^2(\mathbb{R}^n)]^N})^{-1}\|_{\mathcal{B}([L^2_1(\mathbb{R}^n)]^N, [L^2_{-1}(\mathbb{R}^n)]^N)} &\leq C, \\ \lambda \in \mathbb{R}, \mu \in (0, \infty). \end{aligned} \tag{1.3}$$

The first result on a global limiting absorption principle on the entire real line for free, massive Dirac operators $H_0(m) = \alpha \cdot (-i\nabla) + m\beta$, $m > 0$ (cf. (3.4) for details), in the case of dimensions $n = 3$, is due to Iftimovici and Măntoiu [32] in 1999. The first such result for free, massless Dirac operators $H_0 = \alpha \cdot (-i\nabla)$ (cf. (3.2)), again in dimensions $n = 3$, was proved by Saitō and Umeda [46] in 2008 upon relying on quite different methods as part of their study of zero-energy eigenvalues and resonances. Since, apparently, no other global results are available in the massless case, we now fill this gap, and by suitably modifying the approach of Iftimovici and Măntoiu, we prove a global limiting absorption principle (i.e., for energies on the entire real line) for free, massless Dirac operators in all dimensions $n \in \mathbb{N}$, $n \geq 2$. We note, however, that a limiting absorption principle for Dirac operators with continuous matrix-valued short-range potentials in the massless case $m = 0$ (resp., in the massive case $m > 0$) in two and higher dimensions, valid on $(-\infty, 0) \cup (0, \infty)$ (resp., valid on $(-\infty, -m) \cup (m, \infty)$), assuming regular thresholds, has just been proven by Erdoğan, Goldberg, and Green [21] upon relying on a different approach.

It is gratifying that the new results in dimensions $n \in \mathbb{N} \setminus \{3\}$, $n \geq 2$, include, in particular, the two-dimensional case $n = 2$ which is known to have some relevance in applications to graphene.

We emphasize that proving a limiting absorption principle for free (usually, constant coefficient) operators H_0 (resp., $H_0(m)$) is always a first step in proving similar statements (typically, away from essential spectrum thresholds, though) for interacting Hamiltonians $H = H_0 + V$ (resp., $H(m) = H_0(m) + V$) on the basis of a sophisticated perturbative approach. The strategy behind such an approach has been spelled out in great detail, for instance, by Yafaev in [49, Sect. 4.6, 4.7].

While we follow the broad contours outlined in the approach employed by Iftimovici and Măntoiu [32] in the massive case, $m > 0$, there are notable differences in our treatment of the massless case, $m = 0$; of these we note, in particular, the following:

- First, we do not rely on Nelson’s commutator theorem in proving essential self-adjointness of the conjugate operator A introduced in (3.8). Instead, we employ an extension of an essential self-adjointness result for general first-order matrix-valued differential operators (such as, A , upon applying the Fourier transform) going back to Chernoff, [18], but see also [31] for a more modern treatment. Here, we treat only the flat case but permit Lipschitz coefficients instead of the more restrictive

(and more common) smoothness assumption. We expect this topic, to be treated in detail in Section 2, to be of independent interest.

- Second, given our focus on massless Dirac operators in all dimensions $n \geq 2$, we were led to a new conjugate operator A in (3.8) when compared to the massive case discussed in [32]. This causes a variety of additional technical difficulties in Section 3 briefly outlined in the paragraph following (3.10).

- Third, rather than applying Hardy's inequality as in [32], which is only applicable in dimensions $n \geq 3$ (a restriction we wanted to avoid by all means), we now apply Kato's inequality (cf. (3.23), and especially, (3.25)).

Our principal result, the global limiting absorption principle on the real line for H_0 is presented in Section 3 (see Theorem 3.13). This is followed by a standard application to scattering theory for the pair of self-adjoint operators $(H = H_0 + V, H_0)$, with sufficiently weak interactions V (cf. Theorem 3.15), and some remarks that put our results in proper perspective.

For additional (and more general) references in the context of smooth operator theory, limiting absorption principles, completeness of wave operators, and absence of singular continuous spectra, see, for instance, [2], [3], [4], [6], [8, Ch. 17], [9], [10], [11], [12], [20], [22], [23], [24], [27], [28], [29], [36], [37], [39], [41], [42], [43, Sect. XIII.7], [44], [45], [46], [48], [49, Ch. 4], [50, Chs. 0–2], [51]–[53], in particular, global limiting absorption principles for Schrödinger operators can be found in [22], [23], [45].

1.2. A summary of the notation employed: We conclude this introduction by summarizing some of the notation used throughout this paper: Vectors in \mathbb{R}^n are denoted by $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ or $p = (p_1, \dots, p_n) \in \mathbb{R}^n$, $n \in \mathbb{N}$. For $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ we abbreviate

$$\langle x \rangle = (1 + |x|^2)^{1/2}, \quad (1.4)$$

where $|x| = (x_1^2 + \dots + x_n^2)^{1/2}$ denotes the standard Euclidean norm of $x \in \mathbb{R}^n$, $n \in \mathbb{N}$.

The dot symbol, “ \cdot ”, is used in three different ways: First, it denotes the standard scalar product in \mathbb{R}^n ,

$$x \cdot y = \sum_{j=1}^n x_j y_j, \quad x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n. \quad (1.5)$$

Second, we will also use it for n -vectors of operators, $\underline{A} = (A_1, \dots, A_n)$ and $\underline{B} = (B_1, \dots, B_n)$ acting in the same Hilbert space in the form

$$\underline{A} \cdot \underline{B} = \sum_{j=1}^n A_j B_j, \quad (1.6)$$

whenever it is obvious how to resolve the domain issues of the possibly unbounded operators involved. Moreover, for T an operator in some Hilbert space \mathcal{H} and $A = (a_{j,k})_{1 \leq j, k \leq N} \in \mathbb{C}^{N \times N}$ an $N \times N$ matrix with constant complex-valued entries acting in \mathbb{C}^N , $N \in \mathbb{N}$, we will avoid tensor product notation in

$$T \otimes A \text{ in } \mathcal{H} \otimes \mathbb{C}^N \quad (1.7)$$

such that

$$\mathcal{H} \otimes \mathbb{C}^N \text{ is identified with } \mathcal{H}^N = \begin{pmatrix} \mathcal{H} \\ \vdots \\ \mathcal{H} \end{pmatrix}, \quad (1.8)$$

and

$$T \otimes A \text{ is identified with } TA = (Ta_{j,k})_{1 \leq j, k \leq N} = (a_{j,k}T)_{1 \leq j, k \leq N} = AT. \quad (1.9)$$

That is, we interpret $T \otimes A$ as entrywise multiplication, resulting in an $N \times N$ block operator matrix $TA = AT$. Thus, if $\underline{T} = (T_1, \dots, T_n)$, with T_j , $1 \leq j \leq n$, operators in \mathcal{H} , and $\underline{A} = (A_1, \dots, A_n)$, with $A_j \in \mathbb{C}^{N \times N}$, $1 \leq j \leq n$, $N \times N$ matrices in \mathbb{C}^N , we will finally employ the dot symbol also in the form

$$\underline{T} \cdot \underline{A} = \sum_{j=1}^n T_j A_j = \sum_{j=1}^n A_j T_j = \underline{A} \cdot \underline{T}, \quad (1.10)$$

where $T_j A_j = A_j T_j$, $1 \leq j \leq n$, are defined as in (1.9). In the interest of clarity we temporarily underlined vectors of operators (and matrices) such as $\underline{T} = (T_1, \dots, T_n)$; we will refrain from doing so in the bulk of this manuscript.

For X a given set, $A \in X^{N \times N}$, $N \in \mathbb{N}$, represents an $N \times N$ matrix A with entries in X .

Let \mathcal{H}, \mathcal{K} be separable complex Hilbert spaces, $(\cdot, \cdot)_{\mathcal{H}}$ the scalar product in \mathcal{H} (linear in the second argument), $\|\cdot\|_{\mathcal{H}}$ the norm on \mathcal{H} , and $I_{\mathcal{H}}$ the identity operator in \mathcal{H} . If T is a linear operator mapping (a subspace of) a Hilbert space into another, then $\text{dom}(T)$ and $\text{ker}(T)$ denote the domain and kernel (i.e., null space) of T . The closure of a closable operator A is denoted by \bar{A} .

The resolvent set and spectrum of a closed operator T are denoted by $\rho(T)$ and $\sigma(T)$, respectively.

The Banach space of bounded linear operators on a separable complex Hilbert space \mathcal{H} is denoted by $\mathcal{B}(\mathcal{H})$.

For a densely defined closed operator S in \mathcal{H} we employ the abbreviation $\langle S \rangle := (I_{\mathcal{H}} + |S|^2)^{1/2}$, and similarly, if $T = (T_1, \dots, T_n)$, with T_j densely defined and closed in \mathcal{H} , $1 \leq j \leq n$,

$$\langle T \rangle = (I_{\mathcal{H}} + |T|^2)^{1/2}, \quad |T| = (|T_1|^2 + \dots + |T_n|^2)^{1/2}, \quad (1.11)$$

whenever it is obvious how to define $|T_1|^2 + \dots + |T_n|^2$ as a nonnegative self-adjoint operator.

To simplify notation, we will frequently omit Lebesgue measure whenever possible and simply use $L^2(\mathbb{R}^n)$ instead of $L^2(\mathbb{R}^n; dx)$, etc. If no confusion can arise, the identity operator in $L^2(\mathbb{R}^n)$ is simply denoted by I , and I_N represents the identity operator in \mathbb{C}^N , $N \in \mathbb{N}$.

The symbol \mathcal{F} is used to denote the Fourier transform and $\widehat{f} := \mathcal{F}f$. For $N \in \mathbb{N}$, the Fourier transform of functions in $[L^2(\mathbb{R}^n)]^N$ is taken component-wise, that is, if $f = (f_1, \dots, f_N)^{\top} \in [L^2(\mathbb{R}^n)]^N$, then $\widehat{f} = (\widehat{f}_1, \dots, \widehat{f}_N)^{\top}$.

If $\Omega \subseteq \mathbb{R}^n$ is open, then $C_0^\infty(\Omega)$ denotes the set of infinitely differentiable functions on \mathbb{R}^n with compact support in Ω . In addition, $S(\mathbb{R}^n)$ denotes the Schwartz space of rapidly decreasing functions on \mathbb{R}^n , $S'(\mathbb{R}^n)$ the space of tempered distributions, and $W^{k,p}(\mathbb{R}^n)$, $k \in \mathbb{N}$, $p \geq 1$, the standard L^p -based Sobolev spaces.

We abbreviate $\mathbb{C}_\pm = \{z \in \mathbb{C} \mid \pm \operatorname{Im}(z) > 0\}$. The symbol $\lfloor \cdot \rfloor$ denotes the floor function on \mathbb{R} , that is, $\lfloor x \rfloor$ characterizes the largest integer less than or equal to $x \in \mathbb{R}$.

Following a standard practice in Mathematical Physics, we will simplify the notation of operators of multiplication by a scalar or matrix-valued function V and hence use V rather than the more elaborate symbol M_V throughout this manuscript.

2. ESSENTIAL SELF-ADJOINTNESS OF FIRST-ORDER DIFFERENTIAL OPERATORS WITH LIPSCHITZ COEFFICIENTS

We start with a general self-adjointness result for first-order differential expressions which we believe is of independent interest. It will be applied in the subsequent section in connection with the operator A in (3.8).

Let $n, N \in \mathbb{N}$ be fixed. We consider $N \times N$ block operator matrices of bounded operators of multiplication

$$F_j \in \mathcal{B}(L^2(\mathbb{R}^n))^{N \times N}, \quad j \in \mathbb{N}, 1 \leq j \leq n, \quad (2.1)$$

(e.g., F_j can be $N \times N$ matrix-valued operators of multiplication satisfying $F_j = (F_{j,k,\ell})_{1 \leq k, \ell \leq N}$, $F_{j,k,\ell} \in L^\infty(\mathbb{R}^n)$, $1 \leq j \leq n$, $1 \leq k, \ell \leq N$) together with the unbounded, closable operators

$$-i\partial_j : [W^{1,2}(\mathbb{R}^n)]^N \rightarrow [L^2(\mathbb{R}^n)]^N, \quad j \in \mathbb{N}, 1 \leq j \leq n. \quad (2.2)$$

where $\partial_j = \partial/\partial x_j$, $1 \leq j \leq n$. We put

$$\nabla := \begin{pmatrix} \partial_1 \\ \vdots \\ \partial_n \end{pmatrix} : [W^{1,2}(\mathbb{R}^n)]^N \rightarrow \left([L^2(\mathbb{R}^n)]^N \right)^n = \begin{pmatrix} [L^2(\mathbb{R}^n)]^N \\ \vdots \\ [L^2(\mathbb{R}^n)]^N \end{pmatrix}, \quad (2.3)$$

where $\left([L^2(\mathbb{R}^n)]^N \right)^n$ denotes the n -fold direct sum of $[L^2(\mathbb{R}^n)]^N$ with itself. One notes that ∇ is, in fact, densely defined and closed. Define

$$-\Delta := \nabla^* \nabla = - \sum_{j=1}^n \partial_j^2 : [W^{2,2}(\mathbb{R}^n)]^N \rightarrow [L^2(\mathbb{R}^n)]^N. \quad (2.4)$$

The main goal of this section is to prove the essential self-adjointness of a class of first-order differential operators. To make the result precise, we introduce the following set of assumptions, which will remain in effect throughout the remainder of this section.

Hypothesis 2.1. (i) Suppose F_j , $1 \leq j \leq n$ satisfy (2.1).

(ii) Let $\mathcal{D} \subseteq [L^2(\mathbb{R}^n)]^N$ be a core for ∇ with the property that

$$F_j(\mathcal{D}) \subseteq \mathcal{D}, \quad F_j^*(\mathcal{D}) \subseteq \mathcal{D}, \quad j \in \mathbb{N}, 1 \leq j \leq n. \quad (2.5)$$

(iii) Suppose that the commutators

$$[-i\partial_k, F_j], [-i\partial_k, F_j^*] : \mathcal{D} \rightarrow [L^2(\mathbb{R}^n)]^N, \quad j, k \in \mathbb{N}, 1 \leq j, k \leq n, \quad (2.6)$$

extend to bounded operators

$$d_k(F_j), d_k(F_j^*) \in \mathcal{B}\left([L^2(\mathbb{R}^n)]^N\right), \quad j, k \in \mathbb{N}, 1 \leq j, k \leq n, \quad (2.7)$$

respectively.

(iv) Define the symmetric unbounded first-order differential operator

$$L := -i \sum_{j=1}^n F_j \partial_j - i \sum_{j=1}^n \partial_j F_j^* : \mathcal{D} \rightarrow [L^2(\mathbb{R}^n)]^N. \quad (2.8)$$

Proposition 2.2. *Assume Hypothesis 2.1. Then items (ii) and (iii) in Hypothesis 2.1 hold with \mathcal{D} replaced by $[W^{1,2}(\mathbb{R}^n)]^N$. Moreover,*

$$[W^{1,2}(\mathbb{R}^n)]^N \subseteq \text{dom}(\bar{L}). \quad (2.9)$$

Proof. The proof that items (ii) and (iii) in Hypothesis 2.1 hold with the choice $\mathcal{D} = [W^{1,2}(\mathbb{R}^n)]^N$ is standard (cf., e.g., [26, Proposition 2.1]).

One notices that L is symmetric by construction. In particular, L is closable and hence \bar{L} is well-defined. To show that $[W^{1,2}(\mathbb{R}^n)]^N \subseteq \text{dom}(\bar{L})$, define the bounded operators

$$F = (F_1, \dots, F_n) : ([L^2(\mathbb{R}^n)]^N)^n \rightarrow [L^2(\mathbb{R}^n)]^N, \quad (2.10)$$

and

$$(F^*)^\top = (F_1^*, \dots, F_n^*) : ([L^2(\mathbb{R}^n)]^N)^n \rightarrow [L^2(\mathbb{R}^n)]^N. \quad (2.11)$$

Using matrix multiplication one verifies that

$$L = F \cdot (-i\nabla) + (F^*)^\top \cdot (-i\nabla) + \sum_{j=1}^n d_j(F_j^*) : \mathcal{D} \rightarrow [L^2(\mathbb{R}^n)]^N. \quad (2.12)$$

Since F , $(F^*)^\top$, and $d_j(F_j^*)$, $1 \leq j \leq n$, are all bounded and \mathcal{D} is a core for ∇ , one infers

$$[W^{1,2}(\mathbb{R}^n)]^N = \text{dom}(\nabla) \subseteq \text{dom}(\bar{L}), \quad (2.13)$$

establishing (2.9). \square

The proof of the main result of this section relies on the following lemma which yields a useful resolvent-type identity.

Lemma 2.3. *Assume Hypothesis 2.1 and set*

$$R_m := (I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1} : [L^2(\mathbb{R}^n)]^N \rightarrow [L^2(\mathbb{R}^n)]^N, \quad m \in \mathbb{N}. \quad (2.14)$$

Then the following resolvent identity holds:

$$\begin{aligned} (\bar{L}R_m - R_mL^*)g &= -\frac{1}{m} \sum_{j,k=1}^n (\partial_k R_m d_k(F_j) + R_m d_k(F_j) \partial_k) \partial_j R_m g \\ &\quad - \frac{1}{m} \sum_{j,k=1}^n \partial_j (\partial_k R_m d_k(F_j^*) + R_m d_k(F_j^*) \partial_k) R_m g \end{aligned} \quad (2.15)$$

for all $g \in \text{dom}(L^)$ and all $m \in \mathbb{N}$. In particular, the commutator*

$$\bar{L}(I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1} - (I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1}L^* : \text{dom}(L^*) \rightarrow [L^2(\mathbb{R}^n)]^N \quad (2.16)$$

extends to a bounded operator $X_m : [L^2(\mathbb{R}^n)]^N \rightarrow [L^2(\mathbb{R}^n)]^N$ for all $m \in \mathbb{N}$ and

$$\sup_{m \in \mathbb{N}} \|X_m\|_{\mathcal{B}([L^2(\mathbb{R}^n)]^N)} < \infty. \quad (2.17)$$

Proof. The second claim of the lemma follows from the estimate

$$\left\| \partial_j \partial_k R_m \right\|_{\mathcal{B}([L^2(\mathbb{R}^n)]^N)} \leq m, \quad m \in \mathbb{N}, \quad (2.18)$$

and from our assumption that

$$d_k(F_j^*), d_k(F_j) : [L^2(\mathbb{R}^n)]^N \rightarrow [L^2(\mathbb{R}^n)]^N, \quad 1 \leq j, k \leq n, j, k \in \mathbb{N}, \quad (2.19)$$

are bounded. One notes also that

$$X_m^* = -X_m, \quad m \in \mathbb{N}. \quad (2.20)$$

Therefore, we focus on proving identity (2.15).

We first prove identity (2.15) for $f \in \text{dom}(L) = \mathcal{D}$. To this end, let $f \in \mathcal{D}$ be given. The desired identity is a result of the following computation (applying Proposition 2.2 to take care of domain issues):

$$\begin{aligned} \bar{L}R_m f - R_m L f &= -i \sum_{j=1}^n (F_j R_m - R_m F_j) \partial_j f - i \sum_{j=1}^n \partial_j (F_j^* R_m - R_m F_j^*) f \\ &= \frac{i}{m} \sum_{j=1}^n (\Delta R_m F_j R_m - R_m F_j \Delta R_m) (\partial_j f) \\ &\quad + \frac{i}{m} \sum_{j=1}^n \partial_j (\Delta R_m F_j^* R_m - R_m F_j^* \Delta R_m) f \\ &= -\frac{1}{m} \sum_{j,k=1}^n (\partial_k R_m d_k(F_j) R_m + R_m d_k(F_j) \partial_k R_m) (\partial_j f) \\ &\quad - \frac{1}{m} \sum_{j,k=1}^n \partial_j (\partial_k R_m d_k(F_j^*) R_m + R_m d_k(F_j^*) \partial_k R_m) f. \end{aligned} \quad (2.21)$$

We have now shown that identity (2.15) holds for all $f \in \text{dom}(L)$, that is,

$$(\bar{L}R_m - R_m L) f = X_m f, \quad f \in \text{dom}(L). \quad (2.22)$$

For $g \in \text{dom}(L^*)$ one then computes

$$\begin{aligned} ((\bar{L}R_m - R_m L^*)g, f)_{[L^2(\mathbb{R}^n)]^N} &= (g, (R_m L - \bar{L}R_m)f)_{[L^2(\mathbb{R}^n)]^N} \\ &= -(g, X_m f)_{[L^2(\mathbb{R}^n)]^N} \\ &= (X_m g, f)_{[L^2(\mathbb{R}^n)]^N}, \quad f \in \text{dom}(L), \end{aligned} \quad (2.23)$$

where the last equality makes use of (2.20). Since $\text{dom}(L)$ is dense in $[L^2(\mathbb{R}^n)]^N$, this proves identity (2.15). \square

Lemma 2.3, in particular (2.15), is essentially a restatement of the resolvent identity. The slightly complicated form of this identity is due to our assumptions on the coefficients F_j, F_j^* , $1 \leq j \leq n$. Indeed, we do *not* assume that these operators preserve the domain of the Laplacian $\Delta : [W^{2,2}(\mathbb{R}^n)]^N \rightarrow [L^2(\mathbb{R}^n)]^N$. In fact, they are only assumed to preserve the domain of the gradient $\nabla : [W^{1,2}(\mathbb{R}^n)]^N \rightarrow ([L^2(\mathbb{R}^n)]^N)^n$.

With these preparations in place, we are now ready to state and prove the main result of this section.

Theorem 2.4. *Assume Hypothesis 2.1. Then $L : \mathcal{D} \rightarrow [L^2(\mathbb{R}^n)]^N$ is essentially self-adjoint.*

Proof. The operator L is symmetric by construction. To prove that L is essentially self-adjoint it suffices to show that

$$\text{dom}(L^*) \subseteq \text{dom}(\bar{L}). \quad (2.24)$$

To this end, let $g \in \text{dom}(L^*)$. One notes that

$$\lim_{m \rightarrow \infty} (I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1}g = g, \quad (2.25)$$

and, moreover,

$$(I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1}g \in \text{dom}(\bar{L}), \quad (2.26)$$

since $[W^{2,2}(\mathbb{R}^n)]^N = \text{dom}(-\Delta) \subseteq [W^{1,2}(\mathbb{R}^n)]^N \subseteq \text{dom}(\bar{L})$. By Lemma 2.3 one infers

$$\bar{L}(I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1}g = (I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1}L^*g + X_m g, \quad m \in \mathbb{N}, \quad (2.27)$$

so that the sequence

$$\left\{ \bar{L}(I_{[L^2(\mathbb{R}^n)]^N} - m^{-1}\Delta)^{-1}g \right\}_{m \in \mathbb{N}} \quad (2.28)$$

is bounded in the norm of $[L^2(\mathbb{R}^n)]^N$, implying $g \in \text{dom}(\bar{L})$. \square

Remark 2.5. Theorem 2.4 extends Proposition 10.2.11 in Higson and Roe [31] (see also [18]) in the smooth context, while we now permit Lipschitz coefficients F_j in the differential expression L in (2.8). More precisely, assuming smooth coefficients, Higson and Roe prove that any first-order, symmetric differential operator of finite propagation speed acting on the smooth compactly supported sections of a smooth hermitian vector bundle on any complete manifold (without boundary) is essentially self-adjoint. While the results proved here are very similar in nature (although restricted to the flat case), the actual strategy of proof in Theorem 2.4 differs from the one employed in [31] and is inspired by ideas appearing in [33] and [34], [40] in the context of Hilbert C^* -modules. In a broader picture our methods relate to the noncommutative geometry program [19], since our proof is in some sense coordinate-free and can therefore be readily generalized to a much wider array of unbounded operators of the form $\sum_j A_j \cdot D_j + B$ (with D_j and D_k commuting for all $1 \leq j, k \leq n$), provided that an appropriate reference operator, for example $\sum_j D_j^2$, is already well-understood.

For a very recent approach to essential self-adjointness of first-order differential operators with applications to Dirac-type operators we refer to [7] (the approach in [7] is quite different, relying on ellipticity conditions which are not used in our setup).

Needless to stress, self-adjointness is one of the single most important properties of an unbounded operator, because of its implications to the spectrum and to the Borel functional calculus. \diamond

3. A GLOBAL LIMITING ABSORPTION PRINCIPLE FOR FREE, MASSLESS DIRAC OPERATORS

In 1999, Iftimovici and Măntoiu [32] proved a global limiting absorption principle, that is, one on the entire real axis and hence including threshold energies $\pm m$ for

the free, massive Dirac operator $H_0(m)$ with mass $m > 0$ in three dimensions. The first proof of a global limiting absorption principle for massless Dirac operators H_0 in three dimensions is due to Saitō and Umeda [46] in 2008. As no other result on a global limiting absorption principle in the massless case is known to us, we now fill this gap and upon modifying the approach by Iftimovici and Măntoiu for $m > 0$, we treat free, massless ($m = 0$) Dirac operators H_0 in all dimensions $n \in \mathbb{N}$, $n \geq 2$. This includes, in particular, a new result for the case $n = 2$ which is known to be connected to applications to graphene. In this context, we emphasize, however, the recent result by Erdoğan, Goldberg, and Green [21], who prove a limiting absorption principle for Dirac operators with continuous matrix-valued short-range potentials in the massless case $m = 0$ (resp., in the massive case $m > 0$) in two and higher dimensions, valid on $(-\infty, 0) \cup (0, \infty)$ (resp., valid on $(-\infty, -m) \cup (m, \infty)$), assuming regular thresholds.

Here the notion “free” Dirac operator refers to a particular constant coefficient first-order matrix-valued differential operator with vanishing electric (and magnetic) potentials, see (3.2).

To rigorously define the free massless n -dimensional Dirac operators to be studied in this manuscript, we introduce the following basic assumption.

Hypothesis 3.1. *Let $n \in \mathbb{N}$, $n \geq 2$, set $N = 2^{\lfloor (n+1)/2 \rfloor}$, and denote by α_j , $1 \leq j \leq n$, $\alpha_{n+1} := \beta$, $n+1$ anti-commuting self-adjoint $N \times N$ matrices with squares equal to I_N , that is,*

$$\alpha_j^* = \alpha_j, \quad \alpha_j \alpha_k + \alpha_k \alpha_j = 2\delta_{j,k} I_N, \quad 1 \leq j, k \leq n+1. \quad (3.1)$$

Given Hypothesis 3.1, we introduce in $[L^2(\mathbb{R}^n)]^N$ the free massless Dirac operator as follows,

$$H_0 = \alpha \cdot (-i\nabla) = \sum_{j=1}^n \alpha_j (-i\partial_j), \quad \text{dom}(H_0) = [W^{1,2}(\mathbb{R}^n)]^N. \quad (3.2)$$

Employing the relations (3.1), one observes that

$$H_0^2 = -I_N \Delta, \quad \text{dom}(H_0^2) = [W^{2,2}(\mathbb{R}^n)]^N. \quad (3.3)$$

For completeness we also recall that the massive free Dirac operator in $[L^2(\mathbb{R}^n)]^N$ associated with the mass parameter $m > 0$ then would be of the form

$$H_0(m) = H_0 + m\beta, \quad \text{dom}(H_0(m)) = [W^{1,2}(\mathbb{R}^n)]^N, \quad m > 0, \quad \beta = \alpha_{n+1}, \quad (3.4)$$

but we will primarily study the massless case $m = 0$ in this paper.

In the special one-dimensional case $n = 1$, one can choose α_1 to be one of the three Pauli matrices. Similarly, in the massive case, β would typically be a second Pauli matrix (different from α_1).

The main goal of this section is to obtain a uniform limiting absorption principle for the free n -dimensional massless Dirac operator in dimensions $n \geq 2$. The method of proof employed relies on Kato’s inequality (cf. (3.23), not to be confused with his distributional inequality) and the construction of an auxiliary operator A that has a positive commutator with the free massless Dirac operator H_0 .

To set the stage for the definition of A we introduce the following assumption:

Hypothesis 3.2. *Let $\eta : \mathbb{R}^n \rightarrow \mathbb{R}$ denote a radial function of the form*

$$\eta(p) = h(|p|), \quad p \in \mathbb{R}^n, \quad (3.5)$$

where $h : [0, \infty) \rightarrow [0, 1]$ is defined by

$$h(r) := \begin{cases} r, & r \in [0, 1/2), \\ k(r), & r \in [1/2, 1), \\ 1, & r \in [1, \infty), \end{cases} \quad (3.6)$$

and the function $k : [1/2, 1) \rightarrow [0, \infty)$ is nondecreasing and chosen so that $\eta \in C^\infty(\mathbb{R}^n \setminus \{0\})$.

The operator of multiplication by the independent variable x_j in $[L^2(\mathbb{R}^n)]^N$ will be denoted by Q_j and we shall write

$$Q = (Q_1, \dots, Q_n). \quad (3.7)$$

Given Hypotheses 3.1 and 3.2, we introduce in $[L^2(\mathbb{R}^n)]^N$ the operator

$$\begin{aligned} A &= \frac{1}{2} \left[(\alpha \cdot (-i\nabla))(-\Delta)^{-1}\eta(-i\nabla)((-i\nabla) \cdot Q) \right. \\ &\quad \left. + (Q \cdot (-i\nabla))(-\Delta)^{-1}\eta(-i\nabla)(\alpha \cdot (-i\nabla)) \right], \\ \text{dom}(A) &= \mathcal{D}_0(\mathbb{R}^n), \end{aligned} \quad (3.8)$$

where

$$\mathcal{D}_0(\mathbb{R}^n) = \mathcal{F}^{-1} \left([C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N \right) \subset [S(\mathbb{R}^n)]^N. \quad (3.9)$$

That is, $\mathcal{D}_0(\mathbb{R}^n)$ consists of functions whose Fourier transforms have compact support and no support in a neighborhood of $p = 0$.

In addition, we introduce

$$B = \eta(-i\nabla) \in \mathcal{B} \left([L^2(\mathbb{R}^n)]^N \right), \quad (3.10)$$

defined via the spectral theorem.

We emphasize that the necessity of including the factor $\eta(-i\nabla)$ in the definition (3.8) of A considerably complicates matters as at various occasions we will have to consider B^{-1} in combination with other operators. The corresponding massive case, $m > 0$, as treated by Iftimovici and Măntoiu [32], corresponds to the bounded operator $(-\Delta + m^2 I)^{-1}$ instead of $(-\Delta)^{-1}$ in A , and hence does not require the introduction of the term $\eta(-i\nabla)$ in (3.8). Naturally, this considerably influences some technical aspects in the proofs of this section.

For any $n \in \mathbb{N}$, we also introduce the scale of weighted L^2 -spaces,

$$\begin{aligned} L_s^2(\mathbb{R}^n) &= \{f \in L^2(\mathbb{R}^n) \mid \langle Q \rangle^s f \in L^2(\mathbb{R}^n)\} = L^2(\mathbb{R}^n; (1 + |x|^2)^s dx), \\ L_{-s}^2(\mathbb{R}^n) &= [L_s^2(\mathbb{R}^n)]^*, \quad s \geq 0. \end{aligned} \quad (3.11)$$

Proposition 3.3. *Assume Hypotheses 3.1 and 3.2. Then the operator A is essentially self-adjoint on $\mathcal{D}_0(\mathbb{R}^n)$ and*

$$[L_1^2(\mathbb{R}^n)]^N \subseteq \text{dom}(\overline{A}). \quad (3.12)$$

Proof. Considering $\widehat{A} = \mathcal{F}A\mathcal{F}^{-1}$, \widehat{A} reads

$$\begin{aligned} \widehat{A} &= \frac{1}{2} \left[(\alpha \cdot Q)|Q|^{-2}\eta(Q)(Q \cdot (-i\nabla)) + ((-i\nabla) \cdot Q)|Q|^{-2}\eta(Q)(\alpha \cdot Q) \right], \\ \text{dom}(\widehat{A}) &= [C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N. \end{aligned} \quad (3.13)$$

In this context we note that $C_0^\infty(\mathbb{R}^n \setminus \{0\})$, $n \in \mathbb{N}$, $n \geq 2$, is a core for ∇ (see, e.g., [25, p. 97]). Therefore, the operator \widehat{A} is a first-order differential operator of the form introduced in (2.8) with F_j , $1 \leq j \leq n$, defined by

$$F_j(p) = \frac{1}{2}(\alpha \cdot p)|p|^{-2}\eta(p)p_j, \quad p \in \mathbb{R}^n \setminus \{0\}, \quad 1 \leq j \leq n. \quad (3.14)$$

Given Hypothesis 3.2 on η , one can check that F_j leaves the core $C_0^\infty(\mathbb{R}^n \setminus \{0\})$ invariant (in particular, the $|x|$ -behavior of $\eta(Q)$ near $x = 0$ is not felt by functions in $C_0^\infty(\mathbb{R}^n \setminus \{0\})$). Moreover, Hypothesis 3.2 also guarantees that the partial derivatives of F_j are bounded functions, and therefore

$$\overline{[-i\partial_k, F_j]} \in \mathcal{B}\left([L^2(\mathbb{R}^n)]^N\right), \quad 1 \leq j, k \leq n. \quad (3.15)$$

Thus, the assumptions of Hypothesis 2.1 are satisfied, and therefore, Theorem 2.4 implies that the operator \widehat{A} is essentially self-adjoint. The inclusion (3.12) then follows from (2.9). \square

In the next result, we compute the commutator of H_0 with A .

Proposition 3.4. *Assume Hypotheses 3.1 and 3.2, then*

$$i[A, H_0]\psi = B\psi, \quad \psi \in \mathcal{D}_0(\mathbb{R}^n). \quad (3.16)$$

Therefore, for each $\psi_1, \psi_2 \in [L_1^2(\mathbb{R}^n)]^N \cap [W^{1,2}(\mathbb{R}^n)]^N$,

$$(\psi_1, B\psi_2)_{[L^2(\mathbb{R}^n)]^N} = (iH_0\psi_1, \overline{A}\psi_2)_{[L^2(\mathbb{R}^n)]^N} - (i\overline{A}\psi_1, H_0\psi_2)_{[L^2(\mathbb{R}^n)]^N}. \quad (3.17)$$

Proof. We compute the commutator of $\widehat{H}_0 = \alpha \cdot Q$ and

$$\widehat{A} = \frac{1}{2}\left(\widehat{H}_0|Q|^{-2}\eta(Q)(Q \cdot (-i\nabla)) + ((-i\nabla) \cdot Q)\widehat{H}_0|Q|^{-2}\eta(Q)\right) \quad (3.18)$$

on $[C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N$. To this end, one observes that

$$\begin{aligned} [\widehat{H}_0, Q \cdot (-i\nabla)] &= [\widehat{H}_0, (-i\nabla) \cdot Q] \\ &= \sum_{j,k=1}^n [\alpha_j Q_j, Q_k (-i\partial_k)] \\ &= -i \sum_{j=1}^n Q_j \alpha_j [Q_j, \partial_j] \\ &= i\widehat{H}_0 \end{aligned} \quad (3.19)$$

on $[C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N$. As a result, one obtains

$$\begin{aligned} [\widehat{H}_0, \widehat{A}] &= 2^{-1}\widehat{H}_0|Q|^{-2}\eta(Q)[\widehat{H}_0, Q \cdot (-i\nabla)] + 2^{-1}[\widehat{H}_0, (-i\nabla) \cdot Q]\widehat{H}_0|Q|^{-2}\eta(Q) \\ &= (i/2)\widehat{H}_0|Q|^{-2}\eta(Q)\widehat{H}_0 + (i/2)\widehat{H}_0\widehat{H}_0|Q|^{-2}\eta(Q) \\ &= i\eta(Q) \\ &= i\widehat{B} \end{aligned} \quad (3.20)$$

on $[C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N$, where

$$\widehat{B} := \eta(Q). \quad (3.21)$$

\square

Assuming Hypothesis 3.2, the square root $B^{1/2}$ is defined by the spectral theorem via

$$B^{1/2} = \eta(-i\nabla)^{1/2} \in \mathcal{B}\left([L^2(\mathbb{R}^n)]^N\right). \quad (3.22)$$

The next result requires Kato's inequality in \mathbb{R}^n , $n \geq 2$, which is of the form (cf., e.g., [5, p. 19], [30])

$$K_n \int_{\mathbb{R}^n} d^n x |x|^{-1} |f(x)|^2 \leq \int_{\mathbb{R}^n} d^n p |p| |\widehat{f}(p)|^2, \quad f \in \text{dom}((-\Delta)^{1/4}), \quad n \in \mathbb{N}, \quad n \geq 2, \quad (3.23)$$

for some constants $K_n > 0$. In particular,

$$K_n \int_{\mathbb{R}^n} d^n x |x|^{-1} |f(x)|^2 \leq \int_{\mathbb{R}^n} d^n p \langle p \rangle |\widehat{f}(p)|^2, \quad \widehat{f} \in L^2_{1/2}(\mathbb{R}^n), \quad n \in \mathbb{N}, \quad n \geq 2, \quad (3.24)$$

equivalently, and in the form to be used below,

$$K_n \int_{\mathbb{R}^n} d^n p |p|^{-1} |\widehat{f}(p)|^2 \leq \int_{\mathbb{R}^n} d^n x \langle x \rangle |f(x)|^2, \quad f \in L^2_{1/2}(\mathbb{R}^n), \quad n \in \mathbb{N}, \quad n \geq 2. \quad (3.25)$$

Proposition 3.5. *Assume Hypothesis 3.2. Then*

$$B^{-1/2} \in \mathcal{B}\left([L^2_{1/2}(\mathbb{R}^n)]^N, [L^2(\mathbb{R}^n)]^N\right), \quad (3.26)$$

that is, there exists $C_1 > 0$ such that

$$\|B^{-1/2}\psi\|_{[L^2(\mathbb{R}^n)]^N} \leq C_1 \|\psi\|_{[L^2_{1/2}(\mathbb{R}^n)]^N}, \quad \psi \in [L^2_{1/2}(\mathbb{R}^n)]^N. \quad (3.27)$$

Proof. The nonnegative function $\eta(p)^{-1/2}$ of $p = (p_1, \dots, p_n)$ satisfies

$$\eta(p)^{-1/2} \leq 2^{1/2} + \frac{1}{|p|^{1/2}}, \quad p \neq 0, \quad (3.28)$$

so it suffices to note that

$$|-i\nabla|^{-1/2} \in \mathcal{B}\left([L^2_{1/2}(\mathbb{R}^n)]^N, [L^2(\mathbb{R}^n)]^N\right), \quad (3.29)$$

which follows from Kato's inequality in the form of (3.25),

$$\begin{aligned} \|| -i\nabla|^{-1/2}\psi\|_{[L^2(\mathbb{R}^n)]^N} &= \|| |p|^{-1/2}\widehat{\psi}\|_{[L^2(\mathbb{R}^n)]^N} \leq K_n^{-1/2} \|\psi\|_{[L^2_{1/2}(\mathbb{R}^n)]^N}, \\ &\psi \in [L^2_{1/2}(\mathbb{R}^n)]^N, \quad n \geq 2. \end{aligned} \quad (3.30)$$

□

Proposition 3.6. *Assume Hypotheses 3.1 and 3.2. Then there exists a constant $C_2 > 0$ such that*

$$\|B^{-1/2}A\psi\|_{[L^2(\mathbb{R}^n)]^N} \leq C_2 \|\psi\|_{[L^2_1(\mathbb{R}^n)]^N}, \quad \psi \in \text{dom}(A) = \mathcal{D}_0(\mathbb{R}^n). \quad (3.31)$$

Thus, the operator $B^{-1/2}A : \mathcal{D}_0(\mathbb{R}^n) \rightarrow [L^2(\mathbb{R}^n)]^N$ extends to an element

$$B^{-1/2}\overline{A} \in \mathcal{B}\left([L^2_1(\mathbb{R}^n)]^N, [L^2(\mathbb{R}^n)]^N\right). \quad (3.32)$$

Proof. With \widehat{A} defined by (3.18), write

$$\widehat{A} = \frac{1}{2}(\widehat{A}_\ell + \widehat{A}_r), \quad (3.33)$$

where

$$\widehat{A}_\ell = \widehat{H}_0|Q|^{-2}\eta(Q)(Q \cdot (-i\nabla)), \quad (3.34)$$

$$\widehat{A}_r = ((-i\nabla) \cdot Q)\widehat{H}_0|Q|^{-2}\eta(Q), \quad (3.35)$$

on $[C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N$. Then one observes that

$$\begin{aligned} \widehat{A}_r &= -i \sum_{j=1}^n \partial_j Q_j \widehat{H}_0|Q|^{-2}\eta(Q) \\ &= -i \sum_{j=1}^n [\partial_j, Q_j \widehat{H}_0|Q|^{-2}\eta(Q)] + \widehat{A}_\ell. \end{aligned} \quad (3.36)$$

One notes that the operator $Q_j \widehat{H}_0|Q|^{-2}\eta(Q)$ represents the operator $F_j(Q)$, where F_j is defined according to (3.14). Hence, by (3.15) it follows that the commutator $[\partial_j, Q_j \widehat{H}_0|Q|^{-2}\eta(Q)]$ extends to a bounded operator for every $j = 1, \dots, n$. Therefore, the operator \widehat{A}_0 defined by

$$\widehat{A}_0 := -i \sum_{j=1}^n [\partial_j, Q_j \widehat{H}_0|Q|^{-2}\eta(Q)], \quad \text{dom}(\widehat{A}_0) = [C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N, \quad (3.37)$$

extends to a bounded operator

$$\overline{\widehat{A}_0} : [L^2(\mathbb{R}^n)]^N \rightarrow [L^2(\mathbb{R}^n)]^N \quad (3.38)$$

commuting with $\widehat{B} = \eta(Q)$. In particular, one infers that

$$\widehat{A} = \widehat{A}_\ell + \frac{1}{2}\overline{\widehat{A}_0} : [C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N \rightarrow [L^2(\mathbb{R}^n)]^N. \quad (3.39)$$

Writing

$$A_\ell = (\alpha \cdot (-i\nabla))(-\Delta)^{-1}\eta(-i\nabla)((-i\nabla) \cdot Q), \quad (3.40)$$

one obtains

$$\begin{aligned} \|B^{-1/2}A_\ell\psi\|_{[L^2(\mathbb{R}^n)]^N} &= \|(\alpha \cdot (-i\nabla))(-\Delta)^{-1}\eta(-i\nabla)^{1/2}((-i\nabla) \cdot Q)\psi\|_{[L^2(\mathbb{R}^n)]^N} \\ &\leq C_\ell \|\psi\|_{[L^2_1(\mathbb{R}^n)]^N}, \quad \psi \in \mathcal{D}_0(\mathbb{R}^n), \end{aligned} \quad (3.41)$$

for an appropriate constant $C_\ell > 0$, by the (bounded) functional calculus for self-adjoint operators. Finally, one uses (3.41), the fact that $\overline{\widehat{A}_0}$ commutes with \widehat{B} , boundedness of $\overline{\widehat{A}_0}$, and Proposition 3.5, to obtain

$$\begin{aligned} \|B^{-1/2}A\psi\|_{[L^2(\mathbb{R}^n)]^N} &\leq \|B^{-1/2}A_\ell\psi\|_{[L^2(\mathbb{R}^n)]^N} + 2^{-1} \left\| \overline{\widehat{A}_0} \widehat{B}^{-1/2} \widehat{\psi} \right\|_{[L^2(\mathbb{R}^n)]^N} \\ &\leq C_2 \|\psi\|_{[L^2_1(\mathbb{R}^n)]^N}, \quad \psi \in \text{dom}(A), \end{aligned} \quad (3.42)$$

for an appropriate constant $C_2 > 0$. \square

Next, we investigate the commutator of A with B .

Proposition 3.7. *Assume Hypotheses 3.1 and 3.2. Then B leaves the domain of \bar{A} invariant, and there exists $K \in \mathcal{B}([L^2(\mathbb{R}^n)]^N)$, which commutes with B , such that*

$$i[B, \bar{A}]\psi = KB\psi, \quad \psi \in \text{dom}(\bar{A}). \quad (3.43)$$

Thus, $i[B, \bar{A}]$ extends to a bounded operator in $\mathcal{B}([L^2(\mathbb{R}^n)]^N)$.

Moreover, if $\psi_1, \psi_2 \in [L^2_1(\mathbb{R}^n)]^N$, then

$$(\psi_1, KB\psi_2)_{[L^2(\mathbb{R}^n)]^N} = (i\bar{A}\psi_1, B\psi_2)_{[L^2(\mathbb{R}^n)]^N} - (iB\psi_1, \bar{A}\psi_2)_{[L^2(\mathbb{R}^n)]^N}. \quad (3.44)$$

Proof. Using (3.39), one computes

$$\begin{aligned} i[\widehat{B}, \widehat{A}] &= i[\widehat{B}, \widehat{A}_\ell + (1/2)\widehat{A}_0] \\ &= i[\widehat{B}, \widehat{A}_\ell] \\ &= -\widehat{H}_0|Q|^{-2}\eta(Q) \sum_{j=1}^n Q_j[\partial_j, \eta(Q)] \\ &= -\widehat{H}_0\eta(Q)((\nabla\eta)(Q) \cdot Q|Q|^{-2}) \end{aligned} \quad (3.45)$$

on $[C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N$. The claim now follows as

$$-\widehat{H}_0((\nabla\eta)(Q) \cdot Q|Q|^{-2}) : [C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N \rightarrow [L^2(\mathbb{R}^n)]^N \quad (3.46)$$

extends to a bounded operator on $[L^2(\mathbb{R}^n)]^N$ by Hypothesis 3.2 and since the set $[C_0^\infty(\mathbb{R}^n \setminus \{0\})]^N$ is a core for \widehat{A} by definition. \square

Assuming Hypotheses 3.1 and 3.2, define

$$T_\varepsilon^\pm(\lambda, \mu) = H_0 - (\lambda \pm i\mu)I_{[L^2(\mathbb{R}^n)]^N} \mp i\varepsilon B, \quad \lambda \in \mathbb{R}, \mu > 0, \varepsilon \in [0, 1]. \quad (3.47)$$

Proposition 3.8. *Assume Hypotheses 3.1 and 3.2. For each $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$, $\varepsilon \in [0, 1)$, $T_\varepsilon^\pm(\lambda, \mu)$ is a linear homeomorphism from $[W^{1,2}(\mathbb{R}^n)]^N$ to $[L^2(\mathbb{R}^n)]^N$ satisfying*

$$\|T_\varepsilon^\pm(\lambda, \mu)\psi\|_{[L^2(\mathbb{R}^n)]^N}^2 \geq \mu^2\|\psi\|_{[L^2(\mathbb{R}^n)]^N}^2, \quad \psi \in [W^{1,2}(\mathbb{R}^n)]^N. \quad (3.48)$$

Proof. Let $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$, and $\varepsilon \in [0, 1)$. The claim is evident if $\varepsilon = 0$, so we assume from now on that $\varepsilon \neq 0$. To obtain the lower bound in (3.48), let $\psi \in [W^{1,2}(\mathbb{R}^n)]^N$. Since H_0 is self-adjoint and B is nonnegative,

$$\begin{aligned} &\text{Re}\{([H_0 - \lambda I_{[L^2(\mathbb{R}^n)]^N} \mp i\varepsilon B]\psi, \mp i\mu\psi)_{[L^2(\mathbb{R}^n)]^N}\} \\ &= \text{Re}\{\mp i\mu([H_0 - \lambda I_{[L^2(\mathbb{R}^n)]^N}]\psi, \psi)_{[L^2(\mathbb{R}^n)]^N}\} + \text{Re}\{\varepsilon\mu(B\psi, \psi)_{[L^2(\mathbb{R}^n)]^N}\} \\ &= \varepsilon\mu(\psi, B\psi)_{[L^2(\mathbb{R}^n)]^N} \geq 0, \quad \psi \in [W^{1,2}(\mathbb{R}^n)]^N, \end{aligned} \quad (3.49)$$

and therefore,

$$\begin{aligned} &\|T_\varepsilon^\pm(\lambda, \mu)\psi\|_{[L^2(\mathbb{R}^n)]^N}^2 \\ &= \|[H_0 - \lambda I_{[L^2(\mathbb{R}^n)]^N} \mp i\varepsilon B]\psi\|_{[L^2(\mathbb{R}^n)]^N}^2 + \mu^2\|\psi\|_{[L^2(\mathbb{R}^n)]^N}^2 \\ &\quad + 2\text{Re}\{([H_0 - \lambda I_{[L^2(\mathbb{R}^n)]^N} \mp i\varepsilon B]\psi, \mp i\mu\psi)_{[L^2(\mathbb{R}^n)]^N}\} \\ &\geq \mu^2\|\psi\|_{[L^2(\mathbb{R}^n)]^N}^2, \quad \psi \in [W^{1,2}(\mathbb{R}^n)]^N. \end{aligned} \quad (3.50)$$

It is clear that

$$T_\varepsilon^\pm(\lambda, \mu) \in \mathcal{B}\left([W^{1,2}(\mathbb{R}^n)]^N, [L^2(\mathbb{R}^n)]^N\right) \quad (3.51)$$

for each $\lambda \in \mathbb{R}$ and $\mu \in (0, \infty)$, since the operator $i\varepsilon B$ is bounded. Moreover, $T_\varepsilon^\pm(\lambda, \mu)$ is a closed bijection. In fact, (3.48) immediately implies that $T_\varepsilon^\pm(\lambda, \mu)$ is an injection with a closed range. In addition, since $[T_\varepsilon^\pm(\lambda, \mu)]^* = T_\varepsilon^\mp(\lambda, \mu)$, $T_\varepsilon^\pm(\lambda, \mu)$ is a surjection. \square

By Proposition 3.8, $T_\varepsilon^\pm(\lambda, \mu)$ is a linear homeomorphism from $[W^{1,2}(\mathbb{R}^n)]^N$ to $[L^2(\mathbb{R}^n)]^N$ for each $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$, $\varepsilon \in [0, 1)$. Therefore, we define

$$G_\varepsilon^\pm(\lambda, \mu) := (T_\varepsilon^\pm(\lambda, \mu))^{-1} \in \mathcal{B}\left([L^2(\mathbb{R}^n)]^N, [W^{1,2}(\mathbb{R}^n)]^N\right) \subset \mathcal{B}\left([L^2(\mathbb{R}^n)]^N\right), \\ \lambda \in \mathbb{R}, \mu \in (0, \infty), \varepsilon \in [0, 1), \quad (3.52)$$

noting that $G_\varepsilon^\pm(\lambda, \mu)$ are adjoint to one another,

$$(G_\varepsilon^\pm(\lambda, \mu))^* = G_\varepsilon^\mp(\lambda, \mu), \quad \lambda \in \mathbb{R}, \mu \in (0, \infty), \varepsilon \in [0, 1). \quad (3.53)$$

We also define the scalar-valued function

$$F_\varepsilon^\pm(\lambda, \mu; \psi) := (\psi, G_\varepsilon^\pm(\lambda, \mu)\psi)_{[L^2(\mathbb{R}^n)]^N}, \\ \lambda \in \mathbb{R}, \mu \in (0, \infty), \varepsilon \in [0, 1), \psi \in [L^2(\mathbb{R}^n)]^N. \quad (3.54)$$

The following result is a standard application of the second resolvent identity, so we omit the details of its proof.

Proposition 3.9. *Assume Hypotheses 3.1 and 3.2. For each $\psi \in [L^2(\mathbb{R}^n)]^N$, $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$, the function*

$$F^\pm(\lambda, \mu; \psi) : \begin{cases} [0, 1) \rightarrow \mathbb{C}, \\ \varepsilon \mapsto F_\varepsilon^\pm(\lambda, \mu; \psi), \end{cases} \quad (3.55)$$

is smooth on $[0, 1)$ and

$$\frac{dF_\varepsilon^\pm(\lambda, \mu; \psi)}{d\varepsilon} = \pm(\psi, iG_\varepsilon^\pm(\lambda, \mu)BG_\varepsilon^\pm(\lambda, \mu)\psi)_{[L^2(\mathbb{R}^n)]^N}, \quad \varepsilon \in [0, 1). \quad (3.56)$$

Proposition 3.10. *Assume Hypotheses 3.1 and 3.2. $G_\varepsilon^\pm(\lambda, \mu)$ leaves $[L_1^2(\mathbb{R}^n)]^N$ invariant:*

$$G_\varepsilon^\pm(\lambda, \mu)[L_1^2(\mathbb{R}^n)]^N \subset [L_1^2(\mathbb{R}^n)]^N. \quad (3.57)$$

In addition, the following estimates hold:

$$\|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\psi\|_{[L^2(\mathbb{R}^n)]^N}^2 \leq \varepsilon^{-1}|F_\varepsilon^\pm(\lambda, \mu; \psi)|, \quad \psi \in [L^2(\mathbb{R}^n)]^N, \quad (3.58)$$

$$\|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\phi\|_{[L^2(\mathbb{R}^n)]^N} \leq C_1\varepsilon^{-1}\|\phi\|_{[L_{1/2}^2(\mathbb{R}^n)]^N}, \quad \phi \in [L_{1/2}^2(\mathbb{R}^n)]^N, \quad (3.59) \\ \lambda \in \mathbb{R}, \mu \in (0, \infty), \varepsilon \in (0, 1),$$

where C_1 is the same constant as in (3.27).

Proof. To prove the invariance claim, it suffices to show that $G_\varepsilon^\pm(\lambda, \mu)$ preserves the domain of the unbounded self-adjoint operator Q_j for all $1 \leq j \leq n$. Thus, let $j \in \mathbb{N}$, with $1 \leq j \leq n$, be fixed. Note that the sequence of bounded operators

$$\left\{ i(iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1} \right\}_{m=1}^\infty \quad (3.60)$$

converges strongly to the identity operator in $[L^2(\mathbb{R}^n)]^N$. One computes

$$\begin{aligned} & \left[iQ_j (iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1}, G_\varepsilon^\pm(\lambda, \mu) \right] \\ &= G_\varepsilon^\pm(\lambda, \mu) \left[H_0 \mp i\varepsilon B, iQ_j (iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1} \right] G_\varepsilon^\pm(\lambda, \mu) \\ &= G_\varepsilon^\pm(\lambda, \mu) \left[H_0 \mp i\varepsilon B, m(iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1} \right] G_\varepsilon^\pm(\lambda, \mu). \end{aligned} \quad (3.61)$$

In addition, the operator B preserves the domain of Q_j and $[Q_j, B] : \text{dom}(Q_j) \rightarrow [L^2(\mathbb{R}^n)]^N$ extends to a bounded operator $q_j(B) \in \mathcal{B}([L^2(\mathbb{R}^n)]^N)$. Therefore,

$$\begin{aligned} & m \left[H_0 \mp i\varepsilon B, (iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1} \right] \\ &= i\alpha_j (iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1} \\ & \quad \mp i\varepsilon (iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1} q_j(B) (iI_{[L^2(\mathbb{R}^n)]^N} + m^{-1}Q_j)^{-1}, \end{aligned} \quad (3.62)$$

which converges strongly as $m \rightarrow \infty$ to

$$-i\alpha_j \pm i\varepsilon q_j(B). \quad (3.63)$$

The fact that $G_\varepsilon^\pm(\lambda, \mu)$ preserves the domain of Q_j now follows since the righthand side of (3.61) converges strongly to

$$-G_\varepsilon^\pm(\lambda, \mu) [i\alpha_j \mp i\varepsilon q_j(B)] G_\varepsilon^\pm(\lambda, \mu). \quad (3.64)$$

To prove (3.58), let $\psi \in [L^2(\mathbb{R}^n)]^N$, $\lambda \in \mathbb{R}$, and $\mu \in (0, \infty)$ be fixed. Then

$$\begin{aligned} & |F_\varepsilon^\pm(\lambda, \mu; \psi)| \geq \pm \text{Im} \left[(\psi, G_\varepsilon^\pm(\lambda, \mu)\psi)_{[L^2(\mathbb{R}^n)]^N} \right] \\ &= \pm \text{Im} \left[((H_0 - (\lambda \pm i\mu)I_{[L^2(\mathbb{R}^n)]^N} \mp i\varepsilon B) G_\varepsilon^\pm(\lambda, \mu)\psi, G_\varepsilon^\pm(\lambda, \mu)\psi)_{[L^2(\mathbb{R}^n)]^N} \right] \\ &= \mu \|G_\varepsilon^\pm(\lambda, \mu)\psi\|_{[L^2(\mathbb{R}^n)]^N}^2 + \varepsilon \|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\psi\|_{[L^2(\mathbb{R}^n)]^N}^2 \\ &\geq \varepsilon \|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\psi\|_{[L^2(\mathbb{R}^n)]^N}^2, \end{aligned} \quad (3.65)$$

which yields (3.58). Next,

$$\begin{aligned} & |F_\varepsilon^\pm(\lambda, \mu; \phi)| = |(\phi, G_\varepsilon^\pm(\lambda, \mu)\phi)_{[L^2(\mathbb{R}^n)]^N}| \\ &\leq \|B^{-1/2}\phi\|_{[L^2(\mathbb{R}^n)]^N} \|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\phi\|_{[L^2(\mathbb{R}^n)]^N} \\ &\leq C_1 \|\phi\|_{[L^2_{1/2}(\mathbb{R}^n)]^N} \|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\phi\|_{[L^2(\mathbb{R}^n)]^N}, \quad \phi \in [L^2_{1/2}(\mathbb{R}^n)]^N. \end{aligned} \quad (3.66)$$

where the last estimate makes use of Proposition 3.5. Upon combining (3.65) and (3.66), one obtains

$$\begin{aligned} & \varepsilon \|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\phi\|_{[L^2(\mathbb{R}^n)]^N}^2 \leq |F_\varepsilon^\pm(\lambda, \mu; \phi)| \\ &\leq C_1 \|\phi\|_{[L^2_{1/2}(\mathbb{R}^n)]^N} \|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\phi\|_{[L^2(\mathbb{R}^n)]^N}, \end{aligned} \quad (3.67)$$

which then implies

$$\|B^{1/2}G_\varepsilon^\pm(\lambda, \mu)\phi\|_{[L^2(\mathbb{R}^n)]^N} \leq C_1 \varepsilon^{-1} \|\phi\|_{[L^2_{1/2}(\mathbb{R}^n)]^N}, \quad \phi \in [L^2_{1/2}(\mathbb{R}^n)]^N. \quad (3.68)$$

and (3.59) follows. \square

The main goal of this section is to prove a uniform limiting absorption principle for H_0 , namely, given $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$, there exists a $C \in (0, \infty)$ (independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$), such that

$$\|(H_0 - (\lambda \pm i\mu)I_{[L^2(\mathbb{R}^n)]^N})^{-1}\|_{\mathcal{B}([L^2_1(\mathbb{R}^n)]^N, [L^2_{-1}(\mathbb{R}^n)]^N)} \leq C, \quad (3.69)$$

$$\lambda \in \mathbb{R}, \mu \in (0, \infty).$$

To accomplish this, it actually suffices to prove

$$|F_0^\pm(\lambda, \mu; \psi)| \leq C' \|\psi\|_{[L^2_1(\mathbb{R}^n)]^N}^2, \quad \psi \in [L^2_1(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty), \quad (3.70)$$

for some constant $C' \in (0, \infty)$ (independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$), by the next lemma. The latter is surely well-known, but we include a proof for completeness.

Lemma 3.11. *Suppose that \mathcal{H}_+ is a Banach space that embeds continuously and densely into the Hilbert space \mathcal{H} so that \mathcal{H} embeds continuously and densely into $\mathcal{H}_- := \mathcal{H}_+^*$. If $H : \text{dom}(H) \subset \mathcal{H} \rightarrow \mathcal{H}$ is a self-adjoint operator, then*

$$\|(H - (\lambda \pm i\mu)I_{\mathcal{H}})^{-1}\|_{\mathcal{B}(\mathcal{H}_+, \mathcal{H}_-)} \leq C_1, \quad \lambda \in \mathbb{R}, \mu \in (0, \infty), \quad (3.71)$$

for some constant $C_1 \in (0, \infty)$ (independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$) if and only if

$$|(\psi, (H - (\lambda \pm i\mu)I_{\mathcal{H}})^{-1}\psi)_{\mathcal{H}}| \leq C_2 \|\psi\|_{\mathcal{H}_+}^2, \quad \psi \in \mathcal{H}_+, \lambda \in \mathbb{R}, \mu \in (0, \infty), \quad (3.72)$$

for some constant $C_2 \in (0, \infty)$ (independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$).

Proof. The equivalence of (3.71) and (3.72) follows from the estimates

$$\sup_{\psi \in \mathcal{H}_+, \|\psi\|_{\mathcal{H}_+} = 1} |(\psi, T\psi)_{\mathcal{H}}| \leq \|T\|_{\mathcal{B}(\mathcal{H}_+, \mathcal{H}_-)} \leq 4 \sup_{\psi \in \mathcal{H}_+, \|\psi\|_{\mathcal{H}_+} = 1} |(\psi, T\psi)_{\mathcal{H}}|, \quad (3.73)$$

which hold for any $T \in \mathcal{B}(\mathcal{H}_+, \mathcal{H})$. In fact, for $T \in \mathcal{B}(\mathcal{H}_+, \mathcal{H})$, one obtains

$$\begin{aligned} \|T\|_{\mathcal{B}(\mathcal{H}_+, \mathcal{H}_-)} &= \sup_{\psi \in \mathcal{H}_+, \|\psi\|_{\mathcal{H}_+} = 1} \|T\psi\|_{\mathcal{H}_-} \\ &= \sup_{\psi \in \mathcal{H}_+, \|\psi\|_{\mathcal{H}_+} = 1} \left(\sup_{\phi \in \mathcal{H}_+, \|\phi\|_{\mathcal{H}_+} = 1} |(T\psi)(\phi)| \right) \\ &= \sup_{\psi, \phi \in \mathcal{H}_+, \|\psi\|_{\mathcal{H}_+} = \|\phi\|_{\mathcal{H}_+} = 1} |(\phi, T\psi)_{\mathcal{H}}|, \end{aligned} \quad (3.74)$$

and (3.73) follows by the polarization principle. \square

The following lemma will be used in the proof of the global limiting absorption principle for H_0 .

Lemma 3.12 (Lemma 7.A.1 in [4]). *Let $(a, b) \subset \mathbb{R}$ be an open interval and let f , ϕ , and ψ be nonnegative real functions on (a, b) with f bounded and measurable and $\phi, \psi \in L^1((a, b); dx)$. Assume there exist constants $\omega \in [0, \infty)$ and $\theta \in [0, 1)$ with*

$$f(\lambda) \leq \omega + \int_{\lambda}^b dt [\phi(t)f(t)^\theta + \psi(t)f(t)], \quad \lambda \in (a, b). \quad (3.75)$$

Then

$$f(\lambda) \leq e^{\int_{\lambda}^b dt \psi(t)} \cdot \left[\omega^{1-\theta} + (1-\theta) \int_{\lambda}^b ds \phi(s) e^{(\theta-1) \int_s^b dt \psi(t)} \right]^{1/(1-\theta)}, \quad \lambda \in (a, b). \quad (3.76)$$

Given these preparations, everything is finally in place to state and prove the principal result of this section, a global limiting absorption principle for H_0 in all dimensions $n \in \mathbb{N}$, $n \geq 2$:

Theorem 3.13. *Assume Hypotheses 3.1 and 3.2 and let $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$. Then*

$$\left\| (H_0 - (\lambda \pm i\mu)I_{[L^2(\mathbb{R}^n)]^N})^{-1} \right\|_{\mathcal{B}([L_1^2(\mathbb{R}^n)]^N, [L_{-1}^2(\mathbb{R}^n)]^N)} \leq C \quad (3.77)$$

for some constant $C \in (0, \infty)$, independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$. Equivalently,

$$\left\| \langle Q \rangle^{-1} (H_0 - (\lambda \pm i\mu)I_{[L^2(\mathbb{R}^n)]^N})^{-1} \langle Q \rangle^{-1} \right\|_{\mathcal{B}([L^2(\mathbb{R}^n)]^N)} \leq C, \quad \lambda \in \mathbb{R}, \mu \in (0, \infty). \quad (3.78)$$

Consequently, $\langle Q \rangle^{-1}$ is H_0 -Kato-smooth, that is, given $\varepsilon_0 > 0$, for each $f \in [L^2(\mathbb{R}^n)]^N$,

$$\begin{aligned} \sup_{\varepsilon \in (0, \varepsilon_0), \|f\|_{[L^2(\mathbb{R}^n)]^N} = 1} \frac{1}{4\pi^2} \int_{\mathbb{R}} d\lambda \left[\left\| \langle Q \rangle^{-1} (H_0 - (\lambda + i\varepsilon)I_{[L^2(\mathbb{R}^n)]^N})^{-1} f \right\|_{[L^2(\mathbb{R}^n)]^N}^2 \right. \\ \left. + \left\| \langle Q \rangle^{-1} (H_0 - (\lambda - i\varepsilon)I_{[L^2(\mathbb{R}^n)]^N})^{-1} f \right\|_{[L^2(\mathbb{R}^n)]^N}^2 \right] < \infty. \end{aligned} \quad (3.79)$$

Proof. By Lemma 3.11, it suffices to show

$$|F_0^\pm(\lambda, \mu; \psi)| \leq C \|\psi\|_{[L_1^2(\mathbb{R}^n)]^N}^2, \quad \psi \in [L_1^2(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty), \quad (3.80)$$

for some constant $C > 0$.

Choose $\psi_1 = G_\varepsilon^\pm(\lambda, \mu)^* \psi$ and $\psi_2 = G_\varepsilon^\pm(\lambda, \mu) \psi$. On one hand, since $\psi \in [L_1^2(\mathbb{R}^n)]^N$, Proposition 3.10 guarantees that $\psi_1, \psi_2 \in [L_1^2(\mathbb{R}^n)]^N$. On the other hand, $\psi_1, \psi_2 \in [W^{1,2}(\mathbb{R}^n)]^N$ since

$$G_\varepsilon^\pm(\lambda, \mu) \in \mathcal{B}\left([L^2(\mathbb{R}^n)]^N, [W^{1,2}(\mathbb{R}^n)]^N\right). \quad (3.81)$$

Therefore, combining the results of Propositions 3.4 and 3.7, one obtains

$$\begin{aligned} \frac{d}{d\varepsilon} F_\varepsilon^\pm(\lambda, \mu; \psi) &\stackrel{(3.56)}{=} \pm i (G_\varepsilon^\mp(\lambda, \mu) \psi, B G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N} \\ &\stackrel{(3.17)}{=} \pm (H_0 G_\varepsilon^\mp(\lambda, \mu) \psi, \overline{A} G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N} \\ &\quad \mp (\overline{A} G_\varepsilon^\mp(\lambda, \mu) \psi, H_0 G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N} \\ &= \pm (\psi, \overline{A} G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N} \mp (\overline{A} G_\varepsilon^\mp(\lambda, \mu) \psi, \psi)_{[L^2(\mathbb{R}^n)]^N} \\ &\quad - (i\varepsilon B G_\varepsilon^\mp(\lambda, \mu) \psi, \overline{A} G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N} \\ &\quad - (\overline{A} G_\varepsilon^\mp(\lambda, \mu) \psi, i\varepsilon B G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N} \\ &\stackrel{(3.44)}{=} \pm (\overline{A} \psi, G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N} \mp (G_\varepsilon^\mp(\lambda, \mu) \psi, \overline{A} \psi)_{[L^2(\mathbb{R}^n)]^N} \\ &\quad + \varepsilon (K B^{1/2} G_\varepsilon^\mp(\lambda, \mu) \psi, B^{1/2} G_\varepsilon^\pm(\lambda, \mu) \psi)_{[L^2(\mathbb{R}^n)]^N}, \end{aligned} \quad (3.82)$$

$$\psi \in [L_1^2(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty), \varepsilon \in (0, 1).$$

Hence,

$$\begin{aligned} \left| \frac{d}{d\varepsilon} F_\varepsilon^\pm(\lambda, \mu; \psi) \right| &\leq \left\| B^{-1/2} \overline{A} \psi \right\|_{[L^2(\mathbb{R}^n)]^N} \left[\left\| B^{1/2} G_\varepsilon^\pm(\lambda, \mu)^* \psi \right\|_{[L^2(\mathbb{R}^n)]^N} \right. \\ &\quad \left. + \left\| B^{1/2} G_\varepsilon^\pm(\lambda, \mu) \psi \right\|_{[L^2(\mathbb{R}^n)]^N} \right] \\ &\quad + \|K\|_{\mathcal{B}([L^2(\mathbb{R}^n)]^N)} \varepsilon \left\| B^{1/2} G_\varepsilon^\pm(\lambda, \mu)^* \psi \right\|_{[L^2(\mathbb{R}^n)]^N} \left\| B^{1/2} G_\varepsilon^\pm(\lambda, \mu) \psi \right\|_{[L^2(\mathbb{R}^n)]^N} \end{aligned}$$

$$\begin{aligned} &\leq C_4 \varepsilon^{-1/2} |F_\varepsilon^\pm(\lambda, \mu; \psi)|^{1/2} \|\psi\|_{[L_1^2(\mathbb{R}^n)]^N}, \\ &\psi \in [L_1^2(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty), \varepsilon \in (0, 1), \end{aligned} \quad (3.83)$$

where we applied Propositions 3.6 and 3.10, and $C_4 \in (0, \infty)$ is a constant independent of $\lambda \in \mathbb{R}$, $\mu \in (0, \infty)$, and $\varepsilon \in (0, 1)$.

Next, fix $r_0 \in (0, 1)$, and integrate over $[r, r_0] \subset (0, 1)$ in (3.83) to obtain for arbitrary $\psi \in [L_1^2(\mathbb{R}^n)]^N$,

$$\begin{aligned} &C_4 \|\psi\|_{[L_1^2(\mathbb{R}^n)]^N} \int_r^{r_0} d\varepsilon \varepsilon^{-1/2} |F_\varepsilon^\pm(\lambda, \mu; \psi)|^{1/2} \\ &\geq \int_r^{r_0} d\varepsilon \left| \frac{d}{d\varepsilon} F_\varepsilon^\pm(\lambda, \mu; \psi) \right| \\ &\geq \left| \int_r^{r_0} d\varepsilon \frac{d}{d\varepsilon} F_\varepsilon^\pm(\lambda, \mu; \psi) \right| \\ &= |F_{r_0}^\pm(\lambda, \mu; \psi) - F_r^\pm(\lambda, \mu; \psi)| \\ &\geq \left| |F_r^\pm(\lambda, \mu; \psi)| - |F_{r_0}^\pm(\lambda, \mu; \psi)| \right|, \quad \lambda \in \mathbb{R}, \mu \in (0, \infty), r \in (0, r_0), \end{aligned} \quad (3.84)$$

which yields

$$\begin{aligned} |F_r^\pm(\lambda, \mu; \psi)| &\leq |F_{r_0}^\pm(\lambda, \mu; \psi)| + C_4 \|\psi\|_{[L_1^2(\mathbb{R}^n)]^N} \int_r^{r_0} d\varepsilon \varepsilon^{-1/2} |F_\varepsilon^\pm(\lambda, \mu; \psi)|^{1/2}, \\ &\psi \in [L_1^2(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty), r \in (0, r_0). \end{aligned} \quad (3.85)$$

By Lemma 3.12 with $\theta = 1/2$,

$$\begin{aligned} |F_r^\pm(\lambda, \mu; \psi)| &\leq [|F_{r_0}^\pm(\lambda, \mu; \psi)|^{1/2} + C_4 \|\psi\|_{[L_1^2(\mathbb{R}^n)]^N} (r_0^{1/2} - r^{1/2})]^2, \\ &\psi \in [L_1^2(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty), r \in (0, r_0). \end{aligned} \quad (3.86)$$

In addition, by Propositions 3.5 and 3.10,

$$\begin{aligned} |F_{r_0}^\pm(\lambda, \mu; \psi)|^{1/2} &= |(\psi, G_{r_0}^\pm(\lambda, \mu)\psi)_{[L^2(\mathbb{R}^n)]^N}|^{1/2} \\ &\leq \|B^{-1/2}\psi\|_{[L^2(\mathbb{R}^n)]^N}^{1/2} \|B^{1/2}G_{r_0}^\pm(\lambda, \mu)\psi\|_{[L^2(\mathbb{R}^n)]^N}^{1/2} \\ &\leq C_1 r_0^{-1/2} \|\psi\|_{[L_{1/2}^2(\mathbb{R}^n)]^N}, \quad \psi \in [L_{1/2}^2(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty). \end{aligned} \quad (3.87)$$

Therefore,

$$\begin{aligned} |F_r^\pm(\lambda, \mu; \psi)| &\leq [C_1 r_0^{-1/2} \|\psi\|_{[L_{1/2}^2(\mathbb{R}^n)]^N} + C_4 \|\psi\|_{[L_1^2(\mathbb{R}^n)]^N} (r_0^{1/2} - r^{1/2})]^2, \\ &\psi \in [L_1^2(\mathbb{R}^n)]^N, \lambda \in \mathbb{R}, \mu \in (0, \infty), r \in (0, r_0). \end{aligned} \quad (3.88)$$

Finally, taking the limit $r \rightarrow 0^+$ and applying Proposition 3.9, one obtains

$$|F_0^\pm(\lambda, \mu; \psi)| \leq [C_1 r_0^{-1/2} + C_4 r_0^{1/2}]^2 \|\psi\|_{[L_1^2(\mathbb{R}^n)]^N}^2, \quad (3.89)$$

which yields the desired inequality with $C = [C_1 r_0^{-1/2} + C_4 r_0^{1/2}]^2$.

The claim (3.78) is clear from the fact that $\langle Q \rangle^{-s}$ is an isometric isomorphism from $L_t^2(\mathbb{R}^n)$ onto $L_{s+t}^2(\mathbb{R}^n)$, $s, t \in \mathbb{R}$. That $\langle Q \rangle^{-1}$ is H_0 -Kato-smooth follows from (3.78) (cf., e.g., [43, p. 148], [49, p. 134]). \square

Of course, Theorem 3.13 implies the absence of any singular spectrum (i.e., the absence of eigenvalues and singular continuous spectrum) of H_0 ,

$$\sigma_s(H_0) = \emptyset, \quad (3.90)$$

but since H_0 is unitarily equivalent, via Fourier transform, to the operator of multiplication by $\alpha \cdot p$, $p \in \mathbb{R}^n$, purely absolutely continuous spectrum of H_0 was obvious from the outset.

Remark 3.14. In the massive case, where $H_0(m) = H_0 + m\beta$, $m > 0$, a global limiting absorption principle (i.e., for energies on the entire real line) for $H_0(m)$ was proved in dimension $n = 3$ by Iftimovici and Măntoiu [32] in 1999. The corresponding massless case (i.e., for the operator H_0) in dimension $n = 3$ was settled in 2008 by Saitō and Umeda [46] employing entirely different methods in their study of zero eigenvalues and zero-energy resonances of massless Dirac operators. Our result, Theorem 3.13, appears to be new for $n \in \mathbb{N} \setminus \{3\}$, $n \geq 2$. The cases of two and higher-dimensional massless as well as massive Dirac operators with continuous matrix-valued short-range potentials has also just been studied by Erdoğan, Goldberg, and Green [21], employing an entirely different approach. The authors prove a limiting absorption principle in the massless case $m = 0$ (resp., in the massive case $m > 0$) valid on $(-\infty, 0) \cup (0, \infty)$ (resp., valid on $(-\infty, -m) \cup (m, \infty)$), assuming regular thresholds. \diamond

Theorem 3.13 implies existence and completeness (in fact, unitarity) of wave operators for the pair of self-adjoint operators $(H = H_0 + V, H_0)$ for sufficiently “weak” perturbations V of H_0 in the following standard manner (we refer to [49, Ch. 4], especially, [49, Theorem 4.6.1], for details): Consider the self-adjoint matrix-valued potential $V = \{V_{\ell, m}\}_{1 \leq \ell, m \leq N}$ satisfying for some $C \in (0, \infty)$,

$$V \in [L^\infty(\mathbb{R}^n)]^{N \times N}, \quad \|V(x)\|_{\mathbb{C}^{N \times N}} \leq C \langle x \rangle^{-2} \text{ for a.e. } x \in \mathbb{R}^n, \quad (3.91)$$

and define the interacting massless Dirac operator H via

$$H = H_0 + V, \quad \text{dom}(H) = \text{dom}(H_0) = [W^{1,2}(\mathbb{R}^n)]^N. \quad (3.92)$$

One infers that

$$\|\langle x \rangle V \langle x \rangle\|_{\mathbb{C}^{N \times N}} \leq C \text{ for a.e. } x \in \mathbb{R}^n, \quad (3.93)$$

and

$$\begin{aligned} \langle Q \rangle^{-1} (H - zI_{[L^2(\mathbb{R}^n)]^N})^{-1} \langle Q \rangle^{-1} &= \langle Q \rangle^{-1} (H_0 - zI_{[L^2(\mathbb{R}^n)]^N})^{-1} \langle Q \rangle^{-1} \\ &\times [I + \langle Q \rangle V (H_0 - zI_{[L^2(\mathbb{R}^n)]^N})^{-1} \langle Q \rangle^{-1}]^{-1}, \quad z \in \mathbb{C} \setminus \mathbb{R}. \end{aligned} \quad (3.94)$$

Theorem 3.15. *Assume Hypotheses 3.1 and 3.2, (3.91), and define the operator H as in (3.92). If in addition,*

$$\sup_{\lambda \in \mathbb{R}, \mu \in (0, \infty)} \|\langle Q \rangle V (H_0 - (\lambda \pm i\mu)I_{[L^2(\mathbb{R}^n)]^N})^{-1} \langle Q \rangle^{-1}\|_{\mathcal{B}([L^2(\mathbb{R}^n)]^N)} < 1, \quad (3.95)$$

then the wave operators

$$W_\pm(H, H_0) = \text{s-lim}_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0}, \quad W_\pm(H_0, H) = \text{s-lim}_{t \rightarrow \pm\infty} e^{itH_0} e^{-itH}, \quad (3.96)$$

exist and are complete, that is,

$$\begin{aligned} \ker(W_\pm(H, H_0)) &= \ker(W_\pm(H_0, H)) = \{0\}, \\ \text{ran}(W_\pm(H, H_0)) &= \text{ran}(W_\pm(H_0, H)) = [L^2(\mathbb{R}^n)]^N. \end{aligned} \quad (3.97)$$

In fact, they are unitary and adjoint to each other, $W_{\pm}(H, H_0)^* = W_{\pm}(H_0, H)$. In particular, H and H_0 are unitarily equivalent and hence H is spectrally purely absolutely continuous.

Remark 3.16. Since we permit a (sufficiently decaying) matrix-valued potential V in H , this includes, in particular, the case of electromagnetic interactions introduced via minimal coupling, that is, V describes also special cases of the form,

$$H(q, A) := \alpha \cdot (-i\nabla - A) + qI_N = H_0 + [qI_N - \alpha \cdot A], \quad \text{dom}(H(q, A)) = [W^{1,2}(\mathbb{R}^n)]^N, \quad (3.98)$$

where (q, A) represent the electromagnetic potentials on \mathbb{R}^n , with $q : \mathbb{R}^n \rightarrow \mathbb{R}$, $q \in L^\infty(\mathbb{R}^n)$, $A = (A_1, \dots, A_n)$, $A_j : \mathbb{R}^n \rightarrow \mathbb{R}$, $A_j \in L^\infty(\mathbb{R}^n)$, $1 \leq j \leq n$, and for some $C \in (0, \infty)$,

$$|q(x)| + |A_j(x)| \leq C \langle x \rangle^{-2} \text{ for a.e. } x \in \mathbb{R}^n, \quad 1 \leq j \leq n. \quad (3.99)$$

◇

Remark 3.17. Using the notion of (local) strong operator smoothness as described in detail in [50, Sects. 4.4–4.7], the decay rate $\langle \cdot \rangle^{-2}$ in (3.91), (3.99) can be relaxed to $\langle \cdot \rangle^{-\rho}$ for some $\rho > 1$ (cf. [13]) and this then permits situations where H has eigenvalues and hence H is no longer unitarily equivalent to H_0 and spectrally purely absolutely continuous. ◇

Acknowledgments. We are indebted to Andrei Iftimovici for helpful discussions and for valuable hints to the literature on the commutator calculus. We would also like to thank Eduard Tsekanovskii for helpful discussions.

The authors are indebted to the Banff International Research Station for Mathematical Innovation and Discovery (BIRS) for their extraordinary hospitality during the focused research group on *Extensions of index theory inspired by scattering theory* (17frg668), June 18–25, 2017, where part of this work was done.

REFERENCES

- [1] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, Dover, New York, 1972.
- [2] S. Agmon, *Spectral properties of Schrödinger operators and scattering theory*, Ann. Sc. Norm. Sup. Pisa, Ser. 4, **2**, 151–218 (1975).
- [3] D. Aiba, *Absence of zero resonances of massless Dirac operators*, Hokkaido Math. J. **45**, 263–270 (2016).
- [4] W. Amrein, A. Boutet de Monvel, and V. Georgescu, *C_0 -Groups, Commutator Methods and Spectral Theory of N -Body Hamiltonians*, Progr. in Math., Vol. 135, Birkhäuser, Basel, 1996.
- [5] A. Balinsky and W. D. Evans, *Spectral Analysis of Relativistic Operators*, Imperial College Press, London, 2011.
- [6] E. Balslev and B. Helffer, *Limiting absorption principle and resonances for the Dirac operator*, Adv. Appl. Math. **13**, 186–215 (1992).
- [7] L. Bandara and H. Saratchandran, *Essential self-adjointness of powers of first-order differential operators on non-compact manifolds with low-regularity metrics*, J. Funct. Anal. **273**, 3719–3758 (2017).
- [8] H. Baumgärtel and M. Wollenberg, *Mathematical Scattering Theory*, Akademie Verlag, Berlin, 1983.
- [9] M. Ben-Artzi and A. Devinatz, *The limiting absorption principle for partial differential operators*, Memoirs Amer. Math. Soc. **66**, No. 364, 1–70 (1987).
- [10] N. Boussaid and S. Golénia, *Limiting absorption principle for some long range perturbations of Dirac systems at threshold energies*, Commun. Math. Phys. **299**, 677–708 (2010).

- [11] A. Boutet de Monvel-Berthier, D. Manda, and R. Purice, *Limiting absorption principle for the Dirac operator*, Ann. Inst. H. Poincaré, **58**, 413–431 (1993).
- [12] A. Boutet de Monvel and M. Măntoiu, *The method of the weakly conjugate operator*, in *Inverse and Algebraic Quantum Scattering Theory*, B. Apagyi, G. Endrédi, and P. Lévy (eds.), Springer, Heidelberg, 1997.
- [13] A. Carey, F. Gesztesy, G. Levitina, R. Nichols, F. Sukochev, and D. Zanin, *On the limiting absorption principle for massless Dirac operators*, in preparation.
- [14] A. Carey, F. Gesztesy, G. Levitina, D. Potapov, F. Sukochev, and D. Zanin, *On index theory for non-Fredholm operators: a $(1+1)$ -dimensional example*, Math. Nachrichten **289**, 575–609 (2016).
- [15] A. Carey, F. Gesztesy, G. Levitina, and F. Sukochev, *On the index of a non-Fredholm model operator*, Operators and Matrices **10**, 881–914 (2016).
- [16] A. Carey, F. Gesztesy, H. Grosse, G. Levitina, D. Potapov, F. Sukochev, and D. Zanin, *Trace formulas for a class of non-Fredholm operators: A review*, Rev. Math. Phys. **28**, No. 10, (2016), 1630002 (55 pages).
- [17] A. Carey, F. Gesztesy, D. Potapov, F. Sukochev, and Y. Tomilov, *On the Witten index in terms of spectral shift functions*, J. Analyse Math. **132**, 1–61 (2017).
- [18] P. R. Chernoff, *Essential self-adjointness of powers of generators of hyperbolic equations*, J. Funct. Anal. **12**, 401–414 (1973).
- [19] A. Connes, *Noncommutative geometry*, Academic Press, San Diego, 1994.
- [20] T. Daude, *Scattering theory for massless Dirac fields with long-range potentials*, J. Math. Pures Appl. **84**, 615–665 (2005).
- [21] M. B. Erdoğan, M. Goldberg, and W. R. Green, *Limiting absorption principle and Strichartz estimates for Dirac operators in two and higher dimensions*, arXiv:1706.05257.
- [22] M. B. Erdoğan, M. Goldberg, and W. Schlag, *Strichartz and smoothing estimates for Schrödinger operators with large magnetic potentials in \mathbb{R}^3* , J. Eur. Math. Soc. **10**, 507–531 (2008).
- [23] M. B. Erdoğan, M. Goldberg, and W. Schlag, *Strichartz and smoothing estimates for Schrödinger operators with almost critical magnetic potentials in three and higher dimensions*, Forum Math. **21**, 687–722 (2009).
- [24] M. B. Erdoğan and W. R. Green, *The Dirac equation in two dimensions: dispersive estimates and classification of threshold obstructions*, Commun. Math. Phys. **352**, 719–757 (2017).
- [25] W. G. Faris, *Self-Adjoint Operators*, Lecture Notes in Mathematics, Vol. 433, Springer, Berlin, 1975.
- [26] I. Forsyth, B. Mesland, and A. Rennie, *Dense domains, symmetric operators and spectral triples*. New York J. Math. **20**, 1001–1020 (2014).
- [27] V. Georgescu and M. Măntoiu, *On the spectral theory of singular Dirac type Hamiltonians*, J. Operator Th. **46**, 289–321 (2001).
- [28] C. Gérard, *A proof of the abstract limiting absorption principle by energy estimates*, J. Funct. Anal. **254**, 2707–2724 (2008).
- [29] S. Golénia and T. Jecko, *A new look at Mourre’s commutator theory*, Complex Anal. Operator Th. **1**, 399–422 (2007).
- [30] I. Herbst, *Spectral theory of the operator $(p^2 + m^2)^{1/2} - Ze^2/r$* , Commun. Math. Phys. **53**, 285–294 (1977).
- [31] N. Higson and J. Roe, *Analytic K-Homology*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000.
- [32] A. Iftimovici and M. Măntoiu, *Limiting absorption principle at critical values for the Dirac operator*, Lett. Math. Phys. **49**, 235–243 (1999).
- [33] J. Kaad, *Differentiable absorption of Hilbert C^* -modules, connections, and lifts of unbounded operators*, J. Noncommut. Geom. **11**, 1037–1068 (2017).
- [34] J. Kaad and M. Lesch, *A local global principle for regular operators in Hilbert C^* -modules*, J. Funct. Anal. **262**, no. 10, 4540–4569 (2012).
- [35] H. Kalf, T. Okaji, and O. Yamada, *The Dirac operator with mass $m_0 \geq 0$: Non-existence of zero modes and of threshold eigenvalues*, Doc. Math. **20**, 37–64 (2015); Addendum, Doc. Math. **22**, 1–3 (2017).
- [36] T. Kato, *Wave operators and similarity for some non-selfadjoint operators*, Math. Ann. **162**, 258–279 (1966).

- [37] S. T. Kuroda, *An Introduction to Scattering Theory*, Aarhus University Lecture Notes Series, No. 51, 1978.
- [38] A. Kuzhel, *Characteristic Functions and Models of Nonself-Adjoint Operators*, Kluwer Academic Publishers, Boston, 1996.
- [39] M. Măntoiu and M. Pascu, *Global resolvent estimates for multiplication operators*, J. Operator Th. **36**, 283–294 (1996).
- [40] B. Mesland and A. Rennie, *Nonunital spectral triples and metric completeness in unbounded KK -theory*, J. Funct. Anal. **271**, no. 9, 2460–2538 (2016).
- [41] C. Pladdy, Y. Saitō, and T. Umeda, *Resolvent estimates for the Dirac operator*, Analysis **15**, 123–149 (1995).
- [42] C. Pladdy, Y. Saitō, and T. Umeda, *Radiation condition for Dirac operators*, J. Math. Kyoto Univ. **37-4**, 567–584 (1998).
- [43] M. Reed and B. Simon, *Methods of Modern Mathematical Physics. IV: Analysis of Operators*, Academic Press, New York, 1978.
- [44] S. Richard, *Some improvements in the method of weakly conjugate operator*, Lett. Math. Phys. **76**, 27–36 (2006).
- [45] M. Ruzhansky and M. Sugimoto, *Structural resolvent estimates and derivative nonlinear Schrödinger equations*, Commun. Math. Phys. **314**, 281–304 (2012).
- [46] Y. Saitō and T. Umeda, *The zero modes and zero resonances of massless Dirac operators*, Hokkaido Math. J. **37**, 363–388 (2008).
- [47] B. Simon, *Trace Ideals and Their Applications*, Mathematical Surveys and Monographs, Vol. 120, 2nd ed., Amer. Math. Soc., Providence, RI, 2005.
- [48] V. Vogelsang, *Absolutely continuous spectrum of Dirac operators for long-range potentials*, J. Funct. Anal. **76**, 67–86 (1988).
- [49] D. R. Yafaev, *Mathematical Scattering Theory. General Theory*, Amer. Math. Soc., Providence, RI, 1992.
- [50] D. R. Yafaev, *Mathematical Scattering Theory. Analytic Theory*, Math. Surveys and Monographs, Vol. 158, Amer. Math. Soc., Providence, RI, 2010.
- [51] O. Yamada, *On the principle of limiting absorption for the Dirac operator*, Publ. RIMS, Kyoto Univ. **8**, 557–577 (1972/73).
- [52] O. Yamada, *Eigenfunction expansions and scattering theory for Dirac operators*, Publ. RIMS, Kyoto Univ. **11**, 651–689 (1976).
- [53] O. Yamada, *A remark on the limiting absorption method for Dirac operators*, Proc. Japan. Acad. **69**, Ser. A, 243–246 (1993).

MATHEMATICAL SCIENCES INSTITUTE, AUSTRALIAN NATIONAL UNIVERSITY, KINGSLEY ST., CANBERRA, ACT 0200, AUSTRALIA AND SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF WOLLONGONG, NSW, AUSTRALIA, 2522

E-mail address: acarey@maths.anu.edu.au

URL: <http://maths.anu.edu.au/~acarey/>

DEPARTMENT OF MATHEMATICS, BAYLOR UNIVERSITY, ONE BEAR PLACE #97328, WACO, TX 76798-7328, USA

E-mail address: Fritz_Gesztesy@baylor.edu

URL: <http://www.baylor.edu/math/index.php?id=935340>

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF SOUTHERN DENMARK, CAMPUSVEJ 55, DK-5230 ODENSE M, DENMARK

E-mail address: kaad@imada.sdu.dk

SCHOOL OF MATHEMATICS AND STATISTICS, UNSW, KENSINGTON, NSW 2052, AUSTRALIA

E-mail address: g.levitina@student.unsw.edu.au

MATHEMATICS DEPARTMENT, THE UNIVERSITY OF TENNESSEE AT CHATTANOOGA, 415 EMCS BUILDING, DEPT. 6956, 615 McCALLIE AVE, CHATTANOOGA, TN 37403, USA

E-mail address: Roger-Nichols@utc.edu

URL: <http://www.utc.edu/faculty/roger-nichols/index.php>

SCHOOL OF MATHEMATICS AND STATISTICS, UNSW, KENSINGTON, NSW 2052, AUSTRALIA

E-mail address: d.potapov@unsw.edu.au

SCHOOL OF MATHEMATICS AND STATISTICS, UNSW, KENSINGTON, NSW 2052, AUSTRALIA

E-mail address: f.sukochev@unsw.edu.au