

University of Wollongong

## Research Online

---

Faculty of Science, Medicine and Health -  
Papers: part A

Faculty of Science, Medicine and Health

---

1-1-2014

### Progress on understanding atmospheric mercury hampered by uncertain measurements

Daniel A. Jaffe  
*University of Washington*


Seth Lyman  
*Utah State University*


Helen Amos  
*Harvard University*

M Gustin  
*University of Nevada*

Jiaoyan Huang  
*University of Nevada*

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>

 See next page for additional authors

 Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

---

#### Recommended Citation

Jaffe, Daniel A.; Lyman, Seth; Amos, Helen; Gustin, M; Huang, Jiaoyan; Selin, Noelle E.; Levin, Leonard; Ter Schure, Arnout; Mason, Robert P.; Talbot, R; Rutter, Andrew; Finley, Brandon; Jaegle, Lyatt; Shah, Viral; McClure, Crystal; Ambrose, Jesse; Gratz, Lynne; Lindberg, Steven; Weiss-Penzias, Peter; Sheu, Guey-Rong; Feddersen, Dara; Horvat, Milena; Dastoor, Ashu; Hynes, Anthony J.; Mao, Huiting; Sonke, Jeroen E.; Slemr, Franz; Fisher, Jenny A.; Ebinghaus, Ralf; Zhang, Yanxu; and Edwards, Grant, "Progress on understanding atmospheric mercury hampered by uncertain measurements" (2014). *Faculty of Science, Medicine and Health - Papers: part A*. 1946.

<https://ro.uow.edu.au/smhpapers/1946>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: [research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

## Progress on understanding atmospheric mercury hampered by uncertain measurements

### Abstract

Mercury (Hg) is a potent neurotoxin and globally reducing environmental levels is seen as paramount for protecting human and wildlife health. In 2013, many countries finalized the negotiations on, and have now signed, the Minamata Convention on Mercury, which commits participating countries to reduce emissions and use of mercury. Successful implementation of the treaty will require adequate verification through global monitoring.

### Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

### Publication Details

Jaffe, D. A., Lyman, S., Amos, H. M., Gustin, M. S., Huang, J., Selin, N. E., Levin, L., Ter Schure, A., Mason, R. P., Talbot, R., Rutter, A., Finley, B., Jaegle, L., Shah, V., McClure, C., Ambrose, J., Gratz, L., Lindberg, S., Weiss-Penzias, P., Sheu, G., Feddersen, D., Horvat, M., Dastoor, A., Hynes, A. J., Mao, H., Sonke, J. E., Slemr, F., Fisher, J. A., Ebinghaus, R., Zhang, Y. & Edwards, G. (2014). Progress on understanding atmospheric mercury hampered by uncertain measurements. *Environmental Science and Technology* (Washington), 48 (13), 7204-7206.

### Authors

Daniel A. Jaffe, Seth Lyman, Helen Amos, M Gustin, Jiaoyan Huang, Noelle E. Selin, Leonard Levin, Arnout Ter Schure, Robert P. Mason, R Talbot, Andrew Rutter, Brandon Finley, Lyatt Jaegle, Viral Shah, Crystal McClure, Jesse Ambrose, Lynne Gratz, Steven Lindberg, Peter Weiss-Penzias, Guey-Rong Sheu, Dara Feddersen, Milena Horvat, Ashu Dastoor, Anthony J. Hynes, Huiting Mao, Jeroen E. Sonke, Franz Slemr, Jenny A. Fisher, Ralf Ebinghaus, Yanxu Zhang, and Grant Edwards

## **Progress on understanding atmospheric mercury hampered by uncertain measurements**

Daniel A. Jaffe,<sup>1,2\*</sup> Seth Lyman,<sup>3</sup> Helen M. Amos,<sup>4</sup> Mae S. Gustin,<sup>5</sup> Jiaoyan Huang,<sup>5</sup> Noelle E. Selin,<sup>6</sup> Leonard Levin,<sup>7</sup> Arnout ter Schure,<sup>8</sup> Robert P. Mason,<sup>9</sup> Robert Talbot,<sup>10</sup> Andrew Rutter,<sup>11</sup> Brandon Finley,<sup>1</sup> Lyatt Jaeglé,<sup>2</sup> Viral Shah,<sup>2</sup> Crystal McClure,<sup>2</sup> Jesse Ambrose,<sup>1</sup> Lynne Gratz,<sup>1</sup> Steven Lindberg,<sup>12</sup> Peter Weiss-Penzias,<sup>13</sup> Guey-Rong Sheu,<sup>14</sup> Dara Feddersen,<sup>15</sup> Milena Horvat,<sup>16</sup> Ashu Dastoor,<sup>17</sup> Anthony J. Hynes,<sup>18</sup> Huiting Mao,<sup>19</sup> Jeroen E. Sonke,<sup>20</sup> Franz Slemr,<sup>21</sup> Jenny A. Fisher,<sup>22</sup> Ralf Ebinghaus,<sup>23</sup> Yanxu Zhang,<sup>24</sup> and Grant Edwards<sup>25</sup>

<sup>1</sup>School of Science, Technology, Engineering and Mathematics, University of Washington, Bothell, Washington, United States

<sup>2</sup>Department of Atmospheric Sciences, University of Washington, Seattle, Washington, United States

<sup>3</sup>Bingham Entrepreneurship and Energy Research Center, Utah State University, Vernal, Utah, United States

<sup>4</sup>Department of Earth and Planetary Science, Harvard University, Boston, Massachusetts, United States

<sup>5</sup>Department of Natural Resources and Environmental Sciences, University of Nevada, Reno, Nevada, United States

<sup>6</sup>Engineering Systems Division and Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts, United States

<sup>7</sup>Palo Alto, California, United States

<sup>8</sup>Electric Power Research Institute, Palo Alto, California, United States

<sup>9</sup>Department of Marine Sciences & Chemistry, University of Connecticut, Groton, Connecticut, United States

<sup>10</sup>Institute for Climate and Atmospheric Science, University of Houston, Houston, Texas, United States

<sup>11</sup>Department of Chemistry, Carroll University, Waukesha, Wisconsin, United States

<sup>12</sup>Retired

<sup>13</sup>Department of Microbiology and Environmental Toxicology, University of California, Santa Cruz, California, United States

<sup>14</sup>Department of Atmospheric Sciences, National Central University, Jhongli, Taiwan

<sup>15</sup>Department of Chemistry, University of New Hampshire, Durham, New Hampshire, United States

<sup>16</sup>Department of Environmental Sciences, Jozef Stefan Institute, Ljubljana, Slovenia

<sup>17</sup>Air Quality Research Division, Environment Canada, Dorval, Quebec, Canada

<sup>18</sup>Division of Marine and Atmospheric Chemistry, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, Florida, United States

<sup>19</sup>Department of Chemistry, State University of New York, College of Environmental Science and Forestry, Syracuse, New York, United States

<sup>20</sup>Observatoire Midi-Pyrénées, Université Paul Sabatier, Toulouse, France

<sup>21</sup>Atmospheric Chemistry Division, Max Planck Institute for Chemistry, Mainz, Germany

<sup>22</sup>Centre for Atmospheric Chemistry, University of Wollongong, Wollongong, Australia

<sup>23</sup>Department of Environmental Chemistry, Institute of Coastal Research, Geesthacht, Germany

<sup>24</sup>School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts

<sup>25</sup>Faculty of Science, Macquarie University, Sydney, Australia

Mercury (Hg) is a potent neurotoxin and globally reducing environmental levels is seen as paramount for protecting human and wildlife health. In 2013, many countries finalized the negotiations on, and have now signed, the Minamata Convention on Mercury, which commits participating countries to reduce emissions and use of mercury. Successful implementation of the treaty will require adequate verification through global monitoring.

In the last decade, the number of atmospheric measurements of speciated Hg has rapidly multiplied. Mercury data are being collected by monitoring networks in Europe, Canada, East Asia and the United States. These measurements are relatively expensive and mounting evidence suggests they suffer from significant biases. As a community, we are confronted with the question, “Are current analytical methods for Hg adequate to address policy requirements?” We conclude that better analytical techniques are urgently needed.

In the atmosphere Hg occurs as three operationally-defined forms: gaseous elemental mercury (GEM), gaseous oxidized mercury (GOM) and particle-bound mercury (PBM). GOM and PBM are believed to consist of Hg(II) compounds. Oxidation of GEM to GOM and subsequent deposition is thought to be the major pathway to deliver Hg to ecosystems. Thus, GOM provides a critical link between human emissions and ecosystem exposure. A number of laboratory and theoretical chemical investigations have been performed, but at present there is no consensus on what the chemical form(s) of GOM is(are), nor any reliable method to identify the chemical form(s) in the atmosphere. It is likely that more than one form of Hg(II) exists in the atmosphere, depending on its source.

At present the primary method to identify GOM is use of a KCl coated denuder for concentration and collection, followed by thermal desorption and atomic fluorescence to quantify the desorbed GEM. PBM is collected on a regenerable quartz filter downstream of the denuder and is quantified similarly. This method has been adopted by the Canadian Atmospheric Mercury Network, the U.S. Atmospheric Mercury Network and the European Global Mercury Observation System for standard measurements. QA/QC protocols have been developed for these networks, but these protocols do not provide a way to calibrate for GOM or PBM, quantify collection efficiency or quantify measurement interferences.

A number of studies have been published that indicate there are significant problems with the existing methods for GOM and PBM. By using possible GOM compounds, such as  $\text{HgCl}_2$  and  $\text{HgBr}_2$ , it has been shown that the KCl denuder method is subject to interferences from ozone, water vapor and possibly other compounds as well.<sup>1,2</sup> Other studies have found that KCl denuder measurements do not agree with measurements made with alternative collection surfaces. It is also not clear what is the true uncertainty in measurements of PBM, as comparisons with other methods have yielded large discrepancies between commercial, automated PBM measurements and manual methods with alternative samplers.<sup>3</sup> All of these point to substantial problems with our current methods to quantify GOM and PBM.

At present we are not able to quantify the uncertainty in GOM and PBM measurements. This is because (1) we do not know the chemical form being measured, (2) there is no accepted calibration method and (3) we have a limited understanding of interferences. It is not yet clear whether we will be able to correct GOM and PBM data collected with current methods to account for these biases. This will require a more detailed understanding of Hg chemistry and detailed interference tests with reliable field-deployable calibrators.

We acknowledge that some measurements of GOM made with the KCl denuder method have been successfully interpreted, at least qualitatively. Some studies have demonstrated an inverse relationship and approximate “mass conservation” between GOM and GEM. However, these studies do not provide definitive evidence that the KCl denuder method is free from bias. Work investigating ozone interference has found that the ratio of GOM to GEM can be biased low while the total gaseous Hg concentration (GOM + GEM) is approximately conserved.<sup>1</sup> In other words, GOM can decompose to GEM during sampling and be analyzed as GEM. It is also possible that the KCl denuder method works better in some environments than others.

At present, we believe unspiciated Hg measurements (i.e., measurements with no upstream processing prior to sample introduction into an atomic fluorescence analyzer) are well calibrated and the true uncertainty can be reasonably described. There is some uncertainty as to whether current unspiciated measurements capture total gaseous mercury (TGM) or GEM. If desired, total atmospheric mercury (TAM = GEM + GOM + PBM) can be measured using a pyrolyzing inlet. In principle, measurements of TAM should be more accurate than current methods to measure the individual species.

Working with Hg(II) compounds is challenging due to their toxicity, tendency to decompose and stickiness in calibration systems. But without high quality calibrations it is not possible to validate measurements, quantify uncertainties or fully evaluate atmospheric models.<sup>4</sup> Calibration methods for oxidized Hg compounds are improving<sup>5</sup> and these will improve the quality of future interference studies as well as routine measurement studies for GOM and PBM.

In order to have confidence in atmospheric chemical measurements, we must be able to quantify their uncertainty. This includes routine calibration in realistic conditions and rigorous testing for interferences. At present, the existing and planned atmospheric mercury networks have inadequate attention to quality control for GOM and PBM measurements. To resolve these issues, the community needs to:

1. Develop calibration methods for GOM and provide routine calibrations for field instrumentation;
2. Conduct detailed investigations to quantify interferences in the existing GOM methods and develop new methodologies to measure it; and
3. Conduct fundamental research on the chemistry, reaction kinetics and chemical identity of the compounds that make up GOM and PBM in the atmosphere.

We believe these items should be given high priority by the mercury scientific community. To do otherwise impedes scientific progress and environmental monitoring efforts. Space limitations prevent us from including a complete set of relevant work. For a more complete list of relevant publications, please go to <http://atmos.washington.edu/jaffegroup/Hg>

## References

1. Lyman, S. N.; Jaffe, D. A.; Gustin, M. S. Release of mercury halides from KCl denuders in the presence of ozone. *Atmos. Chem. Phys.* **2010**, *10*, 8197–8204; DOI 10.5194/acp-10-8197-2010.
2. Gustin, M. S.; Huang, J.; Miller, M. B.; Peterson, C.; Jaffe, D. A.; Ambrose, J.; Finley, B. D.; Lyman, S. N.; Call, K.; Talbot, R.; Feddersen, D.; Mao, H.; Lindberg, S. E. Do we understand what the mercury speciation instruments are actually measuring? Results of RAMIX. *Environ. Sci. Technol.* **2013** *47* (13), 7295–7306; DOI 10.1021/es3039104.
3. Talbot, R.; Mao, H.; Feddersen, D.; Smith, M.; Kim, S. Y.; Sive, B.; Haase, K.; Ambrose, J.; Zhou, Y.; Russo, R. Comparison of particulate mercury measured with manual and automated methods. *Atmosphere* **2011**, *2*, 1–20; DOI 10.3390/atmos2010001.
4. Kos, G.; Ryzhkov, A.; Dastoor, A.; Narayan, J.; Steffen, A.; Ariya, P. A.; Zhang, L. Evaluation of discrepancy between measured and modelled oxidized mercury species. *Atmos. Chem. Phys.* **2013**, *13*, 4839–4863; DOI 10.5194/acp-13-4839-2013.
5. Finley, B. D.; Jaffe, D. A.; Call, K.; Lyman, S.; Gustin, M.; Peterson, C.; Miller, M.; Lyman, T. Development, testing, and deployment of an air sampling manifold for spiking elemental and oxidized mercury during the Reno Atmospheric Mercury Intercomparison Experiment (RAMIX). *Environ. Sci. Technol.* **2013**; DOI 10.1021/es304185a.