

University of Wollongong

Research Online

Faculty of Engineering and Information
Sciences - Papers: Part A

Faculty of Engineering and Information
Sciences

2006

Jacket matrices constructed from Hadamard matrices and generalized Hadamard matrices

Ken Finlayson

University of Wollongong, kaf03@uow.edu.au

Moon Ho Lee

Chonbuk National University

Jennifer Seberry

University of Wollongong, jennie@uow.edu.au

Mieko Yamada

Kanazawa University

Follow this and additional works at: <https://ro.uow.edu.au/eispapers>



Part of the [Engineering Commons](#), and the [Science and Technology Studies Commons](#)

Recommended Citation

Finlayson, Ken; Lee, Moon Ho; Seberry, Jennifer; and Yamada, Mieko, "Jacket matrices constructed from Hadamard matrices and generalized Hadamard matrices" (2006). *Faculty of Engineering and Information Sciences - Papers: Part A*. 2259.

<https://ro.uow.edu.au/eispapers/2259>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Jacket matrices constructed from Hadamard matrices and generalized Hadamard matrices

Abstract

Jacket matrices constructed from Hadamard matrices and generalized Hadamard matrices.

Keywords

jacket, hadamard, generalized, constructed, matrices

Disciplines

Engineering | Science and Technology Studies

Publication Details

Finlayson, K., Lee, M., Seberry, J. & Yamada, M. (2006). Jacket matrices constructed from Hadamard matrices and generalized Hadamard matrices. *Australasian Journal of Combinatorics*, 35 83-87.

Jacket matrices constructed from Hadamard matrices and generalized Hadamard matrices *

KEN FINLAYSON

*Centre for Computer Security Research
University of Wollongong
N.S.W. 2522
Australia*

MOON HO LEE

*Institute of Information and Communications
Chonbuk National University, Jeonju
Korea*

JENNIFER SEBERRY

*Centre for Computer Security Research
University of Wollongong
N.S.W. 2522
Australia*

MIEKO YAMADA

*Department of Computational Science
Kanazawa University
Kakuma-machi, Kanazawa
Japan*

Abstract

Jacket matrices are matrices $L = (\ell_{ij})$ with inverse $L^{-1} = \frac{1}{n} (\ell_{ij}^{-1})$, where the inverse is over a group G . They have previously been constructed only from $(1, -1)$ Hadamard matrices. In this note, we give constructions for jacket matrices based on generalized Hadamard matrices.

* This work was partially supported by the Institute of Information Technology Assessment, Korea, the Korea Science and Engineering Foundation (KOSEF), Korea and the Australian Academy of Science.

1 Introduction

Let H be a matrix. We define H^\dagger to be the Hermitian conjugate, or the transpose of the matrix with elements the complex conjugate of the corresponding elements of H . When the entries of H are from a group G , we define H^M to be the transpose of the matrix whose elements are the group inverse of the corresponding elements of H .

An Hadamard matrix H of order n is square, with entries ± 1 and satisfies $HH^T = H^T H = nI$. Seberry and Yamada [10] have surveyed Hadamard matrices and the reader is referred there for more details.

In this paper, if $HH^\dagger = H^\dagger H = nI$ then H is a generalized Hadamard matrix. More generally, generalized Hadamard matrices of two types are of interest. The first (see [1, 4]) have entries which are roots of unity; the second (see [2, 3, 8, 9]) have elements from a finite group.

Let p be an odd prime. Let $1, \alpha, \alpha^2, \dots, \alpha^{p-1}$ be the p th roots of unity. A Butson generalized Hadamard matrix [1] $B = (b_{ij})$ of order p is defined as

$$b_{ij} = \begin{cases} 1 & i = 1 \text{ and } 1 \leq j \leq p \\ 1 & j = 1 \text{ and } 1 \leq i \leq p \\ \alpha^{(i-1)(j-1)} & 2 \leq i, j \leq p \end{cases}$$

Then the core C of B is the $(p-1) \times (p-1)$ matrix (b_{st}) , $2 \leq s, t \leq p$. We observe that C, C^\dagger and C^M are symmetric, and that $C^\dagger = C^M$ is a permutation of C .

A jacket matrix (sometimes called a reverse jacket matrix) $L = (\ell_{ij})$ is a matrix of order n with entries from a group G , with inverse $L^{-1} = \frac{1}{n}(\ell_{ij}^{-1})$.

We can use a jacket matrix L in a jacket transform (also called a reverse jacket transform) as follows. For a vector \mathbf{a} of length n , its transform \mathbf{A} is given by $\mathbf{A} = \mathbf{a}L$. The inverse transform is $\mathbf{a} = \mathbf{A}L^{-1} = \frac{1}{n}\mathbf{A}L^M$.

2 Our constructions

Jacket matrices in their original formulation were constructed from $(1, -1)$ Hadamard matrices (see [5–7]). However, it is possible to construct jacket matrices from generalized Hadamard matrices. We present three such constructions. We also give a method of combining such jacket matrices to form larger jacket matrices.

Let A, B, D be symmetric matrices of order $\frac{n-2}{2}$, whose elements are in an Abelian group (including 1). Let e be a column vector whose elements are all 1. Put

$$X = \begin{pmatrix} 1 & e^t & e^t & 1 \\ e & A & B & e \\ e & B & -D & -e \\ 1 & e^t & -e^t & -1 \end{pmatrix}.$$

If X satisfies

$$XX^M = X^M X = nI$$

then X is a jacket matrix.

2.1 Case 1: $A = B = D$

Let

$$A = B = D = \begin{pmatrix} \omega & \omega^2 \\ \omega^2 & \omega \end{pmatrix},$$

where ω is the cube root of unity. Then

$$X = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \omega & \omega^2 & \omega & \omega^2 & 1 \\ 1 & \omega^2 & \omega & \omega^2 & \omega & 1 \\ 1 & \omega & \omega^2 & -\omega & -\omega^2 & -1 \\ 1 & \omega^2 & \omega & -\omega^2 & -\omega & -1 \\ 1 & 1 & 1 & -1 & -1 & -1 \end{pmatrix}$$

is a 6×6 jacket matrix.

2.2 Case 2: Butson Generalized Hadamard matrices

Let B be a Butson generalized Hadamard matrix of order p , p an odd prime. Let C be the core of B , as defined earlier. Let $A = C$, $B = C^M$, $D = -C$. Then

$$X = \begin{pmatrix} 1 & e^t & e^t & 1 \\ e & C & C^M & e \\ e & C^M & -C & -e \\ 1 & e^t & -e^t & -1 \end{pmatrix}$$

is a $2p \times 2p$ jacket matrix. We observe that the $p = 3$ case is a permutation of the jacket matrix in part 2.1.

Theorem 1 *Let p be an odd prime. Then for every order $2p$, there is a jacket matrix whose entries are the p th roots of unity.*

Taking the Kronecker product of X with t copies of $H_2 = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$, $t \geq 1$, gives the following:

Theorem 2 *Let p be an odd prime. Then there are jacket matrices of order $2^{t+1}p$, $t \geq 0$.*

Where the matrix X has a border of ± 1 , the jacket matrices constructed by the Kronecker product will have a t -deep border of $\pm H_2$. We call such a matrix a jacket matrix with t -size border.

2.3 Case 3: Other generalized Hadamard matrices

Theorem 3 *Given a symmetric generalised Hadamard matrix*

$$G = (g_{ij}) = GH(n, \mathbf{G})$$

of order n over the group \mathbf{G} , there exists a jacket matrix of order $2^{t+1}n$, $t \geq 1$.

For example, consider the matrix $GH(6; \mathbb{Z}_3)$

$$G = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & \omega & \omega^2 & \omega^2 & \omega \\ 1 & \omega & \omega^2 & \omega^2 & \omega & 1 \\ 1 & \omega^2 & \omega^2 & \omega & 1 & \omega \\ 1 & \omega^2 & \omega & 1 & \omega & \omega^2 \\ 1 & \omega & 1 & \omega & \omega^2 & \omega^2 \end{pmatrix}$$

Then the core C of G can be used to construct a jacket matrix of order 12, using the construction in part 2.2.

2.4 A general result

Theorem 4 *Let D_1, D_2, \dots, D_k be jacket matrices, where D_i has order $2^{t_i+1}n_i$, $t_i \geq 0$. Then the Kronecker product*

$$D_1 \otimes \dots \otimes D_k \underbrace{\otimes H_2 \dots \otimes H_2}_{\ell \text{ times}}$$

is a jacket matrix with ℓ -size border, of order $2^m \prod_{i=1}^k n_i$, where $m = k + \ell + \sum_{i=1}^k t_i$.

References

- [1] A. T. Butson, Generalized Hadamard matrices, *Proceedings of the American Mathematical Society* 13 (1963), 894–898.
- [2] W. de Launey, Generalised Hadamard matrices whose rows and columns form a group, in *Combinatorial Mathematics X*, volume 1036 of *Lecture Notes in Mathematics*, pp. 154–176, Berlin-Heidelberg-New York, 1983. Springer-Verlag.
- [3] W. de Launey, A survey of generalised Hadamard matrices and difference matrices $D(k, \lambda; g)$ with large k , *Utilitas Math.* 30 (1986), 5–29.
- [4] D. A. Drake, Partial λ -geometries and generalized matrices over groups, *Canad. J. Math.* 31 (1979), 617–627.

- [5] M. H. Lee, The center weighted Hadamard transform, *IEEE Trans. Circuits Syst.* 36 (1989), 1247–1249.
- [6] M. H. Lee, Fast complex reverse jacket transform, In *Proc. 22nd Symp. Information Theory and its Application (SITA 99)*, Yuzawa, Niigata, Japan, Nov. 30–Dec. 3 1999.
- [7] M. H. Lee, B. S. Rajan and J. Y. Park, A generalized reverse jacket transform, *IEEE Trans. Circuits Syst. II*, 48(7) (2001), 684–690.
- [8] V. Mavron and V. D. Tonchev, On symmetric nets and generalized Hadamard matrices from affine designs, *J. Geometry* 67 (2000), 180–187.
- [9] J. Seberry, A construction for generalized Hadamard matrices, *J. Statistical Inference and Planning* 6 (1980), 365–368.
- [10] J. Seberry and M. Yamada, Hadamard matrices, sequences, and block designs, in J. H. Dinitiz and D. R. Stinson, eds., *Contemporary design theory: a collection of surveys*, pp. 431–560. John Wiley & Sons, 1992.

(Received 4 Nov 2004)