



UNIVERSITY  
OF WOLLONGONG  
AUSTRALIA

University of Wollongong  
Research Online

---

Faculty of Engineering and Information Sciences -  
Papers: Part B

Faculty of Engineering and Information Sciences

---

2019

# Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use: Systematic Review

Ting Song

*University of Wollongong, ts783@uowmail.edu.au*

Siyu Qian

*University of Wollongong, sqian@uow.edu.au*

Ping Yu

*University of Wollongong, ping@uow.edu.au*

---

## Publication Details

Song, T., Qian, S. & Yu, P. (2019). Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use: Systematic Review. *JMIR mHealth and uHealth*, 7 (1), e10899-1-e10899-14.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:  
[research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

# Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use: Systematic Review

## Abstract

**Background:** Unhealthy alcohol use (UAU) is one of the major causes of preventable morbidity, mortality, and associated behavioral risks worldwide. Although mobile health (mHealth) interventions can provide consumers with an effective means for self-control of UAU in a timely, ubiquitous, and cost-effective manner, to date, there is a lack of understanding about different health outcomes brought by such interventions. The core components of these interventions are also unclear.

**Objective:** This study aimed to systematically review and synthesize the research evidence about the efficacy of mHealth interventions on various health outcomes for consumer self-control of UAU and to identify the core components to achieve these outcomes.

**Methods:** We systematically searched 7 electronic interdisciplinary databases: Scopus, PubMed, PubMed Central, CINAHL Plus with full text, MEDLINE with full text, PsycINFO, and PsycARTICLES. Search terms and Medical Subject Headings "mHealth," "text message," "SMS," "App," "IVR," "self-control," "self-regulation," "alcohol\*," and "intervention" were used individually or in combination to identify peer-reviewed publications in English from 2008 to 2017. We screened titles and abstracts and assessed full-text papers as per inclusion and exclusion criteria. Data were extracted from the included papers according to the Consolidated Standards of Reporting Trials-EHEALTH checklist (V 1.6.1) by 2 authors independently. Data quality was assessed by the Mixed Methods Appraisal Tool. Data synthesis and analyses were conducted following the procedures for qualitative content analysis. Statistical testing was also conducted to test differences among groups of studies.

**Results:** In total, 19 studies were included in the review. Of these 19 studies, 12 (63%) mHealth interventions brought significant positive outcomes in improving participants' health as measured by behavioral (n=11), physiological (n=1), and cognitive indicators (n=1). No significant health outcome was reported in 6 studies (6/19, 32%). Surprisingly, a significant negative outcome was reported for the male participants in the intervention arm in 1 study (1/19, 5%), but no change was found for the female participants. In total, 5 core components reported in the mHealth interventions for consumer self-control of UAU were context, theoretical base, delivery mode, content, and implementation procedure. However, sound evidence is yet to be generated about the role of each component for mHealth success. The health outcomes were similar regardless of types of UAU, deployment setting, with or without nonmobile cointervention, and with or without theory.

**Conclusions:** Most studies reported mHealth interventions for self-control of UAU appeared to be improving behavior, especially the ones delivered by short message service and interactive voice response systems. Further studies are needed to gather sound evidence about the effects of mHealth interventions on improving physiological and cognitive outcomes as well as the optimal design of these interventions, their implementation, and effects in supporting self-control of UAU.

## Disciplines

Engineering | Science and Technology Studies

---

**Publication Details**

Song, T., Qian, S. & Yu, P. (2019). Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use: Systematic Review. *JMIR mHealth and uHealth*, 7 (1), e10899-1-e10899-14.

---

Review

# Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use: Systematic Review

---

Ting Song<sup>1,2</sup>, MIT; Siyu Qian<sup>1,2</sup>, PhD; Ping Yu<sup>1,2</sup>, PhD

<sup>1</sup>Centre for IT-Enabled Transformation, School of Computing and Information Technology, Faculty of Engineering and Information Sciences, University of Wollongong, Wollongong, Australia

<sup>2</sup>Illawarra Health and Medical Research Institute, University of Wollongong, Wollongong, Australia

**Corresponding Author:**

Ping Yu, PhD

Centre for IT-Enabled Transformation, School of Computing and Information Technology

Faculty of Engineering and Information Sciences

University of Wollongong

Northfields Avenue

Wollongong, 2522

Australia

Phone: 61 2 4221 5412

Fax: 61 2 4221 4170

Email: [ping@uow.edu.au](mailto:ping@uow.edu.au)

---

## Abstract

**Background:** Unhealthy alcohol use (UAU) is one of the major causes of preventable morbidity, mortality, and associated behavioral risks worldwide. Although mobile health (mHealth) interventions can provide consumers with an effective means for self-control of UAU in a timely, ubiquitous, and cost-effective manner, to date, there is a lack of understanding about different health outcomes brought by such interventions. The core components of these interventions are also unclear.

**Objective:** This study aimed to systematically review and synthesize the research evidence about the efficacy of mHealth interventions on various health outcomes for consumer self-control of UAU and to identify the core components to achieve these outcomes.

**Methods:** We systematically searched 7 electronic interdisciplinary databases: Scopus, PubMed, PubMed Central, CINAHL Plus with full text, MEDLINE with full text, PsycINFO, and PsycARTICLES. Search terms and Medical Subject Headings “mHealth,” “text message,” “SMS,” “App,” “IVR,” “self-control,” “self-regulation,” “alcohol\*,” and “intervention” were used individually or in combination to identify peer-reviewed publications in English from 2008 to 2017. We screened titles and abstracts and assessed full-text papers as per inclusion and exclusion criteria. Data were extracted from the included papers according to the Consolidated Standards of Reporting Trials-EHEALTH checklist (V 1.6.1) by 2 authors independently. Data quality was assessed by the Mixed Methods Appraisal Tool. Data synthesis and analyses were conducted following the procedures for qualitative content analysis. Statistical testing was also conducted to test differences among groups of studies.

**Results:** In total, 19 studies were included in the review. Of these 19 studies, 12 (63%) mHealth interventions brought significant positive outcomes in improving participants' health as measured by behavioral (n=11), physiological (n=1), and cognitive indicators (n=1). No significant health outcome was reported in 6 studies (6/19, 32%). Surprisingly, a significant negative outcome was reported for the male participants in the intervention arm in 1 study (1/19, 5%), but no change was found for the female participants. In total, 5 core components reported in the mHealth interventions for consumer self-control of UAU were context, theoretical base, delivery mode, content, and implementation procedure. However, sound evidence is yet to be generated about the role of each component for mHealth success. The health outcomes were similar regardless of types of UAU, deployment setting, with or without nonmobile cointervention, and with or without theory.

**Conclusions:** Most studies reported mHealth interventions for self-control of UAU appeared to be improving behavior, especially the ones delivered by short message service and interactive voice response systems. Further studies are needed to gather sound evidence about the effects of mHealth interventions on improving physiological and cognitive outcomes as well as the optimal design of these interventions, their implementation, and effects in supporting self-control of UAU.

(*JMIR Mhealth Uhealth* 2019;7(1):e10899) doi:[10.2196/10899](https://doi.org/10.2196/10899)

**KEYWORDS**

systematic review; alcohol drinking; self-control; mobile health; mHealth; treatment outcome

## Introduction

### Background

Unhealthy alcohol use (UAU) is one of the major causes of preventable morbidity, mortality, and related behavioral risks around the world [1,2]. Approximately 3.3 million deaths, accounting for 5.9% of global deaths, were caused by alcohol-related problems annually [3]. Nearly 81% of adults in Australia and 70% in Europe consume alcohol [3,4]. UAU contributed to around 70,000 Australian emergency department presentations in 2014 and 2015 and 77,000 Canadian hospitalizations in 2015 and 2016 [5,6]. It might cause allergic reactions, hormonal disturbances, and intoxication [7,8]. Over time, it might cause diseases such as alcoholic hepatitis, diabetes, cardiovascular and cerebrovascular diseases [9], or psychological problems such as depression, obsession, mania, and suicide [10,11]. Once the brain and neurons are anesthetized, a person might lose self-control [12], leading to social problems such as conflicts, unprepared sexual activities, drunk driving, and violence [13,14]. Therefore, UAU is not only a profound public health challenge but also a social concern.

As an umbrella term, UAU covers various degrees of negative effects of alcohol use on people's well-being [15]. According to the severity, there are 2 major types of UAU: risky drinking and alcohol use disorder (AUD) [15,16].

Risky drinking is also known as problematic drinking, harmful alcohol use, risky single-occasion drinking (RSOD), or heavy episodic drinking. It refers to alcohol use that leads to the risk of negative health consequences [16]. It can be measured by the number of standard drinks (SDs) consumed. An SD is defined by the amount of pure alcohol contained in a drink, and it varies among countries [14,16,17]. For example, in Australia, an SD contains 10 g of pure alcohol, in the United Kingdom and Iceland, it contains only 8 g, whereas in Austria it is 20 g [17]. It is deemed risky drinking if alcohol consumption is more than 5 SDs for men and 4 for women on a single occasion [18]. If total weekly alcohol consumption is greater than or equal to 15 SDs for men and 13 for women in the United States or over 14 SDs for men and 9 for women in Sweden, it is also considered as risky drinking [19-21]. Risky drinking can also be measured by scales such as fast alcohol screening test (FAST), alcohol use disorders identification test (AUDIT), and AUDIT for consumption (AUDIT-C) by scoring 3 or higher in FAST [22], over 8 for men and 6 for women in AUDIT [23], or 4 for men and 3 for women in AUDIT-C [24].

The other major type of UAU is AUD. It is a chronically recurrent brain impairment in which compulsive and maladaptive alcohol use results in behavior dysregulation and negative mood once alcohol consumption is ceased [16,25]. Alcohol abuse and alcohol dependence are 2 major representatives for moderate and severe degrees of AUD, respectively [16,25,26]. Consumers with either of them can suffer from adverse consequences. Alcohol abuse, that is, unrestrained alcohol use, can make consumers fail to meet their

major obligations and cause or exacerbate health and social problems [16,27]. More seriously, alcohol dependence, that is, a constant and strong desire for alcohol use without self-control or consideration of health, might result in physical or mental health problems once a large amount of alcohol is consumed over a long period [28]. To be diagnosed with AUD, a person should meet at least two of the 11 criteria listed in the Diagnostic and Statistical Manual of Mental Disorders 5th Edition in 1 year [29].

Mobile health (mHealth), also known as ecological momentary intervention [30,31], refers to the use of mobile devices, such as mobile phones, personal digital assistants, or other wireless devices, to deliver medical or public health services in a timely manner and in real-world living settings [30,32]. Due to limited human resources available for delivering continuous health care services for community-dwelling consumers suffering from chronic diseases, mHealth interventions are increasingly considered by the decision makers as a potential alternative solution in providing same quality but low-cost services [33]. Similarly, there has been a growing interest in using mobile phones to deliver public health interventions to support consumer self-control of UAU.

mHealth interventions are mainly delivered solely or in combination of 3 channels: short message services (SMS) text messaging, apps, and interactive voice response (IVR). SMS text messaging has been used to guide consumers to change alcohol use behavior, for example, to reduce alcohol intake to enable self-control of UAU [19,34]. Apps have been used to monitor consumers' alcohol use and to provide visual feedback about drinking behavior based on statistical analysis of input data. Raising self-awareness can ignite consumers' self-regulation so as to reduce alcohol use [35,36]. IVR has been used to generate audial interactions and to provide automatic answers to consumer queries on UAU [37,38]. Therefore, these 3 delivery channels can all provide effective and efficient interventions for consumer self-control of UAU.

### Objectives

Recent reviews on digital interventions for self-control of UAU focus on the benefit of such interventions on improving health care services. In total, 2 reviews investigated electronic or Web-based interventions and found that despite a small effect, these interventions might improve behavioral outcomes, particularly for the group less likely to access traditional alcohol interventions such as women, youth, and risky drinkers [39,40]. A total of 5 reviews narrowed down the scope on mHealth interventions for self-control of UAU. In total, 2 of them focused on SMS text messaging and found that although the behavioral outcomes were modest, it was still a worthwhile endeavor [41,42]. The other 3 reviews suggested mHealth interventions, especially the ones that can provide personalized feedback, were beneficial for the reduction of UAU with their high fidelity, anonymity, and accessibility [31,43,44]. However, as the mHealth interventions were still nascent in nature, there is still a lack of understanding about how such interventions really

work for changing UAU. Solid evidence about the efficacy of these interventions from empirical field trials is required. Moreover, the other health outcomes, such as physiological and cognitive outcomes, need to be studied. Therefore, this review aimed to synthesize and understand the research evidence about the efficacy of mHealth interventions on various health outcomes for consumer self-control of UAU and to identify the core components to achieve these outcomes.

## Methods

### Study Design

A mixed-methods systematic review was conducted. Literature search and screening followed the preferred reporting items for systematic reviews and meta-analyses [45]. Data extraction was guided by the Consolidated Standards of Reporting Trials-electronic health checklist (V.1.6.1) [46]. The methodological quality of the studies was assessed by the Mixed Methods Appraisal Tool (MMAT) [47]. Data synthesis and analysis followed the principle of realist synthesis [48] and qualitative content analysis [49].

### Literature Search and Screening

The literature search was performed from December 2016 to March 2017 and further refined in August 2018 in 7 electronic interdisciplinary databases: Scopus, PubMed, PubMed Central, CINAHL Plus with full text, MEDLINE with full text, PsycINFO, and PsycARTICLES (see [Multimedia Appendix 1](#)). The following terms and medical subject headings were used individually or in combination to identify the relevant publications: “mHealth,” “text message,” “SMS,” “App,” “IVR,” “self-control,” “self-regulation,” “alcohol\*,” and “intervention.” To ensure adequate coverage, a manual search was also conducted to identify papers from *Journal of Medical Internet Research* and its sister journals. The search was restricted to peer-reviewed journal papers published in English between 2008 and 2017. In addition, the following criteria were used in the selection of papers.

### Inclusion Criteria

The papers were included in which (1) the research focused on supporting consumer self-control of UAU; (2) health intervention was delivered through mobile phone technologies; and (3) the data were collected from empirical randomized controlled trials.

### Exclusion Criteria

The papers were excluded that (1) reported clinical therapy such as injection and medication rather than consumer active participation in the daily self-control of UAU; (2) did not report any alcohol-related health outcome; (3) used the intervention not dealing with UAU or containing Web-based components delivered by desktop or Web-based computer applications; or (4) were review papers, study protocols, conceptual papers, editorials, government reports, or guidelines in the topic area.

### Data Extraction

Data were extracted using a combination of an Endnote X8 and an Excel spreadsheet by 2 authors independently. These included

name(s) of the author(s), year of publication, country of origin, population type, study setting, type of UAU, study type, eligibility, sample size, study arms and grouping, nonmobile cointervention, mHealth intervention theory, delivery mode, mHealth intervention content, implementation procedure, measurement, and outcomes.

### Quality Assessment of the Studies

All studies were assessed using the 4 criteria in section 2 of the MMAT, in terms of (1) randomization or sequence generation, checking if there is a clear description about randomization; (2) allocation concealment, verifying if there is a clear description about blinding; (3) outcome data, confirming if more than 80% outcomes were reported; and (4) attrition, assessing if less than 20% of the participants dropped out. Responses to each criterion were “yes,” “no,” or “can’t tell.”

### Data Synthesis and Analyses

Data were synthesized and analyzed using an inductive method. We reviewed all data that collected and identified similar notions and tagged them with the same code. Thereafter, we grouped the codes with similar meaning into an overarching concept. Concepts with similar meaning were grouped into a category that addresses our research question. The coding and data management were iteratively developed through constant comparison of the similarities and differences among codes.

To explore the initial outcomes about which components really make the intervention works, chi-square testing was conducted to test the relationship between health outcomes with the following 4 parameters: (1) types of UAU, being risky drinking or AUD; (2) with or without nonmobile cointervention; (3) theory-based or not; (4) deployment setting, being clinical, educational, or community based.


## Results

### Search Outcome

The primary search yielded 1345 publications. After removing duplicates, 517 papers remained. Their titles and/or abstracts were manually screened against the inclusion and exclusion criteria. This led to 41 candidate papers. Of these, 20 were excluded after further scrutinizing the full paper (see [Multimedia Appendix 2](#)). Finally, 21 papers were included. Of these, 4 papers were from 2 studies. Suffoletto et al published 2 papers based on the same study population in 2014 and 2015 [50,51], respectively; so did Agyapong et al in 2012 and 2013 [52,53]. Therefore, a total of 19 studies were eligible for review (see [Figure 1](#) and [Multimedia Appendix 3](#)). Among these studies, 58% (11/19) met all 4 MMAT criteria [19,34,37,38,52-59] and 32% (6/19) met 3 criteria [20,21,35,50,51,60,61], indicating high methodological quality in 90% of these studies (see [Multimedia Appendix 4](#)).

### Characteristics of Studies

Although we searched studies published since 2008, all 19 eligible studies were conducted in 2012 and beyond and were from 7 developed countries (see [Multimedia Appendix 5](#)).

**Figure 1.** Literature search and screening process.

Of these, 9 studies (9/19, 47%) were conducted in the United States [19,34,35,38,50,51,55,57,59,62], 8 in Europe [20,21,37,52-54,56,58,61], and only 2 in New Zealand [60,63].

Study arms ranged from 2 to 6. In total, 12 studies (12/19, 63%) were 2-arm trials with an intervention arm and a control arm [21,34,35,52,53,55,56,58-63]. The control-arm participants received (1) no intervention [35,56,58]; (2) nonmobile intervention with the same content through interview [55,62], email [21], and e-booklet [61]; (3) nonalcohol-related content [34,52,53] or only assessment for monitoring purpose [60,63] through the same mobile devices; or (4) different rewarding mechanisms for their abstinence [59]. A total of 5 studies (5/19, 26%) had 3 arms. Of these, 2 added an assessment-only arm besides the intervention and control arms [50,51,57]. Hasin et al employed an arm in which the participants only received intervention through interview [38]. Gajecki et al used 2 intervention arms delivered by 2 different mobile apps in 1 study [20] and 2 intervention arms that started to use the app at different times in another study [54]. In the last 2 studies (2/19,

11%), Andersson conducted a 5-arm trial in which an mHealth intervention was compared with Web-based intervention and nonintervention. Both the mHealth and Web-based interventions had 2 implementation procedures, single and repeated [37]. Muench et al employed a 6-arm design, including 1 nonintervention arm, 1 assessment-only arm, and 4 intervention arms containing different contents [19].


We identified 5 core components of mHealth interventions for UAU: context, theoretical base, content, delivery mode, and implementation procedure and 3 types of potential health outcomes: behavioral, physiological, and cognitive outcome (see Figure 2 and Multimedia Appendix 6).

### Five Core Components of Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use

#### Context

There are 3 types of contexts: participant characteristics, deployment setting, and nonmobile cointervention, which were conducted simultaneously to support the mHealth intervention.

**Figure 2.** In total, 5 core components of mobile health interventions for self-control of unhealthy alcohol use and 3 types of potential health outcomes.



The participants can be categorized into 2 cohorts according to their age: youth group aged below 35 years [20,21,34,37,50,51,54,57,59,60,62,63] and middle and old aged group, aged above 35 years [19,35,38,52,53,55,56,58,61]. They were either risky drinkers [19-21,34,37,38,50,51,57,59-63] or had AUD [20,35,38,52-56,58]. They suffered from comorbidity of depression [52,53], HIV [38,55], drug dependence [55], or smoking [57]. The interventions were deployed in educational settings [20,21,34,37,54,57,60,62,63], clinical settings [35,38,50-53,56,58], and community-based settings [19,55,59,61]. The nonmobile cointervention included social intervention guided by the theory of motivational interviewing [38,50,51,55,57,62] and paper-based intervention in which participants were provided with guidelines for safe alcohol use about the mHealth intervention [19].

### Theoretical Base

In total, 2 types of theories were reported to guide the design and implementation of the mHealth interventions, including behavioral change theories and psychological theories of motivation.

Behavioral change theories included theory of planned behavior [20,21,50,51,56,63], health belief model [19,50,51], social cognitive theory [21,63], theory of reasoned action [50,51], information motivation behavioral model [50,51], cognitive behavioral therapy [57], and social learning theory [19]. Psychological theories of motivation included self-determination

theory [21,35,63], model of action phases [21,63], and contingency management [59]. Notably, although Aharonovich et al did not report any explicit theory applied to their intervention, the design of their app, HealthCall, was theory based [55].

### Delivery Mode

A total of 3 delivery modes were identified: SMS text messaging (12/19, 63%) [19,21,34,50-53,56,58-63], app (5/19, 26%) [20,35,54,55,57], and IVR (2/19, 11%) [37,38]. In total, 6 apps tested in the 5 studies were TeleCoach [54], Brief Alcohol and Smoking Intervention for College Students via Mobile (BASICS-Mobile) [57], HealthCall-S [55], Alcohol-Comprehensive Health Enhancement Support System (A-CHESS) [35], PartyPlanner, and Promillekoll [20]. Only 2 studies described the underlying operating system for these apps [20,57]. PartyPlanner and Promillekoll ran on the Android or iOS [20], and BASICS-Mobile ran on Blackberry, Android, or iOS [57].

### Content

In total, 3 types of content were designed to support the participants' self-control of UAU. They were information [19-21,34,35,37,38,50-58,60-63], motivation [19,21,34,35,37,50-53,55-57,59,61,62], and reminder [19,35,50,51,55,56,58,59].

Informational content included general and personalized information. The general information facilitated the participants



in (1) enriching their knowledge about risks and negative consequences of UAU [19,34,37,50,51,60,61,63], alcohol-related facts [34,37,54,57,61], social drinking norms [50,51,62], and benefits of reducing drinking amount according to safety guidelines [19]; (2) acquiring strategies to control alcohol use [20,34,50,51,54,55,57,58,62], to handle relapse or cravings [52-54,57,61], to manage emotion [54,61], and to reduce intoxication [37]; (3) getting referrals such as alcohol counseling services [55,58], instant library, and weblinks to further alcohol-related information [35]; and (4) conducting recommended actions for self-control of UAU such as tracking and reporting their drinking facts [19,20,35,37,38,50,51,54-56,58,62,63], reasons for drinking or abstinence [38,55], and estimated blood alcohol concentration (eBAC) value [20], mood [38], medication adherence [38], and well-being [38]; introspecting their performance [21]; or simulating a drinking occasion to set personal goal of eBAC and comparing actual eBAC after drinking against this goal [20].

The personalized information helped the participants in (1) providing the tailored feedback according to their responses [19,37,50,51,55,57]; (2) recommending them to set intermittent low-risk drinking goals [61] to replace drinking alcohol by alternative activities [37,57], to celebrate goal attainment [50,51,61], to self-reflect on challenges of UAU [21], to improve the drinking plan, and to reinforce self-control behavior [37,50,51]; and (3) addressing their problems identified at various stages [19].

Motivational content included (1) encouragement messages for reducing alcohol use [21,34,37,50-53,57,61,62], committing to preset drinking goals [19,50,51,55,56] and medical adherence [52,53], and releasing distress [35]; (2) peer support through sharing experiences with others in the anonymous discussion groups [35]; and (3) possible monetary compensation to incentivize participants to submit their valid on-time video [59].

Reminding content facilitated participants in (1) reminding them to remember and fulfill their promises [50,51,55,56,58,59] and (2) warning them about alcohol risks at their risky drinking times [19] or when they were near high-alcohol places detected by global positioning system [35].

### **Implementation Procedure**

The duration of the interventions varied, ranging from 4 days [62], 1 week [37,60,63], 2 weeks [57], 4 weeks [21,37,59], 6 weeks [34], 7 weeks [20], 2 months [38,55,58], 3 months [19,50-54], 6 months [56,61], to 8 months [35].

With regard to the frequency, SMS text messages were sent once [34], twice [50,51,58], or 4 times [21,60] weekly in 5 studies and once [63], twice [52,53], 1 to 3 times [59], or 4 to 6 times [62] daily in 4 studies. The frequency appeared to reduce when the length of the study increased [56,61]. Haug et al sent 1 SMS text message per week in the first 8 weeks and then 1 per fortnight in the remaining 18 weeks [56]. Brendryen et al sent 1 SMS text message per day for 8 weeks, then 1 per week for 4 weeks, and finally 1 per month in the last 2 months [61]~Brendryen, 2014 #8^ . Muench et al sent the SMS text messages with different content at a different frequency, either

once daily for educating participants about alcohol use or once weekly for self-monitoring content and feedback [19]. The participants in Alessi and Petry's study were given a breathalyzer and the corresponding accessories to self-measure breath alcohol concentration (BrAC) and submit a valid real-time video containing the whole self-measuring process to the organizer via SMS text message 1 to 3 times per day at the fixed time interval to prove their abstinence [50]. The intervention-arm participants would be rewarded with more vouchers if their BrAC value was normal. In contrast, the control-arm participants were not rewarded although their BrAC value was normal [50].

In 3 studies in which the interventions were delivered by mobile apps, the frequency of data collection was once daily in Aharonovich et al's study [55] and once weekly in Gustafson et al and Gajecki et al's study [35,54]. In most cases, apps were used in real time according to a participant's preference, typically to receive a certain recommendation once a preset condition was met. For example, Promillekoll could send real-time notification and the corresponding strategies to control alcohol use if a participant's eBAC was over 0.06% [20]. A-CHESS would send an alarm when a participant was near a high-risk alcohol place to be detected by the embedded global positioning system [35].

Andersson divided his intervention arm into 2 subgroups, both receiving the same content but through different delivery modes, either delivering single IVR every day for 1 week or delivering repeated IVR for 4 weeks [37]. Hasin et al requested their intervention-arm participants to spend 1 to 3 min per day to send back their answers to a series of questions, asking their compliance with drinking guidelines on the previous day via a toll-free number [38]. The participants' phone calls were initially answered by the prerecorded IVR in the first 30 days. After evaluating a participant's IVR data, the consultant reset the person's drinking goal for the next 30 days [38].

In terms of timing of delivery, 10 studies reported the fixed time or time slots to deliver the intervention, the rest were flexibly available on demand. The popular days of intervention delivery were Thursday [34,50,51,58,60,63], Saturday [21,34,59,60,63], and Sunday [21,50,51,59,63], mainly at or after 6 pm [19,34,52,53,56,59,60,63].

All studies conducted the baseline assessments. In total, 6 studies conducted an assessment during the intervention period to explore the initial outcome [35,38,54,55,58,61]. The postintervention assessments were conducted in all studies at different time points with different numbers of repetitive measurement. A total of 17 studies conducted 1 assessment immediately after the intervention [19-21,34,35,38,50-61,63]. A total of 8 studies conducted the second assessment 1 month [38,57], 6 weeks [34], 1 academic semester [60,63], 3 months [50-53], and 4 months [35] after the intervention. In total, 2 studies conducted the third assessment 3 [50,51] and 4 months [38] after the intervention. Only 1 study conducted the fourth assessment after 10 months of the intervention [38]. Instead of immediately measuring the outcomes, in 2 studies, the measures were conducted only after 1 month [62]. Of these, 1 study measured the outcome 4 weeks after the intervention for the

single IVR intervention arm and 1 week after the intervention for the repeated IVR intervention arm [37].

## Health Outcomes

### Behavioral Outcome

Behavioral outcome was measured in 18 studies [19,20,34,35,37,38,50-63]. Significant positive outcome was found in 11 of these studies [19,35,37,38,50-54,59-61,63]. These positive outcomes were measured by 1 or more indicators. These included the decreased number of SDs [19,50,51,54,61,63], heavy drinking days [19,35,38,50,51,59], RSOD or binge drinking prevalence [50,51,54], alcohol-related injury prevalence [50,51], and peak eBAC value [37]; increased number of abstinence days [19,52,53,59] or the increased negative affect score in Alcohol Abstinence Self Efficacy Scale [52,53]; and the decreased score in the Alcohol Addiction Severity Index, Drinker Inventory of Consequences [59], or AUDIT [37].

No significant behavioral change was found in 6 studies [34,55-58,62]. In total, 2 studies reported a gender-related behavioral outcome [20,63]. Contrary to the initial objective of reducing UAU, the male participants in the intervention arm significantly increased drinking frequency, whereas no change was found in the female participants and the control arm in 1 study [20]. In the study conducted by Riordan et al, after providing intervention-arm participants with 1-week SMS text messages, the female participants consumed significantly less alcohol 1 week and 1 semester later than their female counterparts in the control arm. However, no intervention effect was found for the male participants [63].

### Physiological Outcome

Physiological outcome was measured in only 1 study via BrAC [59]. Alessi and Petry found a significant improvement in the percentage of negative BrAC in the intervention group but no significant change in the control group [59].

### Cognitive Outcome

Cognitive outcome was measured in 3 studies [21,34,62] and was significantly positive in only 1 study in which the participants' readiness to change UAU behavior in the intervention arm was significantly improved [62]. No significant cognitive change was found in the other 2 studies in terms of motivation to change and self-confidence to resist alcohol [21,34].

## Comparison With the Differences in Health Outcomes Among Different Groups of Studies

Over half of the SMS- and IVR-enabled interventions were effective in reducing alcohol use or increasing readiness to change UAU in 8 out of 12 studies (67%) [19,50-53,59-63] and 2 out of 2 studies (100%) [37,38], respectively. In contrast, app-enabled interventions were only successful in reducing alcohol use in 2 out of 5 studies (40%) [35,54].

Chi-square test did not find any significant differences in health outcomes among groups of studies with different conditions. It suggested that the health outcomes were similar regardless of the types of UAU studied, whether there was nonmobile

cointervention, whether the study was theoretical-based, or which setting it was deployed.

## Discussion

### Principal Findings and Comparison With Previous Work

This study aimed to synthesize and understand the research evidence about efficacy of mHealth interventions on different health outcomes for consumer self-control of UAU and to identify their core components to achieve these outcomes. In total, 19 studies were systematically reviewed and 3 types of health outcomes such as behavioral, physiological, and cognitive outcome and 5 components of these interventions such as context, theoretical base, delivery mode, contents, and implementation procedure were found.

### Health Outcomes

As approximately two-thirds (11/18) of the studies that measured the behavioral outcomes identified a significant positive change [19,35,37,38,50-54,59-61,63]: mHealth interventions appear to be more effective in changing UAU behavior in comparison with the traditional methods. The results could be explained by the information-motivation-behavioral skills model, which suggests that a participant's behavior change is attributed to the provided information, motivation, and improved skills [64]. This is also in accordance with the findings of Regmi et al's review in smoking cessation context where the abstinence days of smoking increase after applying mHealth interventions [65].

Despite the significant 100% positive physiological outcome measured by BrAC, only 5% of the included studies assessed the physiological measurement [59]. There might be two reasons for this. First, people tend to test their biomedical markers in hospital or clinic rather than by themselves as they might lack corresponding skills and it is inconvenient. Second, the corresponding self-testing devices are not cheap and not all research projects can afford them, especially for the projects with a large sample size. Instead, researchers preferred to measure behavioral outcomes because the rough BAC can be calculated simply using Widmark formula once a participant reports his or her alcohol use [66]. In addition, the unique factor to affect the BrAC is alcohol intake. Therefore, the physiological outcome must change when the behavior changes.

For the same reason, as the cognitive outcomes are inconvenient to measure in comparison with behavioral ones, only 16% of the studies [21,34,62] assessed cognitive changes, of which 33% [62] were significantly improved. This might be because cognition can be influenced by various factors, and their measurement can be somewhat subjective and abstract. For example, Mason et al assessed cognitive change using 5 variables: alcohol expectations, readiness to change drinking behavior, importance of change, confidence in ability to change, and intentions to reduce alcohol use. Only the variable of readiness to change drinking behavior was improved [62]. Notably, although all these 3 studies also reported the improvement in behavior changes, it is still not enough to conclude that behavior always changes with cognition. According to the theory of cognitive dissonance proposed by

Leon Festinger, a person can be motivated to reduce own psychological inconsistency and discomfort by changing the behavior [67]. No matter whether the interventions were genuinely accepted by the participants or not, most of them modified their behavior in compliance with the information they received from the interventions to reduce their interconflicts [67]. Therefore, when their cognitive changes were assessed, they might provide the real thoughts, which might not be consistent with the behavior that were displayed.

Complementing the traditional interventions such as face-to-face counseling, in which unhealthy alcohol users' access to treatment was provided in a passive manner within a confined time and location, mHealth interventions open new opportunities for engaging consumers in positive self-control with increased flexibility. The effect of control was improved by continuous tracking and monitoring, interactive communication, or personalized feedback from health care providers anytime, anywhere [30,68,69].

### ***Five Components of Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use***

Participants in most reviewed studies were risky drinkers without documented pathological conditions [19-21,34,37,38,50,51,57,59-63]. We did not find much difference in the intervention outcome between the types of participants, being risky drinker or AUD. This result is consistent with the finding of Blow et al that health outcomes of an intervention are not influenced by the level of severity of alcohol addiction [70]. However, this is contradictory with the findings in the previous review conducted by White et al that e-interventions can be particularly useful for at-risk users [40]. Kazemi et al also seconded that for this population group, mHealth intervention might be the most cost-effective UAU management strategy [71]. The paradox might be caused by the different conditions such as timing and frequency of the interventions or different population types and settings [42].

The gender difference in intervention outcome found in 2 studies [20,63] might be explained by the observation of Hirschi and Gottfredson that men have lower self-control than women [72]. Notably, these 2 studies were both done on young adult students in university settings. This might suggest that it is much more difficult for males in this setting to change their behavior in terms of UAU. First, there are strong social or peer norms in this cohort, which prevent the change of drinking behavior [73], and second, males seem to be less compliant and agreeable than females, and they lack ability to absorb the meaning of the SMS text messages [74,75]. Riordan et al offered some suggestions for improving SMS text messaging for young men and later demonstrated that using more colloquial tone and sending only messages with the potential social consequences of UAU are better for this population [60]. Similar to the finding of Platt et al [76], we did not find any significant relationship between the health outcome and deployment setting.

Although not having any significant impact on health outcomes, cointervention, such as induction or training to enable a participant to confidently use the apps or IVR, is an integral, vital component for a successful mHealth intervention [77,78]. This might explain why more cases of nonmobile cointervention

were reported in interventions delivered via apps (3/5, 60%) and IVR (1/2, 50%). Most likely, the participants were more familiar with SMS text messaging than the other 2 delivery modes; therefore, the cointervention was less reported in the studies delivered by SMS text messaging (3/12, 25%). Notably, the population of the 2 studies in which interventions were delivered via apps without formal reporting of cointervention was university students at a younger age. This might be because of the internet use and mobile phone technologies are popular in this cohort; thus, the app designers did not consider it necessary to provide the students with training to use the app [79].

Behavior change theory provides the foundation for the formation of strategies to incrementally change a consumer's behavior of UAU [80]. Psychological theory of motivation is used to develop motivational strategies to control UAU against psychological craving for alcohol [81]. Although mHealth interventions based on theory can improve instructional design and the effect of self-control of UAU [76], no significant difference in health outcomes was found in this review for the studies based on theory and those otherwise, which is in accordance with the finding of Garnett et al [82]. There might be two reasons to explain this phenomenon. First, from what was described in the Methods, it appears that theory was implicitly applied to the mHealth interventions although a study might not make the claim to be theory based. For example, Bock et al did not report the use of any theory; however, one of the SMS text messages in their intervention "always have an exit plan" indicated the unconscious application of the theory of planned behavior [34]. Second, it takes time to bring in tangible health outcomes for participant's self-control of UAU [52,53].

Almost all SMS- or IVR-enabled interventions were effective in reducing alcohol use or increasing readiness to change except the mobile apps [20,55,57]. This might be because the former 2 types of interventions were delivered proactively, on regular basis, always accessible to the participants regardless of their intention. In contrast, the participants' access to the app-based interventions relied on their self-action of opening the apps, which might not always happen. This is consistent with the findings in Meredith et al's review [83], and it also recommends that the future mHealth apps need push notifications regularly to prompt the active engagement of the users.

Informational content facilitated the participants to develop essential knowledge and skill to build their capacity to change their belief and UAU behavior. It also provided necessary feedback to enable self-awareness of UAU status, which could help execute self-regulation of UAU. Motivational content provided continuous encouragement and peer support through experience sharing to raise the participants' morale in changing UAU behavior. Reminding content provided constant recall to ensure the participants to stay on track in self-control of UAU. Delivery of these 3 types of content is in line with the model of human practical reasoning developed by Michael Bratman [84].

As the length of the reviewed studies was not long enough, ranging from 4 days to 8 months, it is no surprise that there was no obvious improvement in tangible health outcomes in many studies. Longer duration, that is, 6 months or more [35,61],

more frequent delivery [52,53,59,62] and certain techniques such as tangible incentives [62], and assessment during the intervention [61] might help achieve positive outcomes. In contrast, a relatively small sample size, less than 100 [34,55,56] and a short follow-up period, less than 2 months [57], might cause a lack of significant health outcomes for the interventions. However, whether the health outcomes can be influenced by these factors still needs to be verified.

With the same content and implementation procedure, Andersson et al found differences in health outcomes measured by peak eBAC and AUDIT scores with different delivery modes in which the efficacy was better delivered by IVR than the Web [37]. Similarly, with the same delivery mode and implementation procedure but different content, Muench et al also found differences in health outcomes measured by numbers of SDs, heavy drinking days, and abstinent days. The content that highlighted the negative consequences of UAU was significantly more likely to bring about positive health outcomes than the content that emphasized the benefits of UAU abstinence [19]. Furthermore, with the same content and delivery mode, Gajek et al found that the health outcomes measured by SD and drinking frequency were significantly different with different intervals of intervention [54].

Although the first generation of iPhone was released in June 2007, marking the debut of smartphone technology [85], no eligible studies were found before 2012. It appears that using mobile phones to deliver mHealth interventions for UAU was staged in 2012.

### Limitations

The first limitation of this study was that the coverage of the studies might not be exhaustive, because of which our search

was confined to the 7 databases. However, the comprehensiveness of these databases can ensure the representativeness of the trend suggested by this study. The heterogeneity of participant characteristics, intervention, and health outcome measures makes it difficult to compare rigorously the findings among the studies. A lack of homogenous, quantitative measures in the original studies also deemed it impossible to conduct more rigorous meta-analysis. As only peer-refereed journal papers were included to ensure the rigor of this study, there could be a potential risk of reporting bias toward positive findings.

### Conclusions

This systematic review summarized the extant research evidence about the health outcomes of mHealth interventions for consumer self-control of UAU. A total of 3 health outcomes, that is, physiological, behavioral, and cognitive outcomes and 5 core components of these interventions, that is, context, theoretical base, delivery mode, content, and implementation procedure, were synthesized and analyzed. In comparison with the traditional interventions, the evidence to support effectiveness of mHealth interventions for consumer self-control of UAU is modest at best. A majority of studies showed that mHealth interventions brought positive health outcomes in helping unhealthy alcohol users to proactively engage in self-control of their UAU behavior, especially for the ones delivered by SMS text messaging and IVR systems. Sound evidence is yet to be sought about the effects of these interventions in improving the physiological and cognitive outcomes. Further research is needed to gather evidence about the optimal design of mHealth interventions, their implementation, and effects in supporting consumer self-control of UAU.

---

### Acknowledgments

The authors would like to acknowledge and thank Diane Walton for proofreading the paper.

---

### Conflicts of Interest

None declared.

---

### Multimedia Appendix 1

Search and screen records.

[[PDF File \(Adobe PDF File\), 260KB - mhealth\\_v7i1e10899\\_app1.pdf](#)]

---

### Multimedia Appendix 2

Excluded papers with reasons.

[[PDF File \(Adobe PDF File\), 21KB - mhealth\\_v7i1e10899\\_app2.pdf](#)]

---

### Multimedia Appendix 3

List of included studies.

[[PDF File \(Adobe PDF File\), 13KB - mhealth\\_v7i1e10899\\_app3.pdf](#)]

---

## Multimedia Appendix 4

Quality appraisal of the included studies.

[[PDF File \(Adobe PDF File\), 254KB - mhealth\\_v7i1e10899\\_app4.pdf](#)]

## Multimedia Appendix 5

Characteristics of the included studies.

[[PDF File \(Adobe PDF File\), 156KB - mhealth\\_v7i1e10899\\_app5.pdf](#)]

## Multimedia Appendix 6

Five components and three types of health outcomes.

[[PDF File \(Adobe PDF File\), 312KB - mhealth\\_v7i1e10899\\_app6.pdf](#)]

## References

1. Johnson NB, Hayes LD, Brown K, Hoo EC, Ethier KA, Centers for Disease Control and Prevention (CDC). CDC National Health Report: leading causes of morbidity and mortality and associated behavioral risk and protective factors--United States, 2005-2013. *MMWR Suppl* 2014 Oct 31;63(4):3-27. [Medline: [25356673](#)]
2. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012 Dec 15;380(9859):2224-2260 [FREE Full text] [doi: [10.1016/S0140-6736\(12\)61766-8](#)] [Medline: [23245609](#)]
3. World Health Organization. Global Status Report on Alcohol and Health 2014. Geneva, Switzerland: WHO; 2014.
4. Australian Institute of Health and Welfare. Canberra, Australia: AIHW; 2016. Australia's health 2016 URL: <https://www.aihw.gov.au/getmedia/9844cefb-7745-4dd8-9ee2-f4d1c3d6a727/19787-AH16.pdf.aspx?inline=true> [accessed 2018-04-25] [WebCite Cache ID [6yxA3PG2Q](#)]
5. Australian Bureau of Statistics. 2015. National Health Survey: First Results, 2014-15 URL: <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4364.0.55.0012014-15?OpenDocument> [accessed 2018-04-25] [WebCite Cache ID [6yxAESoLa](#)]
6. Canadian Institute for Health Information. Alcohol Harm In Canada: Examining Hospitalizations Entirely Caused by Alcohol and Strategies to Reduce Alcohol Harm. Ottawa, Canada: CIHR; 2017.
7. Rehm J, Gmel G, Sempos CT, Trevisan M. Alcohol-related morbidity and mortality. *Alcohol Res Health* 2003;27(1):39-51 [FREE Full text] [Medline: [15301399](#)]
8. Emanuele N, Emanuele MA. The endocrine system: alcohol alters critical hormonal balance. *Alcohol Health Res World* 1997;21(1):53-64. [Medline: [15706763](#)]
9. Shield KD, Parry C, Rehm J. Chronic diseases and conditions related to alcohol use. *Alcohol Res* 2013;35(2):155-173 [FREE Full text] [Medline: [24881324](#)]
10. Strakowski SM, DelBello MP, Fleck DE, Adler CM, Anthenelli RM, Keck PE, et al. Effects of co-occurring alcohol abuse on the course of bipolar disorder following a first hospitalization for mania. *Arch Gen Psychiatry* 2005 Aug;62(8):851-858. [doi: [10.1001/archpsyc.62.8.851](#)] [Medline: [16061762](#)]
11. Stickley A, Jukkala T, Norström T. Alcohol and suicide in Russia, 1870-1894 and 1956-2005: evidence for the continuation of a harmful drinking culture across time? *J Stud Alcohol Drugs* 2011 Mar;72(2):341-347. [doi: [10.15288/jsad.2011.72.341](#)] [Medline: [21388607](#)]
12. Abernathy K, Chandler LJ, Woodward JJ. Alcohol and the prefrontal cortex. *Int Rev Neurobiol* 2010;91:289-320 [FREE Full text] [doi: [10.1016/S0074-7742\(10\)91009-X](#)] [Medline: [20813246](#)]
13. Gonzalez-Guarda RM, Ortega J, Vasquez EP, De Santis J. La mancha negra: substance abuse, violence, and sexual risks among Hispanic males. *West J Nurs Res* 2010 Feb;32(1):128-148 [FREE Full text] [doi: [10.1177/0193945909343594](#)] [Medline: [19915207](#)]
14. Taylor B, Irving HM, Kanteres F, Room R, Borges G, Cherpitel C, et al. The more you drink, the harder you fall: a systematic review and meta-analysis of how acute alcohol consumption and injury or collision risk increase together. *Drug Alcohol Depend* 2010 Jul 01;110(1-2):108-116 [FREE Full text] [doi: [10.1016/j.drugalcdep.2010.02.011](#)] [Medline: [20236774](#)]
15. Saitz R. Unhealthy alcohol use. *N Engl J Med* 2005 Feb 10;352(6):596-607. [doi: [10.1056/NEJMcp042262](#)] [Medline: [15703424](#)]
16. Holt S, Tetrault J. Unhealthy alcohol use. *Clin Liver Dis* 2016 Aug;20(3):429-444. [doi: [10.1016/j.cld.2016.02.003](#)] [Medline: [27373607](#)]
17. International Alliance for Responsible Drinking. Drinking guidelines: General population URL: <http://www.iard.org/resources/drinking-guidelines-general-population/> [accessed 2018-04-25] [WebCite Cache ID [6yxAjzep9](#)]
18. Dawson DA. Defining risk drinking. *Alcohol Res Health* 2011;34(2):144-156 [FREE Full text] [Medline: [22330212](#)]

19. Muench F, van Stolk-Cooke K, Kuerbis A, Stadler G, Baumel A, Shao S, et al. A randomized controlled pilot trial of different mobile messaging interventions for problem drinking compared to weekly drink tracking. *PLoS One* 2017;12(2):e0167900 [FREE Full text] [doi: [10.1371/journal.pone.0167900](https://doi.org/10.1371/journal.pone.0167900)] [Medline: [28146560](https://pubmed.ncbi.nlm.nih.gov/28146560/)]
20. Gajecki M, Berman AH, Sinadinovic K, Rosendahl I, Andersson C. Mobile phone brief intervention applications for risky alcohol use among university students: a randomized controlled study. *Addict Sci Clin Pract* 2014;9:11 [FREE Full text] [doi: [10.1186/1940-0640-9-11](https://doi.org/10.1186/1940-0640-9-11)] [Medline: [24985342](https://pubmed.ncbi.nlm.nih.gov/24985342/)]
21. Bendtsen M, Bendtsen P. Feasibility and user perception of a fully automated push-based multiple-session alcohol intervention for university students: randomized controlled trial. *JMIR Mhealth Uhealth* 2014;2(2):e30 [FREE Full text] [doi: [10.2196/mhealth.3233](https://doi.org/10.2196/mhealth.3233)] [Medline: [25098296](https://pubmed.ncbi.nlm.nih.gov/25098296/)]
22. Hodgson R, Alwyn T, John B, Thom B, Smith A. The FAST alcohol screening test. *Alcohol Alcohol* 2002;37(1):61-66 [FREE Full text] [Medline: [11825859](https://pubmed.ncbi.nlm.nih.gov/11825859/)]
23. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption--II. *Addiction* 1993 Jun;88(6):791-804. [doi: [10.1111/j.1360-0443.1993.tb02093.x](https://doi.org/10.1111/j.1360-0443.1993.tb02093.x)] [Medline: [8329970](https://pubmed.ncbi.nlm.nih.gov/8329970/)]
24. Bush K, Kivlahan DR, McDonnell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol use disorders identification test. *Arch Intern Med* 1998 Sep 14;158(16):1789-1795. [doi: [10.1001/archinte.158.16.1789](https://doi.org/10.1001/archinte.158.16.1789)] [Medline: [9738608](https://pubmed.ncbi.nlm.nih.gov/9738608/)]
25. Grant BF, Dawson DA, Stinson FS, Chou PS, Kay W, Pickering R. The Alcohol Use Disorder and Associated Disabilities Interview Schedule-IV (AUDADIS-IV): reliability of alcohol consumption, tobacco use, family history of depression and psychiatric diagnostic modules in a general population sample. *Drug Alcohol Depend* 2003 Jul 20;71(1):7-16. [doi: [10.1016/S0376-8716\(03\)00070-X](https://doi.org/10.1016/S0376-8716(03)00070-X)] [Medline: [12821201](https://pubmed.ncbi.nlm.nih.gov/12821201/)]
26. Grant BF, Goldstein RB, Saha TD, Chou SP, Jung J, Zhang H, et al. Epidemiology of DSM-5 alcohol use disorder: results from the national epidemiologic survey on alcohol and related conditions III. *J Am Med Assoc Psychiatry* 2015 Aug;72(8):757-766 [FREE Full text] [doi: [10.1001/jamapsychiatry.2015.0584](https://doi.org/10.1001/jamapsychiatry.2015.0584)] [Medline: [26039070](https://pubmed.ncbi.nlm.nih.gov/26039070/)]
27. Renna F. Alcohol abuse, alcoholism, and labor market outcomes: looking for the missing link. *Ind Labor Relat Rev* 2008 Oct;62(1):92-103. [doi: [10.1177/001979390806200105](https://doi.org/10.1177/001979390806200105)]
28. Lloyd DA, Turner RJ. Cumulative lifetime adversities and alcohol dependence in adolescence and young adulthood. *Drug Alcohol Depend* 2008 Mar 01;93(3):217-226 [FREE Full text] [doi: [10.1016/j.drugalcdep.2007.09.012](https://doi.org/10.1016/j.drugalcdep.2007.09.012)] [Medline: [17980975](https://pubmed.ncbi.nlm.nih.gov/17980975/)]
29. National Institute on Alcohol Abuse and Alcoholism. Alcohol use disorder URL: <https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/alcohol-use-disorders> [accessed 2018-09-23] [WebCite Cache ID 72eyt3EdY]
30. Heron KE, Smyth JM. Ecological momentary interventions: incorporating mobile technology into psychosocial and health behaviour treatments. *Br J Health Psychol* 2010 Feb;15(Pt 1):1-39 [FREE Full text] [doi: [10.1348/135910709X466063](https://doi.org/10.1348/135910709X466063)] [Medline: [19646331](https://pubmed.ncbi.nlm.nih.gov/19646331/)]
31. Morgenstern J, Kuerbis A, Muench F. Ecological momentary assessment and alcohol use disorder treatment. *Alcohol Res* 2014;36(1):101-109 [FREE Full text] [Medline: [26259004](https://pubmed.ncbi.nlm.nih.gov/26259004/)]
32. Kay M, Santos J, Takane M. World Health Organization. 2011. Mhealth: New horizons for health through mobile technologies 2011 URL: [http://apps.who.int/iris/bitstream/handle/10665/44607/9789241564250\\_eng.pdf?sequence=1&isAllowed=y](http://apps.who.int/iris/bitstream/handle/10665/44607/9789241564250_eng.pdf?sequence=1&isAllowed=y) [accessed 2018-04-25] [WebCite Cache ID 6yxBHznr3]
33. Yu P, Wu M, Yu H, Xiao G. The challenges for the adoption of m-health. : IEEE; 2006 Presented at: International Conference on Service Operations and Logistics, and Informatics; June 21-23, 2006; Shanghai, China p. 181-186. [doi: [10.1109/SOLI.2006.329059](https://doi.org/10.1109/SOLI.2006.329059)]
34. Bock BC, Barnett NP, Thind H, Rosen R, Walaska K, Traficante R, et al. A text message intervention for alcohol risk reduction among community college students: TMAP. *Addict Behav* 2016 Dec;63:107-113. [doi: [10.1016/j.addbeh.2016.07.012](https://doi.org/10.1016/j.addbeh.2016.07.012)] [Medline: [27450909](https://pubmed.ncbi.nlm.nih.gov/27450909/)]
35. Gustafson DH, McTavish FM, Chih MY, Atwood AK, Johnson RA, Boyle MG, et al. A smartphone application to support recovery from alcoholism: a randomized clinical trial. *J Am Med Assoc Psychiatry* 2014 May;71(5):566-572 [FREE Full text] [doi: [10.1001/jamapsychiatry.2013.4642](https://doi.org/10.1001/jamapsychiatry.2013.4642)] [Medline: [24671165](https://pubmed.ncbi.nlm.nih.gov/24671165/)]
36. Gonzalez VM, Dulin PL. Comparison of a smartphone app for alcohol use disorders with an internet-based intervention plus bibliotherapy: a pilot study. *J Consult Clin Psychol* 2015 Apr;83(2):335-345. [doi: [10.1037/a0038620](https://doi.org/10.1037/a0038620)] [Medline: [25622202](https://pubmed.ncbi.nlm.nih.gov/25622202/)]
37. Andersson C. Comparison of web and interactive voice response (IVR) methods for delivering brief alcohol interventions to hazardous-drinking university students: a randomized controlled trial. *Eur Addict Res* 2015;21(5):240-252. [doi: [10.1159/000381017](https://doi.org/10.1159/000381017)] [Medline: [25967070](https://pubmed.ncbi.nlm.nih.gov/25967070/)]
38. Hasin DS, Aharonovich E, O'Leary A, Greenstein E, Pavlicova M, Arunajadai S, et al. Reducing heavy drinking in HIV primary care: a randomized trial of brief intervention, with and without technological enhancement. *Addiction* 2013 Jul;108(7):1230-1240 [FREE Full text] [doi: [10.1111/add.12127](https://doi.org/10.1111/add.12127)] [Medline: [23432593](https://pubmed.ncbi.nlm.nih.gov/23432593/)]

39. Dedert EA, McDuffie JR, Stein R, McNiel JM, Kosinski AS, Freiermuth CE, et al. Electronic interventions for alcohol misuse and alcohol use disorders: a systematic review. *Ann Intern Med* 2015 Aug 4;163(3):205-214. [doi: [10.7326/M15-0285](https://doi.org/10.7326/M15-0285)] [Medline: [26237752](https://pubmed.ncbi.nlm.nih.gov/26237752/)]
40. White A, Kavanagh D, Stallman H, Klein B, Kay-Lambkin F, Proudfoot J, et al. Online alcohol interventions: a systematic review. *J Med Intern Res* 2010;12(5):e62 [FREE Full text] [doi: [10.2196/jmir.1479](https://doi.org/10.2196/jmir.1479)] [Medline: [21169175](https://pubmed.ncbi.nlm.nih.gov/21169175/)]
41. Keoleian V, Polcin D, Galloway GP. Text messaging for addiction: a review. *J Psychoactive Drugs* 2015;47(2):158-176 [FREE Full text] [doi: [10.1080/02791072.2015.1009200](https://doi.org/10.1080/02791072.2015.1009200)] [Medline: [25950596](https://pubmed.ncbi.nlm.nih.gov/25950596/)]
42. Riordan B, Flett J, Conner T, Scarf D. Text message interventions for alcohol use: Current research and future directions. In: Gutierrez W, editor. *Alcohol and Drug Abuse: Alcohol Consumption: Patterns, Influences and Health Effects*. Hauppauge, New York: Nova Science Publishers; 2016:187-191.
43. Fowler LA, Holt SL, Joshi D. Mobile technology-based interventions for adult users of alcohol: a systematic review of the literature. *Addict Behav* 2016 Nov;62:25-34. [doi: [10.1016/j.addbeh.2016.06.008](https://doi.org/10.1016/j.addbeh.2016.06.008)] [Medline: [27310031](https://pubmed.ncbi.nlm.nih.gov/27310031/)]
44. Kazemi DM, Borsari B, Levine MJ, Li S, Lamberson KA, Matta LA. A systematic review of the mHealth interventions to prevent alcohol and substance abuse. *J Health Commun* 2017 May;22(5):413-432. [doi: [10.1080/10810730.2017.1303556](https://doi.org/10.1080/10810730.2017.1303556)] [Medline: [28394729](https://pubmed.ncbi.nlm.nih.gov/28394729/)]
45. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009 Jul 21;6(7):e1000097 [FREE Full text] [doi: [10.1371/journal.pmed.1000097](https://doi.org/10.1371/journal.pmed.1000097)] [Medline: [19621072](https://pubmed.ncbi.nlm.nih.gov/19621072/)]
46. Eysenbach G, CONSORT-EHEALTH Group. CONSORT-EHEALTH: improving and standardizing evaluation reports of web-based and mobile health interventions. *J Med Intern Res* 2011;13(4):e126 [FREE Full text] [doi: [10.2196/jmir.1923](https://doi.org/10.2196/jmir.1923)] [Medline: [22209829](https://pubmed.ncbi.nlm.nih.gov/22209829/)]
47. Pluye P, Robert E, Cargo M, Bartlett G, O'Cathain A, Griffiths F, et al. Department of Family Medicine, McGill University. 2011. Proposal: A mixed methods appraisal tool for systematic mixed studies URL: <http://mixedmethodsappraisaltoolpublic.pbworks.com/w/file/attach/84371689/MMAT%202011%20criteria%20and%20tutorial%202011-06-29updated2014.08.21.pdf> [accessed 2018-04-25] [WebCite Cache ID 6yxBYcfF4]
48. Rycroft-Malone J, McCormack B, Hutchinson AM, DeCorby K, Bucknall TK, Kent B, et al. Realist synthesis: illustrating the method for implementation research. *Implement Sci* 2012 Apr 19;7:33 [FREE Full text] [doi: [10.1186/1748-5908-7-33](https://doi.org/10.1186/1748-5908-7-33)] [Medline: [22515663](https://pubmed.ncbi.nlm.nih.gov/22515663/)]
49. Elo S, Kyngäs H. The qualitative content analysis process. *J Adv Nurs* 2008 Apr;62(1):107-115. [doi: [10.1111/j.1365-2648.2007.04569.x](https://doi.org/10.1111/j.1365-2648.2007.04569.x)] [Medline: [18352969](https://pubmed.ncbi.nlm.nih.gov/18352969/)]
50. Suffoletto B, Kristan J, Chung T, Jeong K, Fabio A, Monti P, et al. An interactive text message intervention to reduce binge drinking in young adults: a randomized controlled trial with 9-month outcomes. *PLoS One* 2015;10(11):e0142877 [FREE Full text] [doi: [10.1371/journal.pone.0142877](https://doi.org/10.1371/journal.pone.0142877)] [Medline: [26580802](https://pubmed.ncbi.nlm.nih.gov/26580802/)]
51. Suffoletto B, Kristan J, Callaway C, Kim KH, Chung T, Monti PM, et al. A text message alcohol intervention for young adult emergency department patients: a randomized clinical trial. *Ann Emerg Med* 2014 Dec;64(6):664-72.e4 [FREE Full text] [doi: [10.1016/j.annemergmed.2014.06.010](https://doi.org/10.1016/j.annemergmed.2014.06.010)] [Medline: [25017822](https://pubmed.ncbi.nlm.nih.gov/25017822/)]
52. Agyapong VI, McLoughlin DM, Farren CK. Six-months outcomes of a randomised trial of supportive text messaging for depression and comorbid alcohol use disorder. *J Affect Disord* 2013 Oct;151(1):100-104. [doi: [10.1016/j.jad.2013.05.058](https://doi.org/10.1016/j.jad.2013.05.058)] [Medline: [23800443](https://pubmed.ncbi.nlm.nih.gov/23800443/)]
53. Agyapong VI, Ahern S, McLoughlin DM, Farren CK. Supportive text messaging for depression and comorbid alcohol use disorder: single-blind randomised trial. *J Affect Disord* 2012 Dec 10;141(2-3):168-176. [doi: [10.1016/j.jad.2012.02.040](https://doi.org/10.1016/j.jad.2012.02.040)] [Medline: [22464008](https://pubmed.ncbi.nlm.nih.gov/22464008/)]
54. Gajecki M, Andersson C, Rosendahl I, Sinadinovic K, Fredriksson M, Berman AH. Skills training via smartphone app for university students with excessive alcohol consumption: a randomized controlled trial. *Int J Behav Med* 2017 Dec;24(5):778-788 [FREE Full text] [doi: [10.1007/s12529-016-9629-9](https://doi.org/10.1007/s12529-016-9629-9)] [Medline: [28224445](https://pubmed.ncbi.nlm.nih.gov/28224445/)]
55. Aharonovich E, Stohl M, Cannizzaro D, Hasin D. HealthCall delivered via smartphone to reduce co-occurring drug and alcohol use in HIV-infected adults: a randomized pilot trial. *J Subst Abuse Treat* 2017 Dec;83:15-26 [FREE Full text] [doi: [10.1016/j.jsat.2017.09.013](https://doi.org/10.1016/j.jsat.2017.09.013)] [Medline: [29129192](https://pubmed.ncbi.nlm.nih.gov/29129192/)]
56. Haug S, Lucht MJ, John U, Meyer C, Schaub MP. A pilot study on the feasibility and acceptability of a text message-based aftercare treatment programme among alcohol outpatients. *Alcohol Alcohol* 2015 Mar;50(2):188-194. [doi: [10.1093/alcalc/agu107](https://doi.org/10.1093/alcalc/agu107)] [Medline: [25600249](https://pubmed.ncbi.nlm.nih.gov/25600249/)]
57. Witkiewitz K, Desai SA, Bowen S, Leigh BC, Kirouac M, Larimer ME. Development and evaluation of a mobile intervention for heavy drinking and smoking among college students. *Psychol Addict Behav* 2014 Sep;28(3):639-650. [doi: [10.1037/a0034747](https://doi.org/10.1037/a0034747)] [Medline: [25000269](https://pubmed.ncbi.nlm.nih.gov/25000269/)]
58. Lucht MJ, Hoffman L, Haug S, Meyer C, Pussehl D, Quellmalz A, et al. A surveillance tool using mobile phone short message service to reduce alcohol consumption among alcohol-dependent patients. *Alcohol Clin Exp Res* 2014 Jun;38(6):1728-1736. [doi: [10.1111/acer.12403](https://doi.org/10.1111/acer.12403)] [Medline: [24730528](https://pubmed.ncbi.nlm.nih.gov/24730528/)]
59. Alessi SM, Petry NM. A randomized study of cellphone technology to reinforce alcohol abstinence in the natural environment. *Addiction* 2013 May;108(5):900-909 [FREE Full text] [doi: [10.1111/add.12093](https://doi.org/10.1111/add.12093)] [Medline: [23279560](https://pubmed.ncbi.nlm.nih.gov/23279560/)]

60. Riordan BC, Conner TS, Flett JA, Scarf D. A text message intervention to reduce first year university students' alcohol use: a pilot experimental study. *Digit Health* 2017;3:2055207617707627 [FREE Full text] [doi: [10.1177/2055207617707627](https://doi.org/10.1177/2055207617707627)] [Medline: [29942597](https://pubmed.ncbi.nlm.nih.gov/29942597/)]
61. Brendryen H, Lund IO, Johansen AB, Riksheim M, Nesvåg S, Duckert F. Balance--a pragmatic randomized controlled trial of an online intensive self-help alcohol intervention. *Addiction* 2014 Feb;109(2):218-226. [doi: [10.1111/add.12383](https://doi.org/10.1111/add.12383)] [Medline: [24134709](https://pubmed.ncbi.nlm.nih.gov/24134709/)]
62. Mason M, Benotsch EG, Way T, Kim H, Snipes D. Text messaging to increase readiness to change alcohol use in college students. *J Prim Prev* 2014 Feb;35(1):47-52. [doi: [10.1007/s10935-013-0329-9](https://doi.org/10.1007/s10935-013-0329-9)] [Medline: [24114551](https://pubmed.ncbi.nlm.nih.gov/24114551/)]
63. Riordan BC, Conner TS, Flett JA, Scarf D. A brief orientation week ecological momentary intervention to reduce university student alcohol consumption. *J Stud Alcohol Drugs* 2015 Jul;76(4):525-529. [doi: [10.15288/jsad.2015.76.525](https://doi.org/10.15288/jsad.2015.76.525)] [Medline: [26098027](https://pubmed.ncbi.nlm.nih.gov/26098027/)]
64. Fisher W, Fisher J, Harman J. The information-motivation-behavioral skills model: A general social psychological approach to understanding and promoting health behavior. In: *Social Psychological Foundations of Health and Illness*. Hoboken, New Jersey: Wiley-Blackwell; 2003:82-106.
65. Regmi K, Kassim N, Ahmad N, Tuah N. Effectiveness of mobile apps for smoking cessation: a review. *Tob Prev Cessation* 2017 Apr 12;3(April):12 [FREE Full text] [doi: [10.18332/tpc/70088](https://doi.org/10.18332/tpc/70088)]
66. Brouwer IG. The Widmark formula for alcohol quantification. *S Afr Dent J* 2004 Nov;59(10):427-428. [Medline: [15726763](https://pubmed.ncbi.nlm.nih.gov/15726763/)]
67. Festinger L. *A Theory of Cognitive Dissonance*. Redwood, CA: Stanford University Press; 1962.
68. Gurman TA, Rubin SE, Roess AA. Effectiveness of mHealth behavior change communication interventions in developing countries: a systematic review of the literature. *J Health Commun* 2012;17(Suppl 1):82-104. [doi: [10.1080/10810730.2011.649160](https://doi.org/10.1080/10810730.2011.649160)] [Medline: [22548603](https://pubmed.ncbi.nlm.nih.gov/22548603/)]
69. Mohr DC, Schueller SM, Montague E, Burns MN, Rashidi P. The behavioral intervention technology model: an integrated conceptual and technological framework for eHealth and mHealth interventions. *J Med Intern Res* 2014;16(6):e146 [FREE Full text] [doi: [10.2196/jmir.3077](https://doi.org/10.2196/jmir.3077)] [Medline: [24905070](https://pubmed.ncbi.nlm.nih.gov/24905070/)]
70. Blow FC, Ilgen MA, Walton MA, Czynk EK, McCammon R, Chermack ST, et al. Severity of baseline alcohol use as a moderator of brief interventions in the emergency department. *Alcohol Alcohol* 2009;44(5):486-490 [FREE Full text] [doi: [10.1093/alcalc/agg031](https://doi.org/10.1093/alcalc/agg031)] [Medline: [19692345](https://pubmed.ncbi.nlm.nih.gov/19692345/)]
71. Kazemi DM, Borsari B, Levine MJ, Lamberson KA, Dooley B. REMIT: Development of a mHealth theory-based intervention to decrease heavy episodic drinking among college students. *Addict Res Theory* 2017 Dec 26;26(5):377-385 [FREE Full text] [doi: [10.1080/16066359.2017.1420783](https://doi.org/10.1080/16066359.2017.1420783)]
72. Hirschi T, Gottfredson M. Commentary: testing the general theory of crime. *J Res Crime Delinq* 2016 Aug 18;30(1):47-54. [doi: [10.1177/0022427893030001004](https://doi.org/10.1177/0022427893030001004)]
73. Perkins HW. Social norms and the prevention of alcohol misuse in collegiate contexts. *J Stud Alcohol Suppl* 2002 Mar(14):164-172. [doi: [10.15288/jsas.2002.s14.164](https://doi.org/10.15288/jsas.2002.s14.164)] [Medline: [12022722](https://pubmed.ncbi.nlm.nih.gov/12022722/)]
74. Costa PT, Terracciano A, McCrae RR. Gender differences in personality traits across cultures: robust and surprising findings. *J Pers Soc Psychol* 2001 Aug;81(2):322-331. [doi: [10.1037/0022-3514.81.2.322](https://doi.org/10.1037/0022-3514.81.2.322)] [Medline: [11519935](https://pubmed.ncbi.nlm.nih.gov/11519935/)]
75. Feingold A. Gender differences in personality: a meta-analysis. *Psychol Bull* 1994 Nov;116(3):429-456. [Medline: [7809307](https://pubmed.ncbi.nlm.nih.gov/7809307/)]
76. Platt L, Melendez-Torres GJ, O'Donnell A, Bradley J, Newbury-Birch D, Kaner E, et al. How effective are brief interventions in reducing alcohol consumption: do the setting, practitioner group and content matter? Findings from a systematic review and metaregression analysis. *BMJ Open* 2016 Dec 11;6(8):e011473 [FREE Full text] [doi: [10.1136/bmjopen-2016-011473](https://doi.org/10.1136/bmjopen-2016-011473)] [Medline: [27515753](https://pubmed.ncbi.nlm.nih.gov/27515753/)]
77. Chan SR, Torous J, Hinton L, Yellowlees P. Mobile tele-mental health: increasing applications and a move to hybrid models of care. *Healthcare (Basel)* 2014 May 06;2(2):220-233 [FREE Full text] [doi: [10.3390/healthcare2020220](https://doi.org/10.3390/healthcare2020220)] [Medline: [27429272](https://pubmed.ncbi.nlm.nih.gov/27429272/)]
78. Kirk GD, Himelhoch SS, Westergaard RP, Beckwith CG. Using mobile health technology to improve hiv care for persons living with HIV and substance abuse. *AIDS Res Treat* 2013;2013:194613 [FREE Full text] [doi: [10.1155/2013/194613](https://doi.org/10.1155/2013/194613)] [Medline: [24381751](https://pubmed.ncbi.nlm.nih.gov/24381751/)]
79. Haug S, Paz Castro R, Kowatsch T, Filler A, Dey M, Schaub MP. Efficacy of a web- and text messaging-based intervention to reduce problem drinking in adolescents: results of a cluster-randomized controlled trial. *J Consult Clin Psychol* 2017 Dec;85(2):147-159. [doi: [10.1037/ccp0000138](https://doi.org/10.1037/ccp0000138)] [Medline: [27606700](https://pubmed.ncbi.nlm.nih.gov/27606700/)]
80. Paige SR, Alber JM, Stellefson ML, Krieger JL. Missing the mark for patient engagement: mHealth literacy strategies and behavior change processes in smoking cessation apps. *Patient Educ Couns* 2018 May;101(5):951-955. [doi: [10.1016/j.pec.2017.11.006](https://doi.org/10.1016/j.pec.2017.11.006)] [Medline: [29153592](https://pubmed.ncbi.nlm.nih.gov/29153592/)]
81. Goodwin N, Nelson JA, Ackerman F, Weisskopf T. Global Development And Environment Institute, Tufts University. 2008. Consumption and the Consumer Society URL: [http://www.ase.tufts.edu/gdae/education\\_materials/modules/Consumption\\_and\\_the\\_Consumer\\_Society.pdf](http://www.ase.tufts.edu/gdae/education_materials/modules/Consumption_and_the_Consumer_Society.pdf) [accessed 2018-04-25] [WebCite Cache ID [6yxCU96D8](https://www.webcitation.org/6yxCU96D8)]
82. Garnett C, Crane D, Brown J, Kaner E, Beyer F, Muirhead C, et al. Reported theory use by digital interventions for hazardous and harmful alcohol consumption, and association with effectiveness: meta-regression. *J Med Intern Res* 2018 Feb 28;20(2):e69 [FREE Full text] [doi: [10.2196/jmir.8807](https://doi.org/10.2196/jmir.8807)] [Medline: [29490895](https://pubmed.ncbi.nlm.nih.gov/29490895/)]



83. Meredith SE, Alessi SM, Petry NM. Smartphone applications to reduce alcohol consumption and help patients with alcohol use disorder: a state-of-the-art review. *Adv Health Care Technol* 2015;1:47-54 [FREE Full text] [doi: [10.2147/AHCT.S65791](https://doi.org/10.2147/AHCT.S65791)] [Medline: [27478863](https://pubmed.ncbi.nlm.nih.gov/27478863/)]
84. Bratman M. *Intention, Plans, and Practical Reason*. Stanford, CA: Center for the Study of Language and Information; 1999.
85. West J, Mace M. Browsing as the killer app: explaining the rapid success of Apple's iPhone. *Telecomm Policy* 2010 Jun;34(5-6):270-286. [doi: [10.1016/j.telpol.2009.12.002](https://doi.org/10.1016/j.telpol.2009.12.002)]

## Abbreviations

**A-CHESS:** Alcohol-Comprehensive Health Enhancement Support System  
**AUD:** alcohol use disorder  
**AUDIT:** alcohol use disorders identification test  
**AUDIT-C:** alcohol use disorders identification test for consumption  
**BASICS-Mobile:** Brief Alcohol and Smoking Intervention for College Students via Mobile  
**BrAC:** breath alcohol concentration  
**eBAC:** estimated blood alcohol concentration  
**FAST:** fast alcohol screening test  
**IVR:** interactive voice response  
**mHealth:** mobile health  
**MMAT:** Mixed Methods Appraisal Tool  
**RSOD:** risky single-occasion drinking  
**SD:** standard drink  
**SMS:** short message services  
**UAU:** unhealthy alcohol use

*Edited by G Eysenbach; submitted 29.04.18; peer-reviewed by T Conner, M Bendtsen; comments to author 07.08.18; revised version received 02.10.18; accepted 30.10.18; published 26.01.19*

*Please cite as:*

*Song T, Qian S, Yu P*

*Mobile Health Interventions for Self-Control of Unhealthy Alcohol Use: Systematic Review*

*JMIR Mhealth Uhealth* 2019;7(1):e10899

URL: <http://mhealth.jmir.org/2019/1/e10899/>

doi: [10.2196/10899](https://doi.org/10.2196/10899)

PMID:

©Ting Song, Siyu Qian, Ping Yu. Originally published in JMIR Mhealth and Uhealth (<http://mhealth.jmir.org>), 26.01.2019. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mhealth and uhealth, is properly cited. The complete bibliographic information, a link to the original publication on <http://mhealth.jmir.org/>, as well as this copyright and license information must be included.