University of Wollongong

Research Online

Faculty of Social Sciences - Papers (Archive)

Faculty of Arts, Social Sciences & Humanities

2016

The (possibly negative) effects of physical activity on executive functions: Implications of the changing metabolic costs of brain development

Steven J. Howard

University of Wollongong, stevenh@uow.edu.au

Caylee J. Cook

University of Cape Town

Rihlat Said-Mohamed

University of Witwatersrand

Shane A. Norris

University of Witwatersrand

Catherine E. Draper

University of Cape Town

Follow this and additional works at: <https://ro.uow.edu.au/sspapers>

Part of the [Education Commons](#), and the [Social and Behavioral Sciences Commons](#)

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

The (possibly negative) effects of physical activity on executive functions: Implications of the changing metabolic costs of brain development

Abstract

Background: An area of growth in physical activity research has involved investigating effects of physical activity on children's executive functions. Many of these efforts seek to increase the energy expenditure of young children as a healthy and low-cost way to affect physical, health, and cognitive outcomes.

Methods: We review theory and research from neuroscience and evolutionary biology, which suggest that interventions seeking to increase the energy expenditure of young children must also consider the energetic trade-offs that occur to accommodate changing metabolic costs of brain development. **Results:** According to Life History Theory, and supported by recent evidence, the high relative energy-cost of early brain development requires that other energy-demanding functions of development (ie, physical growth, activity) be curtailed. This is important for interventions seeking to dramatically increase the energy expenditure of young children who have little excess energy available, with potentially negative cognitive consequences. Less energy-demanding physical activities, in contrast, may yield psychosocial and cognitive benefits while not overburdening an underweight child's already scarce energy supply.

Conclusions: While further research is required to establish the extent to which increases in energy-demanding physical activities may compromise or displace energy available for brain development, we argue that action cannot await these findings.

Keywords

physical, development, effects, (possibly, executive, functions; implications, changing, metabolic, costs, brain, negative), activity

Disciplines

Education | Social and Behavioral Sciences

Publication Details

Howard, S. J., Cook, C. J., Said-Mohamed, R., Norris, S. A. & Draper, C. E. (2016). The (possibly negative) effects of physical activity on executive functions: Implications of the changing metabolic costs of brain development. *Journal of Physical Activity and Health*, 13 (9), 1017-1022.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

The (Possibly Negative) Effects of Physical Activity on Executive Functions: Implications of
the Changing Metabolic Costs of Brain Development

Manuscript Type: Review

Keywords: nutrition; neuroscience; psychology; risks of exercise; youth

Abstract Word Count: 200 words

Manuscript Word Count (Text + References): 4,667 words

Date of Revised Manuscript Submission: 1 March 2016

Abstract

Background: An area of growth in physical activity research has involved investigating effects of physical activity on children's executive functions. Many of these efforts seek to increase the energy expenditure of young children as a healthy and low-cost way to affect physical, health, and cognitive outcomes. *Methods:* We review theory and research from neuroscience and evolutionary biology, which suggest that interventions seeking to increase the energy expenditure of young children must also consider the energetic trade-offs that occur to accommodate changing metabolic costs of brain development. *Results:* According to Life History Theory, and supported by recent evidence, the high relative energy-cost of early brain development requires that other energy-demanding functions of development (i.e., physical growth, activity) be curtailed. This is important for interventions seeking to dramatically increase the energy expenditure of young children who have little excess energy available, with potentially negative cognitive consequences. Less energy-demanding physical activities, in contrast, may yield psychosocial and cognitive benefits while not overburdening an underweight child's already scarce energy supply. *Conclusions:* While further research is required to establish the extent to which increases in energy-demanding physical activities may compromise or displace energy available for brain development, we argue that action cannot await these findings.

The (Possibly Negative) Effects of Physical Activity on Executive Functions: Implications of the Changing Metabolic Costs of Brain Development

An area of recent growth in physical activity research has involved investigating the effects of physical activity not just on physical (e.g., motor development) and health outcomes (e.g., body composition), but also cognitive outcomes in children. Particular interest has been directed toward executive functions (EFs), given mounting evidence for their relation to: (i) school readiness, academic learning, and success (e.g., literacy, numeracy, learning more broadly)^{1,2}; (ii) social and emotional outcomes (e.g., theory of mind, moral conduct)³; and (iii) developmental disorders (e.g., ADHD)⁴. EFs refer to cognitive control processes that allow individuals to activate, maintain, and manipulate information in mind (working memory), resist task-irrelevant impulses and distractions (inhibition), and flexibly shift attention between tasks, rules, or requirements (shifting). EFs thus represent an interesting target for intervention in childhood, given that early EF development appears to set the stage for a broad range of later outcomes.

Whereas recent attempts to support and enhance EFs by development researchers have often involved computerized ‘brain training’ programs (which now constitutes a more than \$1 billion industry)⁵, physical activity researchers have sought to evaluate whether, and under what conditions, engaging in physical activity might support the development of EFs. It is postulated that physical activities serve to engage and extend higher-order cognition through their physiological, morphological, and neurochemical effects on the brain.⁶ Research in this area has increasingly focused on the early years of life, as intervening in these formative years may generate a more stable and lasting change.⁷ If a cause and effect relationship exists, according to this line of reasoning, physical activity may thus represent a healthy and low-cost way to – at least indirectly – affect a range of outcomes that extend beyond those

68 already well established in the physical activity literature (e.g., cardiovascular health and
69 academic achievement).⁸⁻¹⁰

70 Research in this area, however, has found at best mixed results. Some studies have
71 shown a modest improvement in EFs after either acute aerobic exercise¹¹ or chronic aerobic
72 exercise.¹² Yet two separate meta-analyses have indicated little to no effect of aerobic activity
73 on subsequent executive functioning.^{13,14} There has been a similar lack of consistent success
74 with resistance training interventions.¹⁵ Despite these findings, promising results have been
75 shown with martial arts,¹⁶ yoga,¹⁷ and some other sports,¹⁸ which have yielded moderate
76 effects on EFs and/or EF-related abilities. On this basis, it has been suggested that *what* is
77 done during physical activity (i.e., physical activity that provides opportunity to engage and
78 challenge EFs) may be the most important factor contributing to group-level effects of
79 physical activity on EFs.^{6,19} In fact, Best states in his critical review, “cognitively-engaging
80 exercise appears to have a stronger effect than non-engaging exercise on children’s executive
81 function” (p. 331).⁶ He further asserts that efforts to understand the link between physical
82 activity and cognition are worthwhile, given the importance of physical activity for the
83 developing mind.

84 What may be overlooked in this emerging line of research, however, are the individual or
85 sub-group level effects of physical activity on EFs and cognitive development more broadly.
86 On the one hand, there are known health benefits of physical activity (e.g., obesity
87 prevention, gross motor competence, positive psychosocial and health outcomes).²⁰⁻²³ Also
88 well established are the negative consequences of unhealthy weight status^{24,25} and sedentary
89 behaviour in childhood (e.g., overweight, unfavourable psychosocial health and cognitive
90 outcomes).²⁶⁻²⁹ Yet insights from neuroscience (PET and MRI data)³⁰ and evolutionary
91 biology (Life History Theory applied to humans)³¹ suggest the need to consider the impact of

the metabolic costs of physical activity in terms of energy expenditure across different stages of life and individuals' metabolic conditions.

The Changing Metabolic Costs of Brain Development

Life History Theory (LHT), which is derived from evolutionary biology and ecology,³¹ seeks to explain the processes, timing, and evolutionary forces that shape the timing and form of life events (e.g., early development, maturation, fertility, mortality).³² More specifically, LHT proposes that survival requires the 'capture' of resources (e.g., energy) from the environment and subsequent 'allocation' of resources (e.g., energy, time) to activities that are essential to maximize fitness (i.e., reproduction and survival). Given that resources available to an individual are finite (often termed an 'energy budget' in the case of energy resources),³² allocation of energy thus must involve trade-offs with other energy-expenditure pursuits. This notion of functional trade-offs to accommodate our finite energy budget is supported by evidence that energy expenditure scales closely to body mass,³³ which suggests a ceiling on the typical energy expenditure for a given body size. While evidence shows humans to have a low energy budget relative to their body mass index, compared to other placental animals, it is notable that this overall energy budget appears to be largely unaffected by physical activity levels or energy input³⁴ (and instead may relate to increased investment in brain structures).

LHT further posits that the allocation of energy and time changes with life history stages (in infancy more energy is allocated to physical and brain growth, whereas in adolescence the energy is more often allocated to reproduction) and circumstances (i.e., ecological factors, health, and social context).^{35,36} In humans, LHT suggests that resources can be allocated to three fundamental functions: maintenance, growth, and reproduction.³⁶ According to the principle of allocation, trade-offs are expected between these functions such that greater resource allocation to one function depletes the resources available for another function.

Particularly relevant to infant, child, and adolescent development are the energetic trade-offs that must occur to ensure physical growth, brain development, the learning of complex cultural practices, and mastery of energy acquisition skills. While it is clear that the pre-adult life-stage is unusually long by primate standards,³⁷ such that a sizeable proportion of pre-adult growth does not occur until adolescence,³⁸ less clear are the evolutionary reasons for such a protracted developmental phase.³⁰ From a life history theory perspective, it has been suggested that the high relative energy-cost of brain development requires that other energy-demanding functions of development be curtailed.^{32,39,40}

In line with this assertion, research by Kuzawa et al. suggests that, in early childhood, a down-regulation in energy devoted to physical growth (e.g., body weight) coincides with an increase in the energy demands for brain development.³⁰ Specifically, using pooled PET and MRI data the researchers found an average pattern that, beginning at around 6 months of age, steady increases in brain glucose demands were accompanied by proportional decreases in body-weight growth. In contrast to previous suggestions of peak brain energy costs at birth,⁴¹ Kuzawa and colleagues found that the glucose demands of the brain peaked at around 4.3 years of age – accounting for approximately 65% of resting metabolic rates and 43% of daily energy requirements – which co-occurred with a low point in body-growth energy demands. After this age, gradual physical growth (i.e., height, weight) was accompanied by decreases in brain energy demands, as both converged toward adult levels.³⁰

In light of these findings, Kuzawa et al. propose that, in humans, physical growth and activity may be attenuated in the early years to increase the energy available for brain development.³⁰ For typically developing children, Kuzawa et al.’s assertion receives support from studies on primates and other non-human animals showing investment in locomotor play carried “sizeable costs” to physical growth^{42,43}, as well as longitudinal human research suggesting physical activity levels may decrease across the preschool years⁴⁴ as children

approach the peak in brain glucose demands. Reasons for this physical activity decline are likely complex and multi-factorial, however, as illustrated by evidence of a continued decline in physical activity after brain energy demands begin to decrease.⁴⁵ Nevertheless, this decrease in physical activity levels coinciding with increasing brain glucose demands suggests a potential compensatory relationship. If true, attenuation of physical-activity-based energy expenditure in the early years may be especially pronounced in children who are already energy-deficient.

Evidence suggesting a more pronounced decline in physical activity levels amongst malnourished children is mixed, however, and is exacerbated by the different aspects of malnourishment that have been used to characterize children in these studies (e.g. underweight, stunting, wasting). Furthermore, some studies have compared undernourished and adequately nourished children in terms of physical activity *abilities*, such as gross motor skills and fitness, and have often found more similarities than differences. For instance, a study in India with children (5-10 years old) with chronic protein energy malnutrition found that malnourished and adequately nourished children were comparable in their motor speed and coordination.⁴⁶ Similarly, a South African study showed similarities in fitness between undernourished and adequately nourished children (7-14 years), with undernourished children actually performing better on some of the parameters, including a 1600m run.⁴⁷

In contrast, studies that have found associations between malnutrition and free-living physical activity *levels* have more commonly found that underweight children are less likely to engage in physical activity⁴⁸ and that stunted children have lower levels of physical activity and activity-based energy expenditure.^{49,50} Although studies have not always found this decrement in physical activity levels in undernourished and stunted children,⁵¹⁻⁵³ these contradictory findings could be a product of differing home contexts (e.g., differences in free living or habitual physical activity, such as household chores or activities). While this issue

is currently speculative and undoubtedly requires further study, such as whether, how and under what conditions the patterns identified by Kuzawa et al.³⁰ might result in individual-level trends, it can be argued that action cannot await these findings. That is, if at least some of this decrement in early childhood physical activity is due to the brain's increased energy requirements, this suggests that interventions seeking to dramatically increase the energy expenditure of young children who have little excess energy available (e.g., underweight or undernourished children) could have negative, rather than positive, cognitive consequences.

Physical Activity and Malnutrition

Undernutrition includes underweight (weight-for-age < -2SD), stunting (height-for-age < -2SD), wasting (weight-for-height < -2SD), and thinness (BMI-for-age < -2SD).⁵⁴ Global prevalence of underweight and stunting in young children has decreased since 1990 (from 25% and 40% to 16% and 26%, respectively, in 2011).⁵⁵ Yet these statistics highlight that undernourished children should also be considered when planning early physical activity interventions, particularly in developing nations (where 17.4% of young children are estimated as underweight), and to a lesser extent in developed nations (in which 2.4% of young children are estimated to be underweight).⁵⁵ That is, if the functional energetic trade-off hypothesis is accurate, there may be potential cognitive repercussions of interventions that aim to significantly increase energy expenditure in children with already low 'energy budgets' (underweight, undernourished), which may divert energy away from brain development functions. This would be particularly problematic given that many of these energy-deficient children are found in disadvantaged circumstances, with already impaired cognitive development (including EFs) as a result of chronic nutrition deficits in early life.^{46,56,57}

This hypothesis does not suggest that physical activity should be curtailed in the early years or that efforts to increase physical activity in this age group are not worthwhile. Rather,

the logical extension of these ideas would suggest that these efforts should carefully consider the unique (and possibly different) needs of the children being targeted. To illustrate, low- and middle-income countries including South Africa,⁵⁸ India,⁵⁹ and Brazil⁶⁰ face a double burden of over- and under-nutrition. These burdens can occur within the same community, as both under- and over-nutrition are associated with social and economic inequalities.^{58,61} In settings such as these, it is therefore essential to promote physical activity as a means of preventing and/or managing overweight and obesity in early childhood and later life. At the same time, it is also important to understand the potential physiological consequences of promoting dramatic energy expenditure increases with early childhood populations in which under-nutrition is more prevalent.

Thus far most early childhood physical activity interventions have not differentiated their approach based on weight status. Instead, studies have tended to introduce a general increase in children's energy expenditure and examine its associated group-level physical benefits.⁶²⁻⁶⁸ Further, in such studies participants' weight status (underweight, normal weight, overweight, obese) is rarely reported, thus making it difficult to estimate the impacts of the intervention on underweight and undernourished children. Instead, participants' body composition is more commonly measured as just one component of an outcome measure such as body mass index (BMI),^{64,69,70} weight-for-height z-score,⁶⁴ or is used as a variable to evaluate initial group equivalence.^{62,63,71} The one identified study reporting weight status of the preschool children involved in a physical activity intervention included the expected proportion of children classified as underweight (2.2%).⁷² It therefore can be expected that other studies may have also included children that have differing 'energy budgets' from their normal and overweight peers, yet the intervention has uniformly sought to increase their energy expenditure through physical activity.

This appears to be an under-investigated issue, however, with no literature found on physical activity interventions specifically targeting undernourished children and/or looking at the *cognitive effects* of increasing physical activity for undernourished, young children. That is, of the studies that do collect cognitive data from children (generally older) after a physical activity intervention,^{11,73,74} few report BMI, and none that could be found looked at the cognitive effects of physical activity for undernourished children. It is important to note that the evidence-based energetic trade-off hypothesis *does not* suggest that physical activity should be discouraged or curtailed in underweight children, or that attempts to promote physical activity in these groups should be avoided. Rather, in such cases physical activity may still play an important role in early development. Yet, according to this hypothesis, the means of intervention must be carefully considered and the approach taken may necessarily require a departure from simply increasing the energy expenditure of these children through energy-demanding physical activity. Less energy-demanding physical activities—for example, those that emphasize gross motor development—may yield important psychosocial⁷⁵⁻⁷⁷ and cognitive benefits,⁷⁸ while not overburdening an underweight child's already scarce supply of energy. Where possible, it may be beneficial for these initiatives to also be accompanied by nutrition interventions for children with low energy budgets, considering the negative effect of poor nutrition on cognitive development.^{46,57}

Conclusions and Future Directions

The implications of the changing metabolic costs of brain development—specifically the down-regulation of energy for physical growth and concomitant up-regulation of energy allocated to brain development in the early years—is an important but under-investigated area. For instance, further research is required to establish the extent to which increases in energy-demanding physical activities may compromise or displace the energy available for human brain development and the extent to which this manifests at the individual level. For instance,

the relatively low energy expenditure of humans relative to body mass³⁴ could be posited as a potential mechanism that could at least partially buffer against this energetic trade-off issue. This possibility appears unlikely, however, in light of available human and primate research. Research with Assamese macaques is at least suggestive in this regard, showing that energy-intensive physical activity may reduce the energy available for physical growth (contrary to previous theorizing that only surplus energy would be used).⁴² Alternatively, rather than a direct cause and effect relationship, there may be a set of common factors that negatively impact early physical activity and cognitive development. These may include poor- or under-nutrition, psychosocial stressors, or other factors associated with socioeconomic deprivation.^{79,80} As such, future research should seek to further understand how changing metabolic costs, brain development, and physical activity interact over the course of development (and especially in the preschool years). This may involve research investigating the association between body composition (e.g., lean and fat mass) and aspects of physical activity (e.g., intensity, type, dose, patterns) in settings in transition where malnutrition is more prevalent. While there has begun empirical research into these questions in disadvantaged areas of South Africa, early presentation of the energetic trade-off hypothesis serves as: (a) a theoretical basis from which to contextualize this empirical data; (b) a prompt for physical activity researchers to consider these hypotheses in their research, in advance of these initial empirical findings; and (c) a stimulus for international evaluations of this hypothesis using existing and new datasets (to expedite replications that otherwise would await publication of initial empirical data). Even in the absence of this empirical work, there is a plausible hypothesis that suggests physical activity initiatives in the early years should consider the varied energy budgets and nutrition needs of children to optimize both group- and individual-level outcomes.

264 **Funding**

265 Support for this paper was provided by the DST/NRF Centre of Excellence in Human

266 Development.

References

1. Blair C, Razza RP. Relating effortful control, executive function, and false belief understanding to emerging math and literacy ability in kindergarten. *Child Dev.* 2007;78:647–663.
2. Bull R, Espy KA, Wiebe SA. Short-term memory, working memory, and executive functioning in preschoolers: Longitudinal predictors of mathematical achievement at age 7 years. *Dev Neuropsychol.* 2008;33:205–228.
3. Riggs NR, Jahromi LB, Razza RP, Dillworth-Bart JE, Müeller U. Executive function and the promotion of social-emotional competence. *J Appl Dev Psychol.* 2006;27:300–309.
4. Casey BJ, Tottenham N, Fossella J. Clinical, imaging, lesion, and genetic approaches toward a model of cognitive control. *Dev Psychobiol.* 2002;40:237–254.
5. Hayden EC. Treating schizophrenia: Game on. *Nature.* 2012;483:24-26.
6. Best JR. Effects of physical activity on children’s executive function: Contributions of experimental research on aerobic exercise. *Dev Rev.* 2010;30:331-351.
7. Wass SV, Scerif G, Johnson MH. Training attentional control and working memory – is younger better? *Dev Rev.* 2012;32:360-387.
8. Donnelly JE, Lambourne K. Classroom-based physical activity, cognition, and academic achievement. *Prev Med.* 2011;52:S36-S42.
9. Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phy.* 2010;7:1-16.
10. Strong WB, Malina RM, Blimkie CJR, et al. Evidence based physical activity for school-age youth. *J Paediatr.* 2005;146:732-737.
11. Hillman CH, Pontifex MB, Raine LB, Castelli DM, Hall EE, Kramer AF. The effect of acute treadmill walking on cognitive control and academic achievement in preadolescent children. *Neurosci.* 2009;159:1044–1054.

- 292 12. Davis CL, Tomporowski PD, Boyle CA, et al. Effects of aerobic exercise on overweight
293 children's cognitive functioning: A randomized controlled trial. *Res Q Exercise Sport*.
294 2007;78:510-519.
- 295 13. Angevaren M, Aufdemkampe G, Verhaar HJ, Aleman A, Vanhees L. Physical activity
296 and enhanced fitness to improve cognitive function in older people without known
297 cognitive impairment. *Cochrane Db*. 2008;3:CD005381.
- 298 14. Smith PJ, Blumenthal JA, Hoffman BM, et al. *Psychosom Med*. 2010;72: 239-252.
- 299 15. Liu-Ambrose T, Nagamatsu LS, Graf P, Beattie L, Ashe MC, Handy TC. Resistance
300 training and executive functions: A 12-month randomized controlled trial. *Arch Intern*
301 *Med*. 2010;170:170-178.
- 302 16. Lakes KD, Hoyt WT. Promoting self-regulation through school-based martial arts
303 training. *J Appl Dev Psychol*. 2004;25:283-302.
- 304 17. Manjunath NK, Telles S. Improved performance in the Tower of London test following
305 yoga. *Indian J Physiol Pharm*. 2001;45:351-354.
- 306 18. Davis CL, Tomporowski PD, McDowell JE, et al. Exercise improves executive function
307 and achievement and alters brain activation in overweight children: A randomized,
308 controlled trial. *Health Psychol*. 2011;30:91-98.
- 309 19. Diamond A. Effects of physical exercise on executive functions: Going beyond simply
310 moving to moving with thought. *Ann Sport Med Res*. 2015;2:1011-1015.
- 311 20. Martinez-Gomez D, Tucker J, Heelan KA, Welk GJ, Eisenmann JC. Associations
312 between sedentary behavior and blood pressure in young children. *Arch Pediatr*
313 *Adolescent Med*. 2009;163:724-730.
- 314 21. Saakslähti A, Numminen P, Varstala V, et al. Physical activity as a preventive measure
315 for coronary heart disease risk factors in early childhood. *Scand J Med Sci Spor*.
316 2004;14:143-149.

22. te Velde SJ, van Nassau F, Uijtdewilligen L. Energy balance-related behaviours associated with overweight and obesity in preschool children: A systematic review of prospective studies. *Obes Rev.* 2012 ;13: 56-74.
23. Timmons BW, LeBlanc AG, Carson V, et al. Systematic review of physical activity and health in the early years (aged 0-4 years). *Appl Physiol Nutr Me.* 2012;37:773-792.
24. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: Public-health crisis, common sense cure. *Lancet.* 2002;360: 473-482.
25. Lobstein T, Baur L, Uauy R. Obesity in children and young people: A crisis in public health. *Obes Rev.* 2004;5:4-85.
26. Byun W, Dowda M, Pate RR. Correlates of objectively measured sedentary behavior in US preschool children. *Pediatrics.* 2011;128:937-945.
27. Carson V, Kuzik N, Hunter S, et al. Systematic review of sedentary behavior and cognitive development in early childhood. *Prev Med.* 2015;78:115-122.
28. Kuhl ES, Clifford LM, Stark LJ. Obesity in preschoolers; Behavioral correlates and directions for treatment. *Obesity.* 2012;20:3-29.
29. LeBlanc AG, Spence JC, Carson V, et al. Systematic review of sedentary behaviour and health indicators in the early years (aged 0-4 years). *Appl Physiol Nutr Me.* 2012;37:753-772.
30. Kuzawa CW, Chugani HT, Grossman LI, Lipovich L, Hof PR, Wildman DE. Metabolic costs and evolutionary implications of human brain development. *PNAS.* 2014;111:13010-13015.
31. Gadgil M, Bossert WH. Life historical consequences of natural selection. *American Naturalist.* 1970;104:1-24.
32. Kaplan HS, Gangestad SW. *Life History Theory and evolutionary psychology.* In: Buss DM, ed. *Handbook of Evolutionary Psychology.* New York: Wiley; 2005:68-95.

33. Keliber M. Body size and metabolism. *Hilgardia*. 1932;6:315-353.
34. Pontzer H, Raichlen DA, Gordon AD, et al. Primate energy expenditure and life history. *PNAS*. 2014;111;1433-1437.
35. Bogin B, Silva MIV, Rios L. Life history trade-offs in human growth: Adaptation or pathology? *Am J Hum Biol*. 2007;19:631-642.
36. Hochberg Z, Albertsson-Wikland K. Evo-devo of infantile and childhood growth. *Pediatr Res*. 2008;64:2-7.
37. Hill K. Life history theory and evolutionary anthropology. *Evol Anthropol*. 1993;2:78–88.
38. Walker R, Hill K, Burger O, Hurtado AM. Life in the slow lane revisited: Ontogenic separation between Chimpanzees and humans. *Am J Phys Anthropol*. 2006;129:577-583.
39. Leonard WR, Robertson, ML. Nutritional requirements and human evolution: A bioenergetics model. *Am J Hum Biol*. 1992;4:179-195.
40. Navarette A, van Schaik CP, Isler K. Energetics and the evolution of human brain size. *Nature*. 2011;480:91-93.
41. Fonseca-Azevedo K, Herculano-Houzel S. Metabolic constraint imposes tradeoff between body and number of brain neurons in human evolution. *PNAS*. 2012;109:18571-18576.
42. Berghänel A, Schülke O, Ostner J. Locomotor play drives motor skill acquisition at the expense of growth: A life history trade-off. *Sci Advances*. 2015;1:e1500451.
43. Théoret-Gosselin R, Hamel S, Côté S. The role of maternal behavior and offspring development in the survival of mountain goat kids. *Oecologia*. 2015;178;175-186.
44. Taylor RW, Williams SM, Farmer VL, Taylor BJ. Changes in physical activity over time in young children: A longitudinal study using accelerometers. *PLOS One*. 2013;8:e81567.

45. Trost SG, Pate RR, Sallis JF, 2002. Age and gender differences in objectively measured physical activity in youth. *Med Sci Sport Exer.* 2013;34:350-355.
46. Kar BR, Rao SL, Chandramouli BA. Cognitive development in children with chronic protein energy malnutrition. *Behav Brain Functions.* 2008;4:31-42.
47. Monyeki MA, Kemper HCG, Koppes LLJ, Twisk JWR. Body composition and physical fitness of undernourished South African rural primary school children. *J Clin Nutr.* 2005;59:877-883.
48. Jones S, Hendricks S, Draper CE. Assessment of physical activity and sedentary behavior at preschools in Cape Town, South Africa. *Child Obes.* 2014;10:501-510.
49. Said-Mohamed R, Bernard JY, Ndzana A-C, Pasquet P. Is overweight in stunted preschool children in Cameroon related to reductions in fat oxidation, resting energy expenditure and physical activity? *PLoS ONE.* 2012;7:e39007–39009.
50. Wilson HJ, Dickinson F, Hoffman DJ, Griffiths PL, Bogin B, Varela-Silva MI. Fat free mass explains the relationship between stunting and energy expenditure in urban Mexican Maya children. *Ann Hum Biol.* 2012;39:432–439.
51. Casale D, Desmond C, Richter L. The association between stunting and psychosocial development among preschool children: A study using the South African Birth to Twenty cohort data. *Child Care Hlth Dev.* 2014;40:900–910.
52. Mamabolo RL, Kruger HS, Lennox A, et al. Habitual physical activity and body composition of black township adolescents residing in the North West Province, South Africa. *Public Health Nutr.* 2007;10:1047–1056.
53. Prista A, Maia JAR, Damasceno A, Beunen G. Anthropometric indicators of nutritional status: Implications for fitness, activity, and health in school-age children and adolescents from Maputo, Mozambique. *Am J Clin Nutr.* 2003;77:952-959.
54. World Health Organization. *WHO child growth standards: length/height-for-age,*

weight-for-age, weight-for-length, weight-for-height and body mass index-for-age:

methods and development. Geneva: WHO; 2006.

55. United Nations Children's Fund, World Health Organization, The World Bank.

UNICEF-WHO-World Bank joint child malnutrition estimates. Geneva: WHO; 2012.

56. Miranda MC, Nobrega FJ, Sato K, Pompeia S, Sinnes EG, Bueno OFA.

Neuropsychology and malnutrition: A study with 7 to 10 years-old children in a poor

community. *Rev Bras Mat Infant.* 2007;7:45-54.

57. Morgan KE. The cognitive effects of chronic malnutrition and environment on working

memory and executive function in children. *Independent Study Project ISP Collection.*

Paper 2053; 2015. Available from: http://digitalcollections.sit.edu/isp_collection/2053

58. Monyeki MA, Awotidebe A, Strydom GL, de Riddler JH, Mamabolo RL, Kemper HCG.

The challenges of underweight and overweight in South African children: Are we

winning or losing the battle? A systematic review. *Int J Environ Res Public Health.*

2015;12:1156-1173.

59. Subramanian SV, Perkins JM, Khan KT. Do burdens of underweight and overweight

coexist among lower socioeconomic groups in India? *Am J Clin Nutr.* 2009;90:369–376.

60. Amorim PRS, de Faria RC, Byrne NM, Hills AP. Physical activity and nutritional status

of Brazilian children of low socioeconomic status: Undernutrition and overweight. *Asia*

Pac J Clin Nutr. 2006;15:217-223.

61. Tzioumis E, Adair LS. Childhood dual burden of under- and over-nutrition in low- and

middle-income countries: A critical review. *Food Nutr Bull.* 2015;35:230-243.

62. Alhassan S, Sirard JR, Robinson TN. The effects of increasing outdoor play time on

physical activity in Latino preschool children. *Int J Pediatr Obes.* 2007;2:153-158.

63. Alhassan S, Nwaokelemeh O, Mendoza A, Shitole S, Whitt-Glover MC, Yancey AK. Design and baseline characteristics of the Short bouts of Exercise for Preschoolers (STEP) study. *BMC Public Health*. 2012;12:582-590.
64. Fitzgibbon ML, Stolley MR, Schiffer L, van Horn L, KauferChristoffel K, Dyer A. Hip-hop to Health Jr. for Latino preschool children. *Obesity*. 2006;14:1616-1625.
65. Harvey-Berino J, Rourke J. Obesity prevention in preschool Native-American children: A pilot study using home visiting. *Obes Res*. 2003;11:606-611.
66. Bayer O, von Kries R, Strauss A, et al. Short- and mid-term effects of a setting based prevention program to reduce obesity risk factors in children: A cluster-randomized trial. *Clin Nutr*. 2009;28:122-128.
67. De Bock F, Genser B, Raat H, Fischer JE, Renz-Polster H. A participatory physical activity intervention in preschools: A cluster randomized controlled trial. *Am J Prev Med*. 2013;45:64-74.
68. Trost SG, Fees B, Dzewaltowski D. Feasibility and efficacy of a “Move and Learn” physical activity curriculum in preschool children. *J Phys Activ Health*. 2008;5:88-103.
69. Eliakim A, Nemet D, Balakirski Y, Epstein Y. The effects of nutritional-physical activity school-based intervention on fatness and fitness in preschool children. *J Pediatr Endocrinol Metab*. 2007;6:711-718.
70. Hannon JC, Brown BB. Increasing preschoolers’ physical activity intensities: An activity-friendly preschool playground intervention. *Prev Med*. 2008;46:532-536.
71. Reilly JJ, Kelly L, Montgomery C, Williamson A, et al. Physical activity to prevent obesity in young children: Cluster randomised controlled trial. *BMJ*. 2006;333:1041-1045.

- 438 72. Bellows LL, Davies PL, Anderson J, Kennedy C. Effectiveness of a physical activity
439 intervention for Head Start preschoolers: A randomized intervention study. *Am J Occup*
440 *Ther.* 2012;67:28-36.
- 441 73. Best JR. Exergaming immediately enhances children's executive function. *Dev Psychol.*
442 2011;48:1501-1510.
- 443 74. Chang Y-K, Liu S, Yu H-H, Lee Y-H. Effect of acute exercise on executive function in
444 children with attention deficit hyperactivity disorder. *Arch Clin Neuropsychol.*
445 2012;27:225-237.
- 446 75. Piek JP, Barrett NC, Smith LM, Rigoli D, Gasson N. Do motor skills in infancy and
447 early childhood predict anxious and depressive symptomology at school age? *Hum*
448 *Movement Sci.* 2010;29:777-786.
- 449 76. Piek JP, Baynam GB, Barrett NC. The relationship between fine and gross motor ability,
450 self-perceptions and self-worth in children and adolescents. *Hum Movement Sci.*
451 2006;25:65-75.
- 452 77. Skinner RA, Piek JP. Psychosocial implications of poor motor coordination in children
453 and adolescents. *Hum Movement Sci.* 2001;20:73-94.
- 454 78. Piek JP, Dawson L, Smith LM, Gasson N. The role of early fine and gross motor
455 development on later motor and cognitive ability. *Hum Movement Sci.* 2008;27:668-681.
- 456 79. Blair C, Granger DA, Willoughby M, et al. Salivary cortisol mediates effects of poverty
457 and parenting on executive functions in early childhood. *Child Dev.* 2011;82: 1970-1984.
- 458 80. Blair C, Raver CC. Child development in the context of adversity: Experiential
459 canalization of brain and behavior. *Am Psychol.* 2012;67:309-318.