



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Social Sciences - Papers

Faculty of Social Sciences

2013

Sociodemographic and health-related predictors of self-reported mammogram, faecal occult blood test and prostate specific antigen test use in a large Australian study

Marianne Weber
Cancer Council, NSW

Michelle Cunich
University of Sydney

David Smith
Cancer Council, NSW

Glenn P. Salkeld
University of Wollongong, gsalkeld@uow.edu.au

Freddy Sitas
Cancer Council, NSW

See next page for additional authors

Publication Details

Weber, M. F., Cunich, M., Smith, D. P., Salkeld, G., Sitas, F. & O'Connell, D. (2013). Sociodemographic and health-related predictors of self-reported mammogram, faecal occult blood test and prostate specific antigen test use in a large Australian study. *BMC Public Health*, 13 (1), 429-1-429-13.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Sociodemographic and health-related predictors of self-reported mammogram, faecal occult blood test and prostate specific antigen test use in a large Australian study

Abstract

Background: While several studies have examined factors that influence the use of breast screening mammography, faecal occult blood tests (FOBT) for bowel cancer screening and prostate specific antigen (PSA) tests for prostate disease in Australia, research directly comparing the use of these tests is sparse. We examined sociodemographic and health-related factors associated with the use of these tests in the previous two years either alone or in combination. **Methods.** Cross-sectional analysis of self-reported questionnaire data from 96,711 women and 82,648 men aged 50 or over in The 45 and Up Study in NSW (2006-2010).

Results: 5.9% of men had a FOBT alone, 44.9% had a PSA test alone, 18.7% had both tests, and 30.6% had neither test. 3.2% of women had a FOBT alone, 56.0% had a mammogram alone, 16.2% had both and 24.7% had neither test. Among men, age and socioeconomic factors were largely associated with having both FOBT and PSA tests. PSA testing alone was largely associated with age, family history of prostate cancer, health insurance status and visiting a doctor. Among women, age, use of hormone replacement therapy (HRT), health insurance status, family history of breast cancer, being retired and not having a disability were associated with both FOBT and mammograms. Mammography use alone was largely associated with age, use of HRT and family history of breast cancer. FOBT use alone among men was associated with high income, living in regional areas and being fully-retired and among women, being fully-retired or sick/disabled.

Conclusions: These results add to the literature on sociodemographic discrepancies related to cancer screening uptake and highlight the fact that many people are being screened for one cancer when they could be screened for two.

Disciplines

Education | Social and Behavioral Sciences

Publication Details

Weber, M. F., Cunich, M., Smith, D. P., Salkeld, G., Sitas, F. & O'Connell, D. (2013). Sociodemographic and health-related predictors of self-reported mammogram, faecal occult blood test and prostate specific antigen test use in a large Australian study. *BMC Public Health*, 13 (1), 429-1-429-13.

Authors

Marianne Weber, Michelle Cunich, David Smith, Glenn P. Salkeld, Freddy Sitas, and Dianne O'Connell

RESEARCH ARTICLE

Open Access

Sociodemographic and health-related predictors of self-reported mammogram, faecal occult blood test and prostate specific antigen test use in a large Australian study

Marianne F Weber^{1*}, Michelle Cunich², David P Smith¹, Glenn Salkeld², Freddy Sitas¹ and Dianne O'Connell^{1,2,3,4}

Abstract

Background: While several studies have examined factors that influence the use of breast screening mammography, faecal occult blood tests (FOBT) for bowel cancer screening and prostate specific antigen (PSA) tests for prostate disease in Australia, research directly comparing the use of these tests is sparse. We examined sociodemographic and health-related factors associated with the use of these tests in the previous two years either alone or in combination.

Methods: Cross-sectional analysis of self-reported questionnaire data from 96,711 women and 82,648 men aged 50 or over in The 45 and Up Study in NSW (2006–2010).

Results: 5.9% of men had a FOBT alone, 44.9% had a PSA test alone, 18.7% had both tests, and 30.6% had neither test. 3.2% of women had a FOBT alone, 56.0% had a mammogram alone, 16.2% had both and 24.7% had neither test. Among men, age and socioeconomic factors were largely associated with having both FOBT and PSA tests. PSA testing alone was largely associated with age, family history of prostate cancer, health insurance status and visiting a doctor. Among women, age, use of hormone replacement therapy (HRT), health insurance status, family history of breast cancer, being retired and not having a disability were associated with both FOBT and mammograms. Mammography use alone was largely associated with age, use of HRT and family history of breast cancer. FOBT use alone among men was associated with high income, living in regional areas and being fully-retired and among women, being fully-retired or sick/disabled.

Conclusions: These results add to the literature on sociodemographic discrepancies related to cancer screening uptake and highlight the fact that many people are being screened for one cancer when they could be screened for two.

Keywords: Cancer screening, Mammography, Faecal occult blood test, Prostate specific antigen test, Sociodemographic characteristics, Socioeconomic status

Background

Prostate, breast and bowel cancer are the most commonly diagnosed cancers in Australia and apart from lung cancer, also account for the highest number of cancer-related deaths [1]. There is clear evidence that early detection of breast and bowel cancer via screening reduces the mortality associated with these diseases [2-4] and Australia has national, government-funded screening programs for both

these cancers (breast since 1992 and bowel since 2006). In contrast, although a mortality benefit has been found for the early detection of prostate cancer using prostate specific antigen (PSA) testing [5], this form of screening is not recommended by the Australian Government because the harms outweigh the benefits [5,6]. However, PSA tests have been subsidised by the Australian Government since 1989 and PSA testing for the early detection of prostate cancer has received widespread media attention in recent years [7]. Thus, PSA testing is effectively used as a *de facto*, albeit disorganised, prostate cancer screening program [8].

* Correspondence: marianne@nswcc.org.au

¹Cancer Research Division, Cancer Council NSW, PO Box 572, Kings Cross, Sydney, NSW 1340, Australia

Full list of author information is available at the end of the article

Among Australian women, participation in the breast cancer screening program is around 55% (in the target age group of 50–69 years) [9]. However, screening mammograms are also commonly accessed through the private sector on a user-pays basis and so the participation rate is more likely to be around 75% [10,11]. In the bowel cancer screening program the participation rate among those invited is around 38.4% (currently only people turning 50, 55 and 65 are invited via a mailed, faecal occult blood test; FOBT) [12]. FOBTs are also available outside the program, but are not widely used and so the overall participation rate in the target age group remains similar to the program participation rate [10]. The prevalence of PSA testing for prostate cancer screening purposes in Australia is unknown but estimated to be quite high. A nationally representative study ten years ago found that 63% of men aged >50 years had ever had a PSA test [13] and this proportion is likely to have risen significantly since then [8,14].

While there have been a few recent studies examining factors that influence mammography, FOBT and PSA test use in Australia [11,14-19], research comparing the use of these different cancer screening tests is sparse. Several studies in the USA have directly compared factors associated with being screened with more than one of these test types [20-25] but only one study (limited to men) has been reported from Australia [26]. These studies have all found that many people are screened for breast and prostate cancer but not for bowel cancer. Although the test types are very different in nature (FOBT is self-administered, mammography is an imaging technique done in a specialist clinic and PSA tests are blood tests ordered by a general practitioner), screening for one cancer type may potentially be a “teachable moment” for screening for another cancer type [27-29].

The objective of our study was to examine factors related to the use of FOBT, mammography and PSA tests either alone or in combination. These factors included individual-level socio-economic indicators as well as demographic characteristics and health-related factors. We used self-reported, cross-sectional data from The 45 and Up Study in New South Wales (NSW; Australia's most populous state) from participants aged 50 years and over, who reported that they had never had cancer. In a previous report, we found that 45 and Up Study participants were more than twice as likely to have a FOBT for bowel cancer if they had also had a mammogram or a PSA test [30]. By investigating the use of a range of cancer screening tests within the same population, commonalities and discrepancies in the determinants of cancer screening uptake can be more clearly identified.

Methods

Study sample

The 45 and Up Study is a population-based cohort study of people aged 45 and over in NSW [31]. The cohort was established with the aim of providing reliable evidence to inform health policy to support Australia's healthy ageing population. Participants were randomly sampled from the Medicare Australia database, Australia's universal health insurance system, which includes all citizens and permanent residents of Australia, some temporary residents and refugees. People aged 80 years and over and residents of regional areas were oversampled by a factor of two. Participants completed a mailed self-administered questionnaire and consent form. The participation rate was 18%, however The 45 and Up study sample has excellent heterogeneity and is reasonably representative of the NSW population; has a response rate comparable to similar studies internationally and in Australia; and is among the most representative large scale cohort studies in the world [32]. This paper uses the baseline cross-sectional data from 232,056 people aged 50 and over who completed the questionnaire between January 2006 and February 2010 (77% completed the questionnaire in 2008). We chose the lower age limit of 50 years because screening for bowel and breast cancer is not recommended for people younger than 50 if they are at normal risk, as is PSA testing for prostate cancer in some Australian guidelines (e.g. the Urological Society of Australia and New Zealand [33]).

Participants who reported ever having had cancer of any type (except non-melanoma skin cancer) were excluded from analysis because many of these individuals would have undergone more frequent surveillance than those without cancer, $n = 39,897$ (17%; 20,502 men and 19,395 women). An additional 7,003 men and 5,797 women were excluded because they failed to provide sufficient information on screening (6.6%). This left 82,648 men and 96,711 women for the main analyses ($n = 179,359$).

The 45 and Up Study was approved by the University of New South Wales Human Research Ethics Committee and the Cancer Council New South Wales Ethics Committee.

Ascertainment of screening use

A self-reported history of bowel cancer screening was ascertained from the question, “Have you ever been screened for colorectal (bowel) cancer? If yes, please indicate which test(s) you had”. We restricted our results to FOBT use rather than colonoscopy or sigmoidoscopy because the National Bowel Cancer Screening Program uses FOBT tests, and colonoscopy/sigmoidoscopy can be used as a diagnostic test as well as a screening test. Breast cancer screening was ascertained from the question “Have you ever been for a breast screening mammogram?”. A history of PSA testing was ascertained from

the question, "Have you ever had a blood test ordered by your doctor to check for prostate disease? (PSA test)". We were not able to distinguish men who had a PSA test for prostate cancer screening from those who may have had a PSA test to investigate disease. However, approximately two thirds of the PSA tests administered Australia-wide in 2008 were for screening [34]. Additionally, we excluded men with a history of prostate cancer, which would have eliminated most men who had PSA tests to monitor disease and we compared the results from the full dataset with a sample of men that excluded those who reported ever having an enlarged prostate (16%). For all tests, respondents were asked to indicate how long ago (in years) they had used each test type and the analysis focussed on tests received in the previous two years.

Ascertainment of socioeconomic, demographic and health characteristics

All factors were obtained from the self-administered questionnaire and are listed in Tables 1 and 2. These included age, education, annual household income from all sources (note, the mean annual Australian income in 2009–2010 for people aged 45 years or older was \$50,490 [35]), married or living with a partner, language spoken at home, country of birth, need help with daily tasks due to illness or disability, psychological distress as measured by the Kessler 10 scale [36], and ever used hormone replacement therapy (HRT; women only). We also examined health insurance status, categorised as having 1) a health care concession card (subsidised care for low income earners), 2) a Department of Veterans' Affairs card (subsidised care for current and past members of the Australian defence force and their spouses), 3) private health insurance without extras (covers ambulance and hospital services only), 4) private health insurance with extras (i.e. covers dental treatments and specialist services such as physiotherapy), or 5) no concession card or health Insurance. It should be noted, however, that all Australians have free universal access to hospital treatment in public hospitals and subsidised out-of-hospital medical treatment and medications. Those with concession cards receive larger subsidies and those with private health insurance are covered (to varying degrees, depending on the policy) for treatment in private hospitals.

The questionnaire also collected information about recent treatment for a number of medical conditions (heart attack, other heart disease, high blood pressure, high blood cholesterol, blood clotting problems, asthma, osteoarthritis, thyroid problems, osteoporosis/low bone density, depression and/or anxiety) and in the absence of a single question about visits to a doctor, these items were combined to get a partial measure of contact with the health system for reasons other than screening.

Analyses

We examined the proportion of men who reported having a PSA test alone, a FOBT alone, neither test, or both tests within the past 2 years by socioeconomic, demographic and health characteristics. Similarly, we compared the proportion of women who reported having a mammogram alone, a FOBT alone, neither test or both tests within the last 2 years by socioeconomic, demographic and health characteristics.

Odds ratios and 95% confidence intervals (CI) corresponding to receiving each test type, both tests, or neither test for each socioeconomic, demographic and health factor were estimated using multinomial logistic regression with a generalised logit link function. Each model included all the factors listed in Table 1 for men and Table 2 for women. Missing values for each covariate were included in the models as a separate term to prevent loss of information (data not shown). We also analysed the data after excluding participants with missing values for any covariate.

Results

Descriptive statistics

Of the 82,648 men included in the main analyses, 5.9% reported having an FOBT alone, 44.9% reported having a PSA test alone, 18.7% reported having both tests, and 30.6% reported having neither test within the previous 2 years. Of the 96,711 women included in the main analyses, 3.2% reported having an FOBT alone, 56.0% reported having a mammogram alone, 16.2% reported having both tests, and 24.7% reported having neither test within the previous 2 years. The prevalence of test use weighted for age and region of residence among men in our sample (according to the NSW population in 2006 [37]), was 23.1% for FOBT and 62.3% for PSA tests. Among women, the weighted prevalence was 17.7% for FOBT and 68.2% for mammography. Table 1 shows the distribution of each socioeconomic, demographic and health-related characteristic by each test combination (FOBT, PSA test, both tests, or neither test) for men and Table 2 shows the distribution of each socioeconomic, demographic and health-related characteristic by each test combination (FOBT, mammogram both tests, or neither test) for women.

Multinomial logistic regression

Figure 1 shows the odds ratios for receipt of FOBT alone, PSA test alone, or both tests compared to men who reported having neither test in the previous 2 years. The model included all the factors listed in Table 1 and all factors were significantly associated with screening test use ($p < .0001$). Figure 2 shows the odds ratios for the receipt of FOBT alone, mammography alone, or both tests compared to women who had neither test in the

Table 1 Characteristics of men in the 45 and Up study who reported ever having a faecal occult blood test (FOBT), prostate specific antigen (PSA) test, both tests, or neither test in the previous 2 years

	n (%) ¹	% FOBT only	% PSA test only	% Both tests	% Neither test
Total n	82,648 (100)	5.9	44.9	18.7	30.6
Age					
50-59	32102 (38.8)	7.1	40.3	16.7	35.9
60-69	27572 (33.4)	5.6	47.6	24.0	22.8
70+	22974 (27.8)	4.8	47.7	14.9	32.5
Family history of cancer					
Bowel	9449 (11.4)	6.4	46.3	17.6	29.7
Prostate	7173 (8.7)	4.5	50.4	23.0	22.1
Bowel and Prostate	1421 (1.7)	5.4	49.6	20.7	24.3
Other Cancer	15816 (19.1)	6.4	44.3	19.6	29.7
None	48789 (59.0)	5.9	43.8	17.8	32.4
Place of residence*					
Major City	38086 (46.1)	5.2	46.0	16.4	32.4
Inner Region	28464 (34.4)	6.3	44.5	20.5	28.7
Outer Region	14498 (17.5)	7.2	42.3	21.4	29.2
Remote or Very Remote	1535 (1.9)	4.7	46.5	14.9	33.9
Highest qualification					
No school certificate or other qualification	8876 (10.7)	4.7	45.7	13.8	35.8
School or intermediate certificate	12535 (15.2)	5.3	47.4	17.5	29.9
Higher school or leaving certificate	7932 (9.6)	5.6	44.3	16.6	33.6
Trade, apprenticeship, certificate or diploma	31105 (37.6)	5.9	45.7	19.4	29.0
University degree or higher	20849 (25.2)	7.1	42.0	21.4	29.5
Employment status					
Full time work	22345 (27.0)	6.7	42.8	17.5	33.0
Part time work	5873 (7.1)	6.5	44.3	20.0	29.3
Self employed	11526 (14.0)	5.9	45.1	18.1	30.8
Partially retired	3935 (4.8)	5.9	46.1	23.4	24.6
Fully retired	32971 (39.9)	5.4	46.5	20.2	27.9
Unemployed	1529 (1.9)	6.3	37.8	9.2	46.6
Unpaid work	646 (0.8)	6.2	45.2	15.2	33.4
Looking after home/family	394 (0.5)	5.8	41.9	11.7	40.6
Sick/disabled	2282 (2.8)	4.4	44.0	11.0	40.7
Other	674 (0.8)	3.9	45.6	15.7	34.9
Income (\$AUD)[†]					
Less than 5,000	982 (1.2)	4.1	43.0	8.3	44.7
5,000-9,000	2953 (3.6)	5.2	43.8	9.2	41.8
10,000-19,000	11543 (14.0)	5.0	45.4	14.1	35.5
20,000-29,000	8545 (10.3)	5.8	46.2	18.4	29.5
30,000-39,000	7231 (8.8)	6.4	44.8	21.5	27.4
40,000-49,000	6745 (8.2)	6.0	44.2	21.7	28.2
50,000-69,000	9695 (11.7)	6.4	44.6	21.9	27.2
70,000 or more	22676 (27.4)	6.9	44.1	20.7	28.3
Prefer not to answer	10331 (12.5)	4.8	46.0	17.4	31.9
Health insurance status					
Health care concession card	14022 (17.0)	5.0	43.5	14.1	37.5
Department of Veterans' Affairs card	1832 (2.2)	5.1	42.3	12.1	40.6

Table 1 Characteristics of men in the 45 and Up study who reported ever having a faecal occult blood test (FOBT), prostate specific antigen (PSA) test, both tests, or neither test in the previous 2 years (Continued)

Private health insurance without extras	41033 (49.7)	6.2	46.6	22.0	25.3
Private health insurance with extras	12142 (14.7)	6.0	46.1	20.2	27.6
No concession card or health Insurance	12141 (14.7)	6.4	40.3	12.8	40.5
Married or living with a partner					
No	15150 (18.3)	5.7	40.9	13.4	40.0
Yes	66802 (80.8)	6.0	45.7	19.9	28.5
Non-English language spoken at home					
No	73979 (89.5)	6.1	44.8	19.7	29.3
Yes	8667 (10.5)	4.2	44.9	9.5	41.4
Been treated by a doctor in the past month[§]					
No	38102 (46.1)	6.5	41.5	18.5	33.5
Yes	35939 (43.5)	5.4	48.4	19.6	26.7
Country of birth					
Australia	59373 (71.8)	6.2	45.5	20.3	28.1
Other English speaking country	11214 (13.6)	6.4	41.2	17.8	34.6
Non-English speaking country	11222 (13.6)	4.3	45.4	11.0	39.2
Need help with daily tasks due to illness or disability					
No	76255 (92.3)	6.1	44.8	19.5	30.0
Yes	3761 (4.6)	4.3	43.8	11.9	40.0
Psychological distress Level					
Well	67398 (81.6)	6.1	44.7	19.7	29.5
Mild	3994 (4.8)	6.0	43.9	16.3	33.9
Moderate	1294 (1.6)	5.0	45.4	14.4	35.2
Severe	1243 (1.5)	5.2	41.8	12.6	40.5

[†] Missing data for each covariate has been omitted for the sake of brevity.

* Using the mean Accessibility/Remoteness Index of Australia (ARIA+).

† Pre-tax annual household income from all sources.

§Treated for heart attack, other heart disease, high blood pressure, high blood cholesterol, blood clotting problems, asthma, osteoarthritis, thyroid problems, osteoporosis/low bone density, depression, or anxiety.

previous 2 years. The model included all the factors listed in Table 2 and all factors were significantly associated with screening test use ($p < .0001$). In order to summarise the results presented in the figures we either highlight the factors with odds ratios (OR) of ~ 2 or more (or ~ 0.5 or less) or factors having the largest independent effects.

Overall, the factors with estimated ORs of ~ 2 or more (or ~ 0.5 or less) for the receipt of both FOBT and PSA tests among men were age (60–69 vs 50–59 years; OR = 1.94, 95% CI 1.83-2.06), annual household income less than \$9,000 (vs. \$70,000+; OR = 0.40, 95% CI 0.35-0.47) or \$5,000 (OR = 0.34, 95% CI 0.26-0.43), being fully-retired (vs. full-time employment; OR = 1.93, 95% CI 1.79-2.07) and having private health insurance (OR = 2.10, 95% CI 1.96-2.25). For having PSA tests alone, the factors with the largest independent effects were age (60–69 vs 50–59 years; OR = 1.80, 95% CI 1.72-1.89), having a family history of prostate cancer (OR = 1.64, 95% CI 1.54-1.75), having private health insurance (OR = 1.60, 95% CI 1.52-1.68) and

being treated by a doctor in the past month (OR = 1.49, 95% CI 1.44-1.55). For use of FOBTs alone among men, the factors associated with the largest independent effects were income less than \$5,000 (vs. \$70,000+; OR = 0.52, 95% CI 0.37-0.73), living in outer regional areas (vs. major city; OR = 1.55, 95% CI 1.42-1.69), and being fully-retired (vs. full-time employment; OR = 1.53, 95% CI 1.37-1.71).

Among women, the factors with ORs of ~ 2 or more (or ~ 0.5 or less) for the receipt of both FOBT and mammograms were age (70+ vs 50–59 years; OR = 0.12, 95% CI 0.11-0.13), use of HRT (OR = 2.41, 95% CI 2.29-2.52), health insurance status: private (OR = 2.38, 95% CI 2.22-2.56) or private plus extras (OR = 2.19, 95% CI 2.02-2.39), family history of breast cancer (OR = 2.05, 95% CI 1.89-2.21), being partially retired (vs. full-time employment; OR = 2.03, 95% CI 1.76-2.35) or fully retired (OR = 2.00, 95% CI 1.84-2.19), and needing help with daily tasks (OR = 0.46, 95% CI 0.41-0.52). For use of mammography alone, the factors with estimated ORs of ~ 2 or more (or ~ 0.5 or less) were age (70+ vs 50–59 years; OR = 0.16,

Table 2 Characteristics of women in the 45 and Up study who reported ever having a faecal occult blood test (FOBT), mammogram, both tests, or neither test in the previous 2 years

	Total n (%) ¹	% FOBT only	% Mammogram only	% Both tests	% Neither test
Total n	96711 (100)	3.2	56.0	16.2	24.7
Age					
50-59	42646 (44.1)	2.5	62.5	17.1	18.0
60-69	30538 (31.6)	2.0	65.4	21.7	11.0
70+	23527 (24.3)	5.9	32.0	7.3	54.7
Family history of cancer					
Bowel	11836 (12.2)	4.1	54.1	16.8	25.0
Breast	9937 (10.3)	2.3	62.5	18.1	17.1
Bowel and Breast	2077 (2.2)	3.0	58.7	17.4	20.9
Other Cancer	20273 (21.0)	2.9	57.1	17.2	22.8
None	52588 (54.4)	3.2	54.7	15.2	27.0
Hormone replacement therapy use					
Ever	40641 (42.0)	2.8	61.5	20.0	15.7
Never	53905 (55.7)	3.4	52.2	13.6	30.8
Place of residence*					
Major City	42402 (43.8)	2.9	55.8	14.3	27.0
Inner Region	34676 (35.9)	3.3	56.1	17.2	23.5
Outer Region	17697 (18.3)	3.3	55.9	18.8	22.0
Remote or Very Remote	1853 (1.9)	3.1	60.4	15.1	21.4
Highest qualification					
No school certificate or other qualification	12720 (13.2)	3.4	52.1	10.5	34.0
School or intermediate certificate	28347 (29.3)	2.9	57.0	15.7	24.5
Higher school or leaving certificate	9297 (9.6)	2.9	55.9	13.9	27.3
Trade, apprenticeship, certificate or diploma	24918 (25.8)	3.5	55.9	18.2	22.4
University degree or higher	19928 (20.6)	3.1	57.7	19.5	19.7
Employment status					
Full time work	16829 (17.4)	2.0	64.0	15.6	18.5
Part time work	16599 (17.2)	2.5	63.8	18.7	15.0
Self employed	6559 (6.8)	2.6	60.7	17.9	18.8
Partially retired	3030 (3.1)	2.9	59.3	22.1	15.8
Fully retired	39603 (41.0)	4.1	48.2	15.8	31.8
Unemployed	2088 (2.2)	3.0	54.6	9.4	33.1
Unpaid work	1952 (2.0)	2.9	59.8	17.0	20.3
Looking after home/family	5914 (6.1)	2.7	58.0	14.7	24.6
Sick/disabled	2004 (2.1)	3.8	54.9	9.2	32.0
Other	1245 (1.3)	2.8	56.0	12.4	28.8
Income (SAUD)[†]					
Less than 5,000	1878 (1.9)	3.0	52.6	10.8	33.7
5,000-9,000	4423 (4.6)	3.4	50.1	9.6	37.0
10,000-19,000	14983 (15.5)	4.3	48.1	12.1	35.6
20,000-29,000	9511 (9.8)	3.7	54.3	17.2	24.9
30,000-39,000	7593 (7.9)	3.3	58.6	18.5	19.7
40,000-49,000	6565 (6.8)	2.9	58.8	19.1	19.3
50,000-69,000	9214 (9.5)	2.7	60.4	19.3	17.6
70,000 or more	17348 (17.9)	2.3	62.5	19.6	15.7
Prefer not to answer	20589 (21.3)	3.1	57.1	16.0	23.8

Table 2 Characteristics of women in the 45 and Up study who reported ever having a faecal occult blood test (FOBT), mammogram, both tests, or neither test in the previous 2 years (Continued)

Health insurance status					
Health care concession card	18482 (19.1)	3.7	49.0	11.2	36.0
Department of Veterans' Affairs card	1095 (1.1)	4.8	32.8	6.6	55.9
Private health insurance without extras	47793 (49.4)	2.9	59.8	19.2	18.2
Private health insurance with extras	14187 (14.7)	3.2	56.2	18.0	22.6
No concession card or health Insurance	13428 (13.9)	3.1	54.8	12.1	30.1
Married or living with a partner					
No	28821 (29.8)	4.0	47.3	11.8	36.9
Yes	67599 (69.9)	2.8	59.7	18.0	19.5
Non-english language spoken at home					
No	88039 (91.0)	3.2	55.9	16.9	24.0
Yes	8671 (9.0)	2.5	56.7	8.6	32.2
Country of birth					
Australia	73198 (75.7)	3.2	56.2	17.2	23.4
Other English speaking country	11848 (12.3)	3.4	55.0	16.6	25.0
Non-English speaking country	10720 (11.1)	2.8	56.0	9.4	31.8
Been treated by a doctor in the past month[§]					
No	39055 (40.4)	2.7	58.6	17.0	21.8
Yes	49328 (51.0)	3.6	53.7	16.1	26.6
Need help with daily tasks due to illness or disability					
No	88048 (91.0)	3.1	57.3	16.9	22.8
Yes	5247 (5.4)	4.1	39.0	8.1	48.7
Psychological distress Level					
Well	74194 (76.7)	3.1	57.2	17.6	22.1
Mild	5366 (5.6)	3.2	57.9	14.1	24.8
Moderate	1706 (1.8)	2.9	53.8	13.0	30.3
Severe	1679 (1.7)	2.9	54.6	10.3	32.3

¹ Missing data for each covariate has been omitted for the sake of brevity.

* Using the mean Accessibility/Remoteness Index of Australia (ARIA+).

† Pre-tax annual household income from all sources.

§Treated for heart attack, other heart disease, high blood pressure, high blood cholesterol, blood clotting problems, asthma, osteoarthritis, thyroid problems, osteoporosis/low bone density, depression, or anxiety.

95% CI 0.15-0.17), use of HRT (OR = 2.08, 95% CI 2.01-2.17), and family history of breast cancer (OR = 1.99, 95% CI 1.87-2.12). For use of FOBTs alone among women the factor with the largest independent effect was employment status: fully retired (OR = 1.93, 95% CI 1.63-2.29) or being sick/disabled (OR = 1.93, 95% CI 1.44-2.58) compared to women in full-time employment.

The estimated odds ratios changed less than 10% when we compared the full dataset with a complete case dataset, and also when comparing the full dataset with a sample that excluded men who had ever had an enlarged prostate.

Discussion

The 45 and Up Study is the largest population-based cohort study in Australia and the results presented here, using uniform outcome measures, suggest that there are

different factors related to cancer screening depending on the type of test. Specifically, among women, having mammography alone was associated most strongly with age and well known health-related factors such as HRT use and family history of breast cancer, whereas having a FOBT alone was associated most strongly with being retired or sick/disabled. Among men, having a PSA test alone was associated most strongly with age, family history of prostate cancer, having private health insurance, and being treated by a doctor in the past month, whereas having a FOBT alone was associated most strongly with higher income, living in outer regional areas and being retired. The factors included in our analysis were factors that have been found to be associated with cancer screening in the past [11,14-19], and indeed, all these factors were significantly related to cancer screening in our study. Only one other study has directly




Figure 1 Odds ratios and 95% confidence intervals for receipt of faecal occult blood test (FOBT) alone, prostate specific antigen (PSA) test alone, or both tests for men in the previous 2 years compared to those who reported having neither test in the 45 and Up study.




Figure 2 Odds ratios and 95% confidence intervals for receipt of faecal occult blood test (FOBT) alone, mammogram alone, or both tests for women in the previous 2 years compared to those who reported having neither test in the 45 and Up study.

compared cancer screening modalities in Australia and that study, restricted to men, also found that PSA testing was more common than bowel screening and that having private health insurance, living with a partner and being white and older were associated with any screening [26].

In Australia, even though national guidelines have advocated bowel cancer screening since 1999 for asymptomatic people from the age of 50 years, a freely available program for bowel cancer screening has only been available to people turning 50 and 55 from 2006 and those turning 50, 55 and 65 from 2008. Prior to 2006, FOBTs were available from 1982 via the Rotary program "Bowelscan" which is run annually throughout most states of Australia at a nominal cost to the individual. In contrast, free population screening for breast cancer has been offered to women aged 50–69 since 1992 and has been widely advocated. PSA testing is not currently recommended as a population-based screening tool for prostate cancer, but is subsidised by the government, has had considerable media attention and is an easy test to administer. Thus, given the circumstances surrounding access to each of these tests in Australia, it is not surprising that FOBT use overall was lower than that for mammography and PSA test use and that having each test is associated with different socioeconomic and demographic factors.

A number of studies have proposed that screening for one cancer type is a potential "teachable moment" for screening for another cancer type, especially given that those having mammograms or PSA tests may be a group that are aware of, and interested in, the benefits of the early detection of cancer [27-29]. The breast cancer screening program in Australia is relatively successful (in terms of uptake and reducing breast cancer mortality [38]), and even though the evidence in favour of PSA testing as a screening tool for prostate cancer remains controversial, many men are being tested regardless. We observed in the 45 and Up Study that 72% of women had a mammogram in the previous 2 years and 63% of men had a PSA test. This suggests that bowel screening rates have the potential to increase markedly from the 22% reported here if promoted through existing health networks. Indeed, in a previous report, we found that 45 and Up Study participants were more than twice as likely to have a FOBT if they had also had a mammogram or a PSA test [30]. Currently, the cancer screening programs in Australia are not integrated. However, decision aids that provide information regarding the harms and benefits of each test side by side may increase the uptake of FOBT – especially for men, since positive messages about screening seem to be deeply ingrained for PSA testing [39].

Socioeconomic factors, especially income, had the strongest effects on being screened for two cancer types.

Specifically, those reporting low income and to a lesser degree, education, were less likely to have had two tests. The national monitoring reports of the screening programs, which are based on area-level socioeconomic classification, report lower participation for low socioeconomic sub-groups in the bowel cancer screening program but less so in the breast screening program [9,12,15]. Factors associated with PSA test use have not been widely explored, but one study in Australia found that socioeconomic factors were not strongly associated with PSA test use [18]. We found that low income groups were less likely to have had a FOBT alone, but not a mammogram alone and to a lesser degree, PSA tests alone. Educational status was related to use of FOBT in both men and women, as well as the combined use of FOBT and PSA test/mammography, but was not related to PSA test or mammogram use alone. This finding suggests that low levels of education were specifically related to low levels of bowel cancer screening. Previous studies have demonstrated that social inequality is a factor for the formation of attitudes towards screening [40,41]. For example, one study has shown that people with low socioeconomic status (SES) viewed bowel cancer screening as less beneficial and more frightening than those with higher SES [41]. Thus, when cost and access are not obvious barriers to cancer screening, encouraging people in low socioeconomic groups to participate in the bowel cancer screening program may require targeted promotional campaigns aimed at addressing negative attitudes towards cancer screening.

Interestingly, living in regional and remote areas was not a barrier to cancer screening in our study. Indeed, people in regional and remote areas were *more* likely to be screened for bowel and breast cancer than those living in a major city (there was no variation in the use of PSA tests alone by place of residence). Australia is a vast country and there are health inequalities across geographic locations, especially in areas of geographic isolation, and a number of reports have demonstrated that people in regional areas have poorer survival from cancer than those in major cities [42,43]. However, our study is in concordance with others showing that both mammogram and FOBT uptake via the national programs is higher in rural areas than in the city [12,15,44]. This suggests that geographic differences in cancer survival are possibly due to differences in access to treatments and/or to socioeconomic discrepancies in survival rather than to delayed diagnosis. Higher levels of cancer screening in regional areas could possibly be due to a greater level of community strength and engagement in regional areas, leading to a greater community awareness of cancer screening. For example, the Breast Screening Program in NSW is largely operated out of a mobile van that travels from town to town. This event is

likely to be much more salient in a small town with a single central shopping centre than in the urban sprawl of the city, and indeed, in some Australian cities the program is run entirely from clinics. Thus, the presence of the mobile van may facilitate a level of awareness of the program in rural and regional areas that may be lacking in urban areas. Moreover, limited research has shown that community strength is greater in rural Australia than in the cities [45], and higher levels of social integration are positively related to cancer screening uptake [46]. It is also possible that, in the case of bowel screening, people living in major cities are more likely to have a colonoscopy than those in regional areas where accessibility to colonoscopy clinics is more limited (although not explored here).

Those who were retired, in part-time work or in unpaid work were more likely to be screened than those in fulltime work. This result is consistent with studies showing that often people state that lack of time is a reason for not participating in bowel screening [47,48]. That is, those in fulltime work have less free time to be screened. There was very little variation in screening uptake among those who reported their employment status as sick/disabled or unemployed compared to those in fulltime work. However, both men and women who reported needing help with daily tasks because of illness or disability had low levels of screening test use. This may be explained in terms of access issues and the ability to get to a mammography clinic/doctor, or to perform a FOBT at home. Alternatively, it may be the case that these participants have more immediate health problems, possibly a shorter life expectancy, and have therefore made a reasonable decision that the small long-term benefits of screening are not relevant for them. However we also found that those who reported being treated by a doctor for a major illness in the past month were significantly more likely to have been screened than those who had not been treated in the past month. This finding is in line with previous reports showing that routine visits to a doctor are associated with cancer screening [29,40,49-51], and highlights the important role for general practitioners in cancer testing, even though they may not be responsible for administering the tests (i.e. in the case of FOBT and mammography) [48,52-56].

This study had a number of limitations. Firstly, cancer screening history was derived from self-report, however a meta-analysis of validation studies on self-reported cancer screening use in the USA found that self-reported versus documented history of screening had reasonably high sensitivity (0.78 for FOBT, 0.95 for mammography, 0.71 for PSA) and specificity (0.90 for FOBT, 0.61 for mammography, 0.73 for PSA) [57]. Secondly, we were not able to identify participants who had a PSA test for monitoring, rather than screening purposes, however by

excluding men who had ever had prostate cancer we were able to eliminate most of these. Additionally, when we excluded men who had ever had an enlarged prostate from the sample, the observed odds ratios changed by less than 10%. Finally, our results may not be representative of the general population because cohort study participants tend to be healthier and more health conscious than non-participants [58] and the participation rate was only 18%. However, like most long-term cohort studies, The 45 and Up Study is designed to provide sufficient heterogeneity for valid comparisons within the cohort, rather than specific estimates of prevalence of exposure in the population [59]. Previous reports have demonstrated that The 45 and Up Study cohort has sufficient spread in the responses to questionnaire items for internal comparisons [31] and that exposure-outcome relationships are very similar to those from a survey of a representative sample from the same population [32]. Potential bias resulting from the "healthy cohort" effect, if it is present, generally leads to more conservative results due to reduced representation of population groups who have more extreme health behaviours, such as those who are mentally or physically ill or marginalised for some other reason. Hence, the associations observed between sociodemographic characteristics and screening may be somewhat closer to the null than might be seen among comparisons of community members who do not tend to participate in studies of this type. Caution must also be exercised when interpreting any negative results.

Conclusions

Overall, this paper adds to the empirical literature on economic and social discrepancies related to cancer screening uptake. Most importantly, our results highlight the fact that many people are being screened for one cancer type when they could be screened for two. Strategies aimed at using one test as a 'teachable moment' for promoting another test may help close the gap in socio-demographic discrepancies in cancer screening to a certain extent.

Abbreviations

CI: Confidence intervals; FOBT: Faecal occult blood test; HRT: Hormone replacement therapy; NSW: New South Wales; OR: Odds ratio; PSA: Prostate specific antigen; SES: Socioeconomic status.

Competing interests

The authors have no competing interests to declare.

Authors' contributions

MW had overall responsibility for the design of this study, data management, statistical analysis and drafting the paper. MC, DS, and DO'C made substantial contributions to the conception and design of the study, interpretation of the data, and critically revised the manuscript drafts. FS and GS made substantial contributions to the conception of the study and critically revised the manuscript drafts. All authors have given final approval of this version to be published.

Acknowledgements

This research was completed using data collected through the 45 and Up Study (www.saxinstitute.org.au). The 45 and Up Study is managed by the Sax Institute in collaboration with major partner Cancer Council NSW; and partners the National Heart Foundation of Australia (NSW Division); NSW Ministry of Health; beyondblue: the national depression and anxiety initiative; Ageing, Disability and Home Care, NSW Department of Family and Community Services; the Australian Red Cross Blood Service; and UnitingCare Ageing. We thank the many thousands of people participating in the 45 and Up Study. Michelle Cunich and Glenn Salkeld were supported by the Screening and diagnostic Test Evaluation Program (STEP) funded by the National Health and Medical Research Council of Australia under program grant number 633003. Marianne Weber, David Smith, Freddy Sitas and Dianne O'Connell are funded by Cancer Council NSW. All authors had full access to all of the data (including statistical reports and tables) in this study.

Author details

¹Cancer Research Division, Cancer Council NSW, PO Box 572, Kings Cross, Sydney, NSW 1340, Australia. ²School of Public Health, The University of Sydney, Sydney, NSW 2006, Australia. ³School of Medicine and Public Health, The University of Newcastle, Newcastle, NSW 2308, Australia. ⁴School of Public Health and Community Medicine, The University of New South Wales, Kensington, NSW 2052, Australia.

Received: 31 October 2012 Accepted: 19 April 2013

Published: 3 May 2013

References

1. Australian Institute of Health and Welfare & Australasian Association of Cancer Registries 2012: *Cancer in Australia: an overview, 2012. Cancer series no. 74. Cat. no. CAN 70.* Canberra: AIHW; 2012.
2. Mandel JS, Church TR, Ederer F, Bond JH: **Colorectal cancer mortality: effectiveness of biennial screening for fecal occult blood.** *J Natl Cancer Inst* 1999, **91**:434–437.
3. Winawer SJ, Flehinger BJ, Schottenfeld D, Miller DG: **Screening for colorectal cancer with fecal occult blood testing and sigmoidoscopy.** *J Natl Cancer Inst* 1993, **85**:1311–1318.
4. Nystrom L, Andersson I, Bjurstam N, Frisell J, Nordenskjold B, Rutqvist LE: **Long-term effects of mammography screening: updated overview of the swedish randomised trials.** *Lancet* 2002, **359**:909–919.
5. Moyer VA: **Screening for prostate cancer: U.S. Preventive services task force recommendation statement.** *Ann Intern Med* 2012, **157**:120–134.
6. Australian Health Technology Advisory Committee: *Prostate cancer screening.* Canberra: Commonwealth Department of Health and Family Services; 1996.
7. MacKenzie R, Chapman S, Holding S, McGeechan K: **'A Matter of faith, not science': analysis of media coverage of prostate cancer screening in Australian news media 2003–2006.** *J R Soc Med* 2007, **100**:513–521.
8. Smith DP, Supramaniam R, Marshall VR, Armstrong BK: **Prostate cancer and prostate-specific antigen testing in New south wales.** *Med J Aust* 2008, **189**:315–318.
9. Australian Institute of Health and Welfare: *BreastScreen australia monitoring report 2009–2010. Cancer series no. 72. Cat. no. CAN 68.* Canberra: AIHW; 2012.
10. Department of Health: *Victorian population health survey.* Melbourne, Victoria: Department of Health; 2008.
11. Birch S, Haas M, Savage E, Van Gool K: **Targeting services to reduce social inequalities in utilisation: an analysis of breast cancer screening in New south wales.** *Aust New Zealand Health Policy* 2007, **4**:12.
12. Australian Institute of Health and Welfare: *National bowel cancer screening program monitoring report: phase 2, July 2008–June 2011. Cancer series no. 65. Cat. no. CAN 61.* Canberra: AIHW; 2012.
13. Holden CA, McLachlan RI, Pitts M, Cumming R, Wittert G, Agius PA, Handelsman DJ, de Kretser DM: **Men in Australia telephone survey (MATEs): a national survey of the reproductive health and concerns of middle-aged and older Australian men.** *Lancet* 2005, **366**:218–224.
14. Baade PD, Youlten DR, Coory MD, Gardiner RA, Chambers SK: **Urban–rural differences in prostate cancer outcomes in Australia: what has changed?** *Medical Journal of Australia* 2011, **194**:293–296.
15. Ward PR, Javanparast S, Matt MA, Martini A, Tsourtos G, Cole S, Gill T, Aylward P, Baratin G, Jiwa M, et al: **Equity of colorectal cancer screening: cross-sectional analysis of national bowel cancer screening program data for south Australia.** *Aust N Z J Public Health* 2011, **35**:61–65.
16. Achat H, Close G, Taylor R: **Who has regular mammograms? effects of knowledge, beliefs, socioeconomic status, and health-related factors.** *Prev Med* 2005, **41**:312–320.
17. Gregory TA, Wilson C, Duncan A, Turnbull D, Cole SR, Young G: **Demographic, social cognitive and social ecological predictors of intention and participation in screening for colorectal cancer.** *BMC Public Health* 2011, **11**:38.
18. Weller D, Pinnock C, Silagy C, Hiller JE, Marshall VR: **Prostate cancer testing in SA men: influence of sociodemographic factors, health beliefs and LUTS.** *Aust N Z J Public Health* 1998, **22**:400–402.
19. Cockburn J, Paul C, Tzelepis F, McElduff P, Byles J: **Screening for bowel cancer among NSW adults with varying levels of risk: a community survey.** *Aust N Z J Public Health* 2002, **26**:236–241.
20. Swan J, Breen N, Graubard BI, McNeel TS, Blackman D, Tangka FK, Ballard-Barbash R: **Data and trends in cancer screening in the united states: results from the 2005 national health interview survey.** *Cancer* 2010, **116**:4872–4881.
21. Carlos RC, Underwood W 3rd, Fendrick AM, Bernstein SJ: **Behavioral associations between prostate and colon cancer screening.** *J Am Coll Surg* 2005, **200**:216–223.
22. Carlos RC, Fendrick AM, Patterson SK, Bernstein SJ: **Associations in breast and colon cancer screening behavior in women.** *Acad Radiol* 2005, **12**:451–458.
23. Carlos RC, Fendrick AM, Ellis J, Bernstein SJ: **Can breast and cervical cancer screening visits be used to enhance colorectal cancer screening?** *J Am Coll Radiol* 2004, **1**:769–776.
24. Swan J, Breen N, Coates RJ, Rimer BK, Lee NC: **Progress in cancer screening practices in the united states: results from the 2000 national health interview survey.** *Cancer* 2003, **97**:1528–1540.
25. Lemon S, Zapka J, Puleo E, Luckmann R, Chasan-Taber L: **Colorectal cancer screening participation: comparisons with mammography and prostate-specific antigen screening.** *Am J Public Health* 2001, **91**:1264–1272.
26. Carriere P, Baade P, Newman B, Aitken J, Janda M: **Cancer screening in Queensland men.** *Med J Aust* 2007, **186**:404–407.
27. Molina L, Carlos RC, Fendrick AM, Abrahamse PH, Dong Q, Bernstein SJ: **Completion of colorectal cancer screening in women attending screening mammography.** *Acad Radiol* 2004, **11**:1237–1241.
28. Red SN, Kassin EC, Williams RM, Penek S, Lynch J, Ahaghotu C, Taylor KL: **Underuse of colorectal cancer screening among men screened for prostate cancer: a teachable moment?** *Cancer* 2010, **116**:4703–4710.
29. Wong YN, Coups EJ: **Correlates of colorectal cancer screening adherence among Men Who have been screened for prostate cancer.** *J Cancer Educ* 2011, **26**:301–307.
30. Weber MF, Banks E, Ward R, Sitas F: **Population characteristics related to colorectal cancer testing in New south wales, Australia: results from the 45 and Up study cohort.** *J Med Screen* 2008, **15**:137–142.
31. Banks E, Redman S, Jorm L, Armstrong B, Bauman A, Beard J, Beral V, Byles J, Corbett S, Cumming R, et al: **Cohort profile: the 45 and up study.** *Int J Epidemiol* 2008, **37**:941–947.
32. Mealing NM, Banks E, Jorm LR, Steel DG, Clements MS, Rogers KD: **Investigation of relative risk estimates from studies of the same population with contrasting response rates and designs.** *BMC Med Res Methodol* 2010, **10**:26.
33. *The early detection of prostate cancer in general practice: supporting patient choice.* <http://www.usanz.org.au/uploads/29168/ufiles/6%20%20PSA%20decision%20card%20041007.pdf>.
34. *Medical benefits schedule statistics.* <http://www.medicareaustralia.gov.au/provider/medicare/mbs.jsp#N1003F>.
35. Australian Bureau of Statistics: *Household income and income distribution 2009–10. Cat No. 6523.0.* Canberra: ABS; 2011.
36. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand SLT, Walters EE, Zaslavsky AM: **Short screening scales to monitor population prevalences and trends in non-specific psychological distress.** *Psychol Med* 2002, **32**:959–976.
37. Australian Bureau of Statistics: *Census of population and housing.* Canberra: Commonwealth of Australia; 2006.
38. Nickson C, Mason KE, English DR, Kavanagh AM: **Mammographic screening and breast cancer mortality: a case–control study and meta-analysis.** *Cancer Epidemiol Biomarkers Prev* 2012, **21**:1479–1488.
39. Gattellari M, Ward JE: **A community-based randomised controlled trial of three different educational resources for men about prostate cancer screening.** *Patient Educ Couns* 2005, **57**:168–182.

40. Cullati S, Charvet-Berard AI, Perneger TV: **Cancer screening in a middle-aged general population: factors associated with practices and attitudes.** *BMC Public Health* 2009, **9**:118.
41. Wardle J, McCaffery K, Nadel M, Atkin W: **Socioeconomic differences in cancer screening participation: comparing cognitive and psychosocial explanations.** *Soc Sci Med* 2004, **59**:249–261.
42. Australian Institute of Health and Welfare and Australasian Association of Cancer Registries (AACR): *Cancer survival in Australia 1992–1997: geographic categories and socioeconomic status.* AIHW cat. no. CAN 17. Canberra: Australian Institute of Health and Welfare (Cancer Series no. 22); 2003.
43. Yu XQ, O'Connell DL, Gibberd RW, Smith DP, Armstrong BK: *Cancer survival, incidence and mortality by area health service in NSW 1994 to 2000.* Sydney: Cancer Council NSW; 2003.
44. O'Byrne AM, Kavanagh AM, Ugoni A, Diver F: **Predictors of non-attendance for second round mammography in an Australian mammographic screening programme.** *J Med Screen* 2000, **7**:190–194.
45. *Change and adjustment in rural Victoria. Presentation to the 13th ministerial rural and regional health forum, 17 August 2007.* http://www.health.vic.gov.au/ruralhealth/downloads/phillips_dpi.pdf.
46. Ye J, Williams SD, Xu Z: **The association between social networks and colorectal cancer screening in American males and females: data from the 2005 health information national trends survey.** *Cancer Causes & Control* : CCC 2009, **20**:1227–1233.
47. Forbes GM, Fritschi L, Mendelson RM, Foster NM, Edwards JT: **Influences on participation in a community-based colorectal neoplasia screening program by virtual colonoscopy in Australia.** *Aust N Z J Public Health* 2004, **28**:283–287.
48. Tong S, Hughes K, Oldenburg BB, Mar CD: **Colorectal cancer screening with faecal occult blood testing: community intention, knowledge, beliefs and behaviour.** *Asia Pac J Public Health* 2006, **18**:16–23.
49. Ioannou GN, Chapko MK, Dornitz JA: **Predictors of colorectal cancer screening participation in the United States.** *Am J Gastroenterol* 2003, **98**:2082–2091.
50. Ross LE, Taylor YJ, Howard DL: **Trends in prostate-specific antigen test use, 2000–2005.** *Public Health Rep* 2011, **126**:228–239.
51. Taylor R, Ivanov O, Page A, Brotherton J, Achat H, Close G: **Predictors of non-attendance from BreastScreen NSW in women who report current mammography screening.** *Aust N Z J Public Health* 2003, **27**:581–587.
52. Janda M, Stanton WR, Hughes K, Del Mar C, Clavarino A, Aitken JF, Tong S, Short L, Leggett B, Newman B: **Knowledge, attitude and intentions related to colorectal cancer screening using faecal occult blood tests in a rural Australian population.** *Asia Pac J Public Health* 2003, **15**:50–56.
53. Page A, Morrell S, Tewson R, Taylor R, Brassil A: **Mammography screening participation: effects of a media campaign targeting Italian-speaking women.** *Aust N Z J Public Health* 2005, **29**:365–371.
54. Slevin TJ, Donnelly N, Clarkson JP, English DR, Ward JE: **Prostate cancer testing: behaviour, motivation and attitudes among western Australian men.** *Med J Aust* 1999, **171**:185–188.
55. Cole SR, Young GP, Byrne D, Guy JR, Morcom J: **Participation in screening for colorectal cancer based on a faecal occult blood test is improved by endorsement by the primary care practitioner.** *J Med Screen* 2002, **9**:147–152.
56. Salkeld GP, Solomon MJ, Short L, Ward J: **Measuring the importance of attributes that influence consumer attitudes to colorectal cancer screening.** *ANZ J Surg* 2003, **73**:128–132.
57. Rauscher GH, Johnson TP, Cho YI, Walk JA: **Accuracy of self-reported cancer-screening histories: a meta-analysis.** *Cancer Epidemiol Biomarkers Prev* 2008, **17**:748–757.
58. Breslow NE, Day N: *Statistical methods in cancer research. Volume II- The design and analysis of cohort studies.* IARC scientific publication No 82. International Agency for Research on Cancer: Lyon, France; 1987.
59. Willett WC, Blot WJ, Colditz GA, Folsom AR, Henderson BE, Stampfer MJ: **Merging and emerging cohorts: not worth the wait.** *Nature* 2007, **445**:257–258.

doi:10.1186/1471-2458-13-429

Cite this article as: Weber et al.: Sociodemographic and health-related predictors of self-reported mammogram, faecal occult blood test and prostate specific antigen test use in a large Australian study. *BMC Public Health* 2013 **13**:429.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

