

PRODUCT SYSTEMS OVER RIGHT-ANGLED ARTIN SEMIGROUPS

NEAL J. FOWLER AND AIDAN SIMS

ABSTRACT. We build upon Mac Lane’s definition of a tensor category to introduce the concept of a product system that takes values in a tensor groupoid \mathcal{G} . We show that the existing notions of product systems fit into our categorical framework, as do the k -graphs of Kumjian and Pask. We then specialize to product systems over right-angled Artin semigroups; these are semigroups that interpolate between free semigroups and free abelian semigroups. For such a semigroup we characterize all product systems which take values in a given tensor groupoid \mathcal{G} . In particular, we obtain necessary and sufficient conditions under which a collection of k 1-graphs form the coordinate graphs of a k -graph.

INTRODUCTION

Product systems were introduced by Arveson in his study of one-parameter semigroups of endomorphisms ([1]). Very roughly, a product system is a family $(E_t)_{t>0}$ of complex Hilbert spaces that is endowed with an associative multiplication such that, for every $s, t > 0$, there is a unitary isomorphism $E_s \otimes E_t \rightarrow E_{s+t}$ which maps the elementary tensor $x \otimes y$ to the product xy . The first discrete analogues of these were studied by Dinh in [3], where the parameter t was constrained to take values in the positive cone of a countable dense subgroup of \mathbb{R} . Product systems over arbitrary semigroups were introduced by Fowler and Raeburn in [8], and the first author has continued this line of investigation in [5] and [6]. Although the papers cited above all focus on the C^* -algebras associated with product systems, here our interest is purely in the algebraic structure of product systems.

This note was inspired by two recent developments. In [7], the notion of a discrete product system was extended to allow for fibers that are right-Hilbert bimodules over a C^* -algebra, thus opening connections with Pimsner’s generalized Cuntz algebras ([15]). Second, in [12] Kumjian and Pask developed the notion of k -graphs, and these have much in common with product systems over the semigroup \mathbb{N}^k . Our first goal is thus to generalize the definition of a product system to encompass these different algebraic structures. We do this in Section 1 by extending Mac Lane’s definition of a monoidal category [13, §VII.1] to that of a *tensor groupoid* \mathcal{G} , and by developing the notion of a product system that takes values in \mathcal{G} . In addition to recovering as product systems the algebraic structures mentioned above, by considering an abelian group G as a tensor groupoid we also obtain as a product system every 2-cocycle of the underlying semigroup S that takes values in G . This suggests that, at least for some tensor groupoids, the set of all product

Received by the editors December 22, 1999 and, in revised form, June 28, 2001.
1991 *Mathematics Subject Classification*. Primary 20F36; Secondary 18B40, 55N20.

©2001 American Mathematical Society

systems over a semigroup S should possess a natural binary operation akin to the multiplication of 2-cocycles in $Z^2(S; G)$. At the end of Section 1 we show that this is indeed the case: if \mathcal{G} is a tensor groupoid which is symmetric in the sense of [13, §XI.1], then one can form the internal tensor product of two product systems over S that take values in \mathcal{G} . The internal tensor product is associative and well-defined on isomorphism classes of product systems, so these isomorphism classes have the natural structure of a semigroup; we denote this semigroup $H^2(S; \mathcal{G})$. Although this notation suggests the existence of a cohomology theory, at this point we have been unable to place product systems in such a framework.

Section 2 is devoted to constructing and classifying product systems over right-angled Artin semigroups; these are semigroups which interpolate between free semigroups and free abelian semigroups. For a given right-angled Artin semigroup P with generating set A , our main result, Theorem 2.1, gives conditions on an A -tuple $(X_a)_{a \in A}$ of objects in \mathcal{G} that allow one to construct a product system over P whose fiber over $a \in A$ is X_a . In Theorem 2.2 we show that our conditions are necessary, and that every product system over P is obtained by our construction. In particular our results allow us to construct and classify k -graphs in terms of the coordinate 1-graphs which generate them; see Remark 2.3. In Proposition 2.8 we use our parameterization to determine when two product systems over P are isomorphic, and in Corollary 2.10 we determine the automorphism group of any product system over P . We close with a description of the semigroup $H^2(P; \mathcal{G})$ when \mathcal{G} is a symmetric tensor groupoid (Proposition 2.11); when \mathcal{G} is an abelian group G , this gives a computation of the second cohomology group $H^2(P; G)$ (Corollary 2.12).

The authors would like to thank the referee for pointing out the connection between our work and Mac Lane’s. The first author would also like to thank Alex Kumjian for several helpful discussions on k -graphs.


1. TENSOR GROUPOIDS AND PRODUCT SYSTEMS

Let \mathcal{G} be a groupoid, regarded as a small category with inverses. We will write $X \in \mathcal{G}$ to denote that X is an object in \mathcal{G} , and $S \in \text{Hom}(X_1, X_2)$ or $S: X_1 \rightarrow X_2$ to denote that S is a morphism from X_1 to X_2 .

We will assume that \mathcal{G} is endowed with the structure of a (relaxed) monoidal category, in the sense of [13, §VII.1]. Thus \mathcal{G} is part of a sextuple $(\mathcal{G}, \otimes, 1_{\mathcal{G}}, \mathcal{B}, \lambda, \rho)$ in which \otimes is a bifunctor $\otimes: \mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$, $1_{\mathcal{G}}$ is a distinguished object in \mathcal{G} , and \mathcal{B} , λ and ρ are natural isomorphisms

$$\begin{aligned} \mathcal{B} &= \mathcal{B}_{X_1, X_2, X_3} : X_1 \otimes (X_2 \otimes X_3) \rightarrow (X_1 \otimes X_2) \otimes X_3, \\ \lambda &= \lambda_X : 1_{\mathcal{G}} \otimes X \rightarrow X, \quad \text{and} \quad \rho = \rho_X : X \otimes 1_{\mathcal{G}} \rightarrow X, \end{aligned}$$

such that $\rho_{1_{\mathcal{G}}} = \lambda_{1_{\mathcal{G}}} : 1_{\mathcal{G}} \otimes 1_{\mathcal{G}} \rightarrow 1_{\mathcal{G}}$, and such that the following two diagrams commute for every $X_1, X_2, X_3, X_4 \in \mathcal{G}$:



and

$$\begin{array}{ccc}
 X_1 \otimes (1_{\mathcal{G}} \otimes X_2) & \xrightarrow{\mathcal{B}_{X_1, 1_{\mathcal{G}}, X_2}} & (X_1 \otimes 1_{\mathcal{G}}) \otimes X_2 \\
 1 \otimes \lambda_{X_2} \downarrow & & \downarrow \rho_{X_1} \otimes 1 \\
 X_1 \otimes X_2 & \xrightarrow{\text{id}_{X_1} \otimes X_2} & X_1 \otimes X_2.
 \end{array}$$

By the corollaries in Sections VII.1 and VII.2 of [13], the canonical isomorphisms supplied by these natural isomorphisms allow us to write expressions such as $X_1 \otimes \cdots \otimes X_k$ without bothering to delineate the order in which adjacent factors should be tensored, and to cancel out any extra factors of $1_{\mathcal{G}}$. We shall take advantage of this notational simplification and make only occasional further references to the natural isomorphisms \mathcal{B} , λ and ρ .

We have used the symbol \otimes for our bifunctor rather than Mac Lane’s more neutral \square since our primary motivating examples truly are tensor products (see Examples 1.5 (2) and (3)). Consequently, we have chosen to expand on Mac Lane’s alternative terminology “tensor category” [13, page 252] and refer to $(\mathcal{G}, \otimes, 1_{\mathcal{G}}, \mathcal{B}, \lambda, \rho)$ (or just \mathcal{G}) as a *tensor groupoid*.

Definition 1.1. Let S be a countable semigroup, and let \mathcal{G} be a tensor groupoid. A *product system over S taking values in \mathcal{G}* is a pair (Y, α) in which Y is a collection $(Y_s)_{s \in S}$ of objects in \mathcal{G} , and α is a collection $(\alpha_{s,t})_{s,t \in S}$ of isomorphisms $\alpha_{s,t}: Y_s \otimes Y_t \rightarrow Y_{st}$ such that

$$(1.1) \quad \alpha_{r,s,t}(\alpha_{r,s} \otimes 1_{Y_t}) \mathcal{B}_{Y_r, Y_s, Y_t} = \alpha_{r,st}(1_{Y_r} \otimes \alpha_{s,t}) \quad \text{for every } r, s, t \in S.$$

If S has an identity e , we require that $Y_e = 1_{\mathcal{G}}$, and that, for each $s \in S$, $\alpha_{e,s}$ and $\alpha_{s,e}$ are implemented by λ_{Y_s} and ρ_{Y_s} , respectively.

As alluded to above we will henceforth suppress the natural equivalence \mathcal{B} . Equation (1.1) then becomes

$$(1.2) \quad \alpha_{r,s,t}(\alpha_{r,s} \otimes 1_{Y_t}) = \alpha_{r,st}(1_{Y_r} \otimes \alpha_{s,t}) \quad \text{for every } r, s, t \in S,$$

where both sides are regarded as isomorphisms from $Y_r \otimes Y_s \otimes Y_t$ to Y_{rst} ; we will write $\alpha_{r,s,t}$ for this isomorphism. More generally:

Notation 1.2. If $k \geq 2$ and $s_1, \dots, s_k \in S$, write α_{s_1, \dots, s_k} for the isomorphism

$$Y_{s_1} \otimes \cdots \otimes Y_{s_k} \rightarrow Y_{s_1 \cdots s_k}$$

obtained by repeatedly applying appropriate isomorphisms $\alpha_{s,t}$ on adjacent factors. For $s \in S$ we define $\alpha_s := 1_{Y_s}$.

A moment’s thought shows that this notation makes sense: for each way of associating the factors in $Y_{s_1} \otimes \cdots \otimes Y_{s_k}$ one can apply appropriate isomorphisms $\alpha_{s,t}$ to obtain a morphism with range $Y_{s_1 \cdots s_k}$, and a straightforward inductive argument using the naturality of \mathcal{B} shows that the canonical isomorphisms supplied by \mathcal{B} carry these morphisms into one another.

Definition 1.3. Two product systems (Y, α) and (Z, β) are *isomorphic* if there is a collection $\psi = (\psi_s)_{s \in S}$ of isomorphisms $\psi_s: Y_s \rightarrow Z_s$ such that, for every $s, t \in S$,

the following diagram commutes:

$$\begin{array}{ccc}
 Y_s \otimes Y_t & \xrightarrow{\alpha_{s,t}} & Y_{st} \\
 \psi_s \otimes \psi_t \downarrow & & \downarrow \psi_{st} \\
 Z_s \otimes Z_t & \xrightarrow{\beta_{s,t}} & Z_{st}.
 \end{array}$$

Remark 1.4. It is often useful to not require that the objects of \mathcal{G} form a set. Thus we will sometimes consider structures $\langle \mathcal{G}, \otimes, 1_{\mathcal{G}}, \mathcal{B}, \lambda, \rho \rangle$ in which \mathcal{G} is a category, all of whose morphisms are invertible. This is merely a convenience; for if (Y, α) is a product system that takes values in such a category \mathcal{G} , then (Y, α) takes values in the tensor groupoid \mathcal{G}' whose objects are all possible tensor products $Y_{s_1} \otimes \cdots \otimes Y_{s_n}$.

Examples 1.5. (1) Let G be an abelian group, considered as the morphisms of a groupoid \mathcal{G} with one object. Since G is abelian, $g \otimes h := gh$ defines a functor $\otimes: \mathcal{G}^2 \rightarrow \mathcal{G}$, and it is easy to see that this gives \mathcal{G} the structure of a tensor groupoid; the lone object in \mathcal{G} is the identity object $1_{\mathcal{G}}$, and λ and ρ are both the identity morphism on $1_{\mathcal{G}}$.

If (Y, α) is a product system over S that takes values in \mathcal{G} , then α is a 2-cocycle on S that takes values in G . Since \mathcal{G} has but one object, the map $(Y, \alpha) \mapsto \alpha$ is a bijection between such product systems and the group $Z^2(S; G)$. Moreover, cocycles are cohomologous if and only if the corresponding product systems are isomorphic, so there is a canonical bijection between the set of isomorphism classes of product systems and the cohomology group $H^2(S; G)$.

(2) The product systems considered in [3], [4], [8], [5], and [6] can be placed in our categorical framework. We will follow the convention of the latter three references and consider product systems over a monoid S , and write e for the identity element in S .

Let \mathcal{G} be the category whose objects are nontrivial separable complex Hilbert spaces, and whose morphisms are intertwining unitary isomorphisms. Let \otimes be the usual Hilbert space tensor product, let $1_{\mathcal{G}} = \mathbb{C}$, and let \mathcal{B} , λ and ρ be the natural equivalences determined by

$$(1.3) \quad \mathcal{B}_{X_1, X_2, X_3}(x_1 \otimes (x_2 \otimes x_3)) = (x_1 \otimes x_2) \otimes x_3 \quad \text{for } x_i \in X_i,$$

$$\lambda_X(z \otimes x) = zx \quad \text{and} \quad \rho_X(x \otimes z) = zx$$

for $x \in X$ and $z \in \mathbb{C}$.

Given a product system (Y, α) over S that takes values in \mathcal{G} , define

$$E := \bigsqcup_{s \in S} \{s\} \times Y_s,$$

define $p: E \rightarrow S$ by $p(s, x) := s$, and define multiplication in E by

$$(s, x)(t, y) := (st, \alpha_{s,t}(x, y)).$$

Then E is a product system over S in the sense of [8], and it is easy to see that this defines a bijective correspondence between (isomorphism classes of) the two different types of product systems.

We can replace \mathcal{G} with a tensor groupoid by limiting the number of objects. For $n \geq 1$ let \mathcal{H}_n be the Hilbert space \mathbb{C}^n , and let $\mathcal{H}_\infty := \ell^2(\mathbb{N})$. Let \mathcal{G}' be the tensor groupoid whose objects are all possible Hilbert space tensor products $\mathcal{H}_{n_1} \otimes \cdots \otimes \mathcal{H}_{n_k}$, and whose morphisms are intertwining unitary operators. Exactly

as above, every product system over S (in the sense of [8]) corresponds to a product system over S taking values in \mathcal{G}' .

(3) The product systems studied in [7] can also be placed in our categorical framework. Let A be a C^* -algebra. A right Hilbert A -module is, roughly speaking, a right A -module X_A which is endowed with an A -valued inner product. If X_A is endowed with a left action of A by adjointable operators, we call X a *right-Hilbert A - A bimodule*. (See [15] and [14] for details.)

Let \mathcal{G} be the category in which the objects are right-Hilbert A - A bimodules, and, for objects $X, Y \in \mathcal{G}$, $\text{Hom}(X, Y)$ is the set of all bimodule isomorphisms $X \rightarrow Y$ that preserve the inner product. As a tensoring functor we use the A -balanced internal tensor product (see [14] for details), and then the bimodule ${}_A A_A$ serves as the identity object $1_{\mathcal{G}}$. The natural equivalence \mathcal{B} is again given by (1.3), and λ and ρ are determined by $\lambda_X(a \otimes_A x) := a \cdot x$ and $\rho_X(x \otimes_A a) = x \cdot a$ for $x \in X$ and $a \in A$. As in the previous example, product systems that take values in this category correspond to the product systems introduced in [7].

(4) Let V be a countable set. We construct a category \mathcal{G} as follows. The objects in \mathcal{G} are triples $X = (E, r_E, s_E)$ in which E is a countable set and r_E and s_E are functions $E \rightarrow V$; we think of V and E as the vertices and edges of a directed graph, with r_E and s_E the range and source maps. We will write r and s rather than r_E and s_E when the domain is clear from context, and we somewhat imprecisely regard E as an object in \mathcal{G} . Elements of $\text{Hom}(E, E')$ are bijections $\varphi: E \rightarrow E'$ such that $s = s \circ \varphi$ and $r = r \circ \varphi$.

Define

$$E_1 \otimes E_2 := \{(f_1, f_2) \in E_1 \times E_2 : r(f_1) = s(f_2)\},$$

with range and source maps

$$r(f_1, f_2) := r(f_2) \quad \text{and} \quad s(f_1, f_2) := s(f_1).$$

Write $f_1 \otimes f_2$ for the edge $(f_1, f_2) \in E_1 \otimes E_2$. For $\varphi_1 \in \text{Hom}(E_1, E'_1)$ and $\varphi_2 \in \text{Hom}(E_2, E'_2)$, define $\varphi_1 \otimes \varphi_2 \in \text{Hom}(E_1 \otimes E_2, E'_1 \otimes E'_2)$ by

$$\varphi_1 \otimes \varphi_2(f_1 \otimes f_2) := \varphi_1(f_1) \otimes \varphi_2(f_2).$$

Then \otimes is a functor $\mathcal{G}^2 \rightarrow \mathcal{G}$. Equation (1.3) again defines a natural isomorphism \mathcal{B} between the functors $\otimes \circ (\text{id}_{\mathcal{G}} \times \otimes)$ and $\otimes \circ (\otimes \times \text{id}_{\mathcal{G}})$. We define the identity object $1_{\mathcal{G}}$ to be the triple $(V, \text{id}_V, \text{id}_V)$, and define the natural isomorphisms λ and ρ by $\lambda_E(s_E(f) \otimes f) := f$ and $\rho_E(f \otimes r_E(f)) := f$ for all $f \in E$.

When S is a monoid with no nontrivial idempotents, product systems over S that take values in this category are related to the k -graphs of Kumjian-Pask [12]. To explain the connection, we first recall the definition of a k -graph. Let Λ be (the morphisms of) a countable small category, and consider S as the morphisms of a small category with one object. A functor $d: \Lambda \rightarrow S$ is said to have the *factorization property* if for every $\lambda \in \Lambda$ and $t_1, t_2 \in S$ with $d(\lambda) = t_1 t_2$, there are unique elements $\lambda_1, \lambda_2 \in \Lambda$ such that $\lambda = \lambda_1 \lambda_2$ and $d(\lambda_1) = t_1, d(\lambda_2) = t_2$. When $S = \mathbb{N}^k$, such a pair (Λ, d) is called a *k -graph*.

Suppose Λ is the set of morphisms of a small category with object set V , and suppose $d: \Lambda \rightarrow S$ has the factorization property. We think of (Λ, d) as a generalized k -graph. For each $t \in S$, define $Y_t := d^{-1}(t)$. With range and source maps the reverse of those inherited from Λ (that is, $r = \text{dom}$ and $s = \text{cod}$), Y_t becomes an

object in \mathcal{G} . For each $t_1, t_2 \in S$, define $\alpha_{t_1, t_2} : Y_{t_1} \otimes Y_{t_2} \rightarrow Y_{t_1 t_2}$ by

$$\alpha_{t_1, t_2}(f_1 \otimes f_2) := f_1 f_2 \quad \text{for } f_1 \in Y_{t_1}, f_2 \in Y_{t_2}.$$

We claim that (Y, α) is a product system over S taking values in \mathcal{G} .

To begin with, note that $\alpha_{t_1, t_2} \in \text{Hom}(Y_{t_1} \otimes Y_{t_2}, Y_{t_1 t_2})$: each such map clearly preserves the range and source maps, and the factorization property is precisely the condition needed to ensure that α_{t_1, t_2} is bijective.

Next we will show that $Y_e = V$, where each $v \in V$ is identified with $1_v \in \Lambda$. Fix $v \in V$. Since v is an idempotent so is $d(v)$, and hence $d(v) = e$; that is, $v \in Y_e$. Now fix $\lambda \in Y_e$. Since $e^2 = e$, the factorization property assures us that there are unique elements $\lambda_1, \lambda_2 \in Y_e$ such that $\lambda = \lambda_1 \lambda_2$. Since $s(\lambda)\lambda = \lambda = \lambda r(\lambda)$, we conclude that $\lambda = s(\lambda) = r(\lambda) \in V$. Thus $Y_e = V$.

Finally, for each $t \in S$ we have

$$\alpha_{e, t}(s(\lambda) \otimes \lambda) = s(\lambda)\lambda = \lambda = \lambda_{Y_t}(s(\lambda) \otimes \lambda) \quad \text{for } \lambda \in Y_t,$$

and similarly $\alpha_{t, e}$ is implemented by ρ_{Y_t} . Thus (Y, α) is a product system over S taking values in \mathcal{G} , as claimed.

Conversely, suppose one is given a product system (Y, α) over S taking values in \mathcal{G} . Let

$$\Lambda := \bigcup_{t \in S} \{t\} \times Y_t,$$

and define $\text{dom}, \text{cod} : \Lambda \rightarrow V$ by $\text{dom}(t, f) := r(f)$ and $\text{cod}(t, f) := s(f)$. Then Λ is the set of morphisms of a countable small category with object set V , in which morphisms are composed according to

$$(t_1, f_1)(t_2, f_2) := (t_1 t_2, \alpha_{t_1, t_2}(f_1 \otimes f_2)).$$

Define $d : \Lambda \rightarrow S$ by $d(t, f) := t$. Then d is a functor, and it satisfies the factorization property because each α_{t_1, t_2} is a bijection.

The procedures outlined above are easily seen to be inverses of one another, and hence product systems over S taking values in \mathcal{G} are essentially the same as generalized k -graphs.

It should be pointed out that this example can be regarded as a special case of Example 1.5 (3), as follows. Suppose that (Y, α) is a product system over S taking values in \mathcal{G} . For each $t \in S$, let X_t be the Cuntz-Krieger bimodule associated with the directed graph Y_t , as in [9, Example 1.2]; X_t is the completion of $C_c(Y_t)$ with respect to a certain norm defined using the range map for Y_t . The embeddings $y \in Y_t \mapsto \delta_y \in X_t$ induce isomorphisms $\beta_{s, t} : X_s \otimes X_t \rightarrow X_{st}$ that make (X, β) into a product system of right-Hilbert $C_0(V)$ - $C_0(V)$ bimodules.

Symmetric tensor groupoids. We now discuss tensor groupoids which are symmetric in the sense of [13, §XI.1]. Let \mathcal{G} be a tensor groupoid and let $F : \mathcal{G}^2 \rightarrow \mathcal{G}^2$ be the “flip” functor which interchanges the order of any pair of objects or morphisms (e.g. $F(X_1, X_2) = (X_2, X_1)$). Suppose there is a natural equivalence \mathcal{F} from \otimes to $\otimes \circ F$; that is, there is a collection of isomorphisms $\mathcal{F}_{X_1, X_2} : X_1 \otimes X_2 \rightarrow X_2 \otimes X_1$ such that

$$(S_2 \otimes S_1) \circ \mathcal{F}_{X_1, X_2} = \mathcal{F}_{Y_1, Y_2} \circ (S_1 \otimes S_2)$$

for all $X_i, Y_i \in \mathcal{G}$, and all $S_i \in \text{Hom}(X_i, Y_i)$. Suppose, furthermore, that for every $X_1, X_2, X_3 \in \mathcal{G}$, the diagram

$$(1.4) \quad \begin{array}{ccc} X_1 \otimes 1_{\mathcal{G}} & \xrightarrow{\mathcal{F}_{X_1, 1_{\mathcal{G}}}} & 1_{\mathcal{G}} \otimes X_1 \\ \rho_{X_1} \downarrow & & \downarrow \lambda_{X_1} \\ X_1 & \xrightarrow{\text{id}_{X_1}} & X_1 \end{array}$$

commutes, and that the following two identities hold:

$$(1.5) \quad \mathcal{F}_{X_1, X_2}^{-1} = \mathcal{F}_{X_2, X_1}$$

and

$$(1.6) \quad \mathcal{F}_{X_1, X_2 \otimes X_3} = (1_{X_2} \otimes \mathcal{F}_{X_1, X_3})(\mathcal{F}_{X_1, X_2} \otimes 1_{X_3}).$$

Taking inverses in (1.6) and using (1.5), one also has

$$(1.7) \quad \mathcal{F}_{X_1 \otimes X_2, X_3} = (\mathcal{F}_{X_1, X_3} \otimes 1_{X_2})(1_{X_1} \otimes \mathcal{F}_{X_2, X_3}).$$

Following [13, §XI.1], we call a tensor groupoid \mathcal{G} that admits a natural equivalence \mathcal{F} with these properties a *symmetric* tensor groupoid.

One can check that this is consistent with Mac Lane’s definition of a symmetric monoidal category: our (1.4) corresponds to [13, §XI.1 (6)], (1.5) corresponds to [13, §XI.1 (8)], and (1.6) and (1.7) correspond to [13, §XI.1 (7)] with the instances of \mathcal{B} suppressed.

Examples 1.6. (1) Let \mathcal{G} be the tensor groupoid associated with an abelian group G , as in Examples 1.5(1). Then \mathcal{G} is a symmetric tensor groupoid: \mathcal{G}^2 has but one object, and assigning the identity element of G to this object gives the desired natural equivalence \mathcal{F} .

(2) Let \mathcal{G}' be the tensor groupoid introduced in Examples 1.5(2). For $X_1, X_2 \in \mathcal{G}$, define $\mathcal{F}_{X_1, X_2}: X_1 \otimes X_2 \rightarrow X_2 \otimes X_1$ by

$$\mathcal{F}_{X_1, X_2}(x_1 \otimes x_2) := x_2 \otimes x_1 \quad \text{for } x_i \in X_i.$$

Then \mathcal{F} is a natural equivalence from \otimes to $\otimes \circ F$, so \mathcal{G}' is a symmetric tensor groupoid.

In a symmetric tensor groupoid, one can take tensor products of product systems:

Proposition 1.7. *Suppose (Y, α) and (Y', α') are product systems over semigroups S and S' , respectively, both taking values in a symmetric tensor groupoid \mathcal{G} . For every $(s, s') \in S \times S'$ define*

$$Z_{(s, s')} := Y_s \otimes Y'_{s'},$$

and for every $(s, s'), (t, t') \in S \times S'$ define $\beta_{(s, s'), (t, t')}: Z_{(s, s')} \otimes Z_{(t, t')} \rightarrow Z_{(ss', tt')}$ by

$$\beta_{(s, s'), (t, t')} := (\alpha_{s, t} \otimes \alpha'_{s', t'})(1_{Y_s} \otimes \mathcal{F}_{Y'_{s'}, Y_t} \otimes 1_{Y'_{t'}}).$$

Then (Z, β) is a product system over $S \times S'$ taking values in \mathcal{G} .

Remark 1.8. We call (Z, β) the *external tensor product* of (Y, α) and (Y', α') .

Proof. We must show that β satisfies the associativity condition (1.2). Suppose $(r, r'), (s, s'), (t, t') \in S \times S'$. Then

$$\begin{aligned}
 & \beta_{(r,r'),(st,s't')}(\mathbf{1}_{Z_{(r,r')}} \otimes \beta_{(s,s'),(t,t')}) \\
 &= (\alpha_{r,st} \otimes \alpha'_{r',s't'}) (\mathbf{1}_{Y_r} \otimes \mathcal{F}_{Y'_{r'}, Y_{st}} \otimes \mathbf{1}_{Y'_{s't'}}) \\
 & \quad (\mathbf{1}_{Y_r \otimes Y'_{r'}} \otimes \alpha_{s,t} \otimes \alpha_{s',t'}) (\mathbf{1}_{Y_r \otimes Y'_{r'} \otimes Y_s} \otimes \mathcal{F}_{Y'_{s'}, Y_t} \otimes \mathbf{1}_{Y'_{t'}}) \\
 (1.8) \quad &= (\alpha_{r,st} \otimes \alpha'_{r',s't'}) (\mathbf{1}_{Y_r} \otimes \alpha_{s,t} \otimes \mathbf{1}_{Y'_{r'}} \otimes \alpha'_{s',t'}) \\
 & \quad (\mathbf{1}_{Y_r} \otimes \mathcal{F}_{Y'_{r'}, Y_s \otimes Y_t} \otimes \mathbf{1}_{Y'_{s'} \otimes Y'_{t'}}) (\mathbf{1}_{Y_r \otimes Y'_{r'} \otimes Y_s} \otimes \mathcal{F}_{Y'_{s'}, Y_t} \otimes \mathbf{1}_{Y'_{t'}}),
 \end{aligned}$$

whereas

$$\begin{aligned}
 & \beta_{(rs,r's'),(t,t')}(\beta_{(r,r'),(s,s')} \otimes \mathbf{1}_{Z_{(t,t')}}) \\
 &= (\alpha_{rs,t} \otimes \alpha'_{r',s',t'}) (\mathbf{1}_{Y_{rs}} \otimes \mathcal{F}_{Y'_{r's'}, Y_t} \otimes \mathbf{1}_{Y'_{t'}}) \\
 & \quad (\alpha_{r,s} \otimes \alpha'_{r',s'} \otimes \mathbf{1}_{Y_t \otimes Y'_{t'}}) (\mathbf{1}_{Y_r} \otimes \mathcal{F}_{Y'_{r'}, Y_s} \otimes \mathbf{1}_{Y'_{s'} \otimes Y_t \otimes Y'_{t'}}) \\
 (1.9) \quad &= (\alpha_{rs,t} \otimes \alpha'_{r',s',t'}) (\alpha_{r,s} \otimes \mathbf{1}_{Y_t} \otimes \alpha'_{r',s'} \otimes \mathbf{1}_{Y'_{t'}}) \\
 & \quad (\mathbf{1}_{Y_r \otimes Y_s} \otimes \mathcal{F}_{Y'_{r'} \otimes Y'_{s'}, Y_t} \otimes \mathbf{1}_{Y'_{t'}}) (\mathbf{1}_{Y_r} \otimes \mathcal{F}_{Y'_{r'}, Y_s} \otimes \mathbf{1}_{Y'_{s'} \otimes Y_t \otimes Y'_{t'}}).
 \end{aligned}$$

Since α and α' each satisfy (1.2), the product of the first two factors in (1.8) is equal to the corresponding product in (1.9). Hence it suffices to show that

$$(\mathcal{F}_{Y'_{r'}, Y_s \otimes Y_t} \otimes \mathbf{1}_{Y'_{s'}}) (\mathbf{1}_{Y'_{r'} \otimes Y_s} \otimes \mathcal{F}_{Y'_{s'}, Y_t}) = (\mathbf{1}_{Y_s} \otimes \mathcal{F}_{Y'_{r'} \otimes Y'_{s'}, Y_t}) (\mathcal{F}_{Y'_{r'}, Y_s} \otimes \mathbf{1}_{Y'_{s'} \otimes Y_t}).$$

By (1.6), the left-hand side of this equation is equal to

$$(\mathbf{1}_{Y_s} \otimes \mathcal{F}_{Y'_{r'}, Y_t} \otimes \mathbf{1}_{Y'_{s'}}) (\mathcal{F}_{Y'_{r'}, Y_s} \otimes \mathbf{1}_{Y_t \otimes Y'_{s'}}) (\mathbf{1}_{Y'_{r'} \otimes Y_s} \otimes \mathcal{F}_{Y'_{s'}, Y_t}),$$

and by (1.7), the right-hand side is equal to

$$(\mathbf{1}_{Y_s} \otimes \mathcal{F}_{Y'_{r'}, Y_t} \otimes \mathbf{1}_{Y'_{s'}}) (\mathbf{1}_{Y_s \otimes Y'_{r'}} \otimes \mathcal{F}_{Y'_{s'}, Y_t}) (\mathcal{F}_{Y'_{r'}, Y_s} \otimes \mathbf{1}_{Y'_{s'} \otimes Y_t}).$$

These last two expressions are obviously equal, and the proof is complete. \square

When $S = S'$, one can restrict the external tensor product to the diagonal to obtain another product system over S :

Definition 1.9. Suppose (Y, α) and (Z, β) are product systems over S taking values in a symmetric tensor groupoid \mathcal{G} . The *internal tensor product* $(Y, \alpha) \otimes (Z, \beta)$ is the product system $(Y \otimes Z, \alpha \otimes \beta)$ defined by

$$(Y \otimes Z)_s := Y_s \otimes Z_s \quad \text{for } s \in S$$

and

$$(\alpha \otimes \beta)_{s,t} := (\alpha_{s,t} \otimes \beta_{s,t}) (\mathbf{1}_{Y_s} \otimes \mathcal{F}_{Z_s, Y_t} \otimes \mathbf{1}_{Z_t}) \quad \text{for } s, t \in S.$$

Lemma 1.10. *The internal tensor product is associative and well-defined on isomorphism classes.*

Proof. Let (Y, α) , (Z, β) and (W, γ) be product systems over S taking values in \mathcal{G} , and suppose $s, t \in S$. Making frequent use of (1.6) and (1.7), we calculate

$$\begin{aligned} ((\alpha \otimes \beta) \otimes \gamma)_{s,t} &= ((\alpha \otimes \beta)_{s,t} \otimes \gamma_{s,t})(1_{Y_s \otimes Z_s} \otimes \mathcal{F}_{W_s, Y_t \otimes Z_t} \otimes 1_{W_t}) \\ &= (\alpha_{s,t} \otimes \beta_{s,t} \otimes \gamma_{s,t})(1_{Y_s} \otimes \mathcal{F}_{Z_s, Y_t} \otimes 1_{Z_t \otimes W_s \otimes W_t}) \\ &\quad (1_{Y_s \otimes Z_s \otimes Y_t} \otimes \mathcal{F}_{W_s, Z_t} \otimes 1_{W_t})(1_{Y_s \otimes Z_s} \otimes \mathcal{F}_{W_s, Y_t} \otimes 1_{Z_t \otimes W_t}) \\ &= (\alpha_{s,t} \otimes \beta_{s,t} \otimes \gamma_{s,t})(1_{Y_s \otimes Y_t \otimes Z_s} \otimes \mathcal{F}_{W_s, Z_t} \otimes 1_{W_t}) \\ &\quad (1_{Y_s} \otimes \mathcal{F}_{Z_s, Y_t} \otimes 1_{W_s \otimes Z_t \otimes W_t})(1_{Y_s \otimes Z_s} \otimes \mathcal{F}_{W_s, Y_t} \otimes 1_{Z_t \otimes W_t}) \\ &= (\alpha_{s,t} \otimes (\beta \otimes \gamma)_{s,t})(1_{Y_s} \otimes \mathcal{F}_{Z_s \otimes W_s, Y_t} \otimes 1_{Z_t \otimes W_t}) \\ &= (\alpha \otimes (\beta \otimes \gamma))_{s,t}. \end{aligned}$$

This gives associativity.

Now suppose that $(\psi_s)_{s \in S}$ is an isomorphism from (Y, α) to (Y', α') , and that $(\varphi_s)_{s \in S}$ is an isomorphism from (Z, β) to (Z', β') . For any $s, t \in S$ we have

$$\begin{aligned} (\psi_{st} \otimes \varphi_{st})(\alpha \otimes \beta)_{s,t} &= (\psi_{st} \otimes \varphi_{st})(\alpha_{s,t} \otimes \beta_{s,t})(1_{Y_s} \otimes \mathcal{F}_{Z_s, Y_t} \otimes 1_{Z_t}) \\ &= (\alpha'_{s,t} \otimes \beta'_{s,t})(\psi_s \otimes \psi_t \otimes \varphi_s \otimes \varphi_t)(1_{Y_s} \otimes \mathcal{F}_{Z_s, Y_t} \otimes 1_{Z_t}) \\ &= (\alpha'_{s,t} \otimes \beta'_{s,t})(1_{Y'_s} \otimes \mathcal{F}_{Z'_s, Y'_t} \otimes 1_{Z'_t})(\psi_s \otimes \varphi_s \otimes \psi_t \otimes \varphi_t) \\ &= (\alpha' \otimes \beta')_{s,t}(\psi_s \otimes \varphi_s \otimes \psi_t \otimes \varphi_t), \end{aligned}$$

so $(\psi_s \otimes \varphi_s)_{s \in S}$ is an isomorphism from $(Y \otimes Z, \alpha \otimes \beta)$ to $(Y' \otimes Z', \alpha' \otimes \beta')$. \square

Remark 1.11. Essentially the same proofs show that the external tensor product is also associative and well-defined on isomorphism classes of product systems.

Motivated by Examples 1.5(1), we write $Z^2(S; \mathcal{G})$ for the set of product systems over S taking values in \mathcal{G} , and $H^2(S; \mathcal{G})$ for the set of isomorphism classes of such product systems. It follows from the previous lemma that when \mathcal{G} is symmetric, the internal tensor product makes both $Z^2(S; \mathcal{G})$ and $H^2(S; \mathcal{G})$ into semigroups.

When \mathcal{G} is the symmetric tensor groupoid associated with an abelian group G (as in Examples 1.5(1) and 1.6(1)), the map $(Y, \alpha) \in Z^2(S; \mathcal{G}) \mapsto \alpha \in Z^2(S; G)$ is an isomorphism of groups, and descends to an isomorphism from $H^2(S; \mathcal{G})$ to $H^2(S; G)$. In the next section we give an explicit description of $H^2(S; \mathcal{G})$ for arbitrary \mathcal{G} in the special case when S is a right-angled Artin semigroup.

2. RIGHT-ANGLED ARTIN SEMIGROUPS

Let Γ be a (non-directed) graph with countable vertex set A . We will assume that Γ is simple; that is, that Γ has no loops (edges from a vertex to itself) or multiple edges. We write $a \leftrightarrow b$ when $a, b \in A$ are joined by an edge in Γ .

Let \mathbb{F}_A be the free group on A , and let $*_{\Gamma}\mathbb{Z}$ be the graph product of $|A|$ copies of \mathbb{Z} ; that is, $*_{\Gamma}\mathbb{Z}$ is the quotient of \mathbb{F}_A by the smallest normal subgroup that contains the commutators $[a, b]$ for which $a \leftrightarrow b$. (See [10] and [11] for details.) Since each of the factors in the graph product is \mathbb{Z} , $*_{\Gamma}\mathbb{Z}$ is a *right-angled Artin group*. Let $\pi: \mathbb{F}_A \rightarrow *_{\Gamma}\mathbb{Z}$ be the canonical quotient map.

Consider the homomorphism $\ell: \mathbb{F}_A \rightarrow \mathbb{Z}$ determined by $\ell(a) = 1$ for $a \in A$. Since every commutator $[a, b]$ belongs to the kernel of ℓ , we have $\ker \pi \subseteq \ker \ell$. Thus ℓ descends to a homomorphism $*_{\Gamma}\mathbb{Z} \rightarrow \mathbb{Z}$, also denoted ℓ , which satisfies $\ell(\pi(a)) = 1$ for each $a \in A$. We call ℓ the *length function*.

Let \mathbb{F}_A^+ be the subsemigroup of \mathbb{F}_A generated by A . Each element $\mu \in \mathbb{F}_A^+$ can be uniquely written as a word in the alphabet A ; we denote by μ_i the i^{th} element of this word, so that $\mu = \mu_1 \cdots \mu_{\ell(\mu)}$ with $\mu_1, \dots, \mu_{\ell(\mu)} \in A$.

Let P be the subsemigroup of $*_{\Gamma}\mathbb{Z}$ which is the image of \mathbb{F}_A^+ under the quotient map π . We call P a *right-angled Artin semigroup*. It is worth bearing in mind the following extreme cases: if Γ has no edges, then P is the free semigroup \mathbb{F}_A^+ , whereas if Γ is the complete graph on A , then P is free abelian.

The remainder of this note is devoted to constructing and classifying product systems over P . Our analysis makes use of a specific section $\delta: P \rightarrow \mathbb{F}_A^+$ of the quotient map π , called the *preferred section*. To define it, we fix once and for all a well-ordering of the vertex set A . (Since A is countable, this does not require the Axiom of Choice: one can simply enumerate the elements of A .) The section δ is defined recursively, starting with $\delta(\pi(a)) := a$ for each $a \in A$. Suppose δ has been defined on all words of length at most k for some $k \geq 1$. Fix $t \in P$ of length $k + 1$, and use the well-ordering of A to define

$$a := \min\{\mu_1 : \mu \in \mathbb{F}_A^+, \pi(\mu) = t\}.$$

Choose any $\mu \in \mathbb{F}_A^+$ such that $\pi(\mu) = t$ and $\mu_1 = a$, and express $\mu = \mu_1\mu'$. Then $t' := \pi(\mu')$ is independent of the choice of μ and has length k , and we define $\delta(t) := a\delta(t')$.

Now suppose \mathcal{G} is a tensor groupoid. For our first theorem, fix a collection $(X_a)_{a \in A}$ of objects in \mathcal{G} , and define

$$X_{\mu} := X_{\mu_1} \otimes \cdots \otimes X_{\mu_{\ell(\mu)}} \quad \text{for } \mu \in \mathbb{F}_A^+.$$

Write 1_{μ} for the identity morphism on X_{μ} .

Suppose $T = (T_{a,b})_{a \leftrightarrow b}$ is a collection of isomorphisms

$$T_{a,b}: X_a \otimes X_b \rightarrow X_b \otimes X_a$$

such that

$$(2.1) \quad T_{a,b}^{-1} = T_{b,a} \quad \text{whenever } a \leftrightarrow b,$$

and, whenever a, b and c form the vertices of a triangle in Γ (i.e., whenever $a \leftrightarrow b, b \leftrightarrow c$ and $c \leftrightarrow a$), the following hexagonal equation is satisfied:

$$(2.2) \quad (T_{b,c} \otimes 1_a)(1_b \otimes T_{a,c})(T_{a,b} \otimes 1_c) = (1_c \otimes T_{a,b})(T_{a,c} \otimes 1_b)(1_a \otimes T_{b,c}).$$

(Both sides of this equation are isomorphisms $X_a \otimes X_b \otimes X_c \rightarrow X_c \otimes X_b \otimes X_a$.)

Our first theorem asserts that such a collection T is all that is necessary to construct a product system over P .

Theorem 2.1. *Fix a well-ordering of A , and let $\delta: P \rightarrow \mathbb{F}_A^+$ be the corresponding preferred section of the quotient map $\pi: \mathbb{F}_A^+ \rightarrow P$. Then there is a unique product system $(Y, \alpha) = (Y^T, \alpha^T)$ over P taking values in the tensor groupoid \mathcal{G} such that*

$$(2.3) \quad Y_t = X_{\delta(t)} \quad \text{for every } t \in P,$$

$$(2.4) \quad \alpha_{s,t} = 1_{\delta(st)} \quad \text{if } \delta(st) = \delta(s)\delta(t), \text{ and}$$

$$(2.5) \quad \alpha_{\pi(a),\pi(b)} = T_{a,b} \quad \text{if } a \leftrightarrow b \text{ and } a > b.$$

Our second theorem asserts that, up to isomorphism, this construction gives all possible product systems over P . It also implies that the well-ordering used in Theorem 2.1 does not affect the isomorphism class of the resulting product system.

Theorem 2.2. *Suppose (Z, β) is a product system over P which takes values in the tensor groupoid \mathcal{G} . Define*

$$(2.6) \quad X_a := Z_{\pi(a)} \quad \text{for } a \in A$$

and

$$(2.7) \quad T_{a,b} := \beta_{\pi(b),\pi(a)}^{-1} \beta_{\pi(a),\pi(b)} \quad \text{for every } a, b \in A \text{ such that } a \leftrightarrow b.$$

Then the collection $(T_{a,b})_{a \leftrightarrow b}$ satisfies (2.1) and (2.2). Moreover, the corresponding product system (Y^T, α^T) given by Theorem 2.1 is isomorphic to (Z, β) via an isomorphism $(\psi_s)_{s \in P}$ such that, for each $a \in A$, $\psi_{\pi(a)}$ is the identity morphism on $Z_{\pi(a)}$.

Notice that if $a \leftrightarrow b$, then $\pi(a)\pi(b) = \pi(b)\pi(a)$, and the isomorphisms $\beta_{\pi(a),\pi(b)}$ and $\beta_{\pi(b),\pi(a)}$ each have range $Z_{\pi(a)\pi(b)}$. Thus the equation (2.7) used to define $T_{a,b}$ makes sense.

Remark 2.3. Before proving these theorems, we give an application to the k -graphs of Kumjian and Pask. Resume the notation of Examples 1.5(4). Let $d: \Lambda \rightarrow \mathbb{N}^k$ be a k -graph, and let $V := d^{-1}(0)$ be the set of objects in Λ . For each $t \in \mathbb{N}^k$, $Y_t := d^{-1}(t)$ is the edge set of a directed graph with vertex set V ; the range and source maps are the reverse of those inherited from Λ . Moreover, since d satisfies the factorization property, for every $t_1, t_2 \in \mathbb{N}^k$ the map $\alpha_{t_1,t_2}: Y_{t_1} \otimes Y_{t_2} \rightarrow Y_{t_1+t_2}$ defined by $\alpha_{t_1,t_2}(f_1 \otimes f_2) = f_1 f_2$ is an isomorphism; composing with α_{t_2,t_1}^{-1} we see that $Y_{t_1} \otimes Y_{t_2}$ and $Y_{t_2} \otimes Y_{t_1}$ are isomorphic.

Given a collection E_1, \dots, E_k of countable directed graphs, each with vertex set V , which satisfy

$$E_i \otimes E_j \cong E_j \otimes E_i \quad \text{for } 1 \leq i, j \leq k,$$

one might thus ask if there is a k -graph $d: \Lambda \rightarrow \mathbb{N}^k$ such that $d^{-1}(e_i)$ is isomorphic to E_i for each i . (Here $\{e_i : 1 \leq i \leq k\}$ is the canonical basis for \mathbb{N}^k .) In [12, Section 6], Kumjian and Pask observed that, when $k = 2$, any isomorphism $\theta: E_2 \otimes E_1 \rightarrow E_1 \otimes E_2$ can be used to construct such a 2-graph. Roughly, the idea is this. For each $t \in \mathbb{N}^2$ let

$$E_t := E_1^{\otimes t_1} \otimes E_2^{\otimes t_2}$$

with the usual range and source maps, and define $\Lambda := \bigsqcup_{t \in \mathbb{N}^k} E_t$. One can use θ in the obvious way to construct isomorphisms $E_s \otimes E_t \rightarrow E_{s+t}$ for every $s, t \in \mathbb{N}^2$, and the resulting binary operation on Λ makes it a small category with object set V . With $d: \Lambda \rightarrow \mathbb{N}^k$ defined by $d(f) := t$ for $f \in E_t$, (Λ, d) is a k -graph.

When $k \geq 3$, things are more complicated. For each pair (i, j) with $1 \leq i < j \leq k$, fix an isomorphism $E_j \otimes E_i \rightarrow E_i \otimes E_j$. Somewhat imprecisely, we write $f \otimes g \mapsto g' \otimes f'$ for each of these isomorphisms, and we also write $g \otimes f \mapsto f' \otimes g'$ for the inverse maps $E_i \otimes E_j \rightarrow E_j \otimes E_i$. Taking $A = \{1, \dots, k\}$ with its usual ordering, Theorems 2.1 and 2.2 say that the analogue of the construction of 2-graphs outlined above yields a k -graph if and only if the following condition holds: whenever $1 \leq i < j < l \leq k$, the composite map

$$(2.8) \quad \begin{aligned} f \otimes g \otimes h &\mapsto f \otimes h' \otimes g' \mapsto h'' \otimes f' \otimes g' \mapsto h'' \otimes g'' \otimes f'' \\ &\mapsto g''' \otimes h''' \otimes f'' \mapsto g''' \otimes f''' \otimes h'''' \mapsto f'''' \otimes g'''' \otimes h'''' \end{aligned}$$

is the identity on $E_l \otimes E_j \otimes E_i$. Moreover, up to isomorphism, every k -graph arises in this manner.

Condition (2.8) holds vacuously when $k = 2$, thus reproducing the result of Kumjian and Pask. It also holds when the vertex matrices M_i of the directed graphs E_i satisfy Robertson and Steger's conditions (H0), (H1a), (H1b) and (H1c) from [16]; that is, if the M_i 's are pairwise commuting $\{0, 1\}$ -matrices such $M_i M_j$ is a $\{0, 1\}$ -matrix whenever $i < j$, and $M_i M_j M_l$ is a $\{0, 1\}$ -matrix whenever $i < j < l$. This is easy to see: under these conditions there are unique isomorphisms $E_j \otimes E_i \rightarrow E_i \otimes E_j$, and (2.8) holds since the identity map is the only automorphism of $E_l \otimes E_j \otimes E_i$. See [12, Examples 1.7] for a more direct translation of the Robertson-Steger conditions into k -graphs.

Note that our result holds for both finite and infinite k , and that one can replace \mathbb{N}^k with an arbitrary right-angled Artin semigroup to obtain a more general result.

To prove Theorems 2.1 and 2.2, we need a few preliminary results. First some notation and terminology. Define an action of the symmetric group S_k on the words of length k in \mathbb{F}_A^+ by

$$\sigma\mu := \mu_{\sigma^{-1}(1)} \cdots \mu_{\sigma^{-1}(k)} \quad \text{for } \sigma \in S_k, \mu \in \mathbb{F}_A^+, \ell(\mu) = k.$$

For $1 \leq i \leq k - 1$, let $\tau_i \in S_k$ be the transposition $(i, i + 1)$; we shall omit the dependence on k , but this should not cause any confusion.

We call τ_i an *allowable transposition for μ* if $\mu_i \leftrightarrow \mu_{i+1}$. Note that since Γ has no loops, τ_i is not an allowable transposition for μ when $\mu_i = \mu_{i+1}$. We call $\sigma \in S_k$ an *allowable permutation for μ* if it can be written as a product $\tau_{i_m} \cdots \tau_{i_1}$ in which τ_{i_j} is an allowable transposition for $\tau_{i_{j-1}} \cdots \tau_{i_1} \mu$ for each j .

Lemma 2.4. *Let σ and ρ be allowable permutations for μ . Then:*

- (1) $\pi(\sigma\mu) = \pi(\mu)$.
- (2) If $i < j$ and $\mu_i \not\leftrightarrow \mu_j$, then $\sigma(i) < \sigma(j)$.
- (3) If $\sigma\mu = \rho\mu$, then $\sigma = \rho$.

Proof. The first assertion follows immediately from the graph product relations upon writing σ as a product of allowable transpositions. For (2), suppose $i < j$ and $\mu_i \not\leftrightarrow \mu_j$. Since the result is obvious for the identity permutation, we may inductively assume that $\sigma = \tau_l \kappa$, where κ is an allowable permutation for μ such that $\kappa(i) < \kappa(j)$ and τ_l is an allowable transposition for $\kappa\mu$. Since τ_l is allowable, we have $(\kappa\mu)_l \leftrightarrow (\kappa\mu)_{l+1}$; that is, $\mu_{\kappa^{-1}(l)} \leftrightarrow \mu_{\kappa^{-1}(l+1)}$. Since $\mu_i \not\leftrightarrow \mu_j$, this implies that either $\kappa(i) \neq l$ or $\kappa(j) \neq l + 1$. From the assumption that $\kappa(i) < \kappa(j)$ we can thus deduce that $\tau_l \kappa(i) < \tau_l \kappa(j)$, and hence $\sigma(i) < \sigma(j)$.

(3) First suppose $\sigma\mu = \mu$. We claim that σ is the identity permutation. If not, then there exists i such that $i < \sigma(i)$. From $\sigma\mu = \mu$ we deduce that $\mu_{\sigma^m(i)} = \mu_{\sigma^{m+1}(i)}$, and hence $\mu_{\sigma^m(i)} \not\leftrightarrow \mu_{\sigma^{m+1}(i)}$, for every m . Repeated applications of (2) yield the contradiction

$$i < \sigma(i) < \sigma^2(i) < \sigma^3(i) < \cdots$$

Now suppose $\sigma\mu = \rho\mu$. Then $\rho^{-1}\sigma$ is an allowable permutation for μ such that $\rho^{-1}\sigma\mu = \mu$, and hence $\sigma = \rho$. \square

We now make use of a result from [10], which, when formulated in the language we have developed, states that if two elements of \mathbb{F}_A^+ have the same image under the quotient map π , then they are connected by a sequence of allowable transpositions; i.e., one is obtainable from the other by an allowable permutation. (See also [11].) Hence for each $\mu \in \mathbb{F}_A^+$ there is an allowable permutation σ for μ such that $\sigma\mu =$

$\delta(\pi(\mu))$. By part (3) of the previous lemma, the permutation σ is unique. Thus we define:

Definition 2.5. For each $\mu \in \mathbb{F}_A^+$, let σ_μ be the unique allowable permutation for μ such that $\sigma_\mu\mu = \delta(\pi(\mu))$.

For each permutation σ let

$$\iota(\sigma) := |\{(i, j) : i < j \text{ and } \sigma(i) > \sigma(j)\}|,$$

the number of inversions in σ .

Lemma 2.6. Let $\mu \in \mathbb{F}_A^+$.

- (1) If ρ is an allowable permutation for μ , then $\sigma_{\rho\mu} = \sigma_\mu\rho^{-1}$.
- (2) If $\sigma_\mu(i) > \sigma_\mu(i + 1)$, then $\iota(\sigma_{\tau_i\mu}) = \iota(\sigma_\mu) - 1$ and $\mu_i \leftrightarrow \mu_{i+1}$.

Proof. (1) By Lemma 2.4(1) we have

$$\sigma_\mu\mu = \delta(\pi(\mu)) = \delta(\pi(\rho\mu)) = \sigma_{\rho\mu}(\rho\mu).$$

Since $\sigma_{\rho\mu}\rho$ is an allowable permutation for μ , (1) now follows from part (3) of Lemma 2.4.

The assumption $\sigma_\mu(i) > \sigma_\mu(i + 1)$ implies that $\iota(\sigma_\mu\tau_i) = \iota(\sigma_\mu) - 1$. The first conclusion of (2) now follows from (1), and the second is an immediate consequence of Lemma 2.4(2). □

The following proposition is our main technical result.

Proposition 2.7. Let $T = (T_{a,b})_{a \leftrightarrow b}$ be a family of isomorphisms

$$T_{a,b}: X_a \otimes X_b \rightarrow X_b \otimes X_a$$

which satisfies (2.1) and (2.2). Fix a well-ordering of the vertex set A and let $\delta: P \rightarrow \mathbb{F}_A^+$ be the corresponding preferred section. Then there is a unique family $(U_\mu)_{\mu \in \mathbb{F}_A^+}$ of isomorphisms $U_\mu: X_\mu \rightarrow X_{\delta(\pi(\mu))}$ which satisfies

$$(2.9) \quad U_\mu = 1_\mu \quad \text{if } \delta(\pi(\mu)) = \mu$$

and

$$(2.10) \quad U_\mu = U_{\tau_i\mu}T_i^\mu \quad \text{whenever } \mu_i \leftrightarrow \mu_{i+1},$$

where T_i^μ is the isomorphism

$$1_{\mu_1} \otimes \cdots \otimes 1_{\mu_{i-1}} \otimes T_{\mu_i, \mu_{i+1}} \otimes 1_{\mu_{i+2}} \otimes \cdots \otimes 1_{\mu_{\ell(\mu)}}: X_\mu \rightarrow X_{\tau_i\mu}.$$

For this family, we have

$$(2.11) \quad U_{\delta(\pi(\mu))\nu}(U_\mu \otimes 1_\nu) = U_{\mu\nu} = U_{\mu\delta(\pi(\nu))}(1_\mu \otimes U_\nu) \quad \text{for all } \mu, \nu \in \mathbb{F}_A^+.$$

Proof. We begin by recursively defining the family $(U_\mu)_{\mu \in \mathbb{F}_A^+}$. If $\iota(\mu) = 0$, define $U_\mu := 1_\mu$. Now let $k \geq 0$, and suppose that we have defined U_μ for every $\mu \in \mathbb{F}_A^+$ such that $\iota(\sigma_\mu) \leq k$. Fix $\mu \in \mathbb{F}_A^+$ such that $\iota(\sigma_\mu) = k + 1$, and let

$$j := \min\{l : \sigma_\mu(l) > \sigma_\mu(l + 1)\}.$$

By Lemma 2.6(2) we have $\iota(\sigma_{\tau_j\mu}) = \iota(\sigma_\mu) - 1 = k$ (so that $U_{\tau_j\mu}$ is defined) and $\mu_j \leftrightarrow \mu_{j+1}$ (so that T_j^μ is defined), so we can define U_μ recursively by

$$U_\mu := U_{\tau_j\mu}T_j^\mu.$$

If $\delta(\pi(\mu)) = \mu$, then $\iota(\sigma_\mu) = 0$, so (2.9) holds by definition of U_μ . We claim that

$$(2.12) \quad U_\mu = U_{\tau_i\mu} T_i^\mu \quad \text{whenever } \sigma_\mu(i) > \sigma_\mu(i+1);$$

again we remark that, by Lemma 2.6(2), the condition $\sigma_\mu(i) > \sigma_\mu(i+1)$ ensures that T_i^μ is defined. Before verifying (2.12), let us show how it implies (2.10). For this, it suffices to show that (2.10) holds whenever $\mu_i \leftrightarrow \mu_{i+1}$ and $\sigma_\mu(i) < \sigma_\mu(i+1)$. Using Lemma 2.6(1), we compute that

$$\sigma_{\tau_i\mu}(i) = \sigma_\mu\tau_i(i) = \sigma_\mu(i+1) > \sigma_\mu(i) = \sigma_\mu\tau_i(i+1) = \sigma_{\tau_i\mu}(i+1).$$

Hence we may apply (2.12) to $\tau_i\mu$ to deduce that

$$U_{\tau_i\mu} = U_\mu T_i^{\tau_i\mu}.$$

Composing both sides on the right with $(T_i^{\tau_i\mu})^{-1} = T_i^\mu$ gives (2.10).

We will verify (2.12) by induction on $\iota(\sigma_\mu)$. If $\iota(\sigma_\mu) = 0$, then $\sigma_\mu = \text{id}$, and (2.12) holds vacuously. Let $k \geq 1$, and suppose inductively that (2.12) holds whenever $\iota(\sigma_\mu) \leq k-1$. Fix $\mu \in \mathbb{F}_A^+$ with $\iota(\sigma_\mu) = k$, and suppose $i \in \{1, \dots, \ell(\mu) - 1\}$ is such that $\sigma_\mu(i) > \sigma_\mu(i+1)$. Let

$$j := \min\{l : \sigma_\mu(l) > \sigma_\mu(l+1)\},$$

so that by definition $U_\mu = U_{\tau_j\mu} T_j^\mu$. Then $j \leq i$, and we consider three cases.

Case 1: $i = j$. Then $U_\mu = U_{\tau_i\mu} T_i^\mu$ holds by definition of U_μ .

Case 2: $i \geq j + 2$.

By Lemma 2.6(2) we have $\iota(\sigma_{\tau_i\mu}) = k - 1$, and using Lemma 2.6(1) we check that

$$\sigma_{\tau_i\mu}(j) = \sigma_\mu\tau_i(j) = \sigma_\mu(j) > \sigma_\mu(j+1) = \sigma_\mu\tau_i(j+1) = \sigma_{\tau_i\mu}(j+1).$$

Hence we may apply (2.12) to $\tau_i\mu$ to obtain

$$U_{\tau_i\mu} = U_{\tau_j\tau_i\mu} T_j^{\tau_i\mu}.$$

Similarly,

$$U_{\tau_j\mu} = U_{\tau_i\tau_j\mu} T_i^{\tau_j\mu}.$$

Since $i \geq j + 2$ we have $\tau_j\tau_i = \tau_i\tau_j$. Moreover, with $\nu := \mu_1 \cdots \mu_{j-1}$, $\lambda := \mu_{j+2} \cdots \mu_{i-1}$ and $\theta := \mu_{i+2} \cdots \mu_{\ell(\mu)}$, we have

$$\begin{aligned} T_j^{\tau_i\mu} T_i^\mu &= (1_\nu \otimes 1_{\mu_{i+1}\mu_i} \otimes 1_\lambda \otimes T_{\mu_j, \mu_{j+1}} \otimes 1_\theta)(1_\nu \otimes T_{\mu_i, \mu_{i+1}} \otimes 1_\lambda \otimes 1_{\mu_j\mu_{j+1}} \otimes 1_\theta) \\ &= (1_\nu \otimes T_{\mu_i, \mu_{i+1}} \otimes 1_\lambda \otimes 1_{\mu_{j+1}\mu_j} \otimes 1_\theta)(1_\nu \otimes 1_{\mu_i\mu_{i+1}} \otimes 1_\lambda \otimes T_{\mu_j, \mu_{j+1}} \otimes 1_\theta) \\ &= T_i^{\tau_j\mu} T_j^\mu. \end{aligned}$$

Thus

$$U_\mu = U_{\tau_j\mu} T_j^\mu = U_{\tau_i\tau_j\mu} T_i^{\tau_j\mu} T_j^\mu = U_{\tau_j\tau_i\mu} T_j^{\tau_i\mu} T_i^\mu = U_{\tau_i\mu} T_i^\mu,$$

as required.

Case 3: $i = j + 1$.

Lemma 2.6(2) gives $\iota(\sigma_{\tau_j\mu}) = k - 1$, and, since $\sigma_\mu(j) > \sigma_\mu(j+1) > \sigma_\mu(j+2)$, we can use Lemma 2.6(1) to check that

$$(2.13) \quad \begin{aligned} \sigma_{\tau_j\mu}(j+1) &= \sigma_\mu\tau_j(j+1) = \sigma_\mu(j) \\ &> \sigma_\mu(j+2) = \sigma_\mu\tau_j(j+2) = \sigma_{\tau_j\mu}(j+2). \end{aligned}$$

Hence we may apply (2.12) to $\tau_j\mu$ to obtain

$$U_{\tau_j\mu} = U_{\tau_{j+1}\tau_j\mu} T_{j+1}^{\tau_j\mu}.$$

By Lemma 2.6(2), (2.13) also implies that $\iota(\sigma_{\tau_{j+1}\tau_j}) = \iota(\sigma_{\tau_j\mu}) - 1 = k - 2$, and using Lemma 2.6(1) we check that

$$\begin{aligned} \sigma_{\tau_{j+1}\tau_j\mu}(j) &= \sigma_\mu\tau_j\tau_{j+1}(j) = \sigma_\mu(j + 1) \\ &> \sigma_\mu(j + 2) = \sigma_\mu\tau_j\tau_{j+1}(j + 1) = \sigma_{\tau_{j+1}\tau_j\mu}(j + 1). \end{aligned}$$

Hence we may apply (2.12) to $\tau_{j+1}\tau_j\mu$ to obtain

$$U_{\tau_{j+1}\tau_j\mu} = U_{\tau_j\tau_{j+1}\tau_j\mu}T_j^{\tau_{j+1}\tau_j\mu}.$$

Thus

$$\begin{aligned} (2.14) \quad U_\mu &= U_{\tau_j\mu}T_j^\mu \\ &= U_{\tau_{j+1}\tau_j\mu}T_{j+1}^{\tau_j\mu}T_j^\mu \\ &= U_{\tau_j\tau_{j+1}\tau_j\mu}T_j^{\tau_{j+1}\tau_j\mu}T_{j+1}^{\tau_j\mu}T_j^\mu. \end{aligned}$$

Since $\sigma_\mu(j) > \sigma_\mu(j + 1) > \sigma_\mu(j + 2)$, Lemma 2.6(2) implies that μ_j, μ_{j+1} and μ_{j+2} form the vertices of a triangle in Γ . Using expansions such as

$$T_j^\mu = 1_\nu \otimes (T_{\mu_j, \mu_{j+1}} \otimes 1_{\mu_{j+2}}) \otimes 1_\theta,$$

(where $\nu := \mu_1 \cdots \mu_{j-1}$ and $\theta := \mu_{j+3} \cdots \mu_{\ell(\mu)}$), the hexagonal equation (2.2) gives

$$T_j^{\tau_{j+1}\tau_j\mu}T_{j+1}^{\tau_j\mu}T_j^\mu = T_{j+1}^{\tau_j\tau_{j+1}\mu}T_j^{\tau_{j+1}\mu}T_{j+1}^\mu.$$

Using this and $\tau_j\tau_{j+1}\tau_j = \tau_{j+1}\tau_j\tau_{j+1}$ in (2.14) gives

$$(2.15) \quad U_\mu = U_{\tau_{j+1}\tau_j\tau_{j+1}\mu}T_{j+1}^{\tau_j\tau_{j+1}\mu}T_j^{\tau_{j+1}\mu}T_{j+1}^\mu.$$

As above, one now verifies that $\iota(\sigma_{\tau_{j+1}\mu}) = k - 1$ and $\sigma_{\tau_{j+1}\mu}(j) > \sigma_{\tau_{j+1}\mu}(j + 1)$, from which (2.12) gives

$$U_{\tau_{j+1}\mu} = U_{\tau_j\tau_{j+1}\mu}T_j^{\tau_{j+1}\mu}.$$

One then verifies that $\iota(\sigma_{\tau_j\tau_{j+1}\mu}) = k - 2$, and that $\sigma_{\tau_j\tau_{j+1}\mu}(j + 1) > \sigma_{\tau_j\tau_{j+1}\mu}(j + 2)$, from which (2.12) gives

$$U_{\tau_j\tau_{j+1}\mu} = U_{\tau_{j+1}\tau_j\tau_{j+1}\mu}T_{j+1}^{\tau_j\tau_{j+1}\mu}.$$

Combining these last two equations with (2.15) gives

$$\begin{aligned} U_\mu &= U_{\tau_{j+1}\tau_j\tau_{j+1}\mu}T_{j+1}^{\tau_j\tau_{j+1}\mu}T_j^{\tau_{j+1}\mu}T_{j+1}^\mu \\ &= U_{\tau_j\tau_{j+1}\mu}T_j^{\tau_{j+1}\mu}T_{j+1}^\mu \\ &= U_{\tau_{j+1}\mu}T_{j+1}^\mu. \end{aligned}$$

This concludes Case 3, and the proof of (2.12) is complete.

For uniqueness, suppose $(V_\mu)_{\mu \in \mathbb{F}_A^+}$ is a different family of isomorphisms $V_\mu : X_\mu \rightarrow X_{\delta(\pi(\mu))}$ which satisfies (2.9) and (2.10). Choose μ with $\iota(\sigma_\mu)$ minimal such that $U_\mu \neq V_\mu$. Since both collections satisfy (2.9) we have $\iota(\sigma_\mu) \geq 1$, and hence there exists i with $\sigma_\mu(i) > \sigma_\mu(i + 1)$. By Lemma 2.6(2) we have $\iota(\sigma_{\tau_i\mu}) < \iota(\sigma_\mu)$, so by minimality $U_{\tau_i\mu} = V_{\tau_i\mu}$. Using (2.10) we obtain the contradiction

$$U_\mu = U_{\tau_i\mu}T_i^\mu = V_{\tau_i\mu}T_i^\mu = V_\mu.$$

Thus the collection is unique.

We now verify (2.11). We will prove that

$$(2.16) \quad U_{\delta(\pi(\mu))\nu}(U_\mu \otimes 1_\nu) = U_{\mu\nu} \quad \text{for all } \mu, \nu \in \mathbb{F}_A^+;$$

the proof of the other half of (2.11) proceeds in a similar fashion. We verify (2.16) by induction on $\iota(\sigma_\mu)$. If $\iota(\sigma_\mu) = 0$, then $\delta(\pi(\mu)) = \mu$, and

$$U_{\delta(\pi(\mu))\nu}(U_\mu \otimes 1_\nu) = U_{\mu\nu}(1_\mu \otimes 1_\nu) = U_{\mu\nu}.$$

Suppose inductively that (2.16) holds whenever $\iota(\sigma_\mu) \leq k - 1$. Let $\nu \in \mathbb{F}_A^+$ be arbitrary, and fix $\mu \in \mathbb{F}_A^+$ satisfying $\iota(\sigma_\mu) = k$. Since $k \geq 1$, there exists i such that $\sigma_\mu(i) > \sigma_\mu(i + 1)$, and Lemma 2.6(2) gives $\iota(\sigma_{\tau_i\mu}) = k - 1$ and $\mu_i \leftrightarrow \mu_{i+1}$. The inductive hypothesis gives that

$$(2.17) \quad U_{\delta(\pi(\tau_i\mu))\nu}(U_{\tau_i\mu} \otimes 1_\nu) = U_{(\tau_i\mu)\nu}.$$

Since $\pi(\tau_i\mu) = \pi(\mu)$,

$$\begin{aligned} U_{\delta(\pi(\mu))\nu}(U_\mu \otimes 1_\nu) &= U_{\delta(\pi(\tau_i\mu))\nu}(U_{\tau_i\mu} T_i^\mu \otimes 1_\nu) && \text{(by 2.10)} \\ &= U_{\delta(\pi(\tau_i\mu))\nu}(U_{\tau_i\mu} \otimes 1_\nu)(T_i^\mu \otimes 1_\nu) \\ &= U_{(\tau_i\mu)\nu}(T_i^\mu \otimes 1_\nu) && \text{(by 2.17)} \\ &= U_{\tau_i(\mu\nu)} T_i^{\mu\nu} \\ &= U_{\mu\nu} && \text{(by 2.10),} \end{aligned}$$

and the proof is complete by induction. □

Proof of Theorem 2.1. We begin by proving the existence of a product system (Y, α) which satisfies the conditions of the theorem. The collection Y is determined by (2.3). Let $(U_\mu)_{\mu \in \mathbb{F}_A^+}$ be the family of isomorphisms $U_\mu: X_\mu \rightarrow X_{\delta(\pi(\mu))}$ given by Proposition 2.7. If $s, t \in P$, then $Y_s \otimes Y_t = X_{\delta(s)} \otimes X_{\delta(t)} = X_{\delta(s)\delta(t)}$ and $Y_{st} = X_{\delta(st)} = X_{\delta(\pi(\delta(s)\delta(t)))}$, so we can define $\alpha_{s,t}: Y_s \otimes Y_t \rightarrow Y_{st}$ by

$$\alpha_{s,t} := U_{\delta(s)\delta(t)}.$$

To see that (Y, α) satisfies the associativity condition (1.2), suppose $r, s, t \in P$. Setting $\mu = \delta(r)\delta(s)$ and $\nu = \delta(t)$ in the first part of (2.11) gives

$$\alpha_{rs,t}(\alpha_{r,s} \otimes 1_{Y_t}) = U_{\delta(rs)\delta(t)}(U_{\delta(r)\delta(s)} \otimes 1_{\delta(t)}) = U_{\delta(r)\delta(s)\delta(t)},$$

and setting $\mu = \delta(r)$ and $\nu = \delta(s)\delta(t)$ in the second part of (2.11) gives

$$\alpha_{r,st}(1_{Y_r} \otimes \alpha_{s,t}) = U_{\delta(r)\delta(st)}(1_{\delta(r)} \otimes U_{\delta(s)\delta(t)}) = U_{\delta(r)\delta(s)\delta(t)}.$$

Thus (1.2) holds, and (Y, α) is a product system.

To check (2.4), suppose $\delta(st) = \delta(s)\delta(t)$. By (2.9) we have $U_{\delta(st)} = 1_{\delta(st)}$, and hence

$$\alpha_{s,t} = U_{\delta(s)\delta(t)} = U_{\delta(st)} = 1_{\delta(st)},$$

giving (2.4). For (2.5), suppose $a, b \in A$ satisfy $a \leftrightarrow b$ and $a > b$. Then $\alpha_{\pi(a),\pi(b)} = U_{\delta(\pi(a))\delta(\pi(b))} = U_{ab} = U_{ba} T_{a,b}$, where the last equality is by (2.10). But $\delta(\pi(ba)) = ba$, so by (2.9) we have $U_{ba} = 1_{ba}$, and hence $\alpha_{\pi(a),\pi(b)} = T_{a,b}$, as required.

For uniqueness, we first establish that

$$(2.18) \quad \alpha_{s_1, \dots, s_k} = U_{\delta(s_1) \dots \delta(s_k)} \quad \text{for all } k \geq 1 \text{ and } s_1, \dots, s_k \in P.$$

(See Notation 1.2.) This equation holds by definition when $k \leq 2$, so suppose inductively that it holds for some $k \geq 2$. Let $s_1, \dots, s_{k+1} \in P$. Then

$$\begin{aligned} \alpha_{s_1, \dots, s_{k+1}} &= \alpha_{s_1 \dots s_k, s_{k+1}}(\alpha_{s_1, \dots, s_k} \otimes 1_{Y_{s_{k+1}}}) && \text{(associativity of } \alpha) \\ &= U_{\delta(s_1 \dots s_k)\delta(s_{k+1})}(U_{\delta(s_1) \dots \delta(s_k)} \otimes 1_{\delta(s_{k+1})}) && \text{(induction)} \\ &= U_{\delta(s_1) \dots \delta(s_{k+1})}, \end{aligned}$$

with the last equality following from (2.11) by setting $\mu = \delta(s_1) \dots \delta(s_k)$ and $\nu = \delta(s_{k+1})$. Hence (2.18) holds for $k + 1$, and inductively for all k .

Now suppose (Y, β) is another product system over P which satisfies (2.3), (2.4) and (2.5). Define

$$V_\mu := \beta_{\pi(\mu_1), \dots, \pi(\mu_{\ell(\mu)})} \quad \text{for } \mu \in \mathbb{F}_A^+.$$

Then V_μ is an isomorphism from $Y_{\pi(\mu_1)} \otimes \dots \otimes Y_{\pi(\mu_k)}$ to $Y_{\pi(\mu)}$, or, equivalently, from X_μ to $X_{\delta(\pi(\mu))}$. We claim that $(V_\mu)_{\mu \in \mathbb{F}_A^+}$ is the unique family of isomorphisms given by Proposition 2.7.

We begin by verifying (2.9). Suppose $\delta(\pi(\mu)) = \mu$. If $\ell(\mu) = 1$, then $V_\mu = 1_\mu$ by definition. Suppose inductively that $V_\mu = 1_\mu$ whenever $\delta(\pi(\mu)) = \mu$ and $\ell(\mu) \leq k$ for some $k \geq 1$. Fix $\mu \in \mathbb{F}_A^+$ such that $\delta(\pi(\mu)) = \mu$ and $\ell(\mu) = k + 1$. Express $\mu = \mu_1\nu$. By definition of $\delta(\pi(\mu))$ we have $\delta(\pi(\mu)) = \mu_1\delta(\pi(\nu))$, and since $\delta(\pi(\mu)) = \mu = \mu_1\nu$, we deduce that $\delta(\pi(\nu)) = \nu$. Since

$$\delta(\pi(\mu_1))\delta(\pi(\nu)) = \mu_1\nu = \mu = \delta(\pi(\mu)) = \delta(\pi(\mu_1)\pi(\nu)),$$

(2.4) gives that $\beta_{\pi(\mu_1), \pi(\nu)} = 1_\mu$. By induction we also have $V_\nu = 1_\nu$, so

$$V_\mu = \beta_{\pi(\mu_1), \pi(\nu)}(1_{\pi(\mu_1)} \otimes V_\nu) = 1_\mu,$$

as required.

We now verify (2.10). First suppose that $a, b \in A$ satisfy $a \leftrightarrow b$. Then

$$V_{ab} = \beta_{\pi(a), \pi(b)} = \begin{cases} T_{a,b} & \text{if } a > b \\ 1_{ab} & \text{if } a < b, \end{cases}$$

so

$$\begin{aligned} V_{ab}T_{b,a} &= \begin{cases} 1_{ba} & \text{if } a > b \\ T_{b,a} & \text{if } a < b \end{cases} \\ &= V_{ba}. \end{aligned}$$

Now suppose $\mu \in \mathbb{F}_A^+$ and $\mu_i \leftrightarrow \mu_{i+1}$. Express $\mu = \nu\mu_i\mu_{i+1}\theta$ with $\nu, \theta \in \mathbb{F}_A^+$. Then

$$\begin{aligned} V_{\tau_i\mu}T_i^\mu &= V_{\tau_i\mu}(1_\nu \otimes T_{\mu_i, \mu_{i+1}} \otimes 1_\theta) \\ &= \beta_{\pi(\nu), \pi(\mu_{i+1}\mu_i), \pi(\theta)}(V_\nu \otimes V_{\mu_{i+1}\mu_i} \otimes V_\theta)(1_\nu \otimes T_{\mu_i, \mu_{i+1}} \otimes 1_\theta) \\ &= \beta_{\pi(\nu), \pi(\mu_i\mu_{i+1}), \pi(\theta)}(V_\nu \otimes V_{\mu_i\mu_{i+1}} \otimes V_\theta) \\ &= V_\mu, \end{aligned}$$

giving (2.10).

By the uniqueness assertion of Proposition 2.7, we have $V_\mu = U_\mu$ for every $\mu \in \mathbb{F}_A^+$, which, together with (2.18), gives

$$\alpha_{\pi(\mu_1), \dots, \pi(\mu_k)} = \beta_{\pi(\mu_1), \dots, \pi(\mu_k)} \quad \text{for every } \mu \in \mathbb{F}_A^+.$$

Now suppose $s, t \in P$. Let $k := \ell(s)$, $l := \ell(t)$, $\mu := \delta(s)$, and $\nu := \delta(t)$. Then

$$\alpha_{s,t}(\alpha_{\pi(\mu_1), \dots, \pi(\mu_k)} \otimes \alpha_{\pi(\nu_1), \dots, \pi(\nu_l)}) = \alpha_{\pi(\mu_1), \dots, \pi(\mu_k), \pi(\nu_1), \dots, \pi(\nu_l)}$$

and

$$\beta_{s,t}(\beta_{\pi(\mu_1), \dots, \pi(\mu_k)} \otimes \beta_{\pi(\nu_1), \dots, \pi(\nu_l)}) = \beta_{\pi(\mu_1), \dots, \pi(\mu_k), \pi(\nu_1), \dots, \pi(\nu_l)},$$

from which we deduce that $\alpha_{s,t} = \beta_{s,t}$. Thus (Y, α) is the unique product system over P which satisfies (2.3), (2.4) and (2.5). \square

Proof of Theorem 2.2. It is obvious that (2.1) holds for the family $(T_{a,b})_{a \leftrightarrow b}$ defined by (2.7). To verify the hexagonal equation (2.2), suppose a, b and c are the vertices of a triangle in Γ . We will show that the following equivalent version of (2.2) is satisfied:

$$(T_{b,a} \otimes 1_c)(1_b \otimes T_{c,a})(T_{c,b} \otimes 1_a)(1_c \otimes T_{a,b})(T_{a,c} \otimes 1_b)(1_a \otimes T_{b,c}) = 1_{abc}.$$

The left-hand side of this equation is

$$\begin{aligned} & (\beta_{\pi(a), \pi(b)}^{-1} \otimes 1_c)(\beta_{\pi(b), \pi(a)} \otimes 1_c)(1_b \otimes \beta_{\pi(a), \pi(c)}^{-1})(1_b \otimes \beta_{\pi(c), \pi(a)}) \\ & (\beta_{\pi(b), \pi(c)}^{-1} \otimes 1_a)(\beta_{\pi(c), \pi(b)} \otimes 1_a)(1_c \otimes \beta_{\pi(b), \pi(a)}^{-1})(1_c \otimes \beta_{\pi(a), \pi(b)}) \\ & (\beta_{\pi(c), \pi(a)}^{-1} \otimes 1_b)(\beta_{\pi(a), \pi(c)} \otimes 1_b)(1_a \otimes \beta_{\pi(c), \pi(b)}^{-1})(1_a \otimes \beta_{\pi(b), \pi(c)}), \end{aligned}$$

which by five applications of (1.2) simplifies to

$$\begin{aligned} & (\beta_{\pi(a), \pi(b)}^{-1} \otimes 1_c)\beta_{\pi(ba), \pi(c)}^{-1}\beta_{\pi(b), \pi(ac)}\beta_{\pi(b), \pi(ca)}^{-1}\beta_{\pi(bc), \pi(a)}\beta_{\pi(cb), \pi(a)}^{-1}\beta_{\pi(c), \pi(ba)} \\ & \beta_{\pi(c), \pi(ab)}^{-1}\beta_{\pi(ca), \pi(b)}\beta_{\pi(ac), \pi(b)}^{-1}\beta_{\pi(a), \pi(cb)}(1_a \otimes \beta_{\pi(b), \pi(c)}). \end{aligned}$$

Since $\pi(ab) = \pi(ba)$, $\pi(bc) = \pi(cb)$ and $\pi(ca) = \pi(ac)$, this in turn collapses to

$$(\beta_{\pi(a), \pi(b)}^{-1} \otimes 1_c)\beta_{\pi(ab), \pi(c)}^{-1}\beta_{\pi(a), \pi(bc)}(1_a \otimes \beta_{\pi(b), \pi(c)}),$$

which by one last application of (1.2) is the identity morphism on X_{abc} , as required.

Let (Y, α) be the product system (Y^T, α^T) associated with this collection T ; that is, $Y_s := X_{\delta(s)}$ and $\alpha_{s,t} := U_{\delta(s)\delta(t)}$, where $(U_\mu)_{\mu \in \mathbb{F}_A^+}$ is the family of isomorphisms $U_\mu: X_\mu \rightarrow X_{\delta(\pi(\mu))}$ given by Proposition 2.7. We will define an isomorphism ψ from (Y, α) to (Z, β) . Fix $s \in P$, let $\mu := \delta(s)$, and let $k := \ell(\mu)$. Then

$$Y_s = X_{\delta(s)} = X_{\mu_1} \otimes \cdots \otimes X_{\mu_k} = Z_{\pi(\mu_1)} \otimes \cdots \otimes Z_{\pi(\mu_k)},$$

so

$$\psi_s := \beta_{\pi(\mu_1), \dots, \pi(\mu_k)}$$

is an isomorphism from Y_s to $Z_{\pi(\mu)} = Z_s$. It remains only to show that

$$\begin{array}{ccc} Y_s \otimes Y_t & \xrightarrow{\alpha_{s,t}} & Y_{st} \\ \psi_s \otimes \psi_t \downarrow & & \downarrow \psi_{st} \\ Z_s \otimes Z_t & \xrightarrow{\beta_{s,t}} & Z_{st} \end{array}$$

commutes for every $s, t \in P$. Let $\mu = \delta(s)$ as above, let $\nu = \delta(t)$, and let $l := \ell(\nu)$. Since $\alpha_{s,t} = U_{\mu\nu}$ and $\beta_{s,t}(\psi_s \otimes \psi_t) = \beta_{\pi(\mu_1), \dots, \pi(\mu_k), \pi(\nu_1), \dots, \pi(\nu_l)}$, it suffices to show that

$$(2.19) \quad \beta_{\pi(\theta_1), \dots, \pi(\theta_m)} = \psi_{\pi(\theta)}U_\theta \quad \text{for every } \theta = \theta_1 \cdots \theta_m \in \mathbb{F}_A^+.$$

We will establish this equation by induction on $\iota(\sigma_\theta)$. If $\iota(\sigma_\theta) = 0$, then $\delta(\pi(\theta)) = \theta$, and (2.9) gives $U_\theta = 1_\theta$. Since $\delta(\pi(\theta)) = \theta$, (2.19) is then immediate from the definition of $\psi_{\pi(\theta)}$. Suppose inductively that (2.19) holds whenever $\iota(\sigma_\theta) \leq k$ for some $k \geq 0$, and fix $\theta \in \mathbb{F}_A^+$ with $\iota(\sigma_\theta) = k + 1$. There exists i such that

$\sigma_\theta(i) > \sigma_\theta(i+1)$, and Lemma 2.6(2) gives that $\theta_i \leftrightarrow \theta_{i+1}$ and $\iota(\sigma_{\tau_i\theta}) = \iota(\sigma_\theta) - 1 = k$. By (2.10) and induction we thus have

$$(2.20) \quad \psi_{\pi(\theta)}U_\theta = \psi_{\pi(\tau_i\theta)}U_{\tau_i\theta}T_i^\theta = \beta_{\pi(\theta_1), \dots, \pi(\theta_{i+1}), \pi(\theta_i), \dots, \pi(\theta_m)}T_i^\theta.$$

To further simplify the right-hand side of this equation, let $\mu := \theta_1 \cdots \theta_{i-1}$, let $\nu := \theta_{i+2} \cdots \theta_m$, and observe that

$$\begin{aligned} & \beta_{\pi(\theta_1), \dots, \pi(\theta_{i+1}), \pi(\theta_i), \dots, \pi(\theta_m)} \\ &= \beta_{\pi(\mu), \pi(\theta_{i+1}\theta_i), \pi(\nu)}(\beta_{\pi(\theta_1), \dots, \pi(\theta_{i-1})} \otimes \beta_{\pi(\theta_{i+1}), \pi(\theta_i)} \otimes \beta_{\pi(\theta_{i+2}), \dots, \pi(\theta_m)}) \end{aligned}$$

and

$$T_i^\theta = 1_\mu \otimes T_{\theta_i, \theta_{i+1}} \otimes 1_\nu.$$

By the definition of $T_{\theta_i, \theta_{i+1}}$ we have

$$\beta_{\pi(\theta_{i+1}), \pi(\theta_i)}T_{\theta_i, \theta_{i+1}} = \beta_{\pi(\theta_i), \pi(\theta_{i+1})},$$

so equation (2.20) simplifies to

$$\begin{aligned} \psi_{\pi(\theta)}U_\theta &= \beta_{\pi(\mu), \pi(\theta_i\theta_{i+1}), \pi(\nu)}(\beta_{\pi(\theta_1), \dots, \pi(\theta_{i-1})} \otimes \beta_{\pi(\theta_i), \pi(\theta_{i+1})} \otimes \beta_{\pi(\theta_{i+2}), \dots, \pi(\theta_m)}) \\ &= \beta_{\pi(\theta_1), \dots, \pi(\theta_m)}. \end{aligned}$$

This completes the induction, and hence the proof of the theorem. □

Proposition 2.8. *Suppose (Y, α) and (Z, β) are product systems over the right-angled Artin semigroup P which take values in the tensor groupoid \mathcal{G} . If $\psi = (\psi_s)_{s \in P}$ is an isomorphism from (Y, α) to (Z, β) , then defining*

$$(2.21) \quad \vartheta_a := \psi_{\pi(a)} \quad \text{for } a \in A$$

gives a collection $\vartheta := (\vartheta_a)_{a \in A}$ of isomorphisms $\vartheta_a: Y_{\pi(a)} \rightarrow Z_{\pi(a)}$ which satisfies

$$(2.22) \quad (\vartheta_b \otimes \vartheta_a)\alpha_{\pi(b), \pi(a)}^{-1}\alpha_{\pi(a), \pi(b)} = \beta_{\pi(b), \pi(a)}^{-1}\beta_{\pi(a), \pi(b)}(\vartheta_a \otimes \vartheta_b)$$

for every $a, b \in A$ such that $a \leftrightarrow b$. Moreover, given any such collection ϑ , there is a unique isomorphism $\psi: (Y, \alpha) \rightarrow (Z, \beta)$ such that $\psi_{\pi(a)} = \vartheta_a$ for every $a \in A$.

Remark 2.9. If Γ has no edges, then P is the free semigroup \mathbb{F}_A^+ , and Theorem 2.1 associates a product system to each collection $(X_a)_{a \in A}$ of objects in \mathcal{G} . Since condition (2.22) is then vacuous, Proposition 2.8 implies that the A -tuple of isomorphism classes of the X_a 's is a complete isomorphism invariant for product systems over \mathbb{F}_A^+ .

Proof of Proposition 2.8. Suppose $\psi = (\psi_s)_{s \in P}$ is an isomorphism from (Y, α) to (Z, β) ; that is, ψ_s is an isomorphism $Y_s \rightarrow Z_s$, and

$$\psi_{st}\alpha_{s,t} = \beta_{s,t}(\psi_s \otimes \psi_t) \quad \text{for all } s, t \in P.$$

Applying this equation with $s = \pi(a)$ and $t = \pi(b)$ gives

$$\beta_{\pi(b), \pi(a)}^{-1}\beta_{\pi(a), \pi(b)}(\psi_{\pi(a)} \otimes \psi_{\pi(b)}) = \beta_{\pi(b), \pi(a)}^{-1}\psi_{\pi(a)\pi(b)}\alpha_{\pi(a), \pi(b)},$$

and applying it with $s = \pi(b)$ and $t = \pi(a)$ gives

$$(\psi_{\pi(b)} \otimes \psi_{\pi(a)})\alpha_{\pi(b), \pi(a)}^{-1}\alpha_{\pi(a), \pi(b)} = \beta_{\pi(b), \pi(a)}^{-1}\psi_{\pi(b)\pi(a)}\alpha_{\pi(a), \pi(b)}.$$

Since $\pi(a)\pi(b) = \pi(b)\pi(a)$, this shows that (2.22) holds for the collection ϑ defined by (2.21).

Conversely, suppose we have a collection $\vartheta = (\vartheta_a)_{a \in A}$ of isomorphisms $\vartheta_a : Y_{\pi(a)} \rightarrow Z_{\pi(a)}$ which satisfies (2.22). Define

$$W_a := Y_{\pi(a)} \quad \text{and} \quad X_a := Z_{\pi(a)} \quad \text{for } a \in A$$

and, for every $a, b \in A$ such that $a \leftrightarrow b$, define

$$S_{a,b} := \alpha_{\pi(b), \pi(a)}^{-1} \alpha_{\pi(a), \pi(b)} : W_a \otimes W_b \rightarrow W_b \otimes W_a$$

and

$$T_{a,b} := \beta_{\pi(b), \pi(a)}^{-1} \beta_{\pi(a), \pi(b)} : X_a \otimes X_b \rightarrow X_b \otimes X_a.$$

By Theorem 2.2, the collections $S = (S_{a,b})_{a \leftrightarrow b}$ and $T = (T_{a,b})_{a \leftrightarrow b}$ satisfy equations (2.1) and (2.2), and the product systems (Y, α) and (Z, β) are isomorphic to (Y^S, α^S) and (Y^T, α^T) , respectively, via isomorphisms which are the identity on the fibers over $\pi(a)$ for $a \in A$. Hence it suffices to construct an isomorphism ψ from (Y^S, α^S) to (Y^T, α^T) such that $\psi_{\pi(a)} = \vartheta_a$ for every $a \in A$.

We begin by observing that, for each $a \in A$, ϑ_a is an isomorphism $W_a \rightarrow X_a$, and that the hypothesis (2.22) can be rewritten as

$$(2.23) \quad (\vartheta_b \otimes \vartheta_a) S_{a,b} = T_{a,b} (\vartheta_a \otimes \vartheta_b) \quad \text{whenever } a \leftrightarrow b.$$

Define

$$\vartheta_\mu := \vartheta_{\mu_1} \otimes \cdots \otimes \vartheta_{\mu_{\ell(\mu)}} : W_\mu \rightarrow X_\mu \quad \text{for } \mu \in \mathbb{F}_A^+$$

and

$$\psi_s := \vartheta_{\delta(s)} \quad \text{for } s \in P.$$

Then $\psi := (\psi_s)_{s \in P}$ is a collection of isomorphisms $\psi_s : Y_s^S \rightarrow Y_s^T$. We claim that

$$(2.24) \quad \psi_{st} \alpha_{s,t}^S = \alpha_{s,t}^T (\psi_s \otimes \psi_t) \quad \text{for } s, t \in P$$

is satisfied, so that ψ is an isomorphism of product systems.

Let $(U_\mu)_{\mu \in \mathbb{F}_A^+}$ and $(V_\mu)_{\mu \in \mathbb{F}_A^+}$ be the families of isomorphisms

$$U_\mu : X_\mu \rightarrow X_{\delta(\pi(\mu))} \quad \text{and} \quad V_\mu : W_\mu \rightarrow W_{\delta(\pi(\mu))}$$

given by Proposition 2.7, so that

$$\alpha_{s,t}^S = V_{\delta(s)\delta(t)} \quad \text{and} \quad \alpha_{s,t}^T = U_{\delta(s)\delta(t)} \quad \text{for } s, t \in P.$$

The equation (2.24) which we aim to verify can then be rewritten as

$$\vartheta_{\delta(st)} V_{\delta(s)\delta(t)} = U_{\delta(s)\delta(t)} \vartheta_{\delta(s)\delta(t)} \quad \text{for all } s, t \in P,$$

so it suffices to show that

$$(2.25) \quad \vartheta_{\delta(\pi(\mu))} V_\mu = U_\mu \vartheta_\mu \quad \text{for all } \mu \in \mathbb{F}_A^+.$$

We establish this by induction on $\iota(\sigma_\mu)$. If $\iota(\sigma_\mu) = 0$, then $\delta(\pi(\mu)) = \mu$, and the equation holds by (2.9). Suppose (2.25) is satisfied whenever $\iota(\sigma_\mu) \leq k - 1$ for some $k \geq 1$. Fix $\mu \in \mathbb{F}_A^+$ with $\iota(\sigma_\mu) = k$. Since $k \geq 1$, there exists i such that $\sigma_\mu(i) > \sigma_\mu(i + 1)$, and by Lemma 2.6(2) we have $\iota(\sigma_{\tau_i \mu}) = \iota(\sigma_\mu) - 1 = k - 1$ and $\mu_i \leftrightarrow \mu_{i+1}$. By (2.10) and induction,

$$(2.26) \quad \vartheta_{\delta(\pi(\mu))} V_\mu = \vartheta_{\delta(\pi(\tau_i \mu))} V_{\tau_i \mu} S_i^\mu = U_{\tau_i \mu} \vartheta_{\tau_i \mu} S_i^\mu.$$

Express $\mu = \nu \mu_i \mu_{i+1} \theta$ with $\nu, \theta \in \mathbb{F}_A^+$. Using expansions such as

$$\vartheta_{\tau_i \mu} = \vartheta_\nu \otimes (\vartheta_{\mu_{i+1}} \otimes \vartheta_{\mu_i}) \otimes \vartheta_\theta$$

and

$$S_i^\mu = 1_\nu \otimes S_{\mu_i, \mu_{i+1}} \otimes 1_\theta,$$

it is easy to see that equation (2.23) gives $\vartheta_{\tau_i \mu} S_i^\mu = T_i^\mu \vartheta_\mu$. Using this in (2.26) and applying (2.10) gives

$$\vartheta_{\delta(\pi(\mu))} V_\mu = U_{\tau_i \mu} \vartheta_{\tau_i \mu} S_i^\mu = U_{\tau_i \mu} T_i^\mu \vartheta_\mu = U_\mu \vartheta_\mu,$$

thus establishing (2.25). □

Using Proposition 2.8 it is easy to characterize the automorphism group of a product system over P .

Corollary 2.10. *Suppose (Y, α) is a product system over the right-angled Artin semigroup P which takes values in the tensor groupoid \mathcal{G} . Then the automorphism group of (Y, α) is isomorphic to the subgroup of $\prod_{a \in A} \text{Aut } Y_{\pi(a)}$ consisting of those A -tuples $(\vartheta_a)_{a \in A}$ which satisfy*

$$\alpha_{\pi(a), \pi(b)}(\vartheta_a \otimes \vartheta_b) \alpha_{\pi(a), \pi(b)}^{-1} = \alpha_{\pi(b), \pi(a)}(\vartheta_b \otimes \vartheta_a) \alpha_{\pi(b), \pi(a)}^{-1}$$

whenever $a \leftrightarrow b$.

The semigroup $H^2(P; \mathcal{G})$. Let \mathcal{G} be a symmetric tensor groupoid. We now describe the structure of the semigroup $H^2(P; \mathcal{G})$ in terms of the collections $T = (T_{a,b})_{a \leftrightarrow b}$ used in Theorem 2.1 to construct product systems (Y^T, α^T) . Consider the composite map

$$T \mapsto (Y^T, \alpha^T) \in Z^2(P; \mathcal{G}) \mapsto [(Y^T, \alpha^T)] \in H^2(P; \mathcal{G}).$$

By Theorem 2.2, this map is surjective and does not depend on the choice of well-ordering of the vertex set A . The following proposition describes the equivalence relation required on the domain to make the map bijective, and then describes the binary operation on the domain which corresponds to multiplication in $H^2(P; \mathcal{G})$.

Proposition 2.11. *Let $(W_a)_{a \in A}$ and $(X_a)_{a \in A}$ be collections of objects in \mathcal{G} , and let $S = (S_{a,b})_{a \leftrightarrow b}$ and $T = (T_{a,b})_{a \leftrightarrow b}$ be collections of isomorphisms*

$$S_{a,b}: W_a \otimes W_b \rightarrow W_b \otimes W_a \quad \text{and} \quad T_{a,b}: X_a \otimes X_b \rightarrow X_b \otimes X_a$$

which satisfy conditions (2.1) and (2.2). Then $[(Y^S, \alpha^S)] = [(Y^T, \alpha^T)]$ as elements of $H^2(P; \mathcal{G})$ if and only if there exists a collection $(\vartheta_a)_{a \in A}$ of isomorphisms $\vartheta_a: W_a \rightarrow X_a$ which satisfies

$$(2.27) \quad (\vartheta_b \otimes \vartheta_a) S_{a,b} = T_{a,b} (\vartheta_a \otimes \vartheta_b) \quad \text{whenever } a \leftrightarrow b.$$

Moreover, multiplication in $H^2(P; \mathcal{G})$ is given by

$$[(Y^S, \alpha^S)][(Y^T, \alpha^T)] = [(Y^{S \otimes T}, \alpha^{S \otimes T})],$$

where $((S \otimes T)_{a,b})_{a \leftrightarrow b}$ is the collection of isomorphisms

$$(S \otimes T)_{a,b}: (W_a \otimes X_a) \otimes (W_b \otimes X_b) \rightarrow (W_b \otimes X_b) \otimes (W_a \otimes X_a)$$

defined by

$$(S \otimes T)_{a,b} := (1_{W_b} \otimes \mathcal{F}_{W_a, X_b} \otimes 1_{X_a})(S_{a,b} \otimes T_{a,b})(1_{W_a} \otimes \mathcal{F}_{X_a, W_b} \otimes 1_{X_b}).$$

Proof. Since $(\alpha_{\pi(b),\pi(a)}^T)^{-1}\alpha_{\pi(a),\pi(b)}^T = T_{a,b}$ whenever $a \leftrightarrow b$, the first assertion follows immediately from Proposition 2.8. For the second, first recall that multiplication in $H^2(P; \mathcal{G})$ is given by the internal tensor product (Definition 1.9), so that

$$[(Y^S, \alpha^S)][(Y^T, \alpha^T)] = [(Y^S \otimes Y^T, \alpha^S \otimes \alpha^T)].$$

We claim that

$$(2.28) \quad (S \otimes T)_{a,b} = (\alpha^S \otimes \alpha^T)_{\pi(b),\pi(a)}^{-1}(\alpha^S \otimes \alpha^T)_{\pi(a),\pi(b)}$$

whenever $a \leftrightarrow b$. This will complete the proof, since it then follows from Theorem 2.2 that the collection $((S \otimes T)_{a,b})_{a \leftrightarrow b}$ satisfies (2.1) and (2.2), and that $(Y^{S \otimes T}, \alpha^{S \otimes T})$ is isomorphic to $(Y^S \otimes Y^T, \alpha^S \otimes \alpha^T)$.

To establish (2.28), first observe that if $a \leftrightarrow b$, then

$$\begin{aligned} & (\alpha_{\pi(b),\pi(a)}^S \otimes \alpha_{\pi(b),\pi(a)}^T)^{-1}(\alpha_{\pi(a),\pi(b)}^S \otimes \alpha_{\pi(a),\pi(b)}^T) \\ &= ((\alpha_{\pi(b),\pi(a)}^S)^{-1}\alpha_{\pi(a),\pi(b)}^S \otimes (\alpha_{\pi(b),\pi(a)}^T)^{-1}\alpha_{\pi(a),\pi(b)}^T) \\ &= S_{a,b} \otimes T_{a,b}. \end{aligned}$$

Using this and Definition 1.9 we thus have

$$\begin{aligned} & (\alpha^S \otimes \alpha^T)_{\pi(b),\pi(a)}^{-1}(\alpha^S \otimes \alpha^T)_{\pi(a),\pi(b)} \\ &= (1_{W_b} \otimes \mathcal{F}_{X_b, W_a} \otimes 1_{X_a})^{-1}(\alpha_{\pi(b),\pi(a)}^S \otimes \alpha_{\pi(b),\pi(a)}^T)^{-1} \\ & \quad (\alpha_{\pi(a),\pi(b)}^S \otimes \alpha_{\pi(a),\pi(b)}^T)(1_{W_a} \otimes \mathcal{F}_{X_a, W_b} \otimes 1_{X_b}) \\ &= (1_{W_b} \otimes \mathcal{F}_{W_a, X_b} \otimes 1_{X_a})(S_{a,b} \otimes T_{a,b})(1_{W_a} \otimes \mathcal{F}_{X_a, W_b} \otimes 1_{X_b}) \\ &= (S \otimes T)_{a,b}, \end{aligned}$$

giving (2.28). □

As a corollary we calculate $H^2(P; G)$ for an arbitrary abelian group G . For this, let E be the set of edges in Γ , and, when $a \leftrightarrow b$, write $e_{a,b}$ for the edge between a and b . Note that $e_{a,b} = e_{b,a}$.

Corollary 2.12. *Fix a well-ordering of A , and let $\delta: P \rightarrow \mathbb{F}_A^+$ be the corresponding preferred section. Then for each function $f: E \rightarrow G$, there is a unique 2-cocycle $\alpha^f \in Z^2(P; G)$ which satisfies*

$$(2.29) \quad \alpha_{s,t}^f = 1_G \quad \text{if } \delta(st) = \delta(s)\delta(t)$$

and

$$(2.30) \quad \alpha_{\pi(a),\pi(b)}^f = f(e_{a,b}) \quad \text{if } a \leftrightarrow b \text{ and } a > b.$$

The resulting map $f \mapsto [\alpha^f]$ is an isomorphism from $\prod_{e \in E} G$ to $H^2(P; G)$.

Proof. Let \mathcal{G} be the tensor groupoid with one object, morphisms G , and tensoring functor $g \otimes h := gh$ for $g, h \in G$. Given a function $f: E \rightarrow G$, define a collection $(T_{a,b})_{a \leftrightarrow b}$ of morphisms by

$$(2.31) \quad T_{a,b} := \begin{cases} f(e_{a,b}) & \text{if } a < b, \\ f(e_{a,b})^{-1} & \text{if } a > b. \end{cases}$$

Equation (2.1) is obviously satisfied, and the hexagonal equation (2.2) holds since G is abelian. Let $(Y^T, \alpha^T) \in Z^2(P; \mathcal{G})$ be the product system given by Theorem 2.1. Then $\alpha^f := \alpha^T$ is the unique element of $Z^2(P; G)$ which satisfies (2.29) and (2.30),

and using this uniqueness property it is easy to see that $f \mapsto [\alpha^f]$ is a group homomorphism from $\prod_{e \in E} G$ to $H^2(P; G)$. Moreover, condition (2.27) in Proposition 2.11 implies that this homomorphism is injective. Since every collection $(T_{a,b})$ arises from a function $f: E \rightarrow G$ according to (2.31), and since $T \mapsto [(Y^T, \alpha^T)]$ is surjective, so is $f \mapsto [\alpha^f]$. \square

Remark 2.13. When Γ has no edges, Corollary 2.12 says that the free semigroup \mathbb{F}_A^+ has trivial second cohomology. When Γ is the complete graph on A , it says that $H^2(\mathbb{N}^k; G)$ is isomorphic to the direct product of $\binom{k}{2}$ copies of G .

REFERENCES

- [1] W. Arveson, *Continuous analogues of Fock space*, Memoirs Amer. Math. Soc. **80** (1989), No. 409. MR **90f**:47061
- [2] J. Crisp and M. Laca, *On the Toeplitz algebras of right-angled and finite-type Artin groups*, J. Austral. Math. Soc., to appear.
- [3] H. T. Dinh, *Discrete product systems and their C^* -algebras*, J. Funct. Anal. **102** (1991), 1–34. MR **93d**:46097
- [4] H. T. Dinh, *On generalized Cuntz C^* -algebras*, J. Operator Theory **30** (1993), 123–135. MR **95m**:46112
- [5] N. J. Fowler, *Compactly-aligned discrete product systems, and generalizations of \mathcal{O}_∞* , International J. Math. **10** (1999), No. 6, 721–738. CMP 2000:02
- [6] N. J. Fowler, *Discrete product systems of finite-dimensional Hilbert spaces, and generalized Cuntz algebras*, preprint.
- [7] N. J. Fowler, *Discrete product systems of Hilbert bimodules*, Pacific J. Math., to appear.
- [8] N. J. Fowler and I. Raeburn, *Discrete product systems and twisted crossed products by semigroups*, J. Funct. Anal. **155** (1998), 171–204. MR **99k**:46118
- [9] N. J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999), 155–181. MR **2001b**:46093
- [10] E. R. Green, *Graph products of groups*, Thesis, The University of Leeds, 1990.
- [11] S. Hermiller and J. Meier, *Algorithms and geometry for graph products of groups*, J. of Algebra **171** (1995), 230–257. MR **96a**:20052
- [12] A. Kumjian and D. Pask, *Higher rank graph C^* -algebras*, New York J. Math **6** (2000), 1–20. MR **2001b**:46102
- [13] S. Mac Lane, *Categories for the working mathematician*, Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998. MR **2001j**:18001
- [14] P. S. Muhly and B. Solel, *Tensor algebras over C^* -correspondences (representations, dilations, and C^* -envelopes)*, J. Funct. Anal. **158** (1998), 389–457. MR **99j**:46066
- [15] M. V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbb{Z}* , Fields Institute Communications **12** (1997), 189–212. MR **97k**:46069
- [16] G. Robertson and T. Steger, *Affine buildings, tiling systems and higher rank Cuntz-Krieger algebras*, J. reine angew. Math. **513** (1999), 115–144. MR **2000j**:46109

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEWCASTLE, NSW 2308, AUSTRALIA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEWCASTLE, NSW 2308, AUSTRALIA