University of Wollongong

Research Online

Faculty of Health and Behavioural Sciences -
Papers (Archive)

Faculty of Science, Medicine and Health

1-1-2000

Global-perspective jitter improves vection in central vision

Stephen A. Palmisano

University of Wollongong, stephenp@uow.edu.au

Barbara Gillam

University of New South Wales

Shane Blackburn

University of New South Wales

Follow this and additional works at: <https://ro.uow.edu.au/hbspapers>

Part of the [Arts and Humanities Commons](#), [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Palmisano, Stephen A.; Gillam, Barbara; and Blackburn, Shane: Global-perspective jitter improves vection in central vision 2000, 57-67.

<https://ro.uow.edu.au/hbspapers/2709>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Global-perspective jitter improves vection in central vision

Abstract

Previous vection research has tended to minimise visual - vestibular conflict by using optic-flow patterns which simulate self-motions of constant velocity. Here, experiments are reported on the effect of adding 'global-perspective jitter' to these displays – simulating forward motion of the observer on a platform oscillating in horizontal and/or vertical dimensions. Unlike non-jittering displays, jittering displays produced a situation of sustained visual - vestibular conflict. Contrary to the prevailing notion that visual - vestibular conflict impairs vection, jittering optic flow was found to produce shorter vection onsets and longer vection durations than non-jittering optic flow for all of jitter magnitudes and temporal frequencies examined. On the basis of these findings, it would appear that purely radial patterns of optic flow are not the optimal inducing stimuli for vection. Rather, flow patterns which contain both regular and random-oscillating components appear to produce the most compelling subjective experiences of self-motion.

Keywords

central, improves, vision, perspective, vection, jitter, global

Disciplines

Arts and Humanities | Life Sciences | Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Palmisano, S. A., Gillam, B. & Blackburn, S. (2000). Global-perspective jitter improves vection in central vision. *Perception*, 29 (1), 57-67.

GLOBAL PERSPECTIVE JITTER IMPROVES VECTION IN CENTRAL VISION

Stephen Palmisano¹, Barbara Gillam² & Shane Blackburn²

¹Human Performance Laboratory, 103 Farquharson Building, 4700 Keele Street, Toronto,
Ontario, M3J 1P3, CANADA.

²School of Psychology, University of New South Wales, POB 1, Kensington, NSW,
2052, AUSTRALIA.

Other Contact Information:

Tel: (416) 736-5659

Fax: (416) 736-5857

Email: Stevep@hpl.crestech.ca

Abstract. Previous vection research has tended to minimise visual-vestibular conflict by utilising optic flow patterns which simulate self-motions of constant velocity. The present experiments examined the effect of adding ‘global perspective jitter’ to these displays - simulating forwards motion of the observer on a platform oscillating in horizontal and/or vertical dimensions. Unlike non-jittering displays, jittering displays produced a situation of sustained visual-vestibular conflict. Contrary to the prevailing notion that visual-vestibular conflict impairs vection, jittering optic flow was found to produce shorter vection onsets and longer vection durations than non-jittering optic flow - for all of jitter magnitudes and temporal frequencies examined. On the basis of these findings, it would appear that purely radial patterns of optic flow are not the optimal inducing stimuli for vection. Rather, flow patterns which contain both regular and random/oscillating components appear to produce the most compelling subjective experiences of self-motion.

1 Introduction

A number of sensory systems are responsible for the perception and control of self-motion - visual, vestibular, proprioceptive, somatosensory and auditory systems may all play a role (Benson 1990; Gibson 1966; Howard 1982; Johansson 1977). Of these modalities, the visual and vestibular systems appear to be particularly important. Research has shown that the visual system can register any type of self-motion on the basis of the optic flow¹ presented to the observer (ie active/passive, linear/rotary, constant velocity/accelerating self-motions - Brandt et al 1973; Lishman and Lee 1973). However, vision appears primarily sensitive to optic flow patterns with low temporal frequencies and/or simulating self-motions of constant velocity (Berthoz et al 1979; Berthoz et al, 1975; Dichgans and Brandt 1978; van Asten et al 1988). Conversely, the vestibular system registers only accelerating self-motions - on the basis of the inertia of the fluid in the semicircular canals and otolith organs (Benson 1990; Howard 1986a). As a result, this modality is primarily sensitive to brief high frequency stimulations (>1Hz; Diener et al 1982; 1984; Melville-Jones & Young, 1978) and is unable to distinguish between traveling at a constant linear velocity and remaining stationary (Lishman and Lee 1973).

Visual and vestibular systems normally provide redundant information about self-motion (Gibson 1966). However, there are situations where they apparently conflict² (for example when a stationary observer views an IMAX movie representing accelerating self-motion). The simplest solution to such a conflict would be for vision to dominate self-motion perception (Lishman and Lee 1973). Consistent with this notion, compelling illusions of self-motion can be induced by visual information alone - referred to asvection. For example, Lee and his colleagues have shown that when subjects are placed

inside a 'swinging room' - where the walls and ceiling swing back and forth - they quickly experience a visual illusion that they themselves are swaying (Lee and Aronson 1974; Lee and Lishman 1975; Lishman and Lee 1973). They argue that such illusory self-motions occur because visual information (indicating self-motion) overrides input from the vestibular, somatosensory and proprioceptive systems (indicating that the observer is in fact stationary).

However, other vection research suggests that visual-vestibular conflicts are not so simply resolved. Psychophysical studies using optic flow patterns which simulated constant velocity (rotary) self-motions found that: (1) stationary observers initially (for the first 3-4s) perceive the flow as entirely due to object motion; (2) they then experience a period of perceived self-acceleration in the opposite direction to the flow; and (3) finally, after about 8-18s exposure, observers perceive the flow as entirely due to self-motion (Brandt et al 1973; Held et al 1975; Young et al 1975). Zacharias and Young (1981) explain this time course in terms of the presence or absence of visual-vestibular conflict - defined as the difference between the *current vestibular signal* and the *expected vestibular signal based on the optic flow*. According to their theory, the optic flow initially produces a visual-vestibular conflict in stationary subjects, since the expected vestibular input is absent (real self-motions of constant velocity are preceded by a brief period of acceleration which would be detected by the vestibular system), resulting in perceived object-motion. This conflict fades quickly and disappears, as the expected vestibular input decreases below detection threshold (real constant velocity self-motions generate negligible vestibular input), resulting in exclusive vection.

Additional support for the notion that visual-vestibular conflict impairs vection is provided by observations that: (1) onset latencies for (circular) vection are shorter when visual and vestibular inputs are initially consistent (eg the observer undergoes an impulse acceleration in the simulated direction of self-motion while viewing a display indicating constant velocity self-motion - Brandt et al 1974; Melcher and Henn 1981; Wong and Frost 1981); (2) conflicting vestibular input can destroy (circular) vection (eg sudden acceleration of the observer in the opposite direction to the simulated self-motion - Teixeira and Lackner 1979; Young et al 1973); (3) (linear) vection along the vertical spinal axis has shorter onset latencies than (linear) vection along the horizontal sagittal axis - the former being thought to generate less visual-vestibular conflict than the latter (Giannopulu and Lepecq 1998); and (4) the lack of any vestibular input during visual displays simulating a roller coaster ride results in rather weak vection (Wann and Rushton 1994).

<INSERT FIGURE 1 ABOUT HERE>

The present experiments further examined the effect of visual-vestibular conflict on vection. Visual displays were of two types: (1) Non-jittering displays simulated forwards self-motion of constant velocity through a three-dimensional cloud of objects (see Figure1B). These radially expanding patterns of optic flow were similar to those used in previous linear vection studies and produced minimal/transient visual-vestibular conflict (eg Andersen and Braunstein 1985; Ohmi and Howard 1988; Palmisano, 1996; Telford et al 1992; Telford and Frost 1993). (2) Jittering displays simulated forwards self-motion of

constant velocity combined with continuous, random horizontal and/or vertical impulse self-accelerations (similar to the effects of ‘camera shake’ - see Figure 1A). These displays were radially expanding patterns of optic flow superimposed with horizontal and/or vertical ‘global perspective jitter’ (at particular points in time, the same randomly determined displacement, modified by perspective, was applied to all points in the flow field)³. Unlike non-jittering displays, jittering displays were designed to produce a situation of sustained visual-vestibular conflict in a stationary observer.

Given the above findings, it was expected that the lack of consistent vestibular input would bias observers to perceive jittering patterns of optic flow as due to object motion rather than self-motion. If so, vection should diminish as the magnitude and temporal frequency of the jitter increases (ie as the mismatch between visual and vestibular information increases).

2 Experiment 1: Does global perspective jitter impairvection?

2.1 *Method*

2.1.1 *Subjects.* Seven male and seven female undergraduate psychology students (aged between 18 and 24 years) participated in this experiment. All had normal or corrected-to-normal vision and had not previously experienced illusions of self-motion in the laboratory.

2.1.2 *Design.* Two independent variables were manipulated in this experiment. (1) *Display Type.* Displays were either jittering or non-jittering patterns of radially expanding optic flow. When present, global perspective jitter occurred along either the horizontal (x) axis, the vertical (y) axis, or both the x- and y- axes. (2) *Display speed.* Each display simulated one of three speeds of forwards self-motion: 2.7m/s, 4m/s and 7m/s. Regardless of their type or speed, all displays simulated self-motion through a three-dimensional cloud of 150 randomly-positioned objects.

2.1.3 *Apparatus.* Displays were generated on an IBM 486-DX (75MHz) personal computer and presented on a superVGA monitor with 1024 H x 768 V pixel resolution. The screen of this monitor subtended a visual angle of 40° H x 32° V when viewed from a chin rest 50cm away. Sincevection has been found to be dominated by the motion of the perceived background (Ohmi and Howard 1988; Telford et al 1992), inducing displays were presented 20cm behind a large black cardboard mask. Kinetic occlusion

always indicated that the display was in the background while the mask was in the foreground. This mask was placed in front of the subject, and two large partitions were placed on either side to restrict his/her vision. Only the monitor could be seen through a square window at the far end of this black viewing booth (1m wide x 2m deep x 2m high). As the inducing displays were viewed monocularly, the subject's right eye was always covered by an eye-patch.

2.1.4 *Visual Displays.* Non-jittering displays simulated forwards self-motion (z-axis) of a constant velocity relative to a three-dimensional cloud of randomly-positioned objects. This was achieved by increasing each object's velocity and total area (0.07° - 1.21°) as it appeared to approach the observer. Jittering displays were identical to non-jittering displays, with the exception that horizontal (x-axis) and/or vertical (y-axis) jitter was added to the optic flow. They simulated the observer moving forwards on a platform which oscillated in horizontal and/or vertical dimensions. The absolute amount of horizontal and/or vertical jitter for each frame was randomly selected from a range of 0 - 1/3 of the simulated forwards displacement. It's direction (left/right for horizontal jitter and up/down for vertical jitter) alternated from frame to frame. This signed jitter was then given the appropriate perspective transformation before it was applied to objects at different simulated locations in depth - ie the jitter component was less for more distant objects - we refer to this as *global perspective jitter*.

Both jittering and non-jittering optic flow consisted of moving green filled-in squares (with a luminance of 3cd/m^2) on a black background (0.03cd/m^2). As objects disappeared off the edge of the screen, they were replaced at the opposite end of space (a simulated

distance of 20m along the z-axis) at the same (x,y) coordinates. To reduce the sensation of their sudden appearance, these objects were initially replaced as dots which were slightly darker (1.6cd/m^2) than the nearer objects. All displays had a frame rate of 30Hz and were symmetrical about both the x- and y-axes.

2.1.5 Procedure. Subjects were told that they would be shown displays of moving objects and that, "Sometimes the objects may appear to be moving towards you; other times you may feel as if you are moving towards the objects. Your task is to press the mouse button down when you feel as if you are moving and hold it down as long as the experience continues. If you don't feel that you are moving then don't press the mouse button" (instructions modified from Andersen and Braunstein, 1985). Subjects were also informed that each display had a fixed duration of 3 minutes and an inter-trial interval of 20 seconds. After two practice trials, the experimental displays were presented in a random order.

2.2 Results

Vection was reported in 166 of the 168 trials (14 subjects responding to 12 stimuli). Of the 2 trials where vection was not induced, 1 had a non-jittering display and the other had a display which jittered in both the x- and y- axes. Separate repeated measures ANOVAs were performed on the onset and duration data. The means are shown in Figures 2A and 2B. Overall, jittering displays were found to produce significantly shorter vection onsets ($F_{1,13} = 11.39$, $p < 0.002$) and significantly longer vection durations ($F_{1,13} = 15.13$, $p < 0.0004$) than non-jittering displays. However, displays with horizontal jitter

did not produce significantly different vection onsets ($F_{1,13} = 0.166$, $p > 0.05$) or durations ($F_{1,13} = 2.055$, $p > 0.05$) from displays with vertical jitter. Similarly, displays which jittered in both directions (horizontal and vertical) did not produce significantly different vection onsets ($F_{1,13} = 0.187$, $p > 0.05$) or durations ($F_{1,13} = 0.857$, $p > 0.05$) from displays which jittered in only one direction.

<INSERT FIGURE 2 ABOUT HERE>

Increasing the simulated speed of forwards self-motion did not significantly effect either vection onsets ($F_{2,26} = 1.034$, $p > 0.05$) or vection durations ($F_{2,26} = 0.240$, $p > 0.05$). No two-way interactions (ie between display type and display speed) reached significance in this experiment.

2.3 *Discussion*

Contrary to prediction, jittering patterns of optic flow were found to induce the most compelling illusions self-motion. Optic flow patterns with global perspective jitter produced shorter vection onsets and longer vection durations than those without. Even when this global perspective jitter occurred in both horizontal and vertical directions (which would have produced the most salient visual-vestibular conflict according to Zacharias and Young's theory), jittering patterns of optic flow were still found to produce more compelling illusions of self-motion. Based on subjects' spontaneous reports⁴, it would appear that both radially expanding and jittering components of the flow were interpreted as self-, not object-, motion. Thus, it would appear that global perspective

jitter was playing a role in the visual perception of self-motion, which overcame any potential impairment produced by its visual-vestibular conflict.

One possible explanation for the advantage of jittering optic flow over non-jittering optic flow, is that the latter, purely radial (expanding) flow, rarely occurs in the real world. Walking, running and even passive transportation usually produce additional random and oscillatory components in the optic flow (Cutting et al 1992). For example, the optic flow presented to a runner will have the following components: (1) a radially expanding component generated by his/her forwards displacement; (2) a vertical sinusoidal component generated by his/her regular up and down displacements; (3) a horizontal sinusoidal component generated by his/her regular side to side displacements; and (4) a random component produced by random horizontal, vertical, and depth displacements (these can become quite significant if the terrain is uneven). Hence, it is possible that jittering optic flow with its additional random (jitter magnitude varied from frame to frame) and oscillatory characteristics (jitter direction - up/down or left/right - alternated from frame to frame) tapped into processes used to perceive self-motion from naturally occurring patterns of optic flow. If so, it might be expected that as jittering optic flow becomes more ecological (ie as jitter range and temporal frequency are reduced), this jitter advantage forvection will increase. This prediction is examined in the following experiments.

3 Experiment 2: Does jitter magnitude effectvection?

Experiment 1 showed that adding global perspective jitter to radially expanding optic flow improves the vection induced in central vision. However, in this experiment, all jittering displays had the same range of random jitter (0-1/3 of the simulated forwards displacement per frame). While jitter of this magnitude could have been produced by real world situations (eg running over uneven terrain, driving a car over an unsealed road, or flying a plane through a region of high turbulence), it is larger than the random flow components accompanying most common self-motions. The present experiment examined whether reducing the range of random jitter (to 0-1/4 or 0-1/5 of the forwards displacement per frame) would increase the ‘jitter advantage’ for vection.

3.1 *Method*

The apparatus, visual displays and procedure were identical to those of experiment 1 - with the sole exception being that two additional jitter ranges were used (0-1/4 and 0-1/5 of the simulated forwards displacement per frame).

3.1.1 *Subjects.* Six male and seven female undergraduate psychology students (aged between 21 and 28 years) participated in this experiment. All had normal or corrected-to-normal vision and had not previously experienced illusions of self-motion in the laboratory.

3.1.2 *Design.* Two independent variables were manipulated in this experiment. (1) *Display Type.* Displays were either jittering or non-jittering patterns of radially

expanding optic flow. When present, global perspective jitter occurred along either the x-axis, the y-axis, or both the x- and y- axes. (2) *Jitter Range*. The absolute amount of horizontal and/or vertical jitter for each frame was randomly selected from one of three ranges: either 0-1/5, 0-1/4, or 0-1/3 of the simulated forwards displacement (all displays simulated a forwards self-motion of 4m/s). This absolute jitter was then altered according to perspective before it was applied to objects at different simulated locations in depth.

3.2 Results

Vection was reported in 155 of the 156 trials (13 subjects responding to 12 stimuli). The only trial that failed to induce vection had a non-jittering display. Separate repeated measures ANOVAs were performed on the onset and duration data. As in experiment 1, jittering displays were found to produce significantly faster vection onsets ($F_{1,12} = 12.225$, $p < 0.001$) and significantly longer vection durations ($F_{1,12} = 24.613$, $p < 0.0001$) than non-jittering displays (see Figures 3A and 3B). While displays with the maximum and middle ranges of jitter (0-1/3 and 0-1/4) did not produce significantly different vection onsets from those with the smallest jitter range (0-1/5) ($F_{1,12} = 0.01$, $p > 0.05$), they did produce significantly longer vection durations ($F_{1,12} = 7.754$, $p < 0.02$). Displays with maximum range of jitter (0-1/3) did not produce significantly different vection onsets ($F_{1,12} = 0.557$, $p > 0.05$) or durations ($F_{1,12} = 0.588$, $p > 0.05$) from those with the middle jitter range (0-1/4). No other main effects or interactions reached significance in this experiment.

<INSERT FIGURE 3 ABOUT HERE>

3.3 *Discussion*

Contrary to the notion that the processes underlying visual self-motion perception might prefer more ecological patterns of optic flow, decreasing the range of the random jitter was found to reduce the ‘jitter advantage’ for vection durations (but not the ‘jitter advantage’ for vection onsets). This finding was also contrary to Zacharias and Young’s (1981) visual-vestibular conflict theory of vection, which predicted that flow patterns with larger ranges of random jitter would impair vection more than those with smaller ranges of random jitter (as the former would produce a greater mismatch between current and expected vestibular signals).

4 Experiment 3: Does jitter frequency effect vection?

Research suggests that visual and vestibular systems are specialized for different types of self-motions - the former being primarily sensitive to low temporal frequency stimulation and constant velocity motion, and the latter being primarily sensitive to high temporal frequency stimulation and brief acceleration (Howard 1986b). While linear vection prefers visual stimulation below 1-0.5 Hz (Berthoz et al 1979; Berthoz et al 1975; Dichgans and Brandt 1978; van Asten et al 1988), vestibular responses to linear acceleration are only elicited at stimulus frequencies above 1 Hz (Diener et al 1982; 1984; Melville-Jones and Young 1978). In the previous experiments, global perspective jitter always occurred 30Hz (ie jitter occurred on each frame of the display). Thus, it is possible that the jitter advantage for vection might be more pronounced for jitter with

lower temporal frequencies (closer to the frequencies preferred by visual self-motion perception). The present experiment examined the effect on vection of reducing jitter frequency from 30 Hz to 1Hz.

4.1 *Method*

The apparatus, visual displays and procedure were identical to those of experiment 2 - with the exception that six temporal frequencies of jitter were used.

4.1.1 *Subjects.* Four male and six female postgraduate psychology students (aged between 23 and 28 years) participated in this experiment. All had normal or corrected-to-normal vision and had not previously experienced illusions of self-motion in the laboratory.

4.1.2 *Design.* Two independent variables were manipulated in this experiment. (1) *Display Type.* Displays were jittering or non-jittering patterns of radially expanding optic flow. (2) *Jitter Frequency.* When present, horizontal and vertical jitter occurred at one of the following temporal frequencies: 1Hz, 2.5Hz, 5Hz, 10Hz, 15Hz or 30Hz. When required, the absolute amount of jitter was randomly selected from the range 0-1/4 of the simulated forwards displacement (all displays simulated a forwards self-motion of 4m/s). This absolute jitter was then altered according to perspective before it was applied to objects at different simulated locations in depth.

4.2 *Results*

Vection was reported in 139 of the 140 trials (10 subjects responding to 14 stimuli). The only trial that failed to induce vection had a non-jittering display. Separate repeated measures ANOVAs were performed on the onset and duration data. Overall, jittering displays were still found to produce significantly shorter vection onsets ($F_{1,9} = 24.137$, $p < 0.0001$) and longer vection durations ($F_{1,9} = 56.326$, $p < 0.0001$) than non-jittering displays (see Figures 4A and 4B). Displays with jitter frequencies of 5 and 10Hz produced significantly longer vection durations than displays with other jitter frequencies ($F_{1,9} = 13.020$, $p < 0.0007$). However, displays with 5 and 10Hz jitter frequencies did not produce significantly different vection onsets from displays with other jitter frequencies ($F_{1,9} = 1.079$, $p > 0.3$).

<INSERT FIGURE 4 ABOUT HERE>

4.3 Discussion

While vection onsets were relatively unaffected by the temporal frequency of the global perspective jitter, jitter frequency did appear to play an important role in vection duration. Specifically, 5-10Hz jitter frequencies were found to produce significantly longer vection durations than higher or lower jitter frequencies. This result would appear to be in stark contrast with the findings of previous studies that linear vection is optimal at low temporal frequencies (ie below 1-0.5Hz - Berthoz et al 1975; 1979). Since reducing the jitter frequency below 5Hz (Experiment 3) and decreasing the range of the random jitter (Experiment 2) both resulted in a diminished jitter advantage, it would appear that jittering displays were not improving vection by mimicking the optic flow patterns produced by naturally occurring self-motions.

5 Conclusions

Contrary to the notion that visual-vestibular conflict impairs vection, the current experiments have demonstrated that jittering patterns of optic flow induce more compelling illusions of self-motion than non-jittering patterns of optic flow (regardless of the direction, range or temporal frequency of the random jitter). It is possible to interpret these findings as support for the notion that visual information dominates the perception of self-motion (ie vestibular information plays a subordinate role in this process - Lishman and Lee 1973; Lee and Lishman 1975). However, it is also possible that any potential impairment produced by visual-vestibular conflict, was obscured by a larger 'jitter advantage'. The latter possibility suggests that the specific characteristics of the visual and vestibular input might determine which source of information dominates the perception of self-motion.

What features of global perspective jitter were responsible for the vection improvements found in the present experiments? One possibility was that global perspective jitter improved vection by obscuring certain artifacts in the computer generated displays. For example, if the visual system was able to resolve the slight relative motions between individual objects in non-jittering displays (produced by their serial - not parallel - production and deletion), this would have reduced/impaired the perception of self-motion (as this would have biased the observer to perceive object-, as opposed to self-, motion). However, the addition of horizontal and/or vertical jitter could have obscured these artifactual motions and hence might have biased the observers - perceiving only the global motion of the objects - towards the perception of self-motion.

This possibility appears remote as jittering and non-jittering displays were designed to avoid 'pixel creep' - both having frame rates of 30Hz.

Global perspective jitter could also have improved vection by reducing adaptation to the optic flow. Consider the time course of the vection induced by a non-jittering pattern of radially expanding optic flow - representing linear self-motion of constant velocity. As the observer adapts to this repetitive and unchanging optic flow, his/her impression of self-motion should continually diminish in magnitude (Denton 1971; Salvatore 1968; Schmidt and Tiffin 1969). However, if a reasonable amount of global perspective jitter is added to the radial flow, it should become more difficult to adapt to this combined flow and hence result in little or no decline in vection over time. Consistent with this account, patterns of optic flow with global perspective jitter were found to produce longer vection durations than those without (experiments 1-3). It should be noted, however, that reduced adaptation alone could not account for the finding that jittering optic flow produced faster vection onsets than non-jittering optic flow - since substantial adaptation to the latter would only occur after vection had been first perceived (eg 30-60s after stimulus onset - Dichgans and Brandt 1978).

A further way that global perspective jitter might have improved vection is by enabling changing-size detectors to extract accurate information about motion in depth from self-motion displays (these contained both motion perspective and changing-size cues to motion in depth). Early research by Regan and Beverley (1978) found evidence for neural mechanisms (changing-size detectors) which are specifically sensitive to changes in retinal image size. However, a later study suggested that these changing-size detectors only provide precise information about motion in depth when the changing-size stimulus

contains frontal plane jitter. Regan and Beverley (1980) noted that during certain trajectories of motion in depth one or other edge of the object remains stationary (or nearly so) as it expands or contracts. They hypothesised that under these circumstances, frontal plane jitter might facilitate the extraction of motion in depth (by removing the restriction of a stationary edge). In different trials, they adapted subjects to stimulus squares which simulated motion along different three-dimensional trajectories: while changing-size cues always simulated the same amount of motion in depth, changing-position cues simulated different amounts of frontal plane motion in different trials. Following this adaptation period, subjects set the stimulus square so that its motion in depth was barely visible. Regan and Beverley found that subjects' final settings were only unbiased by the frontal plane component of adaptation displays when these contained additional (8Hz) frontal plane jitter - suggesting that changing-size detectors require frontal plane jitter if they are to provide accurate estimates of motion in depth. Thus, in the current experiment, jittering patterns of optic flow might have provided additional changing-size based information about the three-dimensional trajectory of the (simulated) self-motion that was not available in non-jittering optic flow, which in turn resulted in a more compelling subjective experiences of self-motion.

In conclusion, the current experiments demonstrate that global perspective jitter can play a significant role in the visual perception of self-motion. Global perspective jitter was found to improve vection induced by optic flow displays, even when corresponding vestibular stimulation was absent. On the basis of these findings, it would appear that purely radial patterns of optic flow are not the optimal inducing stimuli for vection. Rather, it would appear that flow patterns which contain both regular and

random/oscillating components produce the most compelling subjective experiences of self-motion.

References

- Andersen G J, Braunstein M L, 1985 “Induced self-motion in central vision” *Journal of Experimental Psychology: Human Perception & Performance* **11** 122-132
- Benson A J, 1990 “Sensory functions and limitations of the vestibular system” In *Perception and Control of Self-motion* Eds R Warren, A H Wertheim (Hillsdale, New Jersey: Erlbaum) pp 145-170
- Berthoz A, Lacour M, Soechting JF, Vidal PP, 1979 “The role of vision in the control of posture during linear motion” *Progress in Brain Research* **50** 197-209
- Berthoz A, Pavard B, Young LR, 1975 “Perception of linear horizontal self-motion induced by peripheral vision (linearvection)” *Experimental Brain Research* **23** 471-489
- Brandt T, Dichgans J, Büchele W, 1974 “Motion habituation: Inverted self-motion perception and optokinetic after-nystagmus” *Experimental Brain Research* **16** 476-491
- Brandt T, Dichgans J, Koenig E, 1973 “Differential effects of central versus peripheral vision on egocentric and exocentric motion perception” *Experimental Brain Research* **16** 476-491
- Cutting J E, Springer K, Braren P A, Johnson S H, 1992 “Wayfinding on foot from information in retinal, not optical, flow” *Journal of Experimental Psychology: General* **121** 41-72
- Denton G G, 1980 “The influence of visual pattern on perceived speed” *Perception* **9** 393-402

- Dichgans J, Brandt T, 1978 “Visual-vestibular interaction: Effects on self-motion perception and postural control” In *Handbook of Sensory Physiology: Vol. 8. Perception* Eds R Held, H Leibowitz, H L Teuber (New York: Springer-Verlag) pp 755-804
- Diener HC, Dichgans J, Bruzek W, Selinka H, 1982 “Stabilization of human posture during induced oscillations of the body” *Experimental Brain Research* **45** 126-132
- Diener HC, Dichgans J, Guschlbauer B, Mau H, 1984 “The significance of proprioception on postural stabilization as assessed by ischemia” *Brain Research* **296** 103-109
- Gibson J J, 1966 *The senses considered as perceptual systems* (Boston: Houghton Mifflin)
- Giannopulu I, Lepecq J, 1998 “Linear vection chronometry along spinal and sagittal axes in erect man” *Perception* **27** 363-372
- Held R, Dichgans J, Bauer J, 1975 “Characteristics of moving visual scenes influencing spatial orientation” *Vision Research* **15** 357-365
- Howard I P, 1982 *Human visual orientation* (Chichester, Sussex: Wiley) pp 388-398
- Howard I P, 1986a “The vestibular system” In *Handbook of Perception and Human Performance Vol 1: Sensory Processes and Perception* Eds K R Boff, L Kaufman, J P Thomas (New York: Wiley)
- Howard I P, 1986b “The perception of posture, self-motion, and the visual vertical” In *Handbook of Perception and Human Performance Vol 1: Sensory Processes and Perception* Eds K R Boff, L Kaufman, J P Thomas (New York: Wiley)

- Johansson G, 1977 “Studies on the visual perception of locomotion” *Perception* **6** 365-376
- Lee D N, Aronson E, 1974 “Visual proprioceptive control of standing in human infants” *Perception & Psychophysics* **15** 529-532
- Lee D N, Lishman J R, 1975 “Visual proprioceptive control of stance” *Journal of Human Movement Studies* **1** 87-95
- Lishman J R, Lee D N, 1973 “The autonomy of visual kinaesthesia” *Perception* **2** 287-294
- Melcher G A, Henn V, 1981 “The latency of circularvection during different accelerations of the optokinetic stimulus” *Perception & Psychophysics* **30** 552-556
- Melville-Jones G, Young LR, 1978 “Subjective detection of vertical acceleration: A velocity dependent response” *Acta Otolaryngologica* **85** 45-53
- Ohmi M, Howard I P, 1988 “Effect of stationary objects on illusory forward self-motion induced by a looming display” *Perception* **17** 5-12
- Palmisano S, 1996 “Perceiving self-motion in depth: the role of stereoscopic motion and changing-size cues” *Perception & Psychophysics* **58** 1168-1176
- Regan D, Beverley K I, 1978 “Looming detectors in the human visual pathway” *Vision Research* **18** 415-421
- Regan D, Beverley K I, 1980 “Visual responses to changing-size and to sideways motion for different directions of motion in depth: linearization of visual responses” *Journal of the Optical Society of America* **11** 1289-1296
- Riccio GE, Stoffregen TA, 1991 “An ecological theory of motion sickness and postural instability” *Ecological Psychology* **3(3)** 195-240

- Salvatore S, 1968 “Velocity sensing” *Highway Research Record* **292** 79-91
- Schmidt L, Tiffin J, 1969 “Distortion of drivers’ estimates of automobile speed as a function of speed adaptation” *Journal of Applied Psychology* **53** 536-539
- Teixeira R A, Lackner J R, 1979 “Optokinetic motion sickness: Attenuation of visually-induced apparent self-rotation by passive head movements” *Aviation Space and Environmental Medicine* **50** 264-266
- Telford L, Spratley J, Frost B J, 1992 “Linearvection in the central visual field facilitated by kinetic depth cues” *Perception* **21** 337-349
- Telford L, Frost B J, 1993 “Factors affecting the onset and magnitude of linearvection” *Perception & Psychophysics* **53** 682-692
- van Asten WNJC, Gielen CCAM, van der Gon JJD, 1988 “Postural adjustments induced by simulated motion of differently structured environments” *Experimental Brain Research* **73** 371-383
- Wann J, Ruston S, 1994 “The illusion of self-motion in virtual reality environments” *Behavioral and Brain Sciences* **17** 338-340
- Warren W H, Morris M W, Kalish M, 1988 “Perception of translational heading from optical flow” *Journal of Experimental Psychology: Human Perception & Performance* **14** 646-660
- Wong S C P, Frost B J, 1981 “The effect of visual-vestibular conflict on the latency to steady-state visually induced subjective rotation” *Perception & Psychophysics* **30** 228-236
- Young L R, Dichgans J, Murphy R, Brandt T, 1973 “Interaction of optokinetic and vestibular stimuli in motion perception” *Acta oto-laryng* **76** 24-31

Young L R, Oman C M, Dichgans J, 1975 “Influence of head orientation on visually induced pitch and roll sensation” *Aviation, Space and Environmental Medicine* **46** 264-268

Zacharias G L, Young L R, 1981 “Influence of combined visual and vestibular cues on human perception and control of horizontal rotation” *Experimental Brain Research* **41** 159-171

Acknowledgments. Portions of this research were presented at the 1997 Association for Research in Vision and Ophthalmology Conference held in Fort Lauderdale, FL. The authors wish to thank David Regan and William Warren for their feedback and suggestions regarding this article. Correspondence should be addressed to Stephen Palmisano, Human Performance Laboratory, 103 Farquharson Building, 4700 Keele Street, Toronto, Ontario, M3J 1P3, CANADA.

Footnotes

¹Optic flow is defined here as the temporal change in the pattern of light intensities in different directions at the moving point of observation (Gibson 1966; Warren et al 1988).

²The concept of visual-vestibular conflict is not accepted by all researchers. According to Riccio and Stoffregen (1991) there are no situations of sensory conflict, only situations of 'nonredundancy'. Each pattern of multimodal stimulation, regardless of whether it is redundant or nonredundant, represents a specific type of self-motion (ie a possible state of affairs). For example, a nonredundant pattern of multimodal stimulation, which contains visual information that the observer is swaying without corresponding somatosensory information, might specify sway on a nonrigid (as opposed to a rigid) surface.

³'Global perspective jitter' should not be confused with 'global jitter'. While the former represents forwards self-motion on a platform which is oscillating along the horizontal and/or vertical axes, the latter represents forwards self-motion through a sandstorm.

⁴Many subjects spontaneously reported that the experience of self-motion induced by jittering optic flow was similar to walking under the influence of alcohol.

Table and Figure Captions.

A

B

Figure 1: A representation of (A) jittering and (B) non-jittering patterns of radially expanding optic flow.

Figure 1. A representation of (A) jittering and (B) non-jittering patterns of radially expanding optic flow.

Figure 2: The effect of the global perspective jitter on (A) vection onsets and (B) vection durations (Experiment 1). Displays either had no jitter (No), jitter along the x-axis (X), jitter along the y-axis (Y), or jitter along both the x- and y-axes (XY). Error bars represent standard errors of the means.

Figure 2. The effect of global perspective jitter on (A) vection onsets and (B) vection durations (Experiment 1). Displays either had no jitter (No), jitter along the x-axis (X), jitter along the y-axis (Y), or jitter along both the x- and y- axes (XY). Error bars represent standard errors of the means.

Figure 3: The effect of jitter magnitude on (A) vection onsets and (B) vection durations (Experiment 2). Displays either had no jitter (No) or three different jitter ranges (0-1/5, 0-1/4, 0-1/3 of the forwards displacement). The latter jittering displays had jitter along either the x-axis (X), the y-axis (Y), or both the x- and y- axes (XY). Error bars represent the standard errors of the means.

Figure 3. The effect of jitter magnitude on (A) vection onsets and (B) vection durations (Experiment 2). Displays either had no jitter (No) or three different jitter ranges (0-1/5, 0-1/4, 0-1/3 of the forwards displacement). The latter jittering displays had jitter along either the x-axis (X), the y-axis (Y), or both the x- and y- axes (XY). Error bars represent the standard errors of the means.

Figure 4. The effect of jitter temporal frequency on (A) vection onsets and (B) vection durations (Experiment 3). A display either had no jitter (No) or jitter along the x- and y-axes occurring at either 1Hz, 2.5Hz, 5Hz, 10Hz, 15Hz, or 30Hz. Error bars represent the standard errors of the means.

Figure 4. The effect of jitter temporal frequency on (A) vection onsets and (B) vection durations (Experiment 3). A display either had no jitter (No) or jitter along the x- and y-axes occurring at either 1Hz, 2.5Hz, 5Hz, 10Hz, 15Hz, or 30Hz. Error bars represent the standard errors of the means.