

University of Wollongong

## Research Online

---

Faculty of Health and Behavioural Sciences -  
Papers (Archive)

Faculty of Science, Medicine and Health

---

1-1-2012

### Comparison of the DietAdvice website in the clinical trial and primary healthcare contexts

Yasmine Probst

*University of Wollongong, yasmine@uow.edu.au*

Kathryn Boyd

*University Of Newcastle, United Kingdom*

Virva Walkington

*University Of Eastern Finland*

Linda C. Tapsell

*University of Wollongong, ltapsell@uow.edu.au*

Follow this and additional works at: <https://ro.uow.edu.au/hbspapers>



Part of the [Arts and Humanities Commons](#), [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

---

#### Recommended Citation

Probst, Yasmine; Boyd, Kathryn; Walkington, Virva; and Tapsell, Linda C.: Comparison of the DietAdvice website in the clinical trial and primary healthcare contexts 2012, 81.  
<https://ro.uow.edu.au/hbspapers/3196>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: [research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

## Comparison of the DietAdvice website in the clinical trial and primary healthcare contexts

### Abstract

The aim of this study was to evaluate the use of the DietAdvice website, a self-administered online dietary assessment, in the context of a dietary intervention randomized controlled trial [ACTRN 12608000425392]. Data available from a total of n=118 volunteers recruited to the dietary trial and using the website in the clinic at t=0, 3 and 12mo was compared with that obtained from diet histories (DH) and 3-day food record (FR). Fasting blood samples from these time points were also correlated with the macronutrient values obtained. Complete web based records were obtained for 65% of participants at t=0, 83% at t=3 and 75% at t=12mo. At baseline, values for dietary energy (p=0.01), carbohydrate (p=0.036), saturated (p=0.001) and monounsaturated (p=0.014) fat were significantly different for all assessment methods while at 3 months only that for energy (p=0.041) was significantly different. Overall stronger correlations between the macronutrient values from DietAdvice and blood biomarkers were observed at t=12months. In both settings, the data from DietAdvice was most similar to that of the 3-d FR. It appears that trial participants had difficulty using DietAdvice website independently. Compared with data obtained from the primary healthcare setting the limited assistance with the website provided to trial participants may have affected the results.

### Keywords

comparison, healthcare, contexts, clinical, website, trial, dietadvice, primary

### Disciplines

Arts and Humanities | Life Sciences | Medicine and Health Sciences | Social and Behavioral Sciences

### Publication Details

Probst, Y., Boyd, K., Walkington, V. & Tapsell, L. C. (2012). Comparison of the DietAdvice website in the clinical trial and primary healthcare contexts. *Nutrition and Dietetics*, 69 (1), 81.

## **Comparison of the DietAdvice website in the clinical trial and primary healthcare contexts**

**Yasmine Probst<sup>1</sup>**, Kathryn Boyd<sup>2</sup>, Virva Walkington<sup>3</sup>, Linda Tapsell<sup>1</sup>

<sup>1</sup>*University of Wollongong, Australia*

<sup>2</sup>*University of Newcastle, United Kingdom*

<sup>3</sup>*University of Eastern Finland, Finland*

The aim of this study was to evaluate the use of the DietAdvice website, a self-administered online dietary assessment, in the context of a dietary intervention randomized controlled trial [ACTRN 12608000425392]. Data available from a total of n=118 volunteers recruited to the dietary trial and using the website in the clinic at t=0, 3 and 12mo was compared with that obtained from diet histories (DH) and 3-day food record (FR). Fasting blood samples from these time points were also correlated with the macronutrient values obtained. Complete web based records were obtained for 65% of participants at t=0, 83% at t=3 and 75% at t=12mo. At baseline, values for dietary energy (p=0.01), carbohydrate (p=0.036), saturated (p=0.001) and monounsaturated (p=0.014) fat were significantly different for all assessment methods while at 3 months only that for energy (p=0.041) was significantly different. Overall stronger correlations between the macronutrient values from DietAdvice and blood biomarkers were observed at t=12months. In both settings, the data from DietAdvice was most similar to that of the 3-d FR . It appears that trial participants had difficulty using DietAdvice website independently. Compared with data obtained from the primary healthcare setting the limited assistance with the website provided to trial participants may have affected the results.

Contact email - [yasmine@uow.edu.au](mailto:yasmine@uow.edu.au)