

University of Wollongong

Research Online

Faculty of Science, Medicine and Health -
Papers: part A

Faculty of Science, Medicine and Health

1-1-2015

What an anticardiovascular diet should be in 2015

David R. Jacobs Jr

University of Minnesota, Jacobs@epi.umn.edu

Linda C. Tapsell

University of Wollongong, ltapsell@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>



Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Jacobs Jr, David R. and Tapsell, Linda C., "What an anticardiovascular diet should be in 2015" (2015).

Faculty of Science, Medicine and Health - Papers: part A. 3628.

<https://ro.uow.edu.au/smhpapers/3628>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

What an anticardiovascular diet should be in 2015

Abstract

Purpose of review: Given scientific and public debate about optimal diet to prevent cardiovascular disease, and interest in diet and other chronic diseases, we propose that following a few simple dietary principles would reduce chronic disease incidence. **Recent findings:** Nutrition research has been criticized for focusing on individual nutrients and foods, treated like drug therapy. With a few important exceptions, clinical trials of supplemental nutrients have not shown benefit. Although highly specific nutrition information is elusive, diet patterns have provided consistent answers, important for public health. Observational cohort studies have found that some dietary patterns are reported with high reliability over long periods and predict future cardiovascular and other inflammatory-related diseases. Two randomized clinical trials confirmed this finding. There are many common features of Mediterranean and prudent diets, particularly the plant-centered aspect, coupled with variety of foods eaten. A dietary pattern characterized by high fruit, vegetable, legume, whole grain, nut, berry, seed, and fish intakes, and possibly by intakes of dairy, coffee, tea, chocolate, and alcohol (not in excess), but low meat and detrimentally processed foods is associated with reduced incidence of cardiovascular disease and rates of noncardiovascular, noncancer chronic inflammatory-related mortality. **Summary:** A plant-centered diet may be broadly recommended.

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Jacobs Jr, D. R. & Tapsell, L. C. (2015). What an anticardiovascular diet should be in 2015. *Current Opinion in Lipidology*, 26 (4), 270-275.

What an anti-cardiovascular diet should be in 2015

David R. Jacobs, Jr., PhD¹ and Linda C. Tapsell, PhD²

¹Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, Minneapolis, MN 55454, USA

²School of Health Sciences, University of Wollongong, Wollongong, NSW 2522, Australia

Abstract

Purpose of review: Given scientific and public debate about optimal diet to prevent cardiovascular disease, and interest in diet and other chronic diseases, we propose that following a few simple dietary principles would reduce chronic disease incidence.

Recent findings: Nutrition research has been criticized for focusing on individual nutrients and foods, treated like drug therapy. With a few important exceptions, clinical trials of supplemental nutrients have not shown benefit. Although highly specific nutrition information is elusive, diet patterns have provided consistent answers, important for public health. Observational cohort studies have found that some dietary patterns are reported with high reliability over long periods and predict future cardiovascular and other inflammatory-related diseases. Two randomized clinical trials confirmed this finding. There are many common features of Mediterranean and prudent diets, particularly the plant-centered aspect, coupled with variety of foods eaten. A dietary pattern characterized by high fruit, vegetable, legume, whole grain, nut, berry, seed, and fish intakes, and possibly to intakes of dairy, coffee, tea, chocolate, and alcohol (not in excess), but low meat and detrimentally processed foods is associated with reduced incidence of cardiovascular disease and rates of non-cardiovascular, non-cancer chronic inflammatory-related mortality.

Summary: A plant-centered diet may be broadly recommended.

Keywords: chronic disease, cardiovascular disease, diet patterns, plant-centered diet

Introduction

There is substantial scientific and public debate about optimal diet to prevent cardiovascular and other chronic disease. US News and World Report captures this debate annually in “Best Diets”, which in 2015 summarized the rankings by a panel of nutrition professionals of 35 different diet plans¹. The US Dietary Guidelines have made nutrition recommendations, which for many years emphasized reduction in total and saturated fat as one cornerstone, often tending to avoid reference to specific foods^{2,3}. Many people have recommended various forms of low carbohydrate diets (for example, the Atkins diet). Many ways to view “what to eat” have been formalized.

Concurrently, epidemiology has been severely criticized⁴, particularly observational nutritional epidemiology⁵, but also the overwhelming body of detailed studies of nutrition mechanisms⁶. Temple⁶ classified research as complex (“the detailed study of disease mechanisms using such methods as biochemistry and molecular genetics”) or simple (“epidemiology, intervention trials, and analogous studies on animals”) and commented that complex research has received most resources, but more practical value has come from simple research. Archer et al⁷ found that nutritional surveillance in the US is seriously flawed because energy intake is clearly underestimated. They concluded that “the ability to estimate population trends in caloric intake and generate empirically supported public policy relevant to diet-health relationships from U.S. nutritional surveillance is extremely limited”, thereby eliminating all nutritional epidemiology that might be generated from this source. Ioannidis⁵ stated, “Definitive solutions will not come from another million observational papers or a few small randomized trials.” He found a poor track record of observational claims, but lauded two long term randomized clinical trials of dietary patterns^{8,9}.

Some commentators have focused on “nutritionism”^{10,11}, a formulation in which nutrition decisions and policies are focused largely on nutrients, with little or no consideration of the complex nature of foods. Nutritionism at least implicitly supports the financially thriving supplement industry. Several isolated food chemicals have been tested as drugs in randomized clinical trials, with no effect or a harmful effect on long term health¹². A recent clinical trial of niacin plus the anti-flushing agent laropiprant found adverse effects for infection, bleed, and diabetes, among other effects¹³. Exceptions relate to age-related macular degeneration¹⁴ and maternal folate supplementation for prevention of fetal neural tube defects¹⁵ showed benefit. Similarly, viewing diet through the lens of a single nutrient (e.g., low fat, low saturated fat, low carbohydrate) is likely to discard good foods as well as less desirable ones. For example, among highly regarded foods, nuts are high in fat and whole grain foods and all fruits and vegetables are high in carbohydrates. Dairy products are high in saturated fat, but findings related to dairy foods and various outcomes suggest benefit¹⁶.

Therefore concepts about the best diet for long term cardiovascular health have been changing. This review reconsiders diet, focusing on dietary patterns and long term chronic disease outcomes.

A Multi-Dimensional View of Food

Food is complex¹⁷. There are major differences between drug and food research^{17,18}. Most salient is the nature of a drug vs food. Drugs are well-defined, purified compounds which can be unobtrusively consumed and compared to another well-defined regimen, such as a placebo or accepted therapy. Foods are the opposite of well-defined and constant: we eat formerly living organisms that have all the complexity and variety of life forms, nurtured in various ways by farmers, breeders, and genetic modifiers, consumed several times per day, every day, one food balanced against another to maintain energy balance, and affecting chronic disease as some sort of average across foods and time. Many authorities require the internal validity of long term, randomized, clinical trials characteristically used in inferences about drugs before they consider evidence to be of the most persuasive nature. Such trials are difficult at best for study of food, yet it is critical to appraise health value of foods. To this end, several authors have favored reappraisal of the body of nutrition studies with a view towards valid inference; and a more holistic view of nutrition¹⁷⁻²¹, in which dietary policy and even understanding of nutrition itself does not need a meticulous and reductionist breakdown into separate constituent parts¹⁹.

Reductionist science employs syllogisms. An example is the logic underlying avoidance of saturated fat. A syllogism considers 3 propositions, A, B, and C. It asserts that if A being true implies that B is true, and B being true implies that C is true, then it follows without further testing that A being true implies that C is true. The problem of judging the validity of C is reduced to solutions of easier problems. In the particular case, the proposition that intake of saturated fat raises serum cholesterol is strongly supported (A implies B)²²⁻²⁴. The proposition that cholesterol in the serum leads to atherosclerotic disease (B implies C) is also strongly supported by many kinds of studies, for example by presence of cholesterol in atherosclerotic plaque^{25,26}, observational epidemiology^{27,28}, and the success of statin drugs (which dramatically reduce low density lipoprotein cholesterol, among other things) in reducing coronary heart disease and total mortality²⁹. That coronary heart disease reduction also follows serum cholesterol reduction even when total mortality is not reduced³⁰ strengthens the proposition that cholesterol in the serum plays a role in atherosclerotic disease, because it suggests that cholesterol reduction is specific to coronary heart disease, whatever else might eventuate from serum cholesterol lowering. Yet this “efficient” logic is not necessarily correct. Saturated fat is not the only component of saturated fat-containing food; other food components might be important. Many biological changes occur with cholesterol changes, particularly considering the pleiotropic nature of statins, and these also could be important. Direct observation of the association between saturated fat and coronary heart disease did not find the association of saturated fat intake with coronary heart which as expected by the syllogism³¹. Such syllogisms are standard in science, for linking and interpreting chains of evidence, but they should be used cautiously because they are not always right.

In contrast, direct observation of certain dietary patterns with various cardiovascular and other chronic diseases has consistently shown reduced risk³²⁻³⁶, although meta-analysis of specific foods and nutrients was found to be confusing³⁷. This difference in findings possibly relates to the stability of diet patterns over time and reliability of reporting. Jacobs and Orlich¹⁹ pointed out that in most cases diet patterns derived from food frequency questionnaires had high correlation within person over periods of years; two studies reported tracking correlation over 0.5 over 18 and 20 years. Correlations over time of most foods and nutrients are much lower, on the order of 0.2 or less, although a few well-defined foods that are habitually consumed by some

people, such as coffee, tea, or alcohol, have high tracking correlations. In addition, diet patterns integrate over foods eaten and over long periods, whereas individual foods and nutrients do not capture the sum of complex nutrition effects.

Mediterranean and Prudent Diet Patterns That Protect against Chronic Disease

The central focus of diet pattern research has been Mediterranean or prudent diets with various definitions³². In an early article studying middle aged male health professionals, Hu et al.³³ performed principal components analysis and found 2 patterns: “prudent”, with high weightings on green leafy vegetables, dark-yellow vegetables, cruciferous vegetables, other vegetables, legumes, fruit, tomatoes, fish, garlic, poultry, and whole grains; and “western”, with high weightings on red meat, processed meat, refined grains, sweets and desserts, french fries, high-fat dairy products, eggs, high-sugar drinks, snacks, condiments, margarine, potatoes, and butter. The prudent diet was inversely related to incident coronary heart disease, while the western diet was adversely related. This list of foods closely resembles the DASH diet, in which vegetables, fruit, fruit juice, whole grains, nuts, seeds, legumes, and dairy products other than whole milk were increased, and meat, poultry, fish, refined grain, whole milk, sweets, snack foods, fats, oils, and salad dressings were decreased³⁸.

The Prevención con Dieta Mediterránea (PREDIMED) study⁹ instituted a 14 point Mediterranean diet compliance score. Foods favored are olive oil, vegetables, fruit (including natural juices), wine, legumes, fish or shellfish, nuts (including peanuts), chicken, turkey or rabbit meat instead of veal, pork, hamburger or sausage, and sofrito (a sauce made with tomato and onion, leek, or garlic, simmered with olive oil). Foods not favored are red meat, hamburger, or meat products (ham, sausage, etc.), butter, margarine, or cream, sweet/carbonated beverages, and commercial sweets or pastries (not homemade), such as cakes, cookies, biscuits, or custard. Participants randomized to Mediterranean diet groups had about 2 out of 14 points higher score than the randomized control group, starting soon after baseline and throughout the study (Table S1 and Figure S3 in reference 9). Higher baseline scores predicted cardiovascular disease outcomes³⁹.

Mursu et al³⁶ studied various mortality outcomes in the Iowa Women’s Health Study (IWHS) according to two diet scores, the A Priori Diet Quality Score (APDQS)⁴⁰⁻⁴⁵ and the Alternative Healthy Eating Index 2010 (AHEI)⁴⁶.

The APDQS originated in a case control study to differentiate myocardial infarction from healthy controls according to combinations of foods⁴⁰. It was applied in several studies subsequently, showing relationships with demographics⁴¹, several biomarkers⁴²⁻⁴⁵, cognitive function⁴⁶, and fitness⁴⁷. It has been reformulated for different studies, depending on the questionnaire data available. Approximately 40 food groups, covering many aspects of diet, are rated as favorable, adverse or neutral and placed in quantiles according to intake. For example, in the IWHS³⁶, 34 food groups were rated favorable (n=17), adverse (n=10), or neutral (n=7) relative to future chronic disease, based on the literature and expert judgment, then placed in quartiles (or a large nonconsumer group and tertiles among consumers). Mursu et al³⁶ stated: “The positively rated food groups were beans and legumes, beer, coffee, fish, fruit, green vegetables, low-fat dairy, liquor, oil, other vegetables, poultry, seeds, and nuts, soy products, tea,

tomato, whole grains, and wine. The negatively rated food groups were butter, fried foods, fried potatoes, red meat, liver, processed meat, salty snacks, soft drinks, sweets, and whole-fat dairy. The remaining neutrally rated food groups were chocolate, diet soft drinks, eggs, fruit juice, margarine, potatoes, and refined grains.”

Groups were given scores of 0 for lowest quartile of consumption or nonconsumers to 3 for highest consumption grouping. The APDQS was formed by adding these values in a positive direction for beneficially-rated food groups, in a negative direction for adversely-rated food groups, and 0 for neutrally-rated food groups, for a theoretical maximal score of 81. Thus 1 point in the APDQS corresponds to a difference of one higher intake category in beneficially-rated food groups or one lower intake category in adversely-rated food groups. Mean±standard deviation of the APDQS was 38.4±8.2 in the 29,634 women aged 55-69 years and free of self-reported heart disease, diabetes, and cancer at baseline in 1986 and followed for 22 years.

The AHEI-2010⁴⁸ was formulated as an improvement to an earlier similar score that provided dietary guidance related to the US Dietary Guidelines. AHEI-2010 contains 11 components, each of which receives 0-10 points (theoretical maximum 110). Components included vegetables, whole fruit, whole grains, nuts and legumes, red and processed meat, long-chain (n23) fatty acids EPA+DHA, PUFAs, and moderate alcohol intake (all rated favorably) and trans fatty acids, sugar-sweetened beverages and fruit juices, and sodium (all rated adversely). In the IWHS, the mean AHEI-2010 was 40.0±10.4, with correlation of 0.65 between the AHEI-2010 and the APDQS.

Both the APDQS and AHEI-2010 predicted total, cardiovascular, cancer, and non-cardiovascular, non-cancer inflammatory-related mortality, but not other and external causes of mortality. Prediction was numerically strongest for inflammatory-related mortality and weakest for cancer mortality, but statistically significant for all. The risk factor adjusted total mortality rates were 36.9, 34.6, 32.5, and 30.2 per 100 women followed for 22 years across the increasing APDQS quartiles, and similarly for the AHEI-2010. We regard a difference of 36.9 vs 30.2 as of considerable clinical importance in women initially average age 62 years, followed into old age. Furthermore, the women were resurveyed in 2004 (n=15,076) at average age 79 years, and findings were similar or numerically stronger for diet score prediction of all outcomes.

One of the goals of Mursu et al³⁶ was to identify information in each score beyond what was found in the other score. Each score did add some prediction when the other was in the model, but it was difficult to sort out what that information was. More to the point for this review is the commonality between these scores and of these scores with other prudent and Mediterranean diet patterns.

Conclusion

The many diet pattern analyses cited here, though consistent across many settings and having substantial commonality across patterns studied, are observational and therefore considered insufficient in themselves to make causal statements about diet protecting against future chronic disease. However, coupled with the randomized findings of the Lyon Diet-Heart Study⁸ and PREDIMED⁹, we assert that certain diet patterns can prevent or delay a substantial amount of

currently existing chronic disease, particularly cardiovascular and non-cardiovascular, non-cancer inflammatory-related diseases. Such diet patterns are plant-centered, at the same time minimizing intake of nutritionally poor plant foods such as sugar, refined grains and highly processed and very salty foods. Intake of plant seeds, including whole grains, nuts, berries, coffee, and chocolate, is promising⁵⁰. The recommended prudent/Mediterranean diet patterns are highly varied, as can be gleaned from the small increments to the APDQS from 34 widely varying food groups. Apparently there are many ways to achieve a protective diet pattern.

Excellent dietary patterns are not necessarily vegetarian, although much of the meat in the diet is rated poorly in all patterns. Meat is less sustainable than other foods, given the energy cost of growing meat products of all types⁵¹. Fish is highly recommended across all diet patterns, but the fish stock is substantially polluted with heavy metals and other industrial chemicals; this is an ongoing problem^{52,53}. These are difficulties for the future.

We may not ever know the more specific details that have been so earnestly sought in many, many nutrition studies. Indeed, the diet pattern approach to nutrition recommendations taken here leaves many questions unanswered. For example, alcohol intake may be over-endorsed in the APDQS, since points are given for each of wine, beer, and liquor separately and with no upper limit for excessive intake, as opposed to the moderate amount of alcohol recommended in the AHEI-2010 or the wine with meals recommended in PREDIMED. There is evidence that would support rating chocolate as favorable rather than neutral⁵⁴. Furthermore, it is not easy to know about separate effects of food groups (as in APDQS) or dietary principles (as in AHEI-2010). For example, prediction of CVD events in PREDIMED from the 14 individual score components is not always consistent with findings for prediction from the whole score, but individual components do not take the whole diet into account and are subject to confounding with the rest of the diet³⁹. The treatment groups themselves in PREDIMED⁹ and the Lyon Diet Heart Study⁸ are not specific, but rather are highly multi-faceted. In PREDIMED it is not possible to fully sort out whether the benefit was from substituting extra virgin olive oil for the customarily consumed and less expensive refined olive oil, from the nut mixture, or from the overall Mediterranean diet. Whether findings would have been different if a different control regimen had been used is not discernible from the PREDIMED trial.

However, the public and public health researchers do not necessarily need to know details in order to eat and recommend diet patterns that improve public health. The small contribution of any one food to the total score enhances robustness of the scores and supports enjoyment of a varied diet as a major aspect of eating. Thus Pollan's² 7 word aphorism and our doubling of it to 14 words¹⁹, "eat food, mostly plants, not too much, in colorful variety, maximizing nutrients per bite" is a simple approach to healthy and enjoyable eating, which will also tend to enhance sustainability of the food system.

Key Points

1. Much of the dietary advice over the past decades to prevent cardiovascular disease has focused on reduction of saturated fat intake; coronary heart disease mortality decreased over this period. The low saturated fat message altered meat and dairy intake. Nevertheless, this message appears not to be correct in all respects and to have been too restrictive. It missed important focuses on nutritionally-rich plant food intake.

2. Reductionist science operates through syllogisms, which can be misleading. In the case of saturated fat, it fostered a view of nutrition that was unnecessarily restrictive. For dietary supplements, it fostered growth of the dietary supplement industry, which promotes numerous products which have been shown in long-term randomized clinical trials to have no effect or even adverse effects on chronic disease risk.
3. There are many excellent approaches to a diet that prevents chronic disease, but such approaches do not depend on total fat or total carbohydrate intake.
4. Diets that are associated with reduced chronic disease risk and have been successful in long-term randomized clinical trials are characterized by high fruit, vegetable, legume, whole grain, nut, berry, seed, unrefined unsaturated oils, and fish intakes, and more speculatively to intakes of dairy, coffee, tea, chocolate, and alcohol (not in excess), but are low in meat and detrimentally processed foods.

Acknowledgements: D. R. J. is supported by diverse grants from the National Institutes of Health and the Environmental Protection Agency in the USA and is a member of the Scientific Advisory Board of the California Walnut Commission. L. C. T. is supported by grants from the Australian Research Council and the Illawarra Health and Medical Research Institute and is an unpaid member of the Scientific Advisory Board of the California Walnut Commission, and a member of the Science Advisory Committee of the McCormick Science Institute. There was no specific funding received for this paper. D. R. J. wrote the first draft of this paper. L. C. T. performed the critical review and editing.

References

1. US News and World Report. Best Diets 2015. <http://health.usnews.com/best-diet> Accessed January 26, 2015
2. US Department of Agriculture National Agricultural Library. Historical Dietary Guidelines. <http://fnic.nal.usda.gov/dietary-guidance/dietary-guidelines/historical-dietary-guidance> Accessed January 26, 2015
3. US Department of Health and Human Services. History of Dietary Guidelines for Americans. <http://www.health.gov/dietaryguidelines/history.htm> Accessed January 26, 2015
4. Taubes G. Epidemiology faces its limits. *Science* 1995;269:164–9.
5. Ioannidis JP. Implausible results in human nutrition research. *BMJ* 2013;347:f6698.
6. Temple NJ. Nutrition and disease: challenges of research design. *Nutrition*. 2002 Apr;18(4):343-7.
7. Archer E, Hand GA, Blair SN (2013) Validity of U.S. Nutritional Surveillance: National Health and Nutrition Examination Survey Caloric Energy Intake Data, 1971–2010. *PLoS ONE* 8(10): e76632.
8. De Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999;99:779-85.

9. Estruch R, Ros E, Salas-Salvadó J, Covas M-I, Corella D, Arós F, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 2013;368:1279-90.
10. Scrinis G. *Nutritionism: the science and politics of dietary advice*. New York, NY: Columbia University Press, 2013.
11. Pollan M. *In defense of food: an eater's manifesto*. New York, NY: The Penguin Press, 2008.
12. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis. *JAMA* 2007;297:842-57
13. HPS2-THRIVE Collaborative Group, Landray MJ, Haynes R, Hopewell JC, Parish S, Aung T, Tomson J, Wallendszus K, Craig M, Jiang L, Collins R, Armitage J. Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med*. 2014 Jul 17;371(3):203-12.
14. Age-Related Eye Disease Study Research Group. A randomized, placebo-controlled, clinical trial of high-dose supplementation with vitamins C and E, beta carotene, and zinc for age-related macular degeneration and vision loss: AREDS report no. 8. *Arch Ophthalmol*. 2001;119(10):1417-1436.
15. Kim YI. Folic acid fortification and supplementation—good for some but not so good for others. *Nutr Rev* 2007;65:504-11.
16. Weng LC, Steffen LM, Szklo M, Nettleton J, Chambless L, Folsom AR. A diet pattern with more dairy and nuts, but less meat is related to lower risk of developing hypertension in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) study. *Nutrients*. 2013 May 21;5(5):1719-33.
17. Jacobs DR Jr, Tapsell LC, Temple NJ. Food synergy: the key to balancing the nutrition research effort. *Public Health Rev* 2012;33:507-29.
18. Satija A, Yu E, Willett WC, Hu FB. Understanding Nutritional Epidemiology and Its Role in Policy. *Adv Nutr* 2015;6:1-14.
19. Jacobs DR Jr, Orlich MJ. Diet pattern and longevity: do simple rules suffice? A commentary. *Am J Clin Nutr*. 2014 May 28;100(Supplement 1):313S-319S.
20. Fardet A, Rock E. Toward a new philosophy of preventive nutrition: from a reductionist to a holistic paradigm to improve nutritional recommendations. *Adv Nutr*. 2014 Jul 14;5(4):430-46.
21. Mozaffarian D, Ludwig DS. Dietary guidelines in the 21st century--a time for food. *JAMA*. 2010 Aug 11;304(6):681-2.
22. Keys A, Anderson JT, Grande F. Serum cholesterol response to changes in the diet, IV: particular saturated fatty acids in the diet. *Metabolism* 1965;14:776-84.
23. Mensink RP, Katan MB. Effect of dietary trans fatty acids on high-density and low-density lipoprotein cholesterol levels in healthy subjects. *N Engl J Med* 1990;323:439-45.

24. Hegsted DM. A look back at lessons learned and not learned. *J Nutr* 1994;124(suppl):1867S–70S.
25. Sabine J: Cholesterol. New York, Marcel Dekker, Inc, 1977
26. Wissler RW. New insights into the pathogenesis of atherosclerosis as revealed by PDAY. *Pathobiological Determinants of Atherosclerosis in Youth*. Atherosclerosis. 1994 Aug;108 Suppl:S3-20.
27. Jacobs D, Blackburn H, Higgins M, Reed D, Iso H, McMillan G, Neaton J, Nelson J, Potter J, Rifkind B, et al. Report of the Conference on Low Blood Cholesterol: Mortality Associations. *Circulation*. 1992 Sep;86(3):1046-60.
28. Prospective Studies Collaboration, Lewington S, Whitlock G, Clarke R, Sherliker P, Emberson J, Halsey J, Qizilbash N, Peto R, Collins R. Blood cholesterol and vascular mortality by age, sex, and blood pressure: a meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. *Lancet*. 2007 Dec 1;370(9602):1829-39.
29. Ebrahim S, Taylor FC, Brindle P. Statins for the primary prevention of cardiovascular disease. *BMJ*. 2014 Jan 27;348:g280.
30. Gould AL, Rossouw JE, Santanella NC, et al. Cholesterol reduction yields clinical benefit A new look at old data. *Circulation* 1995;91:2274-82.
31. Hu FB, Stampfer MJ, Manson JE, Rimm E, Colditz GA, Rosner BA, Hennekens CH, Willett WC. Dietary fat intake and the risk of coronary heart disease in women. *N Engl J Med*. 1997 Nov 20;337(21):1491-9.
32. Trichopoulou A, Martínez-González MA, Tong TY, Forouhi NG, Khandelwal S, Prabhakaran D, Mozaffarian D, de Lorgeril M. Definitions and potential health benefits of the Mediterranean diet: views from experts around the world. *BMC Med*. 2014 Jul 24;12:112.
33. Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC. Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr* 2000;72:912–21.
34. Stradling C, Hamid M, Taheri S, Thomas GN. A review of dietary influences on cardiovascular health: part 2: dietary patterns. *Cardiovasc Hematol Disord Drug Targets*. 2014;14(1):50-63.
35. Martinez-Gonzalez MA, Bes-Rastrollo M. Dietary patterns, Mediterranean diet, and cardiovascular disease. *Curr Opin Lipidol*. 2014 Feb;25(1):20-6.
36. Mursu J, Steffen LM, Meyer KA, Duprez D, Jacobs DR Jr. Diet quality indexes and mortality in postmenopausal women: the Iowa Women's Health Study. *Am J Clin Nutr*. 2013 Aug;98(2):444-53.
37. Berciano S, Ordovás JM. Nutrition and cardiovascular health. *Rev Esp Cardiol (Engl Ed)*. 2014 Sep;67(9):738-47.
38. Jacobs DR Jr, Gross MD, Steffen L, Steffes MW, Yu X, Svetkey LP, Appel LJ, Vollmer WM, Bray GA, Moore T, Conlin PR, Sacks F. The effects of dietary patterns on urinary

- albumin excretion: results of the Dietary Approaches to Stop Hypertension (DASH) Trial. *Am J Kidney Dis.* 2009 Apr;53(4):638-46.
39. Schröder H, Salas-Salvadó J, Martínez-González MA, Fito M, Corella D, Estruch R, Ros E. Baseline adherence to the Mediterranean diet and major cardiovascular events: Previsión con Dieta Mediterránea trial. *JAMA Intern Med.* 2014 Oct;174(10):1690-2.
 40. Lockheart MS, Steffen LM, Rebnord HM, Fimreite RL, Ringstad J, Thelle DS, Pedersen JI, Jacobs DR Jr. Dietary patterns, food groups and myocardial infarction: a case-control study. *Br J Nutr* 2007;98:380-7.
 41. Sijtsma FP, Meyer KA, Steffen LM, Shikany JM, Van Horn L, Harnack L, Kromhout D, Jacobs DR Jr. Longitudinal trends in diet and effects of sex, race, and education on dietary quality score change: the Coronary Artery Risk Development in Young Adults study. *Am J Clin Nutr* 2012;95:580-6.
 42. Jacobs DR Jr, Sluik D, Rokling-Andersen MH, Anderssen SA, Drevon CA. Association of 1-y changes in diet pattern with cardiovascular disease risk factors and adipokines: results from the 1-y randomized Oslo Diet and Exercise Study. *Am J Clin Nutr* 2009;89:509-17.
 43. Meyer KA, Sijtsma FP, Nettleton JA, Steffen LM, Van Horn L, Shikany JM, Gross MD, Mursu J, Traber MG, Jacobs DR Jr. Dietary patterns are associated with plasma F2-isoprostanes in an observational cohort study of adults. *Free Radic Biol Med* 2013;57:201-9.
 44. Sijtsma FP, Meyer KA, Steffen LM, Van Horn L, Shikany JM, Odegaard AO, Gross MD, Kromhout D, Jacobs DR Jr. Diet quality and markers of endothelial function: the CARDIA study. *Nutr Metab Cardiovasc Dis.* 2014 Jun;24(6):632-8.
 45. Nettleton JA, Schulze MB, Jiang R, Jenny NS, Burke GL, Jacobs DR Jr. A priori-defined dietary patterns and markers of cardiovascular disease risk in the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr* 2008;88:185-94.
 46. Zhu N, Jacobs DR, Meyer KA, He K, Launer L, Reis JP, Yaffe K, Sidney S, Whitmer RA, Steffen LM. Cognitive Function In a Middle Aged Cohort Is Related to Higher Quality Dietary Pattern 5 and 25 Years Earlier: The CARDIA Study. *J Nutr Health Aging.* 2015;19(1):33-8.
 47. Shikany JM, Jacobs DR Jr, Lewis CE, Steffen LM, Sternfeld B, Carnethon MR, Richman JS. Associations between food groups, dietary patterns, and cardiorespiratory fitness in the Coronary Artery Risk Development in Young Adults study. *Am J Clin Nutr.* 2013 Dec;98(6):1402-9.
 48. Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, Stampfer MJ, Willett WC. Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr* 2012;142:1009-18.
 49. McCullough ML, Feskanich D, Stampfer MJ, Giovannucci EL, Rimm EB, Hu FB, Spiegelman D, Hunter DJ, Colditz GA, Willett WC. Diet quality and major chronic

- disease risk in men and women: moving toward improved dietary guidance. *Am J Clin Nutr* 2002;76:1261–71.
50. Ros E, Hu FB. Consumption of plant seeds and cardiovascular health: epidemiological and clinical trial evidence. *Circulation*. 2013 Jul 30;128(5):553-65.
 51. Sabaté J, Soret S. Sustainability of plant-based diets: back to the future. *Am J Clin Nutr*. 2014 Jun 4;100(Supplement 1):476S-482S.
 52. Nøstbakken OJ, Hove HT, Duinker A, Lundebye AK, Berntssen MH, Hannisdal R, Lunestad BT, Maage A, Madsen L, Torstensen BE, Julshamn K. Contaminant levels in Norwegian farmed Atlantic salmon (*Salmo salar*) in the 13-year period from 1999 to 2011. *Environ Int*. 2015 Jan;74:274-80.
 53. Ruzzin J, Bethune C, Goksøyr A, Hylland K, Lee DH, Jacobs DR Jr, Carpenter DO. Comment on "Contaminant levels in Norwegian farmed Atlantic salmon (*Salmo salar*) in the 13-year period from 1999 to 2011" by Nøstbakken et al. *Environ Int*. 2015 Jan 16. pii: S0160-4120(15)00007-0. doi: 10.1016/j.envint.2015.01.003. [Epub ahead of print]
 54. Latham LS, Hensen ZK, Minor DS. Chocolate--guilty pleasure or healthy supplement? *J Clin Hypertens (Greenwich)*. 2014 Feb;16(2):101-6.