



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers

Faculty of Science, Medicine and Health

2016

Vegetarian and omnivorous nutrition - comparing physical performance

Joel Craddock

University of Wollongong, jcc256@uowmail.edu.au

Yasmine Probst

University of Wollongong, yasmine@uow.edu.au

Gregory E. Peoples

University of Wollongong, peoples@uow.edu.au

Publication Details

Craddock, J. C., Probst, Y. C. & Peoples, G. E. (2016). Vegetarian and omnivorous nutrition - comparing physical performance. *International Journal of Sport Nutrition and Exercise Metabolism*, 26 (3), 212-220.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Vegetarian and omnivorous nutrition - comparing physical performance

Abstract

Humans consuming vegetarian-based diets are observed to have reduced relative risk for many chronic diseases. Similarly, regular physical activity has also been shown to assist in preventing, and reducing the severity of these conditions. Many people, including athletes, acknowledge these findings and are adopting a vegetarian-based diet to improve their health status. Furthermore, athletes are incorporating this approach with the specific aim of optimising physical performance. To examine the evidence for the relationship between consuming a predominately vegetarian-based diet and improved physical performance a systematic literature review was performed using the SCOPUS database. No date parameters were set. The keywords; 'vegetarian* OR vegan* AND sport* OR athlete* OR training OR performance OR endurance' were used to identify relevant literature. Included studies; (i) directly compared a vegetarian-based diet to an omnivorous/mixed diet, (ii) directly assessed physical performance, not biomarkers of physical performance, (iii) did not use supplementation emulating a vegetarian diet. Reference lists were hand searched for additional studies. Seven randomised controlled trials and one cross-sectional study met the inclusion criteria. No distinguished differences between vegetarian-based diets and omnivorous mixed diets were identified when physical performance was compared. Consuming a predominately vegetarian-based diet did not improve nor hinder performance in athletes. However, with only 8 studies identified, with substantial variability amongst the studies' experimental designs, aims and outcomes, further research is warranted

Publication Details

Craddock, J. C., Probst, Y. C. & Peoples, G. E. (2016). Vegetarian and omnivorous nutrition - comparing physical performance. *International Journal of Sport Nutrition and Exercise Metabolism*, 26 (3), 212-220.

Vegetarian and omnivorous nutrition – Comparing physical performance

Joel Craddock¹, Dr Yasmine Probst¹, Dr Greg E Peoples².

School of Medicine, University of Wollongong, Northfields Avenue,
Wollongong, NSW 2522, Australia.

Corresponding author and requests for reprints:

Joel Craddock

School of Medicine, University of Wollongong, Northfields Avenue,
Wollongong, NSW 2522, Australia.

Email: jcc256@uowmail.edu.au

Phone: +61422369633

Running Title – Vegetarian-based nutrition and physical performance

Vegetarian and omnivorous nutrition – Comparing physical performance

Abstract: Humans consuming vegetarian-based diets are observed to have reduced relative risk for many chronic diseases. Similarly, regular physical activity has also been shown to assist in preventing, and reducing the severity of these conditions. Many people, including athletes, acknowledge these findings and are adopting a vegetarian-based diet to improve their health status. Furthermore, athletes are incorporating this approach with the specific aim of optimising physical performance. To examine the evidence for the relationship between consuming a predominately vegetarian-based diet and improved physical performance a systematic literature review was performed using the SCOPUS database. No date parameters were set. The keywords; vegetarian* OR vegan* AND sport* OR athlete* OR training OR performance OR endurance' were used to identify relevant literature. Included studies; (i) directly compared a vegetarian-based diet to an omnivorous/mixed diet, (ii) directly assessed physical performance, not biomarkers of physical performance, (iii) did not use supplementation emulating a vegetarian diet. Reference lists were hand searched for additional studies. Seven randomised controlled trials and one cross-sectional study met the inclusion criteria. No distinguished differences between vegetarian-based diets and omnivorous mixed diets were identified when physical performance was compared. Consuming a predominately vegetarian-based diet did not improve nor hinder performance in athletes. However, with only 8 studies identified, with substantial variability amongst the

studies' experimental designs, aims and outcomes, further research is warranted.

Keywords – Vegan, vegetarian, sport

Introduction

A range of vegetarian diets exist, all of which are typically plant-based and are often classified on the exclusion or inclusion of animal, or animal derived products. Vegan, pescovegetarian, ovo-vegetarian, lacto-vegetarian and lacto-ovo-vegetarian (LOV) diets are examples of vegetarian-based diets in which fruits, vegetables, grains, nuts, seeds and legumes represent a high proportion of dietary intake compared to meat and dairy products (Venderly & Campbell, 2006). Table 1 provides an overview of common vegetarian diets. Reduction in coronary heart disease, hypertension, diabetes mellitus, obesity and even some cancers have been observed in participants following vegetarian-based diets (Barnard et al., 2015; Olrich & Fraser., 2014; Ornish et al.,1998; Schmidt et al.,1997). Diets of this nature are typically higher in oligo and polysaccharides, fibre, fruits, vegetables, antioxidants and phytochemicals while lower in saturated fat and cholesterol compared to omnivorous diets (Venderley & Campbell, 2006). Some athletes have adopted a vegetarian diet to acquire the health benefits associated, but also believe the diet may assist in achieving appropriate carbohydrate intake, weight management and other performance enhancing advantages (Fuhrman & Ferreri, 2010). Physical performance is a broad term, however, in the context of this review will include; strength, speed, endurance and power, whilst excluding other traditional components such as balance and

[Insert Table 1 here]

flexibility. Although mechanisms linking a diet high in plant-based foods to improved physical performance are limited, there are three plausible theories described in the literature. Firstly, it has been hypothesised that a vegetarian diet may enhance an athlete's performance due to the high carbohydrate intake leading to improved glycogen stores in the body (Barr & Rideout 2004; Ferreira et al., 2006). Secondly, the increased phytochemicals and antioxidants consumed in vegetarian-based diets may also help reduce oxidative stress associated with prolonged exercise and improve general immunity (Trapp et al, 2010). Thirdly, it is widely accepted that intramuscular acidity can limit high-intensity exercise (Carr et al, 2011). A relationship has been established linking oral supplements, namely sodium bicarbonate and sodium citrate, to altered blood alkalosis levels. When ingested these buffers have been shown to have an ergogenic effect on high-intensity acute exercise (Carr et al, 2011). Conversely, ingested acidic supplements can be ergolytic. Evidence suggests consuming a vegetarian diet will have an alkaline effect on acid-base levels compared to non-vegetarians due to the high fruit and vegetable intake, whilst being lower in animal based proteins (Hietavala et al, 2015; Deriemaeker et al, 2010). Although a long-term vegetarian-based diet may not have the same effect as an acute sodium bicarbonate supplement, it is plausible, that a small 're-setting' change in the homeostatic baseline may be approached when a sustained vegetarian-based diet is followed, leading to a potential increase in physical performance. Despite these promising notions, there remains concern that a sub-optimal vegetarian

diet may increase risk for micronutrient deficiencies and reduce muscle creatine concentrations leading to submaximal performance (Barr & Rideout, 2004). Studies connecting vegetarian diets to improved health are well-established (Barnard et al., 2015; Olrich & Fraser, 2014; Ornish et al.,1998; Schmidt et al.,1997), however, the evidence for this phenomenon to be transferred to improved physical performance in athletes is less clear. The aim of this study was to examine the evidence for the relationship between consumption of a vegetarian-based diet and improved physical performance by conducting a systematic literature review. Due to a vegetarian-based diet theoretically increasing muscle glycogen, cell alkalinity, immunity, while reducing oxidative stress, it is hypothesised that this diet may improve physical performance in athletes.

Methods

Study protocol

A systematic literature review (NHMRC, 2000) was conducted in January 2015 using the SCOPUS database with no date exclusions. The search used the following keywords; vegetarian* OR vegan* AND sport* OR athlete* OR training OR performance OR endurance' in article, keywords or abstract. A search for unpublished literature was not performed though reference lists of the included publications were examined for additional relevant studies. National Health Medical Research Council's levels of evidence were applied to the included studies.

Study selection

Inclusion of studies met the following requirements. The studies (i) directly compared a plant-based diet (e.g. ovo-vegetarian, LOV or vegan) to a typical omnivorous/mixed diet, (ii) directly assessed physical performance, not solely biomarkers of physical performance (immune biomarkers were exempt from this inclusion criterion as physical detection of immunity is difficult to measure). The inclusion criterion was created to assess diet and its effect on performance rather than other external factors such as supplementation and lifestyle factors. It was important for physical performance to be measured, as biomarkers alone may not translate into effects on physical performance. Studies that met the following exclusion criteria were omitted: (i) studies with key words – pregnancy, non-human, high performance liquid chromatography (excluded within database search limits) (ii) journal articles not published in English, (iii) studies examining the relationship between diet *and lifestyle factors* on physical performance. This review was only concerned with diet and its effect on physical performance, not the effect of other lifestyle factors, (iv) published conference papers, short surveys, letters, notes, editorials, articles in press, book series, erratum and conference proceedings. (v) participants taking supplementation to emulate a vegetarian diet.

Data extraction included information on the publication year, study design/quality, number of participants, total sample size, population type, dropouts, intervention, diet, study results/conclusions (Table 2). Study quality was assessed using the quality criteria checklist of the Evidence Analysis Library (<http://www.anddeal.org/>)

of the Academy of Nutrition and Dietetics (2012).

Results

The literature search identified 327 studies of which eight articles met the inclusion criteria.

[Insert figure 1 here]

The eight included studies were varied with respect to population, intervention period, diet composition, and primary objectives including attribute of physical performance (Table 2). For instance, several papers examined muscular power and strength (Campbell et al., 1999; Haub et al., 2005; Wells et al., 2003), four assessed anaerobic and aerobic performance (Baguet et al., 2011; Hanne et al., 1986; Hietavala et al., 2012; Raben et al., 1992) whilst one investigated immune parameters (Richter et al., 1991) in relation to a vegetarian-based diet. In addition, most papers used different physical testing and/or biomarkers. The following sections are structured according to the type of physical performance being analysed, although there was some cross over between studies.

[Insert Table 2]

Resistance (strength/power) training

Three studies examined the difference between a LOV diet and a typical beef-containing western diet and its effect on Resistance Training (RT) in elderly men

(Campbell et al., 1999; Haub et al., 2005; Wells et al., 2003). The studies were unified regarding muscular strength. All three studies found no significant difference in muscular strength or power between the LOV groups and the omnivorous groups except in Wells et al. (2003) where the LOV group displayed a significant increase in strength for knee extensions ($p < 0.01$), yet both groups revealed significant improvements in muscular strength and power. Campbell et al. (1999) did, however, report resistance training induced changes in whole body composition ($p = 0.014$) and an increase in mean type II muscle fibre area size between groups ($p = 0.005$). Similarly, Wells et al. 2003, described haemoglobin and haematocrit were significantly increased in the meat group ($p < 0.01$) though this did not affect strength testing.

Anaerobic and aerobic performance

Four studies were identified relating a vegetarian-based diet to either anaerobic and/or aerobic performance. Hietavala et al. (2012) revealed that a low protein vegetarian diet had no significant effect on exercise time to exhaustion, but oxygen consumption was significantly higher at 40%, 60% and 80% of maximum oxygen consumption compared to a mixed diet (2.03 ± 0.25 vs. 1.82 ± 0.21 l/min, $p=0.035$; 2.86 ± 0.36 vs. 2.52 ± 0.33 l/min, $p < 0.001$ and 4.03 ± 0.50 vs. 3.54 ± 0.58 L/min, $p < 0.001$; respectively). Venous blood pH, strong ion difference, partial pressure of CO_2 , HCO_3^- , was also measured with no significant difference between diets. Comparably, Baguet et al. (2011) found that anaerobic performance improvement (repeated sprint ability test) was not different between the diet groups. Hanne et al. (1986) assessed both anaerobic and aerobic

capacity between vegetarian and non-vegetarian athletes. No significant difference in aerobic performance, as measured by predicted maximum oxygen consumption and Rating of Perceived Exertion (RPE) was observed. Likewise, no significant differences between groups were measured using the Wingate test to assess anaerobic performance (Table 2).

Raben et al. (1992) reported no significant differences between a LOV diet and maximum oxygen consumption, maximal voluntary contraction, endurance performance or muscle glycogen concentrations compared to a mixed diet (both diets controlled for carbohydrate 57%, protein 14% and fat 29%). A significant decrease in fasting serum testosterone was observed over the six week intervention period in the vegetarian groups diet (median 21.1nmol^{-1} to 13.7nmol^{-1} , $p < 0.05$), where no change was observed in the mixed diet. This did not have an effect on any physical performance parameters.

Immune function

Richter et al. (1991) reported that the immune parameters; blood mononuclear cells, and natural killer cells did not differ between a vegetarian and mixed diet after aerobic exercise. Similarly, Phytohaemagglutinin (PHA) and Purified Protein Derivative (PPD; tuberculin) showed no significant differences between dietary groups.

Discussion

This review is the first to explore an exclusive vegetarian-based diet and its

effects on physical performance using a rigorous systematic approach. Earlier investigations have focused on components of a vegetarian diet and performance, or supplementation emulating a vegetarian diet and performance, but none have examined the diet holistically, the way individuals or athletes would typically eat. Due to the limited evidence pool, and the disparate outcomes of performance tested, evaluating the association between a vegetarian-based diet and improved physical performance in athletes was immeasurable. This did not align with the primary hypothesis that a vegetarian diet would improve physical performance in athletes.

Nieman (1999) similarly reviewed vegetarian diets and possible links to improved physical performance. Seventeen scientific papers were assessed by Nieman prior to 1999, with neither a beneficial or a detrimental effect reported. Of the eight papers, which were reviewed in this investigation, all were unified with Nieman's findings. The vegetarian-based diet did not improve nor hinder physical performance. It is noteworthy to declare that all references used in Nieman's 1999 paper were hand searched for inclusion in this systematic review. No additional articles were included. Nieman's study was not extracted in the methodology, and therefore not included in the results of the review. This occurred due to the keywords used by Nieman (Exercise, endurance, athlete, carbohydrate, meat, iron, protein, creatine, vegetarian diet, humans) being broad with more focus on food groups and macronutrients.

Due to limited studies and dissimilar performance measurements, there may still be some merit to the hypothesis forecasting a vegetarian-based diet increasing

performance due to increased muscle glycogen, cell alkalinity and immunity, while reducing oxidative stress. This is particularly true for reducing oxidative stress as no trials were found on the subject.

Strength and Power

The three papers examining a LOV diet were unified, identifying that both the control and LOV groups significantly improved muscular strength and power equally during the study period. All three studies used elderly men as subjects concluding there is no difference between LOV diets and omnivorous diets in RT in elderly men. However, this may not be representative of the larger population. Wider studies are required to confidently consolidate their findings with the inclusion of both genders, and a range of ages. In two out of the three studies texturised vegetable protein was used frequently in the LOV diets with breakfast patties, grillers, chick[pea] patties and veggie dogs highly prominent (Haub et al., 2005; Wells et al., 2003). Products such as these often contain food colour, pH modifier, surface-active substrates, emulsifiers and surfactants (Asgar et al., 2010). Ideally, the aim of this study was to investigate a more whole food vegetarian-based diet. Research limiting the use of texturised vegetable protein products is warranted to more adequately align a plant-based dietary intake, and its response to resistive training.

Anaerobic and aerobic performance

Four papers were identified analysing a vegetarian diet and its effect on endurance and/or aerobic performance. These studies exhibited some

heterogeneity with three of the papers reporting on maximal aerobic capacity, two papers reporting on anaerobic performance and one also including isometric strength performance. No significant differences were observed between dietary intake and physical performance.

As only four studies with small participant pools were identified, it is imprudent to make a judgement on the effect of a vegetarian diet regarding this type of physical performance. The studies were consistent, however, revealing no significant differences between dietary groups in short, middle or endurance performance. This should only serve as a preliminary statement with further research required. This is particularly true with Baguet et al. (2011) issuing both the vegetarian and non vegetarian groups 1g/day of creatine monohydrate to reduce a creatine deficiency in the vegetarian diet group^{*}. Some studies, such as that of Bembien & Lamont (2005), have linked creatine to improved anaerobic performance. Although Baguet et al. (2011) were analysing carnosine concentration, the creatine supplementation may have skewed the results, at least for the applicability of this review. In the study by Baguet et al. (2003), baseline measurements between vegetarians and non vegetarians revealed lower total creatine concentration ($p < 0.05$). If creatine is implicated in improved performance, and vegetarians have reduced concentrations to non-vegetarians, creatine supplementation may be particularly influential in performance results. Supplementing with creatine eliminates it as a variable, enabling the specific focus of carnosine; which has been hypothesised to increase performance,

* Creatine monohydrate was used across both dietary groups to eliminate it as a variable. As both groups supplemented with creatine, it was included.

however, greatly limits the findings to address the relationship between a vegetarian diet and performance in short to middle distance athletes. From this study, it can be supposed, that there is no difference in carnosine concentrations between the two dietary groups (Baguet et al., 2011), however, any links to physical performance must be questioned due to the creatine supplementation. Hietavla et al. (2012) interestingly found that although there was no overall difference between the dietary groups' acid-base status or overall effect on maximum oxygen capacity, cycling efficiency decreased in the LOV group. This would not be a desirable effect for any athlete, which deserves to be explored further. Three of the studies assessing anaerobic and aerobic performance used short treatment periods of vegetarian consumption (Baguet et al; 5 weeks: Raben et al; 6 weeks: Hietavala et al: 4 days). The only study which was included where a vegetarian diet was adopted for an extended period of time, was that of Hanne et al. (1986). The participants in this study, were vegetarian for a minimum of two years. This timeframe would be more suitable to assess metabolic alterations. However, the sample size was small (39 vegetarians) and the investigation did not implement a randomised control study, but a cross sectional assessment. A larger number of participants, longer treatment times, studies without additional supplementation and a greater number studies are needed to confidently make a conclusion about a vegetarian-based diet and its effect on anaerobic an aerobic performance.

Immune parameters and performance

It has been suggested that due to the wealth of phytochemicals, antioxidants and

plethora of micronutrients in vegetarian-based diets, immune function may be improved in the vegetarian population (Nieman, 1988). This was not observed in the single study identified comparing immune status between the two dietary groups (Richter et al., 1991). The treatment groups in this particular study were subjected to a macro energy controlled, 57% Carbohydrate, 14% Protein, 29% Lipids LOV (16% animal derived protein, 84% vegetable protein) or mixed western diet (67% animal derived protein, 33% vegetable protein) diet for a total of 6 weeks. This duration is perhaps lacking the duration for full effects of a vegetarian/vegan diet to become apparent. A larger body of research with an extended duration of vegetarian consumption is needed before this can be concluded.

Mechanisms

Mechanisms other than those hypothesised to discriminate between a vegetarian-based diet and a mixed diet were proposed in some of the studies. Raben et al. (1992) for example hypothesised a decrease in sex serum testosterone due to a vegetarian-based diet. Raben et al suggested non-heme iron may not be absorbed as readily as heme iron and increased fibre intake may reduce the bioavailability of some nutrients, causing implications to the heavily training athlete. This was found not to be the case as the study revealed low level sex serum hormones in the vegetarian group but no changes in physical performance.

Baget et al. (2011) and Hietavala et al. (2012) investigated the relationship

between vegetarian-based diets and their effect on acid-base balance. Heiteva et al. (2012) found no significant difference in venous blood pH, strong ion difference or total concentration of weak acids (Atot), suggesting a low protein, vegetarian-based diet (Protein 10.1%, Carbohydrate 58.7%, Fat 24.7%) may not optimise acid-base balance and thus improve physical performance. Baget et al. (2011) predominately focused on carnosine and its buffering capacity. Lacto-ovo vegetarians revealed lower total creatine concentration ($p < 0.05$) compared to non-vegetarian participants, however, no difference in performance was observed, again suggestive of a vegetarian-based diet being ineffective at providing some sort of buffering effect.

Limitations

A limitation with the body of evidence is that all of the RCT's used extremely short periods of dietary intervention ranging from 4 days to 12 weeks. Changes in stored nutrient concentration could take much longer than this period. For instance, a recent study revealed that the most notable decline in vitamin B-12 in vegan participants occurred between 24 months and 60 months (Madry et al., 2012). The results from the present literature review only offer understanding into the short-term effects of a vegetarian-based diet, which may be useful for acute purposes, such as leading into a competition or race, however; does not address long-term effects. This is significant, as athletes following a vegetarian-based diet would typically do so for extended periods.

Additionally, many of the included papers lacked information on the

standardisation of dietary intake between groups and/or lacked detail about dietary compliance. Jeacocke and Burke (2010), note the possible impact poor dietary control can have on the outcome of a study. This is of particular interest in this review as the sample sizes were small, thereby likely to exaggerate the results of inadequate dietary standardisation between groups.

Whilst this systematic literature review has provided new insights into the effects of vegetarian-based nutrition and physical performance in athletes via a highly rigorous and structured review, some improvements could be made. The present SLR's search criterion encompasses dietary factors and effects on *total* physical performance. Refined search parameters focusing on *specific* domains of physical performance may uncover additional studies within that domain. Furthermore, as limited research was identified exploring vegetarian-based nutrition and physical performance, including cross-sectional studies comparing performance biomarkers may have increased the strength of this review.

Conclusion

Currently, the evidence for consuming a predominately vegetarian-based diet and improved athletic performance is lacking. In the eight studies which were identified in this review, however, the vegetarian-based diet did not improve performance, nor did it hinder it. There appeared to be no differences at least acutely between a vegetarian-based diet and an omnivorous diet in muscular power, muscular strength, anaerobic or aerobic performance. Many limitations were identified within the body of evidence including the total body of evidence

being very small (7 trials and 1 cross sectional analysis), the body of evidence experimental outcomes varied significantly, typically short dietary treatment times were administered (all but one study used treatments of 4 days – 12 weeks), resistance training focused only on elderly men and supplementation was used in one of the trials. More trials are needed to address the limitations and provide stronger evidence towards vegetarian-based diets and their effects on physical performance in athletes. It would be recommended that future research meets high level RCT design with strict vegetarian-based dietary intervention lasting six months or greater to determine the association between a vegetarian-based diet and physical performance.

Acknowledgements

No funding has been received for the preparation of this manuscript. The authors declare that there are no conflicts of interest that are directly relevant to the content of this review. The study was designed by Joel Craddock and Yasmine Probst; data were collected and analyzed by Joel Craddock; data interpretation and manuscript preparation were undertaken by Joel Craddock, Yasmine Probst and Greg Peoples. All authors approved the final version of the paper.

Reference List

Academy of Nutrition and Dietetics, (2012). *Evidence Analysis Manual: Steps in the Academy Evidence Analysis Process*. Available at <http://www.andeanal.org/> Accessed 10/12/2014.

Asgar, M. A., Fazilah, A., Huda, N., Bhat, R., & Karim, A. A. (2010). Nonmeat protein alternatives as meat extenders and meat analogs. *Comprehensive Reviews in Food Science and Food Safety*, 9(5), 513–529.

- Baguet, A., Everaert, I., De Naeyer, H., Reyngoudt, H., Stegen, S., Beeckman, S., Achten, E., Vanhee, L., Volckaert, A., Petrovic, M., Taes, Y., & Derave, W. (2011). Effects of sprint training combined with vegetarian or mixed diet on muscle carnosine content and buffering capacity. *European Journal of Applied Physiology*, 111(10), 2571–2580.
- Barnard, ND, Levin, SM, & Yokoyama, Y 2015, 'Research: A Systematic Review and Meta-Analysis of Changes in Body Weight in Clinical Trials of Vegetarian Diets', *Journal of the Academy of Nutrition and Dietetics*, vol. 115, pp. 954-969
- Barr, S I., & Rideout, C A., (2004). Nutritional considerations for vegetarian athletes. *Nutrition*, 20(7-8), 696–703.
- Bemben, M G., & Lamont, H S. (2005). Creatine supplementation and exercise performance: Recent findings. *Sports Medicine*, 35(2), 107–125.
- Bloomer, R J., & Goldfarb, A. H. (2004). Anaerobic exercise and oxidative stress: A review. *Canadian Journal of Applied Physiology*, 29(3), 245–263.
- Campbell, W W., Barton Jr., M L., Cyr-Campbell, D., Davey, S L., Beard, J L., Parise, G., & Evans, W J. (1999). Effects of an omnivorous diet compared with a lactoovo vegetarian diet on resistance-training-induced changes in body composition and skeletal muscle in older men. *American Journal of Clinical Nutrition*, 70(6), 1032–1039.
- Carr, A J., Hopkins, W G., & Gore, C J., (2011). Effects of Acute Alkalosis and Acidosis on Performance, 41(10), 801–814.
- Deriemaeker, P., Aerenhouts, D., Hebbelinck, M., & Clarys, P, (2010). Nutrient based estimation of acid-base balance in vegetarians and non-vegetarians. *Plant Foods for Human Nutrition (Dordrecht, Netherlands)*, 65(1), 77–82.
- Ferreira, L G., Burini, R C., & Maia, A F., (2006). Vegetarian diets and sports performance . *Dietas Vegetarianas E Desempenho Esportivo*, 19(4), 469–477.
- Fisher-Wellman, K., & Bloomer, R. J. (2009). Acute exercise and oxidative stress: A 30 year history. *Dynamic Medicine*, 8(1).
- Fuhrman, J., & Ferreri, D. M. (2010). Fueling the Vegetarian (Vegan) Athlete. *Current Sports Medicine Reports*, 9(4), 233–241.
- Hanne, N., Dlin, R., & Rotstein, A. (1986). Physical fitness, anthropometric and metabolic parameters in vegetarian athletes. *Journal of Sports Medicine and Physical Fitness*, 26(2), 180–185.

Haub, M D., Wells, A M., & Campbell, W.W. (2005). Beef and soy-based food supplements differentially affect serum lipoprotein-lipid profiles because of changes in carbohydrate intake and novel nutrient intake ratios in older men who resistive-train. *Metabolism: Clinical and Experimental*, 54(6), 769–774.

Hietavala, E.-M., Puurtinen, R., Kainulainen, H., & Mero, A. A. (2012). Low-protein vegetarian diet does not have a short-term effect on blood acid-base status but raises oxygen consumption during submaximal cycling. *Journal of the International Society of Sports Nutrition*, 9(50).

Hietavala, E., Stout, J. R., Hulmi, J. J., Suominen, H., Pitkänen, H., Puurtinen, R., & Mero, A. A. (2015). Effect of diet composition on acid-base balance in adolescents, young adults and elderly at rest and during exercise. *European Journal Of Clinical Nutrition*, 69(3), 399-404

Jeacocke, N. A., & Burke, L. M. (2010). Methods to Standardize Dietary Intake Before Performance Testing. *International Journal Of Sport Nutrition & Exercise Metabolism*, 20(2), 87-103.

Lap Tai, L., & Joan, S. (2014). Beyond Meatless, the Health Effects of Vegan Diets: Findings from the Adventist Cohorts. *Nutrients*, 6(6), 2131-2147

Madry, E., Lisowska, A., Grebowiec, P., & Walkowiak, J. (2012). The impact of vegan diet on B-12 status in healthy omnivores: Five-year prospective study. *Acta Scientiarum Polonorum, Technologia Alimentaria*, 11(2), 209–212.

NHMRC. (2000) How to use the evidence: *assessment and application of scientific evidence, Handbook series on preparing clinical practice guidelines*. Retrieved from https://www.nhmrc.gov.au/_files_nhmrc/publications/attachments/cp69.pdf

Nieman, D. C. (1988). Vegetarian dietary practices and endurance performance. *American Journal of Clinical Nutrition*, 48(3 SUPPL.), 754–761.

Nieman D C., (1998). Physical fitness and vegetarian diets: is there a relation? *American Journal of Clinical Nutrition*, 70(3), 570-575.

Olrich, M J., & Fraser G E., (2014). Vegetarian diets in the Adventist Health Study 2: a review of initial published findings. *American Journal of Clinical Nutrition*, 100(1), 353S-358S.

Ornish, D., Scherwitz, L. W., Billings, J. H., Lance Gould, K., Merritt, T. A., Sparler, S., Armstrong, W T., Ports, T A., Kirkeeide, R L., Hogeboom, C., & Brand, R. J. (1998). Intensive lifestyle changes for reversal of coronary heart disease. *Journal of the American Medical Association*, 280(23), 2001–2007.

Philip, J T. (2013). Nutritional Update for Physicians: Plant-Based Diets. *The Permanente Journal*, 17(2), 61–66.

Raben, A., Kiens, B., Richter, E. A., Rasmussen, L. B., Svenstrup, B., Micic, S., & Bennett, P. (1992). Serum sex hormones and endurance performance after a lacto-ovo vegetarian and a mixed diet. *Medicine and Science in Sports and Exercise*, 24(11), 1290–1297.

Richter, E A., Kiens, B., Raben, A., Tvede, N., & Pedersen, B. K. (1991). Immune parameters in male athletes after a lacto-ovo vegetarian diet and a mixed western diet. *Medicine and Science in Sports and Exercise*, 23(5), 517–521.

Schmidt, T., Wijga, A., Von Zur Muhlen, A., Brabant, G., & Wagner, T. O. (1997). Changes in cardiovascular risk factors and hormones during a comprehensive residential three month kriya yoga training and vegetarian nutrition. *Acta Physiologica Scandinavica, Supplement*, 161(640), 158–162.

Szeto, Y. T., Kwok, T. C. Y., & Benzie, I. F. F. (2004). Effects of a long-term vegetarian diet on biomarkers of antioxidant status and cardiovascular disease risk. *Nutrition*, 20(10), 863–866.

Trapp, D., Knez, W., Sinclair, W. (2010). Could a vegetarian diet reduce exercise-induced oxidative stress? A review of the literature. *Journal of Sports Sciences*, 28(12), 1261–1268.

Venderley, A. M., & Campbell, W. W. (2006). Vegetarian diets: Nutritional considerations for athletes. *Sports Medicine*, 36(4), 293–305.

Wells, A M., Haub, M D., Fluckey, J., Williams, D K., Chernoff, R., & Campbell, W. W. (2003). Comparisons of vegetarian and beef-containing diets on hematological indexes and iron stores during a period of resistive training in older men. *Journal of the American Dietetic Association*, 103(5), 594–601.

2 **Tables**3 Table 1. Classification of dietary patterns[^]

Type of diet	Definition	Red meat	Poultry	Fish	Dairy	Eggs
Non-Vegetarian						
Mixed	Eats red meat, poultry, fish, dairy and eggs.	✓	✓	✓	✓	✓
Vegetarian-Based						
Pesco-Lacto-Ovo Vegetarian (PL)	Eats fish, dairy and eggs, but not red meat or poultry.	✗	✗	✓	✓	✓
Lacto-ovo vegetarian	not red meat, poultry or fish.	✗	✗	✗	✓	✓
Lacto-vegetarian	Eats dairy not red meat, poultry, fish, dairy or eggs.	✗	✗	✗	✓	✗
Ovo-vegetarian	meat, poultry, fish, dairy or eggs.	✗	✗	✗	✗	✓
Pesco-Vegetarian	Eats fish but not red meat, poultry eggs or dairy.	✗	✗	✓	✗	✗
Vegan	Eats no red meat, poultry, fish, dairy or eggs.	✗	✗	✗	✗	✗

20
21 [^] Adapted from Lap Tai & Joan, 2014

Table 2. Body of Evidence – Summary table of included journal articles with American Diabetes Association quality rating template results included.

Study Description	Study Quality	Population	Intervention	Dietary Group	Study Results/Conclusion
Anaerobic and aerobic performance					
Baguet et al (2011)	Pseudo RCT, Level III-1*	n = 20 Healthy, Non-vegetarian participants. 11 males, 9 Females.	5 weeks, sprint training program -Mixed diet -Vegetarian Diet (*Both groups supplemented with creatine).	<u>LOV</u> Energy 9321kJ (P= 13.13%, CHO = 55.08%, F = 28.00%, EtOH = 3.79%) <u>Mixed</u> Energy 9693kJ (P = 15.78%, CHO = 54.55%, F = 26.02%, EtOH = 3.67%)	No performance difference in repeated sprint ability test between LOV diet and mixed diet.
Hietavala et al (2012)	RCT, Level II**	n = 9 Healthy, recreationally active men. No mention of prior eating habits.	Cross over design w/ 16 day washout period 4 day vegetarian diet 4 day normal diet.	<u>Low protein vegetarian diet</u> Energy 1046 kJ, (P = 10.1%, CHO = 58.7%, Fat = 24.7% (Limited grain and dairy) <u>Normal Diet</u> Energy 11687 kJ (P = 17.6%, CHO = 49.8%, Fat = 28.1 %)	Oxygen consumption was significantly higher at 40, 60 and 80% of maximum oxygen capacity (cycle ergometer). Suggestive of poorer exercise economy in vegetarian diet. No further effect on maximal aerobic performance.
Raban et al (1992) ^^	RCT Level II**	n = 8 Endurance trained male athletes. Non-vegetarians before study.	Cross over design with 4 week washout period, 6 week LOV diet 6 week non-vegetarian diet.	<u>Both Diets Included:</u> 57% E Carbohydrate, 14% E Protein, 29% E Lipids <u>LOV</u> Protein composition (16% animal derived protein, 84% vegetable protein) <u>Mixed western</u> Protein composition (67% animal protein, 33% vegetable protein)	No difference in maximal oxygen uptake (graded ergometer cycle or treadmill test) or maximal voluntary contraction (measured with strain gauge) between groups.
Hanne et al al (1989)	Level III-2***	n = 98 49 vegan, lacto vegetarian or LOV 49 Non vegetarians.	NA - Cross sectional cohort.	<u>Vegetarian-based</u> > 2 years (Vegan, lacto vegetarian or LOV) <u>Mixed</u> - Non-vegetarian based diet	No difference in anaerobic (Wingate anaerobic test) or maximal oxygen uptake (cycle stress test) performance.
Strength and Power					
Haub et al (2005) ^	RCT, Level II**	n =21 Healthy men aged 59 - 78 BMI 24-33kg/m2.	12-week RT program 3days/week	<u>LOV</u> 0.6g/protein/kg/day from TVP Energy 9.37MJ (P= 1.17g/kg/day, CHO = 274g/day, F = 85g/day) <u>LOV + Beef</u> 0.6/g/protein/kg/day from beef Energy 9.09MJ (P= 1.10g/kg/day, CHO =	No difference in strength (repetitions until fatigue) or power gains (1 rep max) between a LOV diet + soy or LOV diet +Beef. No difference between groups for upper body or lower body power output.

				282g/day, F = 73g/day)	
Campbell et al (1999)	Pseudo RCT, Level III-1*	n = 19 Sedentary men (51-69 y.o) overweight to moderately obese aged	12 week resistance training program. Group 1 (Mixed Diet) Group 2 (LOV).	<u>LOV</u> (self-selected) Energy ~10.3 MJ (52% E Carbohydrate, 13% E Protein, 34% E Lipid) <u>Mixed</u> - (habitual unrestricted) (provided 50% of total dietary protein from meat) Energy ~ 8.6 MJ (50% E Carbohydrate, 16% E Protein, 32% E Lipid)	No difference in strength (1 Rep Max) with RT between groups for any of the exercises performed.
Wells et al (2003) ^	RCT Level II**	n = 21 Healthy men aged 59 - 78 BMI 24-33kg/m2.	12-week RT program 3days/week	<u>LOV</u> - 0.6g/protein/kg/day from TVP Energy 9.37MJ (P= 1.17g/kg/day, CHO = 274g/day, F = 85g/day) <u>LOV + Beef</u> - 0.6g/protein/kg/day from beef Energy 9.09MJ (P= 1.10g/kg/day, CHO = 282g/day, F = 73g/day)	No differences in strength (1 Rep Max) between groups in all but one exercise. Vegetarian group had a larger increase in strength for knee extensions (p <0.01).
Immune Parameters					
Richter et al (1991) ^^	RCT Level II**	n = 8 8 Endurance trained male athletes (4 cyclists, 1 runner, 1 rower, 2 mixed) Non-vegetarians before the study.	Cross over design with 4 week washout period, 6 week LOV diet 6 week non-vegetarian diet.	<u>Both Diets Included:</u> 57% E Carbohydrate, 14% E Protein, 29% E Lipids <u>LOV</u> Protein composition (16% animal derived protein, 84% vegetable protein) <u>Mixed western</u> Protein composition (67% animal protein, 33% vegetable protein)	No difference in composition or concentration in in-vivo function of human blood mononuclear cells between a meat rich mixed diet, or a LOV diet.

* Pseudo RCT, Level III-I - A study of test accuracy with: an independent, blinded comparison with a valid reference standard, among non-consecutive patients with a defined clinical presentation.

** RCT Level II - A study of test accuracy with: an independent, blinded comparison with a valid reference standard, among consecutive patients with a defined clinical presentation.

*** Level III-2 - A comparative study with concurrent controls: Non-randomised, experimental trial, cohort study, case-control study, interrupted time series with a control group.

^ Same experiment/data set - assessed different parameters of physical activity.

^^ Same experiment /data set - assessed different parameters of physical activity.

Figures




Figure 1. The PRISMA flowchart showing the initial and final number of studies obtained.