University of Wollongong

Research Online

Faculty of Science, Medicine and Health -
Papers: part A

Faculty of Science, Medicine and Health

1-1-2013

A review of observations and models of dynamic topography

Nicolas Flament

University of Sydney, nflament@uow.edu.au

Michael Gurnis

California Institute of Technology

R Dietmar Muller

University of Sydney

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>

Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Flament, Nicolas; Gurnis, Michael; and Muller, R Dietmar, "A review of observations and models of dynamic topography" (2013). *Faculty of Science, Medicine and Health - Papers: part A*. 4274.
<https://ro.uow.edu.au/smhpapers/4274>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

A review of observations and models of dynamic topography

Abstract

The topography of Earth is primarily controlled by lateral differences in the density structure of the crust and lithosphere. In addition to this isostatic topography, flow in the mantle induces deformation of its surface leading to dynamic topography. This transient deformation evolves over tens of millions of years, occurs at long wavelength, and is relatively small (<2 km) in amplitude. Here, we review the observational constraints and modeling approaches used to understand the amplitude, spatial pattern, and time dependence of dynamic topography. The best constraint on the present-day dynamic topography induced by sublithospheric mantle flow is likely the residual bathymetry calculated by removing the isostatic effect of oceanic lithospheric structure from observed bathymetry. Increasing knowledge of the thermal and chemical structure of the lithosphere is important to better constrain present-day mantle flow and dynamic topography. Nevertheless, at long wavelengths (>5000 km), we show that there is good agreement between published residual topography fields, including the one described here, and present-day dynamic topography predicted from mantle flow models, including a new one. Residual and predicted fields show peak-to-peak amplitudes of roughly ± 2 km and a dominant degree two pattern with high values for the Pacific Ocean, southern Africa, and the North Atlantic and low values for South America, western North America, and Eurasia. The flooding of continental interiors has long been known to require both larger amplitudes and to be temporally phase-shifted compared with inferred eustatic changes. Such long-wavelength inferred vertical motions have been attributed to dynamic topography. An important consequence of dynamic topography is that long-term global sea-level change cannot be estimated at a single passive margin. As a case study, we compare the results of three published models and of our model to the subsidence history of well COST-B2 offshore New Jersey. The $<400>\pm 45$ m amount of anomalous subsidence of this well since 85 Ma is best explained by models that predict dynamic subsidence of the New Jersey margin during that period. Explicitly including the lithosphere in future global mantle flow models should not only facilitate such comparisons between model results and data, but also further constrain the nature of the coupling between the mantle and the lithosphere.

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Flament, N., Gurnis, M. & Muller, R. (2013). A review of observations and models of dynamic topography. *Lithosphere*, 5 (2), 189-210.

A review of observations and models of dynamic topography

Nicolas Flament¹, Michael Gurnis², and R. Dietmar Müller¹

¹ *EarthByte Group, School of Geosciences, University of Sydney NSW 2006, Australia*

² *Seismological Laboratory, California Institute of Technology, Pasadena CA, USA*

ABSTRACT

The topography of the Earth is primarily controlled by lateral differences in the density structure of the crust and lithosphere. In addition to this *isostatic* topography, flow in the mantle induces deformation of its surface leading to *dynamic topography*. This transient deformation evolves over tens of millions of years, occurs at long wavelength and is relatively small (< 2 km) in amplitude. Here, we review the observational constraints and modeling approaches used to understand the amplitude, spatial pattern and time-dependence of dynamic topography. The best constraint on the present-day dynamic topography induced by sub-lithospheric mantle flow is likely the residual bathymetry calculated by removing the isostatic effect of oceanic lithospheric structure from observed bathymetry. Increasing knowledge of the thermal and chemical structure of the lithosphere is important to better constrain present-day mantle flow and dynamic topography. Nevertheless, at long wavelengths ($> 5,000$ km), we show that there is good agreement between published residual topography fields, including one described here, and present-day dynamic topography predicted from mantle flow models, including a new one. Residual and predicted fields show peak-to-peak amplitudes of

roughly ± 2 km and a dominant degree two pattern with high values for the Pacific Ocean, southern Africa and the North Atlantic and low values for South America, western North America and Eurasia. The flooding of continental interiors has long been known to require both larger amplitudes and to be temporally phase-shifted compared with inferred eustatic changes. Such long wavelength inferred vertical motions have been attributed to dynamic topography. An important consequence of dynamic topography is that long-term global sea level change cannot be estimated at a single passive margin. As a case study, we compare the results of three published models and of our model to the subsidence history of well COST-B2 offshore New Jersey. The $< 400 \pm 45$ m of anomalous subsidence of this well since 85 Ma are best explained by models that predict dynamic subsidence of the New Jersey margin during that period. Explicitly including the lithosphere in future global mantle flow models should not only facilitate such comparisons between model results and data, but also further constrain the nature of the coupling between the mantle and the lithosphere.

INTRODUCTION

Knowledge of the effect of mantle flow on surface topography has considerably increased over the last 30 years. The rapid improvement in computational algorithms and computing resources has facilitated the modeling of global mantle flow at increasing resolution, that now achieve Earth-like convective vigor. The dramatic expansion of global and regional seismic data sets and numerical methods have lead to commensurate improvements in the resolution and global coverage of mantle tomographic images (e.g. Romanowicz, 2008). In turn these have lead to an improved understanding of the thermo-chemical structure of the mantle (Grand, 2002; Masters et al., 2000; Simmons et

al., 2007). Independently, global plate reconstructions that are used to constrain the evolution of mantle flow and to interpret seismic images of the mantle have likewise improved and such reconstructions are now available back to the Triassic in million year increments (Seton et al., 2012). As a consequence, plate kinematics and seismic tomography can now be assimilated into both forward and backward mantle flow models. Exploiting such improvements, a range of known long-wavelength vertical motions of continental plates have been attributed to mantle flow and hence used as constraints on the spatial character, amplitude and time-dependence of mantle flow. Examples include the Cenozoic uplift of Southern Africa (Gurnis et al., 2000), the late Cenozoic uplift of the Colorado Plateau (Moucha et al., 2009), the Cretaceous subsidence and subsequent uplift of the interior of North America (Liu et al., 2008; Mitrovica et al., 1989), the tilt of northern South America to the East during the Miocene (Shephard et al., 2010), the tilt of Australia since the late Cretaceous (DiCaprio et al., 2009) and the vertical motions of the Slave and Kaapvaal Cratons since the Paleozoic (Zhang et al., 2012). An important concept that derives from such studies is that there is likely “no such thing as a stable continental platform” (Moucha et al., 2008a) which in turn implies that long-term global sea level (eustasy) change cannot be defined based on the detailed analysis of the stratigraphic record in a single area, as has been argued previously (e.g. Miller et al., 2005).

Nevertheless, multiple challenges still limit our understanding of global dynamic topography, including different definitions of the meaning of the terms “*dynamic topography*”, inaccurate estimates of present-day dynamic topography in the absence of a detailed global model of the structure of the lithosphere, and contradicting model

predictions of vertical motions for the same area (Moucha et al., 2008a; Müller et al., 2008b; Spasojevic et al., 2008).

In this review, we first attempt to clarify the definition of dynamic topography. We then explore observational constraints on present-day and time-dependent dynamic topography. For the present-day, we compare three published residual topography fields (Kaban et al., 2003; Panasyuk and Hager, 2000; Steinberger, 2007) and a residual topography field that we derive, and show that they are in broad agreement. We then carry out a non-exhaustive review of analytical and numerical models of dynamic topography, both instantaneous and time-dependent, with an aim of not only showing the progressive improvements of these models but also to expose the strengths and limits of each. As part of this process, we compare the present-day dynamic topography predicted by four published models (Conrad and Husson, 2009; Ricard et al., 1993; Spasojevic and Gurnis, in press; Steinberger, 2007) and by the model that we introduce in this study. This comparison shows that the predicted dynamic topography fields are in good agreement at long wavelength, although the predicted amplitudes differ from one model to another. The residual topographies and modeled dynamic topographies are in broad agreement, although there are important regional mismatches. For example, east and southeast Asia are highs in most residual topography fields but they are also dynamic topography lows. We attribute this discrepancy to the difficulty to constrain residual topography with limited extents of dated oceanic lithosphere. In addition, the northeast Pacific present day residual topography disagrees with modeled changes in dynamic topography, which shows the limitations in constraining a process that occurs over tens of millions of years using a snapshot of residual topography. We next discuss the

consequences of dynamic topography for long-term sea level change. Although the global effect of dynamic topography on long-term sea level change, dominated by changes in the volume of ocean basins, is secondary (< 100 m; Conrad and Husson, 2009; Spasojevic and Gurnis, in press), its regional effects are important. To illustrate this, we carry out a detailed analysis of the anomalous subsidence of COST-B2 well offshore New Jersey that illustrates that this margin has subsided by up to 400 ± 45 m since 85 Ma. We show that this dynamic subsidence is consistent with the model presented in this study and with two previously published models (Müller et al., 2008b; Spasojevic et al., 2008). However, given the large uncertainty associated with the dynamic models, the detailed sea level curve derived from the stratigraphy of the New Jersey margin cannot be corrected for dynamic effects, which implies that it should not be used as long-term global sea level curve. From an observational point of view, a detailed knowledge of the thermal and chemical structure of the lithosphere is key to constraining present-day dynamic topography. From a modeling point of view, explicitly including the lithosphere in future global mantle flow model should not only facilitate the comparison between modeling results and the geological record, but also make dynamic topography an additional constraint on the nature of the coupling between the mantle and the lithosphere.

DYNAMIC TOPOGRAPHY

Dynamic topography is that topography due to flow within the mantle, an idea that was first proposed by Pekeris (1935). It is ‘dynamic’ because the mass anomalies driving the density are moving and this contrasts with isostatic topography in which the

mass anomalies are in a state of quasi-equilibrium. Due to plate tectonics, the topography of the Earth consists of continents and oceans with an average elevation difference of ~ 4.5 km. This first-order difference in elevation is of isostatic origin and results from the different average densities and thicknesses of continental and oceanic crust and lithosphere. The long-known poor correlation between the ocean-continent function and the long wavelength geoid (Kaula, 1972) supports this view. If the topography of the Earth was not affected by any dynamic processes, flat continents would stand ~ 4.5 km above flat abyssal plains. This is clearly not the case, and the topography of the Earth results from an array of dynamic processes operating at different scales in time and space. In the continents, tectonic convergence results in mountain belts such as the Himalayas and the Andes, and tectonic divergence results in rifts, such as the East African or the Salton Sea in California. The process leading to these topographic features is clearly dynamic, but even in active and geologically recent tectonic zones the topography is largely compensated isostatically: the continental crust is thicker than average in mountain belts, and thinner than average in rifts (Mooney et al., 1998).

In the oceans, the bathymetry is primarily determined by cooling and thickening of oceanic lithosphere as it moves away from spreading centers (Langseth et al., 1966; McKenzie, 1967). While it is ultimately dynamic in origin, the bathymetry of the young ocean floor (< 80 Ma) is well explained by combining the principles of isostasy and of the cooling of a semi-infinite half space (Davis and Lister, 1974). Ocean floor older than ~ 80 Ma appears to flatten compared to this simple and elegant physical model, although whether “normal” old seafloor flattens is subject to some debate (Hillier, 2010; Korenaga and Korenaga, 2008). The reason for the apparent shallowing of old ocean floor is the

subject of ongoing debate, and proposed explanations include thermal rejuvenation by hot spots (Smith and Sandwell, 1997), small-scale convection (e.g. Afonso et al., 2008) dynamic support by mantle flow (Kido and Seno, 1994; Zhang et al., 2012) and departure from the topography predicted by thermal boundary layer theory due to radioactive heat production within the mantle (Jarvis and Peltier, 1982). In the absence of consensus regarding the physical process explaining this flattening, the age-depth relationship of the ocean floor can still be accommodated in empirical “plate models” (e.g. Crosby and McKenzie, 2009; Parsons and Sclater, 1977). As in the continents, the topography of the ocean floor is affected by variations in crustal thickness, notably in areas affected by magmatism such as oceanic plateaus. Similarly, variations in lithospheric temperature and magmatic processes also affect continental topography, however there is no simple tectonic model to describe the topography of continents due to their long and complex tectonic history.

Given the number of dynamic processes that affect topography, it is not surprising that the definition of “*dynamic topography*”, which generally describes the effect of mantle convection on surface topography, does not have a single, agreed-upon definition. As discussed above, the bathymetry of the ocean floor is compatible with that of the thermal boundary layer of the convecting mantle and in this respect the subsidence of oceanic lithosphere with age should be included in the definition of dynamic topography. However, the dynamics of the continental lithosphere are more complex and cannot be explained by boundary layer theory. For this reason, defining dynamic topography globally as the difference between observed topography and isostatic crustal topography (e.g. Forte et al., 1993) results in continents dynamically depressed by up to 3 km, which

is inconsistent with observations (Gurnis, 1993a). This discrepancy is in part due to the non negligible isostatic contribution of the continental lithosphere (in the past referred to as tectosphere; Jordan, 1975) to topography. Given the relatively poor global constraints on the complex structure and density of the continental lithosphere (Zoback and Mooney, 2003), dynamic topography is usually defined as that topography originating from sources beneath the upper thermal boundary layer of mantle convection (e.g. Le Stunff and Ricard, 1995; Panasyuk and Hager, 2000). We use this latter definition in the present paper, keeping in mind that it ignores the many lithospheric-scale dynamic processes that affect topography, including those described above.

PRESENT-DAY CONSTRAINTS ON LONG-WAVELENGTH DYNAMIC TOPOGRAPHY: RESIDUAL TOPOGRAPHY

Constraining dynamic topography requires removing the isostatic contribution of the sediments, ice, crust and lithosphere from the observed topography. The remaining topography is called residual topography (Crough, 1983). In the oceans, a choice must be made between correcting either for the physical half-space cooling model (as in Davies and Pribac, 1993; Gurnis et al., 2000) or for a plate model accounting for the apparent flattening of the ocean floor with increasing age (Kaban et al., 2003; Le Stunff and Ricard, 1995; Steinberger, 2007). The former approach implies that mantle support may contribute to the observed flattening of old ocean floor (Kido and Seno, 1994), whereas the latter requires some other physical process to operate. In the continents, a common approach is to remove the isostatic topography due to the continental crust (Forte et al., 1993; Steinberger, 2007). However, correcting for the isostatic contribution of the

continental crust alone can result in continents being either systematically lowered or elevated by one to two kilometers, which is inconsistent with observation (Gurnis, 1993a). Zoback and Mooney (2003) showed that there is no simple relationship between continental crustal thickness and surface elevation, and that isostatic models based on crustal thickness alone systematically over-predict continental elevations, mainly because Archean continental lithosphere is depleted and of intrinsically lower density (Jordan, 1975). Several authors have proposed residual topography maps that remove the isostatic contribution of both the continental crust and continental lithosphere (Kaban et al., 2003; Le Stunff and Ricard, 1995; Panasyuk and Hager, 2000). Such calculations require assumptions relative to the thermo-chemical state of the continental lithosphere, which result in relatively large uncertainties (Kaban et al., 2003; Le Stunff and Ricard, 1995; Panasyuk and Hager, 2000) and underline the need for a better knowledge of the thermal and chemical structure of the lithosphere in order to constrain mantle flow models.

The long wavelength residual topography has been calculated by removing the mean continental elevation from continents (following Gurnis et al., 2000; Lithgow-Bertelloni and Silver, 1998), because the thermal and chemical structures of the continental lithosphere are not well known. In the oceans, the plate cooling model of Crosby and McKenzie (2009) was removed from the observed bathymetry after accounting for isostatic loading by sediments using the method of Sykes (1996) and the global sediment thickness map of Divins (2003). We consider that continents extend to the edge of the continental shelf (magenta contours in Fig. 1A), which roughly corresponds to the -200 m elevation contour (Harrison et al., 1983), and we calculate the equal area arithmetic mean continental elevation as 529 m. The areas between the edge

209 of the continental shelf and ocean floor *per se* as defined by (Müller et al., 2008a) were
210 arbitrarily set to a uniform elevation of -200 m. The resulting residual topography was
211 low-pass filtered using a filter tapered for wavelengths between 2,000 and 5,000 km, to
212 simplify comparison with published estimates of residual topography expanded in
213 spherical harmonics to degree 12 (Kaban et al., 2003; Panasyuk and Hager, 2000;
214 Steinberger, 2007; Fig.1B, C, and D). The residual topography contains an isostatic
215 component evident in high values in the Himalayas and Andes (with present crustal
216 thicknesses greater than 50 km (Mooney et al., 1998) shown as yellow contours, Fig. 1A).
217 In the oceans, the residual topography highs are closely associated with hot-spot tracks
218 (Cazenave et al., 1989; Schroeder, 1984) and Phanerozoic large igneous provinces (LIPs;
219 denoted by green contours in Fig. 1A). Therefore, part of the oceanic residual
220 topography is due to the thicker crust of oceanic plateaus and at least partially
221 isostatically compensated. Ideally, a correction for the thickness of the oceanic crust
222 should be carried out to estimate the topography associated with mantle support. Such
223 studies have been carried out regionally (e.g. Crosby and McKenzie, 2009; Winterbourne
224 et al., 2009) but not yet globally. Finally, the residual topography high over the Canada
225 Basin in the Arctic Ocean (Fig. 1A) can be attributed to the absence of sediment
226 thickness data.

227 Despite being based on different isostatic models of the lithosphere, the four
228 residual topography fields (Fig. 1) agree overall. For half-wavelengths greater than
229 ~ 3,300 km, the peak-to-peak amplitude is ± 2 km, and the root-mean-square amplitude
230 ~ 0.5 km (Table 1). The peak-to-peak amplitude is smallest for the model of Panasyuk
231 and Hager (2000), and largest for the model presented in this study (Table 1) due to the

absence of correcting for isostasy in elevated areas (Fig. 1A). The root-mean-square amplitude is smallest for the model of (Panasyuk and Hager, 2000), and largest for the model of (Kaban et al., 2003 ; Table 1). In terms of pattern, in the continents, features common to all four models include the residual topography lows within central Europe, eastern North America, and western South America, and the residual topography highs within southern and eastern Africa and western North America. In the oceans, the most persistent features between models are the Australian-Antarctic Discordance (AAD) that depresses the south east Indian mid-ocean ridge, the Argentine Basin in the southern South Atlantic, the residual topography low under the western part of the central Atlantic ocean, the large-scale, low-amplitude residual topography low under the northeast Pacific, and the highs under the central and western Pacific (Darwin Rise), the areas offshore southern Africa and around Iceland and the northern mid-Atlantic ridge.

TIME-DEPENDENT CONSTRAINTS ON DYNAMIC TOPOGRAPHY

In principle, dynamic topography can occur over all wavelengths from 10 km to greater than 1,000 km but since crustal and lithospheric density and thickness variations and plate flexure exerts a strong control on topography for wavelengths less than 100 km, most attempts to deconvolve it from the geological record have focused on transient long wavelengths changes that cannot be well explained without invoking dynamic topography. In this section, we first focus on the geological expression of long-wavelength dynamic topography that is the main focus of this paper before giving a brief overview of short-wavelength dynamic topography.

Geological expressions of long-wavelength dynamic topography

Long-wavelength dynamic topography is recorded by lithospheric plates moving over the constantly changing mantle flow and is thus ephemeral over million years to tens of millions of years. As a result, subsidence events are generally followed by uplift and vice versa. Indeed, the fact that dynamic topography is transient over tens of millions of years means that it must be recorded over long time periods over which the preservation potential of the rock record is poor. The very nature of the long-wavelength, low-amplitude dynamic topography signal makes it difficult to isolate it in the geological record that is dominated by shorter wavelength, larger amplitude signals such as lithospheric extension or shortening. One further restriction on the preservation of dynamic topography comes from the geological record itself that is biased towards areas that have been subsiding for long periods of time and have a good preservation potential, as opposed to uplifting areas that are subject to erosion and have a poor preservation potential. However, strata deposited in a dynamic topography depression also have low long-term preservation potential because dynamic topography is transient. Thus, ancient subduction-related dynamic topography is most likely to be represented by unconformities (Burgess et al., 1997). As a consequence, the stratigraphic record is the most accessible archive of the dynamic topography signal, and continental interiors that have not undergone deformation for hundreds of million years are the most promising places to search for the dynamic topography signal (Gurnis, 1992).

Stratigraphic record of continental interiors: examples from North America and Australia

The western interior Cretaceous seaway of North America that covered ~ 40% of the continent and resulted in > 1 km of sediments deposited over ~ 1,000 km (Bond, 1976; Liu and Nummedal, 2004) presents a robust long-wavelength signal. Analysis of Cretaceous sediments in North America suggests that the observed flooding would require a sea-level rise of 310 m, resulting in accumulation of 700 m of sediments (Bond, 1976), but the inferred Cretaceous isopachs (Cook and Bally, 1975) are significantly thicker. Late Cretaceous subsidence was followed by Tertiary uplift with a maximum tilt amplitude of 3 km over ~ 1,400 km (Mitrovica et al., 1989). During the Late Cretaceous, when the sedimentary wedge was thickest, the depositional system was characterized by an eastward migration of a depocenter from Utah to Wyoming (Weimer, 1970; Fig. 2A). The dominant characteristic is a westward downward tilt with an internal depocenter that migrated eastward at 5 cm/yr within North America (Liu et al., 2012; Fig. 2B-F; Liu et al., 2011). Besides this dominating Late-Cretaceous to Tertiary signal, Heller et al. (2003) identified three thin but widespread alluvial conglomeratic units that recorded post-Paleozoic episodes of wide spread tilting across the U.S. Cordillera that have been interpreted in terms of dynamic topography. Finally, a combination of stratigraphic, thermochronologic and provenance constraints showed that the south western corner of the U.S. tilted to the east at ~ 80 Ma and then to the west at ~ 55 Ma over scales of ~ 500 km with an amplitude of ~ 1 km (Wernicke, 2011).

Episodes of anomalous long wavelength motions have also been inferred for Australia that has experienced little orogenic activity since the mid-Mesozoic and is

299 relatively flat. During the Cretaceous, the marine inundation of Australia and eustasy
300 were out of phase. Maximum flooding of Australia occurred in the late Aptian to early
301 Albian (120 to 110 Ma) (Struckmeyer and Brown, 1990), when large regions of the
302 continent experienced marine inundation. Australia became progressively exposed during
303 the Late Cretaceous, reaching a flooding minimum in the Campanian (80 to 70 Ma) when
304 global sea level was near a maximum. This record of anomalous relative sea level
305 suggests that Australia has experienced several episodes of vertical motion, including a
306 large-scale vertical translation of the entire continent and a localized subsidence event
307 (Bond, 1978; Russell and Gurnis, 1994; Veevers and Conaghan, 1984). Late Cretaceous
308 vertical motions appear to have been mostly confined to the Eromanga and Surat basins
309 (Gallagher and Lambeck, 1989). The Eromanga Basin recorded ~ 500 m of tectonic
310 subsidence in 10 million years (Myr) at around 100 Ma (Gallagher and Lambeck, 1989).
311 Apatite fission track analysis suggests that the Cretaceous strata of the Surat Basin and
312 basins on the eastern margin were subsequently eroded (Gallagher et al., 1994), which is
313 consistent with the tilting up toward the east of Cretaceous marine strata (Gallagher et al.,
314 1994; Russell and Gurnis, 1994). Near the end of the Cretaceous, Australia was about
315 250 m higher than it is today (Bond, 1978; Russell and Gurnis, 1994; Veevers and
316 Conaghan, 1984). During the Cenozoic, Australia moved northward after it separated
317 from Antarctica and tilted down to the northeast (DiCaprio et al., 2009; Sandiford, 2007).
318 This tilt must have had an amplitude of 300 m since the Eocene to reconcile interpreted
319 patterns of marine incursion with a predicted topography accounting for global sea level
320 variations (DiCaprio et al., 2009). On the southern margin, long wavelength dynamic
321 topography was enhanced by at least 250 m of shorter wavelength anomalous subsidence,

consistent with the passage of the margin over a north–south elongated, 500 km wide, anomaly of dynamic topography approximately fixed with respect to the mantle (DiCaprio et al., 2009). The present-day position of this depth anomaly is aligned with the Australian-Antarctic Discordance (cf. Whittaker et al., 2010).

Stratigraphic record of continental margins

In addition to continental interiors, some methods have been developed to estimate dynamic topography in areas that have undergone recent tectonic deformation. For instance, post-rift anomalous subsidence, the difference between observed post-rift thermal subsidence and that predicted by an analytic model of lithospheric cooling, has been attributed to dynamic topography on the Queensland Plateau on the northeastern margin of Australia (Müller et al., 2000), in the South China Sea (Xie et al., 2006), between New Zealand and Antarctica (Sutherland et al., 2010), and at the New Jersey margin (see discussion).

Constraints on long-term dynamic uplift

The previous section shows that the stratigraphic record is a powerful constraint on dynamic topography in areas that have experienced long-term subsidence, especially in areas where long-term subsidence has been followed by uplift. Different methods are required to estimate dynamic topography in areas experiencing long-term dynamic uplift, with southern Africa being a well-known example. Both the timing and amplitude of uplift are generally difficult to constrain (Gurnis et al., 2000), due to the poor preservation potential in elevated areas and to the large uncertainty of paleo-altimetry proxies, typically a few hundred meters. As a result, dynamic uplift may be constrained

indirectly by estimating the timing and amplitude of rock uplift. In this respect, thermochronologic methods allow the estimation of the amount of exhumation by constraining both the amplitude and timing of the cooling of uplifted rocks. For instance, indirect estimates of vertical motions via thermochronology have been made for Eromanga Basin in Australia (Gallagher et al., 1994), the Slave Craton (Ault et al., 2009) and Kaapvaal Craton (Flowers and Schoene, 2010) using apatite (U–Th)/He (AHe) or apatite fission-track (AFT) thermochronology. One limit of thermochronologic methods, however, is the non-uniqueness of thermochronologic modeling (Redfield, 2010), which underlines the need to complement it with other, independent, observations (Green et al., 2011).

A recent approach to constraining dynamic uplift consists in “inverting” topographic river profiles by parameterizing erosion in both time and space (Pritchard et al., 2009), which is somewhat different from conventional geomorphology in which uplift is parameterized to estimate erosion. This method, which has been used to estimate the uplift of the Bié dome, Angola over the last 40 Myr (Pritchard et al., 2009). In this case, the method constrains the uplift history of single watersheds over $\leq 1,000$ km and ≤ 30 Myr, and is thus complementary to the stratigraphic and thermochronological records that provide information over longer time scales and larger spatial extents. An alternative method is that of Guillaume et al. (2009), who estimated the tilt of Miocene terraces to estimate the Neogene uplift of central and Eastern Patagonia, possibly due to the opening of a slab window. The main difficulty of this method is the dating of terraces.

Geological expressions and models of short-wavelength dynamic topography

Mantle-driven dynamic processes may also affect topography at shorter wavelengths (100 km to a few hundred kilometers). A major challenge when studying dynamic topography at such wavelengths is to isolate the signal from the contributions of isostasy and elastic flexure that have similar, kilometer-scale, amplitudes. The most vivid expression of dynamic topography is found in oceanic trenches that are typically < 200 km wide yet several kilometers deep (Husson et al., 2012; Zhong and Gurnis, 1994).

Delamination and convective instability of the continental lithosphere are short-wavelength dynamic processes with the potential for a strong topographic expression. A compelling empirical case is that of the renewed uplift of the southern Sierra Nevada Mountains since the mid-Pliocene (Clark et al., 2005) and localized subsidence within the adjacent Great Valley (Saleeby et al., 2012), which has been related to instability of the continental lithosphere evident in a prominent high seismic velocity anomaly that extends from the base of the crust to about 200 km depth (Boyd et al., 2004; Zandt et al., 2004). Other examples of vertical motions associated with continental lithospheric instability include Quaternary uplift in Baja, California (Mueller et al., 2009), Late Cenozoic uplift of the Willowa Mountains in eastern Oregon, U.S.A. (Hales et al., 2005), and uplift of Miocene marine sediments in the Central Anatolian Plateau, Turkey (Cosentino et al., 2012).

Smaller-scale modes of convection may influence surface topography. A prominent mode evident in models of mantle convection are small, low amplitude

instabilities that travel along the base of the cold, top thermal boundary layer of mantle convection (e.g. Huang et al., 2003). Such traveling convective instabilities might be responsible for vertical motions on a 2-20 Myr time scale evident in sequence stratigraphic analysis of passive continental margins (Petersen et al., 2010). This raises the possibility that such localized small-scale convection could be responsible for high frequency, second to third-order sea level fluctuations. Whether the second and third-order sea level changes evident on passive margins are global or local is controversial (e.g. Hallam, 1992; Lovell, 2010). The upwarped edges of the Colorado Plateau in the western United States, as well as the localization of late Neogene–Quaternary magmatism and steep upper-mantle seismic velocity gradients, are potentially indicative of this mode of sub-lithospheric, small-scale convection (van Wijk et al., 2010).

Although the uplift and high elevation of the Colorado Plateau (e.g. Flowers, 2010) has been interpreted as part of the overall long-wavelength Cenozoic uplift of the western U.S. (see above), it might be viewed as a shorter wavelength component of dynamic topography. The vertical motions evident from uplift of Mesozoic marine strata, unroofing from low-temperature thermochronology (Flowers et al., 2008) and reorganizations of drainage patterns (Wernicke, 2011) might be associated with a long wavelength signal associated with the removal of the negatively buoyant Farallon slab (Liu and Gurnis, 2010) or with the upwelling or warm mantle that extends from the top of the lower mantle into the upper mantle (Moucha et al., 2008b, 2009), or with a combination of both processes.

Locations of active hotspot volcanism correspond to swells of residual topography that have a wavelength of several hundred kilometers and an elevation < 2 km (Crosby

and McKenzie, 2009; Crough, 1983). The surface expression of mantle plumes likely extends beyond this main swell. For instance, lateral magmatic underplating and pulses of hot material in the asthenosphere associated with the Iceland plume have been proposed to explain transgression and regression patterns over < 10 Myr observed in the North Atlantic (Lovell, 2010). A Palaeocene–Eocene landscape off the northwest coast of Europe was recently reconstructed using a three-dimensional seismic dataset on the continental shelf north of Scotland (Hartley et al., 2011; Shaw-Champion et al., 2008). Data suggest that this landscape was lifted above sea level in a series of three discrete steps of 200–400 m each, and that it was reburied after ~ 1 Myr of subaerial exposure, which points to transient uplift and subsidence. These transient vertical motions have been interpreted as a pulse of warm asthenosphere that moves in a thin low viscosity channel radially away from the Iceland plume (Rudge et al., 2008).

MODELING LONG-WAVELENGTH DYNAMIC TOPOGRAPHY

Several approaches have been developed to model regional and global dynamic topography, including models that only consider the present day (instantaneous models), if they consider the time domain at all, time-dependent models, and either a forward or inverse matching of present-day mantle structure. In addition, there has been a variety of studies which have looked at generic issues associated with the physics of dynamic topography and their connection with the rock record (Burgess and Gurnis, 1995; Mitrovica and Jarvis, 1985). Models either use or predict present-day quantities (such as the geoid, mantle seismic structure, surface and/or core-mantle boundary dynamic topography, the rate of change of surface topography, and plate motions) to infer mantle

properties. Inverse models and models that use data assimilation can be compared to fewer observations since assimilated data cannot be used to validate the models. This section constitutes an overview of a number of regional and global scale models of dynamic topography, including their strengths and limitations (Table 2), while also looking at some generic issues associated with how physical properties influence dynamic topography, time-dependence of dynamic topography and its connection to the stratigraphic record.

Instantaneous flow models of present-day dynamic topography

Dynamic topography, h , is the deflection of the earth's surface in response to the normal stresses arising from flow in the mantle, or

$$\sigma_{rr} = \Delta\rho gh \quad (1)$$

where σ_{rr} is the total normal stress at the level surface of mantle flow model (in which an otherwise free surface would be deflected by h), $\Delta\rho$ the density difference between the mantle and either air or water, and g the acceleration due to gravity. Pekeris (1935) was first to point out that surface deformation should result from mantle convection. He advanced a hypothesis of thermal convection in Earth's interior, and pointed out that the surface of the convecting mantle would be pushed upwards over rising currents and pulled downwards over sinking currents, later verified by numerous numerical studies (e.g. McKenzie et al., 1974). In order to link mantle structure, flow, the geoid and surface topography, Parsons and Daly (1983), Ricard et al. (1984) and Richards and Hager (1984) developed analytical solutions, essentially Green's function solutions, for viscous

flow in either Cartesian or spherical domains with simply layered viscosities. This idea can be expressed as

$$h = C\Delta T \quad (2)$$

where ΔT is a temperature scale and C is a function of depth that specifies the depth distribution of temperature (density variations) and lateral and radial variations in viscosity. C is not dependent on the absolute value for viscosity, as first demonstrated by Morgan (1965). An example of the dependence of C on depth, or normalized surface deformation “kernel”, is given in Fig. 3 for a case of the analytical model of Hager and Clayton (1989) in which mantle viscosity is radially distributed in four layers ($\eta_{lith} = 10$; $\eta_{asth} = 1/30$; $\eta_{UM} = 1$; $\eta_{LM} = 1$). The kernels show the efficiency with which density heterogeneities induce dynamic topography decreases with depth, and that this effect is more pronounced at shorter wavelengths (Fig. 3). The surface deformation kernel depends on the assumed viscosity structure. The strong decrease of the deformation kernels with increasing wavenumber results from the combined effect of a low viscosity asthenosphere below the lithosphere and then the subsequent increase in viscosity with increasing depth from the upper mantle through the transition zone and into the lower mantle (Hager and Clayton, 1989), a common feature of proposed mantle viscosity structures (Mitrovica and Forte, 2004; Paulson et al., 2007; Steinberger and Calderwood, 2006).

Hager (1984) used the approach given by Eq. 2 with a thermal model of slabs consistent with Benioff zone seismicity and the observed geoid to constrain a minimum 30-fold increase of viscosity from the upper to lower mantle. The approach was then

elaborated on using a mantle density structure scaled from global seismic tomography inversions (Hager et al., 1985), arguing that chemically stratified mantle convection could be ruled out because it produced a pattern of dynamic topography with broad subsidence over Africa. To fit the observed geoid, Hager et al. (1985) revised the minimum viscosity increase from upper to lower mantle to a factor of 10. In contrast, Lithgow-Bertelloni and Silver (1998) fit the residual topography of southern Africa with the dynamic topography predicted by an instantaneous mantle flow model with a lower mantle 50 times more viscous than the upper mantle. These differences in best-fit mantle viscosity structures are not surprising considering that the two flow models are based on different tomography models and that Hager et al. (1985) included all seismic-velocity anomalies as density anomalies in their calculation of surface dynamic topography whereas Lithgow-Bertelloni and Silver (1998) calculated the surface dynamic topography resulting from density heterogeneities below 325 km depth.

Steinberger (2007) and Conrad and Husson (2009) derived the global dynamic topography predicted by instantaneous numerical mantle flow models (Fig. 4B and 4D). These two models, hereafter referred to as S07 and CH09, are broadly similar in that they are based on a mantle density structure derived from S-wave tomography with the assumption that all lateral variations in seismic velocities are thermal, while ignoring density contrasts in the uppermost mantle (220 km in S07; 300 km in CH09). Important differences between these two models include the tomographic inversion, the scaling factor to convert seismic velocity to density anomaly, the surface velocity boundary conditions and the viscosity structure. In addition, S07 is a semi-analytical model using a spherical harmonics approach (Hager and O'Connell, 1979) with a horizontal resolution

of ~ 650 km whereas CH09 is a numerical model using a finite-element approach, *CitcomS* (Zhong et al., 2000), with a horizontal resolution of ~ 100 km. These differences illustrate the large number of free parameters and boundary conditions inherent to mantle flow models.

Despite the major differences between these two models, the present-day dynamic topography fields they predict are overall similar (Figs. 4B and 4D), and compatible with proposed residual topography fields (Fig. 1). Both predicted dynamic topographies show highs within the Pacific, and southern and eastern Africa. The main dynamic topography lows extend under Central and South America, and from Australasia to eastern Europe with a minimum under southeast Asia. The main differences in terms of patterns are the dynamic topography lows in the central South Atlantic, in northwest Africa and in the eastern US predicted by S07 but not by CH09. All three of these dynamic topography lows are observed in residual topography fields (Fig. 1), although the amplitudes are substantially larger in S07 compared to the CH09 and residual fields (Fig. 1). In addition, neither S07 nor CH09 reproduces the residual topography low in the northeast Pacific Ocean.

An important difference between S07 and CH09 is the amplitude of the calculated dynamic topography (Table 1), which has been the subject of a long-standing debate (e.g. Gurnis, 1990). The dynamic topography predicted by model S07 (CH09) presents a peak-to-peak amplitude of ± 3 km (± 1.5 km) and a root-mean-square amplitude of ~ 0.9 km (~ 0.5 km; Table 1). For comparison, the peak-to-peak amplitude of the residual topography fields shown in Fig. 1 is approximately ± 2 km, and their root-mean-square amplitude ~ 0.5 km (Table 1). Therefore, the dynamic topography predicted by

model CH09 (S07) is 25 % smaller (50 % larger) than that of residual topography. The main parameters explaining the amplitude difference between the models are the scaling factor from seismic to density anomaly, greater in model S07, the depth above which density contrasts are ignored, 80 km shallower in S07, and the low-viscosity asthenosphere present in CH09 and not in S07. Steinberger (2007) showed that including the effect of phase transitions in such models only reduces the root-mean-square amplitude of dynamic topography by $\sim 5\%$. Other important effects to investigate include accounting for chemical heterogeneity when converting seismic velocity anomalies to density anomalies (Simmons et al., 2007) and the effect of lateral viscosity variations (Zhong and Davies, 1999). At long wavelengths, lateral variations in viscosity might be small compared to the uncertainty of tomography models, but it is expected to be significant in time-dependent mantle flow models that are largely driven by the evolution of thermal heterogeneity (Moucha et al., 2007).

Still focusing on instantaneous models, the rate of change of dynamic topography provides complementary constraints. Since dynamic topography depends on how fast mantle buoyancy moves in the mantle, Gurnis et al. (2000) showed that

$$\dot{h} = \frac{C'}{\eta} \Delta T^2 \quad (3)$$

where $\dot{h}$ is the rate of change of dynamic topography, C' is a constant related to C in Eq. 2 and therefore to the distribution of buoyancy and viscosity, while η is a reference absolute value of viscosity and ΔT is a temperature scale. In other words, in contrast to just dynamic topography, which does not depend on the absolute viscosity (Morgan, 1965), the rate of change does. These ideas were captured in a numerical model of global,

present-day mantle flow that forward predicted present-day uplift rates. From models matching the constraints on uplift rate of southern Africa, (Gurnis et al., 2000) argued that the mid-lower mantle was 0.2 % less dense than average - much less than would be expected if shear velocities were solely governed by temperature - but with a viscosity increase of a factor of 10 from upper to lower mantle, which again shows that best-fit parameters vary largely between models.

Time-integrated and time-dependent flow models of past and present-day dynamic topography

The previous section shows that present-day dynamic topography can be estimated using the present-day density structure of the mantle as constrained by tomography models. However, a major challenge in estimating the evolution of dynamic topography is that the past density structure of the mantle is not directly known. Several strategies using data assimilation have been developed over the last two decades to model the evolution of mantle flow and of dynamic topography. These strategies include forward, backward and adjoint mantle flow models based on the history of subduction and plate motions given by global tectonic reconstructions, or on mantle tomography, or on a combination of both.

Models of dynamic topography based on subduction history

Analytical models.

Present-day dynamic topography. Ricard et al. (1993) developed a time-dependent analytical model of mantle flow in which subduction history (Lithgow-Bertelloni and Richards, 1998; Lithgow-Bertelloni et al., 1993) was imposed over the last

200 Myr. In this model, slabs are treated as “stokeslets” sinking vertically at a constant rate and slowing down as they enter the lower mantle, while neglecting thermal diffusion. The best-fit model of Ricard et al. (1993) to the observed geoid was obtained for a lower mantle 40 times more viscous than the upper mantle. This estimate based on a forward model is closer to the analytical result of Hager (1984; lower mantle 30 times more viscous than upper mantle) than the best-fit model to the geoid of Hager et al. (1985) with a lower mantle only 10 times more viscous than the upper mantle. Again, such variations in results are not surprising given the differences between the three approaches. The present-day dynamic topography predicted by the model of Ricard et al. (1993) is shown in Fig. 4C. It has a minimum of -1.6 km, a maximum of ~ 600 m and a root-mean square amplitude of ~ 430 m (Table 1). The skew of this dynamic topography field towards negative values is due to the model being driven by slabs, while the return mantle flow is entirely passive: negative dynamic topography is of larger amplitude and covers a smaller area than positive dynamic topography (Fig. 4C). As a consequence, the amplitude of negative dynamic topography predicted by this model is about 20% smaller than negative residual topography (Fig. 1), whereas positive dynamic topography is only 30% of positive residual topography. In terms of spatial distribution, the dynamic topography lows predicted by the forward model of Ricard et al. (1993) under Central/South America and southeast Asia match that predicted by the inverse models of Steinberger (2007) and Conrad and Husson (2009). However, the dynamic topography low under the Gulf of Alaska and Bering Sea predicted by Ricard et al. (1993) is not predicted by inverse models, and it is only observed in one of the four residual

topography fields (Fig. 1), which suggests an overestimation of the amount of subducted material in that area in the plate model used by Ricard et al. (1993).

Evolution of dynamic topography. Gurnis (1993a) developed time-dependent models of mantle flow based on the analytical formulation proposed by Richards and Hager (1984) using the Phanerozoic subduction history of Scotese et al. (1993) to investigate the influence of dynamic topography and its time derivative on continental flooding. This study revealed that a significant part of the observed Phanerozoic continental flooding could be attributed to dynamic subsidence above active subduction zones, suggesting that dynamic topography affects global sea level change at long wavelengths, and therefore that Phanerozoic stratigraphy can be used to constrain mantle flow. Building on this approach, Lithgow-Bertelloni and Gurnis (1997) used a series of five instantaneous global flow models throughout the Cenozoic to reproduce the overall observed trend of flooding and exposure of North America, Australia and Indonesia, although the amplitude of model uplift and subsidence were greater than that inferred from stratigraphic constraints.

Forward numerical models. Gurnis et al. (1998) developed numerical regional mantle convection models with imposed plate kinematics to investigate the evolution of the dynamic topography of the Australian plate since the Cretaceous. Gurnis et al. (1998) used the location of subduction taken from plate reconstructions (Müller et al., 1997) to build a synthetic initial temperature field for their forward models. They reproduced the regional Cretaceous flooding of Australia deduced from paleo-geographic reconstructions in which it had long been known that eastern Australia achieved peak flooding with shallow marine seas about 20 million earlier than inferred peak global sea level (see

above; Veevers and Conaghan, 1984). Essentially, in the Early Cretaceous Australia drifted eastward over the long-lived subduction zone bounding Gondwanaland, which caused widespread marine inundation in the central and eastern part of the continent. Sediments filled the accommodation space as the continent slowly rose following termination of subduction such that when global sea level peaked in the Late Cretaceous there was no remaining accommodation space, resulting in an absence of any major inundation. Gurnis et al. (1998) also showed that the position and history of the equally anomalous Australian-Antarctic Discordance (AAD; Fig. 1), a prominent present-day north-south residual topography low, could also be fit with the same model. The study is notable for two reasons. First, this rather straightforward computational approach linking plate tectonics and thermal convection showed that an entire cycle of dynamic topography could be followed from subsidence to uplift. As we shall see, most other regional applications can only follow part of the entire history (either the subsidence or the uplift phase). This is not a criticism of modeling methods, but reflect fundamental limitations of the geological record. In addition, the study reinforced the need to follow the history of vertical motions in the plate frame of reference, that is tracking the history of a point on a plate that moves with respect to the mantle. This is essential for the detailed comparison of models to geological proxies of vertical motions.

DiCaprio et al. (2010) tracked the evolution of dynamic topography for the Australian plate in a kinematically-driven forward model with assimilation of plate velocities in one million year increments in which a high-resolution regional model was embedded in a global model. This model, starting at 50 Ma with subduction zones inferred from plate reconstructions, reproduced the observed fast Neogene subsidence of

the Australian northeast shelf that would have contributed to the demise of carbonate reefs (DiCaprio et al., 2010). DiCaprio et al. (2010) were also able to exploit new paleogeographic tools that allowed the plate boundaries to smoothly evolve (Gurnis et al., 2012), essential for time-dependent convection models. Zhang et al. (2012) proposed a kinematically-driven forward model extending back to 450 Ma and of average global surface resolution ~ 100 km. The plate reconstruction used in this model consists of 34 stages over the last 450 Ma, and the initial condition was obtained by running the first reconstruction stage for 150 Myr. The vertical motions predicted by this model for the Kaapvaal and Slave Cratons were in agreement to the first-order with the burial histories predicted by thermochronology models.

A new model. We introduce a forward model in which we assimilate the thermal and kinematic field associated with plate reconstructions (Seton et al., 2012) with continuously closing plates (Gurnis et al., 2012) prepared using the software *GPlates* (Boyden et al., 2011). This model is essentially the extension of the approach of Ricard et al. (1993), Lithgow-Bertelloni et al. (1993) and Lithgow-Bertelloni and Richards (1998) to “a full three-dimensional convection problem constrained by plate motions” (Ricard et al., 1993). The velocity field of plate reconstructions is continuously imposed as surface boundary condition by linear interpolation between reconstruction stages defined every million year. The temperature field of the oceanic lithosphere is assimilated using paleo-oceanic ages (Seton et al., 2012) and the half-space cooling model (e.g. Davis and Lister, 1974). One limitation of global mantle convection models with imposed plate velocities is that they result in substantial advective thickening of slabs instead of well-defined one-sided subduction (Table 2). To overcome this

659 limitation, we assimilate the temperature field in the proximity of the subduction zones
660 given by the plate reconstructions. The thermal structure of slabs is derived from the age
661 of the subducting lithosphere using a half-space cooling formulation and assuming a slab
662 dip angle of 45° . This analytical solution is assimilated in the dynamic model within
663 350 km of subduction zones using a blending function. In addition, we ensure that
664 subduction zones appearing during the model are progressively inserted into the upper
665 mantle. The method, ensures that the slab buoyancy flux in the upper mantle is consistent
666 with the plate reconstructions, and avoids unrealistic slab advective thickening (Bower et
667 al., in preparation). The model, solved using *CitcomS* (Zhong et al., 2000), consists of
668 12×128^3 elements and presents an average surface resolution of ~ 50 km. The initial
669 condition (at 200 Ma) consists of a global temperature field (non-dimensional
670 background of 0.5) with cold slabs inserted to a depth of 1750 km and an assimilated cold
671 surface thermal boundary layer as described above (see Zehirovic et al., 2012 for a cross-
672 section of the initial condition under the Tethys). The temperature-dependence of
673 viscosity is implemented using an Arrhenius law with activation energy of 100 kJ/mol
674 (33 kJ/mol) in the upper (lower) mantle, which limits viscosity variations to two orders of
675 magnitude, and in our reference case the lower mantle reference viscosity is 100 times
676 larger than that of the upper mantle (see Zehirovic et al., 2012 for the resulting viscosity
677 profile). The surface dynamic topography is computed by restarting the calculation every
678 5 Myr with no-slip surface boundary conditions to suppress the effect of the surface
679 traction imposed by plate velocities (see Thoraval and Richards, 1997 for a discussion of
680 the effect of model boundary conditions on predicted dynamic topography) and after
681 removing the uppermost 350 km of the mantle, depth to which subduction zones are

682 assimilated.

683 The resulting present-day dynamic topography field (Fig. 4A), broadly similar to
684 that of Ricard et al. (1993; Fig. 4D), has a degree-two pattern with a belt of low dynamic
685 topography extending from Antarctica through the Americas to the Arctic and broadening
686 beneath the whole Eurasian continent. In this subduction-driven model, the dynamic
687 topography highs correspond to passive upwelling under the Pacific Ocean, southern
688 Africa and the Atlantic Ocean. This dynamic topography field is in overall agreement
689 with the residual topography in terms of pattern (Fig. 1) and amplitude (between -2.2 km
690 and +1.6 km; Table 1). Although the mean dynamic topography is zero, the modeled
691 highs have lower amplitude than lows because upwelling is passive. The model features
692 dynamic topography lows under central Europe, eastern North America, and western
693 South America, and a dynamic topography high under southern Africa, all evident in the
694 residual topography (Fig. 1). The dynamic topography under the Argentine Basin and the
695 western United States has the expected sign but its amplitude is lower than expected. The
696 AAD is also of lower amplitude than residual topography, and it is shifted to the west.
697 This discrepancy is tempered by the fact that only about half the excess depth of the AAD
698 is likely dynamic, the rest being due to thin oceanic crust within the AAD (Gurnis et al.,
699 1998; Whittaker et al., 2010). In addition, the positive dynamic topography predicted
700 under the Northeast Pacific Ocean is at odds with the large-scale, low-amplitude residual
701 topography low (Fig. 1). Finally, the residual topography highs associated with plumes
702 in eastern Africa, the western Pacific (Darwin Rise) and Iceland are not reproduced by
703 the model that only captures large-scale, passive mantle upwelling.

704

Time-dependent models of dynamic topography based on mantle-tomography

Backward advection. In order to investigate the time-dependence of mantle flow and dynamic topography, a natural extension of instantaneous flow models consists of backward advecting the mantle density structure derived from tomography. The process essentially consists in changing the sign of gravity in the momentum equation and setting the non-reversible thermal diffusivity to zero in the energy equation (Steinberger and O'Connell, 1997). Some caution is needed since the method accumulates light material at the core mantle boundary, and dense material towards the surface, which results in a decrease of information on the density structure of the mantle back in time. A further limit of backward advection is that thermal diffusion cannot be simply reversed (Ismail-Zadeh et al., 2004). However, simple backward integration is adequate for long wavelengths where advection dominates over diffusion. The first study to model dynamic topography using this approach was that of Conrad and Gurnis (2003) who proposed that the dynamic topography high currently located under southern Africa migrated from eastern Africa over the last 36 Ma. They also pointed out the importance of thermal boundary layers in the backward advection process, and used a half-space cooling model to assimilate oceanic ages in the top thermal boundary layer. This backward advection scheme, in which plate velocities are also assimilated, was reversible for ages up to between 50 and 75 Ma (Conrad and Gurnis, 2003).

Moucha et al. (2008a) estimated dynamic topography both at the surface, following the definition of Forte et al. (1993), and beneath the lithosphere, similarly to the definition used here, from a backward advection model in which data were assimilated in several ways. Firstly, the tomography model used by Moucha et al. (2008a) is a hybrid

tomography/instantaneous mantle flow model that assimilates free-air gravity anomaly, the “residual” topography field corrected for the isostasy of the crust, present-day plate velocities from DeMets et al. (1990), and the excess ellipticity of the core-mantle boundary (Simmons et al., 2007). Secondly, the viscosity profile is constrained using the same convection related observables as above, as well as data related to glacial isostatic adjustment (Mitrovica and Forte, 2004). Thirdly, although plate motions are predicted by the flow rather than driving the flow in the formulation of Moucha et al. (2008a), mantle flow is backward advected using boundary conditions consistent with a no-net-rotation tectonic plate model. Aside from the data assimilation, the model of Moucha et al. (2008a) does not seem to include any specific treatment of thermal boundary layers, important in backward mantle advection schemes (Conrad and Gurnis, 2003), nor of lateral viscosity variations or temperature-dependence of viscosity, important in time-dependent flow models (Moucha et al., 2007). Based on this model, Moucha et al. (2008a) proposed that the New Jersey margin has been dynamically uplifted by 100-200 m over the last 30 Myr, and its conjugate margin in northwest Africa has been uplifted by ~ 100 m during the same period. The important consequence of this work is that sea level curves derived from the well-documented stratigraphy of the New Jersey margin (Miller et al., 2005; Watts and Thorne, 1984) likely contain a dynamic component and cannot be used alone to derive eustatic sea level curves. We will return to how different models fit the New Jersey constraint in the Discussion.

Adjoint models. To overcome the shortcoming of backward advection in dealing with conductive thermal boundary layer, adjoints of the governing equations can be introduced (Bunge et al., 2003; Ismail-Zadeh et al., 2004; Liu and Gurnis, 2008). The

method consists in refining the assumed initial condition by iteratively solving the forward and backward adjoint and minimizing a cost function (in this case between seismic tomography and forward predicted temperature). The main advantage of this method is that it uses a forward solution of the coupled equations of motion and energy while converging upon the known tomographic structure. A significant limitation is the computational cost as a single inversion requires multiple forward and adjoint solutions of the equations and is substantially more expensive than a backward run (Bunge et al., 2003; Table 2).

Liu et al. (2008) and Spasojevic et al. (2009) used a variant of this method to constrain mantle properties finding adjoint models (starting from present day seismic tomography) that best fit the flooding history of North America (Fig. 5A). In these models, the mantle structure is scaled from shear wave seismic tomography (Grand, 2002) between 250 km and 2,400 km depth, and the plate velocities of continuously closing plates (Gurnis et al., 2012). In addition, the flat-lying, continuous Cretaceous Farallon slab could only be backward advected (Fig. 5C) by imposing a viscous stress guide (Liu et al., 2008; Spasojevic et al., 2009). The dynamic topography (Fig. 5B) and its rate of change were used to predict the spatial pattern of marine inundation (Fig 5A). This was then compared to the observed flooding in the western interior seaway and to subsidence rates deduced from borehole data, respectively. The scaling factor between mantle tomography and density structure was constrained by minimizing the misfit between the amplitude of observed and predicted subsidence. The viscosity of the upper mantle was constrained by minimizing the misfit between observed and predicted rates of subsidence. The best fit between model output and subsidence-related observables was

obtained for an upper mantle reference viscosity of 1×10^{21} Pa s and a lower mantle reference viscosity of 1.5×10^{22} Pa s (Liu et al., 2008; Spasojevic et al., 2009). Significantly, the inferred upper mantle viscosity is within the range inferred from postglacial rebound studies (Cathles, 1975; Mitrovica and Forte, 2004). Finally, the best fitting model from (Liu et al., 2008) successfully reproduced the observed craton-ward migration of the prominent depocenter within the Cretaceous section of the interior of North America (Fig. 2C-F).

Hybrid models. Spasojevic and Gurnis (in press) recently proposed a hybrid formulation in which the density structure of the lower mantle is entirely derived from backward advection of a seismic tomography model, whereas in the upper mantle the positively buoyant anomalies are backward advected from seismic tomography and the negatively buoyant subduction zones are imposed from tectonic reconstructions and oceanic ages (Müller et al., 2008a). Spasojevic and Gurnis (in press) carried out a series of forward instantaneous flow models, hereafter referred to as SG12, from 90 Ma in 10 Myr increment based on these density structures. They compared the predicted change in global dynamic topography for several models with different viscosity structures to the uplift and subsidence maps deduced from paleo-geographic reconstructions (Blakey, 2008; Smith et al., 1994) and noted that different models compare best to data in different areas and suggested that the radial viscosity structure of the mantle is not laterally uniform.

This dynamic topography field of Spasojevic and Gurnis (in press; Fig. 4E) is similar to that proposed by Conrad and Husson (2009) which is expected since model CH09 is instantaneous and model SG12 carries no information from the past to the

present. Both models are also based on the same tomography model, S20RTS. The main difference between the two models is the amplitude of dynamic topography lows that is ~ 1.5 times larger in the model of Spasojevic and Gurnis (in press), which reflects (i) that the uppermost 300 km of the mantle have been ignored in CH09 whereas the 250 km uppermost km have been removed in SG12 and (ii) that the scaling factor to convert seismic velocity anomaly to density anomaly is depth-independent and equal to $0.15 \text{ g cm}^{-3} \text{ km}^{-1} \text{ s}$ in CH09 whereas it varies with depth between 0.05 and $0.3 \text{ g cm}^{-3} \text{ km}^{-1} \text{ s}$ in SG12.

DISCUSSION

Comparison between present-day global residual and dynamic topography

The good first-order agreement between four global residual topography fields (Fig. 1), derived independently and based on different assumptions, is encouraging. The most prominent and robust residual topography features are (i) the lows extending from central Europe to southeast Asia, from eastern North America to western South America, the Australian-Antarctic Discordance (AAD), the Argentine Basin, and (ii) the highs under southern and eastern Africa, Iceland and the Darwin Rise.

The global dynamic topography predicted by five numerical mantle flow models (Fig. 4), based on two distinct methods and a hybrid of those, are also in overall agreement with each other, and with the residual topography (Fig. 1). All of these dynamic topography fields, with the exception of Steinberger (2007), are to some extent skewed towards negative values (Table 1), reflecting that negative dynamic topography

819 anomalies are larger in amplitude but cover smaller areas than positive dynamic
820 topography anomalies (since the mean of each field is zero). The wavelength of model
821 dynamic topography is overall larger than that of proposed residual topographies (Fig. 1).
822 The model of Steinberger (2007), based on a compilation of mantle tomography models,
823 presents shorter wavelengths than models by Conrad and Husson (2009) and Spasojevic
824 and Gurnis (in press) based on S20RTS (Ritsema et al., 2004) (Fig. 4). For instance, the
825 dynamic topography high under Iceland and the Azores (Conrad et al., 2004) is distinct in
826 the model of Steinberger (2007), subdued in the model of Conrad and Husson (2009) and
827 absent in Spasojevic and Gurnis (in press) (Fig. 4). Dynamic topography lows in the
828 equatorial South Atlantic ocean are only captured by the model of Steinberger (2007)
829 albeit with a larger amplitude than suggested by residual topography. With more robust
830 processing techniques, especially with full waveform seismic tomography, future mantle
831 tomography models will improve our understanding of global dynamic topography down
832 to $\sim 1,000$ km scale, whereas present global studies are limited to a resolution of $\sim 3,000$ -
833 $5,000$ km (Fig. 4).

834 Both synthetic time-dependent forward mantle flow models (Figs. 4A and C) and
835 instantaneous models based on mantle tomography (Figs. 4B, D and E) predict dynamic
836 topography lows under the continents, with the notable exception of Africa south of the
837 equator and of western North America (out of the five models only that of Ricard et al.
838 (1993) does not predict positive dynamic topography for western North America).
839 Interestingly, all mantle flow models predict a dynamic topography low under southeast
840 Asia that is more pronounced than suggested by residual topography studies, and negative
841 dynamic topography along the east coast of Asia where the residual topography is

positive (Figs. 1 and 4). One limit to constraining residual topography in southeast Asia and along the east coast of Asia is the small extent of oceanic crust, limited to the South China, Sulu, Celebes and Banda Seas in southeast Asia and to the Sea of Japan and southernmost Sea of Okhotsk along the east coast of Asia (see COB contours in Fig. 1A). Because we have assigned a uniform elevation of - 200 m to areas between oceans and continents, our proposed residual topography features a regional residual topography low in southeast Asia and along the east coast of Asia (Fig. 1A) whereas other studies (Figs. 1B, C and D) in which these areas are treated as continental suggest positive residual topography in the Gulf of Thailand, Borneo and the east coast of Asia. The consistency of the dynamic topography low under southeast Asia and the east coast of Asia in both instantaneous flow models (Figs. 4B, D, and E) and forward mantle flow models (Figs. 4A, and C) suggests that the seismically fast mantle is reproduced by forward models. Given the poor control on residual dynamic topography in these regions, dynamic topography models for south and east Asia should be compared to time-dependent observables (e.g. Xie et al., 2006) and to mantle tomography in the case of synthetic forward models. However, southeast Asia is the region with the largest extent of shallow marine seas and such shallow seas are strongly correlated with predicted present-day dynamic topography lows (Gurnis, 1993b).

All mantle flow models predict positive dynamic topography for the entire Pacific Ocean (Fig. 4), at odds with residual topography studies that consistently predict a large-scale, low-amplitude low in the northeast Pacific (Fig. 1). An anti-correlation between residual gravity and residual topography is known for this region (Crosby and McKenzie, 2009), which suggests a dynamic origin for the residual topography low, but there is no

fast seismic anomaly structure under the northeast Pacific (Crosby and McKenzie, 2009). Indeed, Spasojevic et al. (2010) attributed the gravity lows under the northeast Pacific to mid- to upper-mantle upwelling above slab graveyards, thereby explaining the absence of large fast seismic velocity anomalies directly under this region and suggesting that the northeast Pacific could be presently uplifting. Perhaps this is a region with negative residual topography that is experiencing dynamic uplift (on its way up after being drawn down). Therefore, the mismatch between model dynamic topography and present-day residual topography in the northeast Pacific illustrates the limitation in using a snapshot of residual topography to constrain dynamic topography that occurs over time scales of several tens of million years. This underlines the concept that the time evolution of dynamic topography is arguably more important than its present day sign and amplitude, and therefore that quality time-dependent proxies should be favored as model constraints.

Finally, the position of the AAD is well reproduced by instantaneous mantle flow models based on tomography (Figs. 4B, D, and E) but only at long wavelengths by forward models (Figs. 4A and C).

Consequences of dynamic topography for long-term sea level change

Arguably the most important consequence of dynamic topography studies is their influence on eustatic and relative sea level. In this section, we first discuss the effect of global dynamic topography on global sea level and then turn to the regional effects of dynamic topography.

Global dynamic topography and global sea level change

Changes in dynamic topography could have a first order influence on global sea level (Gurnis, 1992). Using a simple model of mantle flow associated with slabs, Gurnis (1993c) computed that the shifting motion of continents over the evolving long wavelength dynamic topography would give rise to a 100-200 m eustatic change with two highs in the mid-Paleozoic and late Mesozoic. Husson and Conrad (2006) developed a model based on boundary layer theory to propose that changes in dynamic topography associated with geologically rapid (≤ 20 Myr) changes in mean global tectonic velocities would only affect eustatic sea level by ~ 20 m, which is relatively small in comparison to eustatic changes of several hundred meters associated with changes in the age-distribution of the ocean floor (e.g. Hays and Pitman, 1973; Müller et al., 2008b; Spasojevic and Gurnis, in press). Husson and Conrad (2006) proposed that the dynamic effect of longer-term (10^8 yr) change in tectonic velocities on sea level could be of up to ~ 80 m.

Extrapolating from the present-day rate of change of dynamic topography, Conrad and Husson (2009) considered the effect on eustasy of continental motion across dynamic topography gradients during the aggregation and dispersal of a supercontinent. The most significant effect would occur during the dispersal phase, when fragments move away from the upwelling expected beneath a mature supercontinent (Coltice et al., 2007; Gurnis, 1988) towards dynamic topography lows associated with subduction zones. Conrad and Husson (2009) estimated that dynamic topography (Fig. 4D) currently triggers a net uplift of the ocean basins thereby contributing to a positive sea level offset of $\sim 90 \pm 20$ m. In addition, they carried out a 1-Myr-long forward model of mantle flow

910 based on the present-day mantle structure and plate motions to propose a current rate of
911 sea level rise induced by dynamic topography of < 1 m/Myr.

912 Spasojevic and Gurnis (in press) simultaneously calculated dynamic topography,
913 the shifting of present-day continental isostatic topography with the plates and the
914 changing age-distribution of the ocean floor, while forcing the water load to fit the
915 evolving geoid in a discrete series of ten instantaneous mantle flow models between
916 90 Ma and present. Based on these calculations, they estimated that sea level has risen by
917 between ~ 110 m and ~ 270 m since 90 Ma because of global dynamic topography,
918 including ~ 110 -200 m caused by sea floor uplift and < 70 m due to continental
919 subsidence. Thus, dynamic topography influences global sea level by offsetting (by
920 about one third) the decrease in sea level imposed by changes in the age-distribution of
921 the ocean floor. Dynamic topography should therefore not be neglected in studies of
922 long-term global sea level change. In computational models, it is possible to create a
923 global sea level curve by averaging the height of all continents with respect to sea level.
924 Things are more complex in reality because large amplitude swings in dynamic
925 topography could have a disproportionate influence on the stratigraphically inferred
926 record of sea level change. This is particularly true on a regional scale, at which both
927 dynamic and tectonic topography often dominates the stratigraphic signal.

928 *Case Study: Has the New Jersey margin been dynamically uplifting or subsiding since*
929 *the Cretaceous?*

930 Another important consequence of dynamic topography studies is that sea level
931 curves defined at a single “stable” passive continental margin (or indeed at any location)

932 may not be representative of eustasy if the margin has been experiencing differential
933 vertical dynamic motion. The most prominent example is the sea level curve proposed by
934 Miller et al. (2005) based on the careful backstripping of five wells from the New Jersey
935 coastal plains which predicts a much smaller decrease in sea level since the Late
936 Cretaceous than other global sea level curves (for details see Müller et al., 2008b;
937 Spasojevic et al., 2008). Recent dynamic topography studies all suggest that the New
938 Jersey margin, although considered tectonically stable, has undergone significant vertical
939 motions driven by mantle flow since the Late Cretaceous (Moucha et al., 2008a; Müller
940 et al., 2008b; Spasojevic et al., 2008). Based on their backward advection model
941 described above, Moucha et al. (2008a) proposed that the New Jersey margin has been
942 dynamically uplifting by ~ 100 to 200 m over the last 30 Ma. In contrast, Spasojevic et
943 al. (2008) argued for dynamic subsidence of the New Jersey margin by > 300 m since
944 55 Ma, including ~ 100 to 200 m over the last 30 Ma, based on the adjoint model of Liu
945 et al. (2008) described above. Finally, Müller et al. (2008b) estimated that the New
946 Jersey margin has subsided by 105 to 385 m since 70 Ma using backward advection
947 models based on three different tomography models. Müller et al. (2008b) considered
948 two types of models: “pure” backward advection models and “modified” backward
949 advections models in which upwellings are continued upward to 220 km depth and
950 downwellings are removed from the uppermost 220 km. For the last 30 Ma, their “pure”
951 backward advection models predict between ~ 50 m and ~ 150 m of margin subsidence
952 while their modified backward advection models predict between 0 and ~ 70 m of margin
953 subsidence, and ~ 90 m of subsidence between 30 Ma and 8 Ma followed by 40 m of
954 uplift since 8 Ma for the tomography model of Grand (2002).

Comparison between observed tectonic subsidence and predicted dynamic topography for well COST-B2. To shed some light on this controversy, we focus on the tectonic subsidence history of the COST B-2 well offshore New Jersey (Scholle, 1977) that we compare to the evolution of dynamic topography predicted by published models and by the model presented in this study.

Subsidence of well COST-B2. Figure 6 shows the tectonic subsidence of well COST B-2 with respect to present-day obtained by backstripping the stratigraphy (courtesy M. Kominz) for two end-member cases of sea level change. Well COST-B2 shows a marked acceleration in post-rift subsidence at ~ 16 Ma (Fig. 6A; APPENDIX 1) that is also evident in strain rate inversion (Fig. 6B; APPENDIX 1) and anomalous subsidence (Fig. 6C; APPENDIX 1) analysis. In addition, both strain rate inversion and anomalous subsidence analysis reveal a phase of anomalous subsidence between 85 and 15 Ma (Figs. 6B, 6C and 9). The sign of this anomalous subsidence depends on the eustatic sea level curve used in backstripping, with changing (constant) sea level resulting in subsidence (uplift), respectively (Figs. 6C and 9). The hypothesis of constant sea level is difficult to reconcile with the growth of ice sheets that would have triggered a sea level drop by 54 m over the last 33 Myr (Miller et al., 2005), and with the increase in the volume of ocean basins that would have caused a sea level drop of at least 120 m since 85 Ma (Müller et al., 2008a; Spasojevic and Gurnis, in press). This suggests that well COST-B2 has been subsiding rather than uplifting over the last 85 Myr, which is in agreement with the subsidence of New Jersey paleo-shorelines by 50-200 m since the Eocene (Spasojevic et al., 2008). Given the time scale (tens of Myrs), this anomalous

977 subsidence could result from dynamic subsidence induced by long-wavelength mantle
978 flow.

979 *Comparison with the dynamic topography predicted by the model from this study.*

980 Figure 7 shows the evolution of dynamic topography predicted by our forward model at
981 151 Ma, 100 Ma, 49 Ma, and present-day along with the coastlines and subduction zones
982 reconstructed through time with respect to the mantle. The predicted two-degree pattern
983 of the dynamic topography field is stable through time, and in this subduction-driven
984 model the main changes in dynamic topography are the location, width and amplitude of
985 dynamic topography lows. Well COST-B2 (Figs. 1, 4 and 7) is initially located over a
986 dynamic topography high and drifts towards the broadening dynamic topography low
987 associated with the old Farallon slab as North America drifts to the west (Figs. 7B and
988 C). Our model predicts that well COST-B2 is currently located on the northeast edge of a
989 mild dynamic topography low (Fig. 7D), which is consistent with other dynamic
990 topography models (Fig. 4) and in broad agreement with estimates of residual topography
991 (Fig. 1), in which the topography low is more pronounced. In contrast to well COST-B2,
992 our model predicts that the synthetic well located on the conjugate margin in Western
993 Sahara would have remained on a dynamic topography high since 200 Ma (Fig. 7). For
994 well COST-B2, the predicted evolution of dynamic topography shows an initial uplift by
995 ~ 150 m in two stages (200 - 170 Ma and 140 – 120 Ma), followed by 850 m of dynamic
996 subsidence since 120 Ma (Fig. 8). This predicted dynamic subsidence of well COST-B2
997 since 120 Ma is consistent with the anomalous strain and with the anomalous subsidence
998 discussed above (Figs. 6 and 9). The model predicts 580 m of dynamic subsidence since
999 85 Ma, which is ~ 30 % greater than the anomalous subsidence deduced from

backstripping using the sea level curve of Haq et al. (1987; Fig. 8; Appendix 1) and only 30 m of subsidence over the last 15 Ma, which is four to six times smaller than the anomalous subsidence for that period (Fig. 9, Appendix 1). As for the conjugate margin, our model predicts that it was uplifted by 250 m from 200 Ma to 110 Ma and that it has subsided by 170 m since 90 Ma (Fig. 8). This predicted subsidence is five times less than that of well COST-B2 (Fig. 8), which confirms that the margin of north west Africa has not been subject to as much post-rift vertical motion as the New Jersey margin (Moucha et al., 2008a). Unfortunately, little data are available to estimate the vertical motions of northwest Africa, particularly for the conjugate margin of New Jersey. Nevertheless, the subsidence of northwest Africa predicted by our models is to first-order compatible with the marine transgression since 40 Ma deduced from paleo-geographic analysis for that area (Spasojevic and Gurnis, in press) and with the moderate anomalous subsidence of the central Atlantic margin in the Senegal Basin (Brun and Lucazeau, 1988). In addition, the relative vertical stability of northwest Africa above a dynamic topography high predicted by our models is in agreement with the thin onshore Cenozoic stratigraphic thickness (~ 70 m) of the Senegal Basin (e.g. Swezey, 2009).

The dynamic subsidence inferred over tens of Myrs for well COST-B2 and from paleo-shoreline analysis in New Jersey appears to be in contradiction with recent uplift reported further south on the east coast of North America (Rowley et al., 2011). Dowsett and Cronin (1990) reported recent uplift along an Early Pliocene wave-cut scarp in South Carolina and North Carolina (see Fig. 7 for location) that Rowley et al. (2011) extended from Georgia to Virginia. Together, these studies suggest up to ~ 50-65 m of uplift since ~ 3.5-3.0 Ma over ~ 500 km in an area located several hundred kilometers south of well

COST-B2. The evolution of dynamic topography predicted by our model at one of the sections reported by Dowsett and Cronin (1990) shown on Fig. 8 is overall similar to that predicted for well COST-B2. An important difference, however, is that the North Carolinian location is predicted to have been dynamically uplifted by ~ 20 m over the last 25 Ma, a period during which well COST-B2 is predicted to have subsided by ~ 50 m. This result underlines the difficulty of identifying the evolution of dynamic topography in the geological record, since a post mid-Pliocene uplift of relatively small amplitude could be more prominent in the present-day landscape than the dynamic subsidence of several hundred meters predicted from the early Aptian to the late Eocene. Determining spatial and temporal trends of dynamic topography in details along the east coast of North America requires further work.

Comparison with the dynamic topography predicted by previous models. To put our results for well COST-B2 in perspective, we compare them in Fig. 9 with the results of three previous studies of dynamic topography for the same area all using the same definition of mantle-flow-driven dynamic topography, excluding the dynamics of the thermal boundary layer (Moucha et al., 2008a; Müller et al., 2008b; Spasojevic et al., 2008). We use the results of Moucha et al. (2008a) for surface dynamic topography calculated using mantle density heterogeneities deeper than 200 km. These extend back to 30 Ma and were integrated over a ~ 450 km x 550 km area offshore New Jersey that includes the location of well COST-B2. The results of Spasojevic et al. (2008) extend back to 50 Ma and were sampled at a point location ~ 150 km west of well COST-B2, and the results of Müller et al. (2008b) extend back to 70 Ma and were sampled at the location of well COST-B2. Given the long wavelength nature of dynamic topography,

the results of each of these studies can be considered representative of the New Jersey margin and compared to our results. Two cases are presented for each model considered, including a second case for our model in which the lower mantle reference viscosity is 50 times larger than that of the upper mantle (this ratio is 100 in the reference case). Modeling results are also compared to the anomalous subsidence deduced from Fig. 6C. Note that this direct comparison implies that post-rift subsidence is only affected by cooling of the lithosphere and dynamic topography, which may be an oversimplification. The Neogene phase of anomalous subsidence (Fig. 9) is difficult to reconcile with the results of Moucha et al. (2008a) and with model “ngrand m.b.a” of Müller et al. (2008b) that predict uplift since 30 Ma and since 10 Ma, respectively. All other models predict subsidence throughout the Neogene, although the amplitude is too small except for case 2 of Spasojevic et al. (2008). The models of Moucha et al. (2008a) predict uplift between 30 Ma and 10 Ma, which is consistent with backstripping for constant sea level, whereas all other models predict subsidence prior to 15 Ma, which is consistent with backstripping using the sea level curve of Haq et al. (1987). In addition, the uplift predicted by the model of Moucha et al. (2008a) is consistent with the geologically-inferred uplift from Virginia to Georgia (Dowsett and Cronin, 1990; Rowley et al., 2011), although the inferred uplift rate is three to six times larger than the modeled uplift rate. As discussed above, sea level has fallen by at least 170 m over the last 85 Ma because of a combination of decreasing volume of ocean basins and ice sheet growth in the Cenozoic. As a consequence, we favor models that predict Cenozoic dynamic subsidence of the New Jersey margin. However, the large uncertainty associated with the dynamic models that predict between 160 m and 500 m of subsidence over the last 50 Ma (Fig. 9) make it

difficult to estimate long-term eustatic sea level from the high quality-record of the New Jersey margin by correcting for dynamic topography (Kominz et al., 2008; Müller et al., 2008b). Long-term eustatic sea level change may be estimated from the global stratigraphic record (Haq et al., 1987), a synthesis of the flooding record from continental platforms around the world (Bond, 1978), or by modeling the change of volume of ocean basins through time (Kominz, 1984; Müller et al., 2008b; Spasojevic and Gurnis, in press), and should be corrected for the global effect of dynamic topography (Conrad and Husson, 2009; Spasojevic and Gurnis, in press).

The contrasting evolution of dynamic topography predicted by different models for the New Jersey margin (Fig. 9) calls for explanation. The adjoint model of Spasojevic et al. (2008) and the forward model presented here are based on numerical approaches different to the backward advection models of Müller et al. (2008b) and of Moucha et al. (2008a). In addition, lateral viscosity variations that influence the evolution of thermal heterogeneity in the mantle (Moucha et al., 2007) are taken into account via the temperature dependence of viscosity in the model presented here and in that of Spasojevic et al. (2008), whereas they are ignored in the models of Moucha et al. (2008a) and of Müller et al. (2008b) which only have radial variations in viscosity. Furthermore, the present model and that of Spasojevic et al. (2008) consider a simple two-layer viscosity structure for the mantle (Paulson et al., 2007) whereas the models of Moucha et al. (2008a) and of Müller et al. (2008b) each consider about 20 viscosity layers throughout the mantle (Mitrovica and Forte, 2004; Steinberger and Calderwood, 2006). These different viscosity structures affect the depth-dependence of surface dynamic topography via the viscosity ratio between the different layers (Hager and Clayton, 1989;

Fig. 3; Eq. 2). For instance, decreasing the viscosity of the lower mantle by a factor of two in our model increases the predicted subsidence since 85 Ma by ~ 15 % (Fig. 9).

The models of Moucha et al. (2008a) and of Müller et al. (2008b) are overall similar in design, yet give different results for the New Jersey margin. The main differences between these two backward advection models include the mantle tomography employed, the conversion from seismic velocity to density, the viscosity structure of the mantle and the treatment of thermal boundary layers – essentially the four main steps of estimating dynamic topography (Braun, 2010). The scaling factor from seismic velocity to density clearly has an effect on the amplitude of the predicted absolute dynamic topography. A constant scaling is applied to the whole mantle by Müller et al. (2008b), but a radially and laterally variable scaling in the tomographic inversion of Simmons et al. (2007) is used by Moucha et al. (2008a). The effect of this spatially variable scaling factor on the spatio-temporal evolution of dynamic topography predicted by Moucha et al. (2008a) is likely to be significant. As for mantle tomography itself, Müller et al. (2008b) predicted Tertiary subsidence of the New Jersey margin using three different S-wave tomography models. The results of Müller et al. (2008b) also confirm that the treatment of thermal boundary layers is important in backward advection models (Conrad and Gurnis, 2003), as shown by the difference between their “pure” and “modified” models in Fig. 9. The “modified” model predicts 150 m less dynamic subsidence during the Tertiary than the “pure” model, presumably due to the removal of shallow downwellings in the “modified” model. The “modified” model also predicts uplift for the last 10 Myr, likely associated with the upward continuation of upwellings, at odds with the subsidence of the New Jersey margin. The lithosphere is thus key not only

to estimating present-day residual topography, as discussed previously, but also to modeling time-dependent dynamic topography. Finally, the main difference between the viscosity structures used by Moucha et al. (2008a) and Müller et al. (2008b) is the 30-km-thick viscosity notch just above the 670-km phase transition (Mitrovica and Forte, 2004) for which the viscosity is 300 times (5 times) lower than the reference viscosity in model V1 (model V2). This narrow low viscosity zone, potentially related to transformational superplasticity or to an internal thermal boundary layer (Pari and Peltier, 1995), should at least partially decouple the upper mantle from the lower mantle. Such a decoupling should act to suppress the effect of the subducting Farallon slab that is imaged close to the top of the lower mantle under the present-day east coast of North America (Liu et al., 2008; Müller et al., 2008b). Thus Cenozoic subsidence of the New Jersey margin, as opposed to the uplift predicted by Moucha et al. (2008a), appears mainly dependent on the absence or presence of a low-viscosity notch just above the lower mantle under North America.

Dynamic topography as a constraint on mantle flow and on the coupling between mantle and lithosphere

The previous section illustrates the need for detailed comparisons between the evolution of dynamic topography predicted by models and that deduced from the geological record, as well as the complexity and uncertainty of dynamic topography models. The comparison of model predictions to the geological record is made complex by the large wavelength nature of the dynamic topography signal and the relative scarcity of high-quality constraints. The New Jersey example discussed above illustrates the importance of the assumptions made when interpreting the geological record.

Nevertheless, such comparisons are valuable since they constitute constraints for the viscosity structure of the mantle, and for the response of the lithosphere to mantle flow.

From an observational point of view, increasing amounts of geophysical and geochemical data and their compilation in databases will over time improve the knowledge of the thermal and chemical structure of the crust and lithosphere, which are critical to constraining present-day dynamic topography. Global databases of stratigraphic data will also be important to re-appraise global sea level change and constrain the evolution of dynamic topography. From a modeling point of view, the comparison between the evolution of topography deduced from the geological record and the dynamic topography predicted by mantle flow models can be made more direct. The usual, indirect, approach consists of comparing the model topography after removing the isostatic component (thermal boundary layer) to the estimated dynamic component of the topography deduced from the geological record. Increasing computing power will soon make it possible to explicitly model the deformation of the continental lithosphere in global mantle flow models. This would permit the simultaneous estimation of isostatic and dynamic topographies, and would facilitate the comparison between modeling results and data. Explicitly modeling both the continental and oceanic lithospheres in global mantle flow models would also open up the opportunity to study the response of the viscoelastic lithosphere to mantle stresses. Because of the computational cost of the finite-element viscoelastic treatment of global mantle flow, viscoelastic studies of sea level have largely been limited to the post-glacial rebound problem (e.g. Paulson et al., 2007). Increasing computational power and new numerical method will make calculations possible that include the full range of rheological and structural variations

from hundreds of meters to global scales (Stadler et al., 2010). This would improve our understanding of topography at intermediate wavelengths (400-800 km) at which both small-scale convection and elastic flexure occur.

CONCLUSIONS

Considerable progress has been made over the last 30 years on the observational constraints on and the modeling of dynamic topography. In addition to constraining the nature of mantle convection and mantle rheology, studies of dynamic topography have led to a new interpretation of the stratigraphic record and of long-term sea level change. Current global models and observations of dynamic topography are in broad agreement at very large scale ($\sim 10,000$ km) but differ in some areas and at smaller scales ($< 5,000$ km). Increasing computing resources and modeling techniques provide the opportunity to develop higher-resolution models that will be more readily comparable to geological data. In parallel, acquiring a better understanding of the global structure of the lithosphere through data acquisition and compilation would improve the constraint on present-day global dynamic topography and therefore on mantle properties.

ACKNOWLEDGEMENTS

We thank C. Conrad, Y. Ricard, S. Spasojevic and B. Steinberger for sharing their results, M. Kominz for the updated COST-B2 well data, and L. Husson for discussions. This paper benefited from comments by two anonymous reviewers. NF was supported by Statoil, MG by the National Science Foundation (through EAR-0810303) and the Caltech

Tectonics Observatory (by the Gordon and Betty Moore Foundation) and RDM by Australian Research Council grant FL0992245. Computer simulations were carried out on the Sun Constellation VAYU Cluster of the Australian National Computational Infrastructure. Figs. 1, 4 and 7 were prepared using the Generic Mapping Tools (Wessel and Smith, 1998). The original *CitcomS* software was obtained from CIG, Computational Infrastructure for Geodynamics (<http://geodynamics.org>). This is Contribution Number 216 of the Tectonics Observatory, Caltech.

APPENDIX 1. TECTONIC SUBSIDENCE OF WELL COST-B2

Backstripping

The updated stratigraphy of well COST-B2 was backstripped using the software of White (1993) for constant sea level and for the eustatic curve proposed by Haq et al. (1987) that features sea level variations of up to 250 m since the Paleozoic. Both tectonic subsidence curves show ~ 2.5 km of tectonic subsidence during the main rifting phase between 210 Ma and ~ 175 Ma, followed by ~ 1.5 km of thermal subsidence. Considering the large error bars on paleo-water depth associated with the two anomalies in thermal subsidence at ~ 85 Ma and ~ 35 Ma, the thermal subsidence phase is overall smooth until ~ 16 Ma when there is a marked acceleration in subsidence.

Strain rate inversion

In addition to backstripping, we carry out a strain rate inversion analysis of the backstripped data following the method of (White, 1993). The best-fit subsidence curves obtained from the inversion are shown in Fig. 6A, and the strain rate history is shown in Fig. 6B for both sea level curves. In both cases, the strain rate history describes a main

1206 first rifting phase between 210 Ma and ~ 165 Ma followed by a relatively minor rifting
1207 phase between 155 Ma and 135 Ma. The inversion predicts an increase in strain rate
1208 from ~ 15 Myr, of amplitude larger than the apparent second rift phase at ~ 145 Ma
1209 (Fig. 6B). In addition, the inversion using the eustatic curve of Haq et al. (1987) predicts
1210 protracted strain between ~ 85 Ma and 15 Ma.

1211 **Forward modeling of tectonic subsidence**

1212 We carry out forward modeling of pure shear lithospheric stretching following the
1213 method of Jarvis and McKenzie (1980) in order to estimate the anomalous subsidence at
1214 well COST-B2. These forward models do not take into account flexure or depth-
1215 dependent extension. We assume a single rifting phase between 210 Ma and 175 Ma,
1216 lithosphere initially 160 km thick, continental crust initially 31 km thick, and varying
1217 stretching factors between 1.8 and 2.2. Although the solution is non-unique, these
1218 parameters give a satisfactory fit to the observed tectonic subsidence (Fig. 6A).

1219 **Estimation of anomalous subsidence**

1220 We estimate the amount of anomalous subsidence for well COST-B2 by subtracting the
1221 thermal subsidence predicted by the forward models from the observed data (Fig. 6C).
1222 For each sea level curve, we use the models that best fits the early stages of post-rift
1223 subsidence (130-110 Ma), namely a stretching factor of 1.9 for the curve of Haq et al.
1224 (1987) and of 2.0 for constant sea level (Fig. 6C). On Fig. 6C, positive (negative) values
1225 reflect anomalous subsidence (uplift) for which the model subsidence is shallower
1226 (deeper) than the backstripped data. Note the large error bars associated with paleo-water
1227 depth, particularly for the period between 85 Ma and 35 Ma for which the uncertainty is

1228 ± 230 m (Fig. 6). We limit our analysis to the relative changes in anomalous subsidence
 1229 that exceed this uncertainty. The case with eustatic sea level change displays 175 ± 70 m
 1230 of relative uplift between 125 Ma and 85 Ma, then 400 ± 45 m of relative subsidence
 1231 since 85 Ma, including 230 ± 110 m since 15 Ma. The case for constant sea level shows
 1232 115 ± 70 m of relative uplift between 125 Ma and 85 Ma and $130 \text{ m} \pm 65$ m of relative
 1233 subsidence since 10 Ma. Note that in this latter case the relative subsidence of 60 ± 45 m
 1234 since 85 Ma is of smaller amplitude than the Neogene phase of subsidence, which implies
 1235 anomalous uplift between 85 Ma and 10 Ma. The large differences in the evolution of
 1236 anomalous subsidence for the case with constant sea level and for the case accounting for
 1237 sea level change illustrate the importance of the sea level curve used when backstripping
 1238 stratigraphic data.

1239 TABLES

1240

TABLE 1. AMPLITUDE OF RESIDUAL AND DYNAMIC TOPOGRAPHY FIELDS

Model	Minimum (m)	Maximum (m)	RMS* (m)
<u>Residual topography (Fig. 1)</u>			
A. This study	-1638	3158	556
B. <i>Steinberger (2007)</i>	-1557	2105	474
C. <i>Panasyuk and Hager (2000)</i>	-1704	1119	439
D. <i>Kaban et al. (2003)</i>	-2053	2144	613
<u>Dynamic topography (Fig. 4)</u>			
A. This study	-2235	1653	760
B. <i>Steinberger (2007)</i>	-2778	3039	909
C. <i>Ricard et al. (1993)</i>	-1605	611	430
D. <i>Conrad and Husson (2009)</i>	-1550	1450	480
E. <i>Spasojevic and Gurnis (2012)</i>	-3652	883	884
*Root mean square amplitude			

1241

TABLE 2. STRENGTHS AND LIMITATIONS OF DYNAMIC TOPOGRAPHY MODELS

Strengths	Limitations
<u>Analytical and semi-analytical models</u>	
Computationally cheap	Radial viscosity variations only Thermal diffusivity neglected
<u>Numerical models</u>	
<i>Forward models</i>	
Computationally relatively cheap	Dependent on synthetic initial condition
Can achieve large dimensional time	Kinematically driven
Can achieve high resolution	May result in unrealistic slab advection
Can be compared to mantle tomography	Upwelling is usually passive
<i>Backward advection</i>	
Computationally relatively cheap	Thermal diffusivity is neglected
Can achieve high resolution	Limited to a few tens of million years
Consistent with the present-day density structure of the mantle	Thermal boundary layers require special treatment Usually kinematically driven
<i>Adjoint models</i>	
Consistent with the present-day density structure of the mantle	Computationally expensive
Thermal diffusivity is accounted for	Usually kinematically driven
<i>Hybrid models</i>	
Computationally relatively cheap	Not fully time dependent
Consistent with the present-day density structure of the mantle	Kinematically driven Thermal diffusivity neglected in the backward advection

1242

1243 **FIGURE CAPTIONS**

1244 Figure 1. Graphical comparison between four residual topography fields calculated
 1245 independently. All four models account for the flattening of old ocean floor (“plate
 1246 model”) with water-loaded oceans. Continents are air-loaded, and the mean residual

topography is set to zero for comparison with dynamic topography models. In the continents, A was calculated by removing the mean continental elevation (529 m), B was obtained by removing the isostatic contribution of the crust only, updated from (Steinberger, 2007) using the age grids of (Müller et al., 2008a), C and D were obtained by removing the isostatic contribution of both the continental crust and the continental lithosphere. The thin and thick black lines are the coastlines and plate boundaries, respectively. In A, the white contours are the continent-ocean boundary from Müller et al. (2008a), the magenta contours are the edge of the continents (-200 m; Harrison et al., 1983), the green contours outline Phanerozoic large igneous provinces and the yellow contours indicate continental crust thicker than 50 km. The red star is the location of well COST-B2 offshore New Jersey. Mollweide projection. Data in B, C and D courtesy B. Steinberger.

Figure 2. Depocenter migration within Cretaceous western interior basin of North America. A. Map and cross-section of deltaic depocenters in the central Rocky Mountain area. 1: pre-Campanian; 2: Campanian and Early Maastrichtian; 3: Maastrichtian. Modified from Figure 15 of Weimer (1970). B. Predicted depth of the Farallon slab and main tectonic elements (STB, dark shading: Sevier thrust belt; WIB, light shading: Western Interior Basin; FAR: Farallon plate). C-F: cumulative (black curves) and incremental (blue bands) residual subsidence from backstripped Late Cretaceous strata, compared to the cumulative dynamic subsidence (pink curves) predicted by the adjoint model of Liu et al. (2008), at different time intervals across section w-x in B. B-F modified from Figure 3 of Liu et al. (2011).

Figure 3. Normalized surface deformation kernel showing the relative deformation induced by a surface mass placed at a given depth in the mantle for spherical harmonic degrees 2 (solid line), 4 (long dashes) and 8 (short dashes). A value of unity implies perfect isostatic compensation. Values on the right-hand side axis indicate the relative, non-dimensional viscosity of each of the four layers (lithosphere, asthenosphere, upper mantle, lower mantle) considered in this case. The limit between upper and lower mantle is shown in gray. Modified from Figure 24 of Hager and Clayton (1989).

Figure 4. Graphical comparison of the present-day dynamic topography (water-loaded in the oceans, air-loaded in the continents, and averaging to zero) predicted by five distinct models. A and C show the dynamic topography fields predicted by time-dependent mantle flow models, both numerical (A) and analytical (C). B, D and E show the dynamic topography field predicted by present-day instantaneous mantle flow models in which the density structure is based on a seismic tomographic model of the interior of the Earth. See text for details. The red star in all panels shows the location of well COST-B2. Mollweide projection. Data in B. courtesy B. Steinberger, C. courtesy Y. Ricard, D. courtesy C. Conrad and E. courtesy S. Spasojevic.

Figure 5. Predictions of the adjoint models of Spasojevic et al. (2009). Flooding (A) and dynamic topography (B) of North America since 100 Ma in a North-America-fixed reference frame. C. Cross-sections at 41°N of the velocity and temperature fields. The curves above the section show plate velocities (red) and dynamic topography (blue) along the profile. Black triangles denote a well in the area of Denver that moves with the continent. Note that the bottom-right cross section is for 90 Ma while the dynamic

topography and flooding predictions are at 100 Ma. Modified from Figures 4 and 12 of Spasojevic et al. (2009).

Figure 6. Backstripped tectonic subsidence, evolution of strain rate and anomalous subsidence for well COST B-2 (location shown on Figs. 1, 4 and 7). A. Backstripped, water-loaded tectonic subsidence plotted as black (sea level curve of Haq et al., 1987) and gray (no sea level correction) circles with associated water depth error bars. The forward-modeled tectonic subsidence is shown in gray dashed curves for stretching factors between 1.8 and 2.2. The modeled, best-fit tectonic subsidence from strain rate inversion is shown as bold blue (sea level curve of Haq et al., 1987) and red (no sea level correction) curves, and the associated time-dependent strain rates are shown in B. C. Anomalous subsidence taken as the difference between the subsidence deduced from backstripping and the subsidence predicted by forward modeling. The reference model is for a stretching factor of 1.9 when sea level changes are taken into account (black curve, circles and error bars) and 2.0 when they are not (gray curve, circles and error bars). Negative values indicate anomalous uplift and positive values indicate anomalous subsidence.

Figure 7. Evolution of air-loaded dynamic topography predicted by our model, shown in 50 Myr intervals with reconstructed coastlines (thin black lines), and subduction zones (teethed thick gray lines). Stars show reconstructed locations of: well COST-B2 offshore New Jersey (large, red star) and a conjugate virtual well in Western Sahara (large, orange star), location 4 of Dowsett and Cronin (1990) in North Carolina (small, green star) and a conjugate virtual well in Senegal (small, cyan star).

Figure 8. Dynamic topography sampled through time at the reconstructed locations shown in Fig. 6. The dashed part of the curve is not shown in Fig. 6. The inset shows dynamic topography with respect to present for the two locations in New Jersey (NJ) and North Carolina (NC). Sen: Senegal; WS: Western Sahara.

Figure 9. Comparison between the dynamic topography predicted by four distinct time-dependent models (two cases are shown for each model) for the New Jersey margin and the anomalous uplift and subsidence at well COST-B2 since 85 Ma. The plain (dotted) gray field shows the anomalous subsidence and associated error relative to the present day based on backstripping with (without) eustatic sea level change.

REFERENCES CITED

- Afonso, J., Zlotnik, S., and Fernández, M., 2008, Effects of compositional and rheological stratifications on small-scale convection under the oceans: Implications for the thickness of oceanic lithosphere and seafloor flattening: *Geophysical Research Letters*, v. 35, p. L20308.
- Ault, A. K., Flowers, R. M., and Bowring, S. A., 2009, Phanerozoic burial and unroofing history of the western Slave craton and Wopmay orogen from apatite (U–Th)/He thermochronometry: *Earth and Planetary Science Letters*, v. 284, no. 1, p. 1-11.
- Blakey, R. C., 2008, Gondwana paleogeography from assembly to breakup—A 500 my odyssey: Resolving the Late Paleozoic Ice Age in Time and Space: *Geological Society of America Special Paper*, v. 441, p. 1-28.
- Bond, G., 1976, Evidence for continental subsidence in North America during the late Cretaceous global submergence: *Geology*, v. 4, p. 557-560.
- , 1978, Speculations on real sea-level changes and vertical motions of continents at selected times in the Cretaceous and Tertiary periods: *Geology*, v. 6, p. 247-250.
- Bower, D. J., Gurnis, M., and Flament, N., in preparation, Assimilating lithosphere and slab history in 4-D dynamic Earth models: *Physics of the Earth and Planetary Interiors*.
- Boyd, O. S., Jones, C. H., and Sheehan, A. F., 2004, Foundering lithosphere imaged beneath the Southern Sierra Nevada, California, USA: *Science*, v. 305, p. 660–662.
- Boyden, J., Müller, R., Gurnis, M., Torsvik, T., Clark, J. A., Turner, M., Ivey-Law, H., Watson, R., and Cannon, J., 2011, Next-generation plate-tectonic

1347 reconstructions using GPlates: Geoinformatics: Cyberinfrastructure for the
 1348 Solid Earth Sciences, p. 95-113.
 1349 Braun, J., 2010, The many surface expressions of mantle dynamics: Nature
 1350 Geoscience, v. 3, no. 12, p. 825-833.
 1351 Brun, M. V. L., and Lucazeau, F., 1988, Subsidence, extension and thermal history of
 1352 the West African margin in Senegal: Earth and Planetary Science Letters, v.
 1353 90, no. 2, p. 204-220.
 1354 Bunge, H. P., Hagelberg, C., and Travis, B., 2003, Mantle circulation models with
 1355 variational data assimilation: inferring past mantle flow and structure from
 1356 plate motion histories and seismic tomography: Geophysical Journal
 1357 International, v. 152, no. 2, p. 280-301.
 1358 Burgess, P. M., and Gurnis, M., 1995, Mechanisms for the formation of cratonic
 1359 stratigraphic sequences: Earth and Planetary Science Letters, v. 136, p. 647-
 1360 663.
 1361 Burgess, P. M., Gurnis, M., and Moresi, L., 1997, Formation of sequences in the
 1362 cratonic interior of North America by interaction between mantle, eustatic,
 1363 and stratigraphic processes: Geological Society of America Bulletin, v. 109,
 1364 no. 12, p. 1515-1535.
 1365 Cathles, L. M., 1975, The Viscosity of the Earth's Mantle, Princeton, New Jersey,
 1366 Princeton University Press, 386 p.:
 1367 Cazenave, A., Souriau, A., and Dominh, K., 1989, Global coupling of Earth surface
 1368 topography with hotspots, geoid and mantle heterogeneities: Nature, v. 340,
 1369 no. 6228, p. 54-57.
 1370 Clark, M. K., Maheo, G., Saleeby, J., and Farley, K. A., 2005, The nonequilibrium
 1371 landscape of the southern Sierra Nevada, California: GSA Today, v. 15, no. 9,
 1372 p. 4-10.
 1373 Coltice, N., Phillips, B. R., Bertrand, H., Ricard, Y., and Rey, P., 2007, Global warming
 1374 of the mantle at the origin of flood basalts over supercontinents: Geology, v.
 1375 35, no. 5, p. 391-394.
 1376 Conrad, C. P., and Gurnis, M., 2003, Seismic tomography, surface uplift, and the
 1377 breakup of Gondwanaland: Integrating mantle convection backwards in time:
 1378 Geochemistry Geophysics Geosystems, v. 4, no. 3, p. 1031.
 1379 Conrad, C. P., and Husson, L., 2009, Influence of dynamic topography on sea level
 1380 and its rate of change: Lithosphere, v. 1, no. 2, p. 110-120.
 1381 Conrad, C. P., Lithgow-Bertelloni, C., and Loudon, K. E., 2004, Iceland, the Farallon
 1382 slab, and dynamic topography of the North Atlantic: Geology, v. 32, no. 3, p.
 1383 177-180.
 1384 Cook, T. D., and Bally, A. W., 1975, Stratigraphic Atlas of North and Central America,
 1385 Princeton University Press, 272 p.:
 1386 Cosentino, D., Schildgen, T. F., Cipollari, P., Faranda, C., Gliozzi, E., Hudáčeková, N.,
 1387 Lucifora, S., and Strecker, M. R., 2012, Late Miocene surface uplift of the
 1388 southern margin of the Central Anatolian Plateau, Central Taurides, Turkey:
 1389 GSA Bulletin, v. 124, no. 1-2, p. 133-145.
 1390 Crosby, A. G., and McKenzie, D., 2009, An analysis of young ocean depth, gravity and
 1391 global residual topography: Geophysical Journal International, v. 178, no. 3, p.
 1392 1198-1219.

1393 Crough, S. T., 1983, Hotspot swells: Annual Reviews of Earth and Planetary Science,
1394 v. 11, p. 165-193.

1395 Davies, G. F., and Pribac, F., 1993, Mesozoic seafloor subsidence and the Darwin Rise,
1396 past and present, The Mesozoic Pacific: Geology, Tectonics, and Volcanism,
1397 Volume 77, American Geophysical Union, p. 39-52.

1398 Davis, E. E., and Lister, C. R. B., 1974, Fundamentals of ridge crest topography: Earth
1399 and Planetary Science Letters, v. 21, no. 4, p. 405-413.

1400 DeMets, C., Gordon, R., Argus, D., and Stein, S., 1990, Current plate motions:
1401 Geophysical Journal International, v. 101, no. 2, p. 425-478.

1402 DiCaprio, L., Gurnis, M., and Müller, R. D., 2009, Long-wavelength tilting of the
1403 Australian continent since the Late Cretaceous: Earth and Planetary Science
1404 Letters, v. 278, no. 3-4, p. 175-185.

1405 DiCaprio, L., Müller, R. D., and Gurnis, M., 2010, A dynamic process for drowning
1406 carbonate reefs on the northeastern Australian margin: Geology, v. 38, no. 1,
1407 p. 11-14.

1408 Divins, D. L., 2003, Total Sediment Thickness of the World's Oceans & Marginal Seas:
1409 Boulder, CO, NOAA National Geophysical Data Center.

1410 Dowsett, H. J., and Cronin, T. M., 1990, High eustatic sea level during the middle
1411 Pliocene: Evidence from the southeastern US Atlantic Coastal Plain: Geology,
1412 v. 18, no. 5, p. 435-438.

1413 Flowers, R. M., 2010, The enigmatic rise of the Colorado Plateau: Geology, v. 38, no.
1414 7, p. 671-672.

1415 Flowers, R. M., and Schoene, B., 2010, (U-Th)/He thermochronometry constraints on
1416 unroofing of the eastern Kaapvaal craton and significance for uplift of the
1417 southern African Plateau: Geology, v. 38, no. 9, p. 827-830.

1418 Flowers, R. M., Wernicke, B. P., and Farley, K. A., 2008, Unroofing, incision and uplift
1419 history of the southwestern Colorado Plateau from apatite (U-Th)/He
1420 thermochronometry: GSA Bulletin, v. 120, p. 571-587.

1421 Forte, A. M., Peltier, W. R., Dziewonski, A. M., and Woodward, R. L., 1993, Dynamic
1422 surface topography: A new interpretation based upon mantle flow models
1423 derived from seismic tomography: Geophysical Research Letters, v. 20, no. 3,
1424 p. 225-228.

1425 Gallagher, K., Dumitru, T. A., and Gleadow, A. J. W., 1994, Constraints on the vertical
1426 motion of eastern Australia during the Mesozoic: Basin Research, v. 6, p. 77-
1427 94.

1428 Gallagher, K., and Lambeck, K., 1989, Subsidence, sedimentation and sea-level
1429 changes in the Eromanga Basin, Australia: Basin Research, v. 2, p. 115-131.

1430 Grand, S. P., 2002, Mantle shear-wave tomography and the fate of subducted slabs:
1431 Philosophical Transactions of the Royal Society of London. Series A:
1432 Mathematical, Physical and Engineering Sciences, v. 360, no. 1800, p. 2475-
1433 2491.

1434 Green, P. F., Japsen, P., Chalmers, J. A., and Bonow, J. M., 2011, Thermochronology,
1435 erosion surfaces and missing section in West Greenland: Journal of the
1436 Geological Society, v. 168, no. 4, p. 817-830.

1437 Guillaume, B., Martinod, J., Husson, L., Roddaz, M., and Riquelme, R., 2009, Neogene
 1438 uplift of central eastern Patagonia: Dynamic response to active spreading
 1439 ridge subduction?: *Tectonics*, v. 28, no. 2, p. TC2009.
 1440 Gurnis, M., 1988, Large-scale mantle convection and the aggregation and dispersal of
 1441 supercontinents: *Nature*, v. 332, no. 6166, p. 695-699.
 1442 Gurnis, M., 1990, Bounds on global dynamic topography from Phanerozoic flooding
 1443 of continental platforms: *Nature*, v. 344, no. 6268, p. 754-756.
 1444 Gurnis, M., 1992, Long-term controls on eustatic and epeirogenic motions by mantle
 1445 convection: *GSA Today*, v. 2, p. 141-157.
 1446 Gurnis, M., 1993a, Comment on "Dynamic surface topography: A new interpretation
 1447 based upon mantle flow models derived from seismic tomography"; by A. M.
 1448 Forte, W. R. Peltier, A. M. Dziewonski and R. L. Woodward: *Geophysical*
 1449 *Research Letters*, v. 20, no. 15, p. 1663-1664.
 1450 Gurnis, M., 1993b, Depressed continental hypsometry behind oceanic trenches: A
 1451 clue to subduction controls on sea level change: *Geology*, v. 21, p. 29-32.
 1452 -, 1993c, Phanerozoic marine inundation of continents driven by dynamic
 1453 topography above subducting slabs: *Nature*, v. 364, p. 589-593.
 1454 Gurnis, M., Mitrovica, J. X., Ritsema, J., and van Heijst, H.-J., 2000, Constraining mantle
 1455 density structure using geological evidence of surface uplift rates: The case of
 1456 the African Superplume: *Geochemistry Geophysics Geosystems*, v. 1, no. 7, p.
 1457 1-31.
 1458 Gurnis, M., Müller, R. D., and Moresi, L., 1998, Cretaceous vertical motion of Australia
 1459 and the Australian Antarctic Discordance: *Science*, v. 279, no. 5356, p. 1499-
 1460 1504.
 1461 Gurnis, M., Turner, M., Zahirovic, S., DiCaprio, L., Spasojevic, S., Müller, R. D., Boyden,
 1462 J., Seton, M., Manea, V. C., and Bower, D. J., 2012, Plate tectonic
 1463 reconstructions with continuously closing plates: *Computers & Geosciences*,
 1464 v. 38, no. 1, p. 35-42.
 1465 Hager, B. H., 1984, Subducted slabs and the geoid: constraints on mantle rheology
 1466 and flow: *Journal of Geophysical Research - Solid Earth*, v. 89, no. B7, p. 6003-
 1467 6015.
 1468 Hager, B. H., and Clayton, R. W., 1989, Constraints on the structure of mantle
 1469 convection using seismic observations, flow models, and the geoid, *in* Peltier,
 1470 W. R., ed., *Mantle Convection, Plate Tectonics and Global Dynamics*: New
 1471 York, Gordon and Breach Science Publishers, p. 657-763.
 1472 Hager, B. H., Clayton, R. W., Richards, M. A., Comer, R. P., and Dziewonski, A. M., 1985,
 1473 Lower mantle heterogeneity, dynamic topography and the geoid: *Nature*, v.
 1474 313, no. 6003, p. 541-545.
 1475 Hager, B. H., and O'Connell, R. J., 1979, Kinematic models of large-scale flow in the
 1476 Earth's mantle: *Journal of Geophysical Research*, v. 84, no. B3, p. 1031-1048.
 1477 Hales, T. C., Abt, D. L., Humphreys, E. D., and Roering, J. J., 2005, A lithospheric
 1478 instability origin for Columbia River flood basalts and Willowa Mountains
 1479 uplift in northeast Oregon: *Nature*, v. 438, p. 842-845.
 1480 Hallam, A., 1992, *Phanerozoic sea level changes*, New York, NY, Columbia University
 1481 Press, 266 p.:

1482 Haq, B. U., Hardenbol, J., and Vail, P. R., 1987, Chronology of fluctuating sea levels
 1483 since the Triassic: *Science*, v. 235, no. 4793, p. 1156-1167.
 1484 Harrison, C. G. A., Miskell, K. J., Brass, G. W., Saltzman, E. S., and Sloan, J. L., II, 1983,
 1485 Continental hypsography: *Tectonics*, v. 2, no. 4, p. 357-377.
 1486 Hartley, R. A., Roberts, G. G., White, N., and Richardson, C., 2011, Transient
 1487 convective uplift of an ancient buried landscape: *Nature Geoscience*, v. 4, p.
 1488 562-565.
 1489 Hays, J. D., and Pitman, W. C., 1973, Lithospheric Plate Motion, Sea Level Changes
 1490 and Climatic and Ecological Consequences: *Nature*, v. 246, no. 5427, p. 18-22.
 1491 Heller, P. L., Dueker, K., and McMillan, M. E., 2003, Post-Paleozoic alluvial gravel
 1492 transport as evidence of continental tilting in the U.S. Cordillera: *Geological*
 1493 *Society of America Bulletin*, v. 115, p. 1122-1132.
 1494 Hillier, J., 2010, Subsidence of "normal" seafloor: Observations do indicate
 1495 "flattening": *Journal of Geophysical Research*, v. 115, no. B3, p. B03102.
 1496 Huang, J., Zhong, S., and van Hunen, J., 2003, Controls on sub-lithospheric small-scale
 1497 convection: *J. Geophys. Res.*, v. 108, no. B8.
 1498 Husson, L., and Conrad, C. P., 2006, Tectonic velocities, dynamic topography, and
 1499 relative sea level: *Geophysical Research Letters*, v. 33, no. 18, p. L18303.
 1500 Husson, L., Guillaume, B., Funicello, F., Faccenna, C., and Royden, L. H., 2012,
 1501 Unraveling topography around subduction zones from laboratory models:
 1502 *Tectonophysics*, v. 526-529, p. 5-15.
 1503 Ismail-Zadeh, A., Schubert, G., Tsepelev, I., and Korotkii, A., 2004, Inverse problem of
 1504 thermal convection: numerical approach and application to mantle plume
 1505 restoration: *Physics of the Earth and Planetary Interiors*, v. 145, no. 1-4, p.
 1506 99-114.
 1507 Jarvis, G. T., and McKenzie, D. P., 1980, Sedimentary basin formation with finite
 1508 extension rates: *Earth and Planetary Science Letters*, v. 48, no. 1, p. 42-52.
 1509 Jarvis, G. T., and Peltier, W., 1982, Mantle convection as a boundary layer
 1510 phenomenon: *Geophysical Journal Of the Royal Astronomical Society*, v. 68,
 1511 no. 2, p. 389-427.
 1512 Jordan, T. H., 1975, The continental tectosphere: *Reviews of Geophysics and Space*
 1513 *Physics*, v. 13, no. 3, p. 1-12.
 1514 Kaban, M. K., Schwintzer, P., Artemieva, I. M., and Mooney, W. D., 2003, Density of the
 1515 continental roots: compositional and thermal contributions: *Earth and*
 1516 *Planetary Science Letters*, v. 209, no. 1-2, p. 53-69.
 1517 Kaula, W. M., 1972, Global gravity and tectonics, *in* Robertson, E. C., ed., *The Nature*
 1518 *of the Solid Earth*: New York, N.Y., McGraw-Hill, p. 385-405.
 1519 Kido, M., and Seno, T., 1994, Dynamic topography compared with residual depth
 1520 anomalies in oceans and implications for age-depth curves: *Geophysical*
 1521 *Research Letters*, v. 21, no. 8, p. 717-720.
 1522 Kominz, M., Browning, J., Miller, K., Sugarman, P., Mizintseva, S., and Scotese, C.,
 1523 2008, Late Cretaceous to Miocene sea-level estimates from the New Jersey
 1524 and Delaware coastal plain coreholes: An error analysis: *Basin Research*, v.
 1525 20, no. 2, p. 211-226.

1526 Kominz, M. A., 1984, Oceanic ridge volumes and sea-level change: An error analysis,
 1527 *in* Schlee, J., ed., Interregional unconformities and hydrocarbon accumulation,
 1528 AAPG Memoir, p. 109-127.
 1529 Korenaga, T., and Korenaga, J., 2008, Subsidence of normal oceanic lithosphere,
 1530 apparent thermal expansivity, and seafloor flattening: Earth and Planetary
 1531 Science Letters, v. 268, no. 1, p. 41-51.
 1532 Langseth, M. G., Jr., Le Pichon, X., and Ewing, M., 1966, Crustal structure of the mid-
 1533 ocean ridges: 5. Heat flow through the Atlantic Ocean floor and convection
 1534 currents: Journal of Geophysical Research, v. 71, no. 22, p. 5321-5355.
 1535 Le Stunff, Y., and Ricard, Y., 1995, Topography and geoid due to lithospheric mass
 1536 anomalies: Geophysical Journal International, v. 122, no. 3, p. 982-990.
 1537 Lithgow-Bertelloni, C., and Gurnis, M., 1997, Cenozoic subsidence and uplift of
 1538 continents from time-varying dynamic topography: Geology, v. 25, no. 8, p.
 1539 735-738.
 1540 Lithgow-Bertelloni, C., and Richards, M. A., 1998, The dynamics of Cenozoic and
 1541 Mesozoic plate motions: Reviews of Geophysics, v. 36, no. 1, p. 27-78.
 1542 Lithgow-Bertelloni, C., Richards, M. A., Ricard, Y., O'Connell, R. J., and Engebretson, D.
 1543 C., 1993, Toroidal-poloidal partitioning of plate motions since 120 Ma:
 1544 Geophysical Research Letters, v. 20, no. 5, p. 375-378.
 1545 Lithgow-Bertelloni, C., and Silver, P. G., 1998, Dynamic topography, plate driving
 1546 forces and the African superswell: Nature, v. 395, no. 6699, p. 269-272.
 1547 Liu, L., and Gurnis, M., 2008, Simultaneous inversion of mantle viscosity and initial
 1548 conditions from an adjoint of mantle convection: Journal of Geophysical
 1549 Research, v. 113.
 1550 -, 2010, Dynamic subsidence and uplift of the Colorado Plateau: Geology, v. 38, p.
 1551 663-666.
 1552 Liu, L., Spasojević, S., and Gurnis, M., 2008, Reconstructing Farallon Plate Subduction
 1553 Beneath North America Back to the Late Cretaceous: Science, v. 322, no.
 1554 5903, p. 934-938.
 1555 Liu, S., and Nummedal, D., 2004, Late Cretaceous subsidence in Wyoming:
 1556 Quantifying the dynamic component: Geology, v. 32, p. 397-400.
 1557 Liu, S., Nummedal, D., and Gurnis, M., 2012, Depocenter shifts and unconformities
 1558 within Late Cretaceous sediments across central Utah and Colorado, USA
 1559 reveals the interplay between mantle dynamics, flexural subsidence and
 1560 eustatic sea levels: Lithosphere - to be submitted.
 1561 Liu, S., Nummedal, D., and Liu, L., 2011, Migration of dynamic subsidence across the
 1562 Late Cretaceous United States Western Interior Basin in response to Farallon
 1563 plate subduction: Geology, v. 39, no. 6, p. 555-558.
 1564 Lovell, B., 2010, A pulse in the planet: regional control of high-frequency changes in
 1565 relative sea level by mantle convection: Journal of the Geological Society, v.
 1566 167, no. 4, p. 637-648.
 1567 Masters, G., Laske, G., Bolton, H., and Dziewonski, A., 2000, The relative behaviour of
 1568 shear velocity, bulk sound speed, and compressional velocity in the mantle:
 1569 Implications for chemical and thermal structure, *in* Karato, S., Forte, A. M.,
 1570 Liebermann, R. C., Masters, G., and Stixrude, L., eds., Earth's Deep Interior:

1571 Mineral Physics and Tomography From the Atomic to the Global Scale:
1572 Washington, D.C., Am. Geophys. Un., p. 63-87.

1573 McKenzie, D. P., 1967, Some Remarks on Heat Flow and Gravity Anomalies: Journal
1574 of Geophysical Research, v. 72, no. 24, p. 6261-6273.

1575 McKenzie, D. P., Roberts, J. M., and Weiss, N. O., 1974, Convection in the Earth's
1576 mantle: towards a numerical simulation: Journal of Fluid Mechanics, v. 62, p.
1577 465-538.

1578 Miller, K. G., Kominz, M. A., Browning, J. V., Wright, J. D., Mountain, G. S., Katz, M. E.,
1579 Sugarman, P. J., Cramer, B. S., Christie-Blick, N., and Pekar, S. F., 2005, The
1580 Phanerozoic record of global sea-level change: Science, v. 310, no. 5752, p.
1581 1293-1298.

1582 Mitrovica, J., and Forte, A., 2004, A new inference of mantle viscosity based upon
1583 joint inversion of convection and glacial isostatic adjustment data: Earth and
1584 Planetary Science Letters, v. 225, no. 1, p. 177-189.

1585 Mitrovica, J. X., Beaumont, C., and Jarvis, G. T., 1989, Tilting of continental interiors
1586 by the dynamical effects of subduction: Tectonics, v. 8, no. 5, p. 1079-1094.

1587 Mitrovica, J. X., and Jarvis, G. T., 1985, Surface deflections due to transient
1588 subduction in a convecting mantle: Tectonophysics, v. 120, p. 211-237.

1589 Mooney, W. D., Laske, G., and Masters, T. G., 1998, CRUST 5.1: A global crustal model
1590 at 5× 5: Journal of Geophysical Research, v. 103, no. B1, p. 727-747.

1591 Morgan, W. J., 1965, Gravity anomalies and convection currents: 1. A sphere and
1592 cylinder sinking beneath the surface of a viscous fluid: Journal of Geophysical
1593 Research, v. 70, p. 6175-6187.

1594 Moucha, R., Forte, A., Mitrovica, J., and Daradich, A., 2007, Lateral variations in
1595 mantle rheology: implications for convection related surface observables and
1596 inferred viscosity models: Geophysical Journal International, v. 169, no. 1, p.
1597 113-135.

1598 Moucha, R., Forte, A. M., Mitrovica, J. X., Rowley, D. B., Quéré, S., Simmons, N. A., and
1599 Grand, S. P., 2008a, Dynamic topography and long-term sea-level variations:
1600 There is no such thing as a stable continental platform: Earth and Planetary
1601 Science Letters, v. 271, no. 1, p. 101-108.

1602 Moucha, R., Forte, A. M., Rowley, D. B., Mitrovica, J. X., Simmons, N. A., and Grand, S.
1603 P., 2008b, Mantle convection and the recent evolution of the Colorado
1604 Plateau and the Rio Grande Rift valley: Geology, v. 36, p. 439-442.

1605 -, 2009, Deep mantle forces and the uplift of the Colorado Plateau: Geophys. Res.
1606 Lett, v. 36, no. (19), p. 1-6.

1607 Mueller, K., Kier, G., Rockwell, T., and Jones, C. H., 2009, Quaternary rift flank uplift of
1608 the Peninsular Ranges in Baja and southern California by removal of mantle
1609 lithosphere: Tectonics, v. 28, p. TC5003.

1610 Müller, R. D., Lim, V. S. L., and Isern, A. R., 2000, Late Tertiary tectonic subsidence on
1611 the northeast Australian passive margin: response to dynamic topography?:
1612 Marine Geology, v. 162, no. 2-4, p. 337-352.

1613 Müller, R. D., Roest, W. R., Royer, J. Y., Gahagan, L. M., and Sclater, J. G., 1997, Digital
1614 isochrons of the world's ocean floor: Journal of Geophysical Research, v. 102,
1615 no. B2, p. 3211-3214.

1616 Müller, R. D., Sdrolias, M., Gaina, C., and Roest, W. R., 2008a, Age, spreading rates,
 1617 and spreading asymmetry of the world's ocean crust: *Geochemistry*
 1618 *Geophysics Geosystems*, v. 9, no. 4, p. Q04006.
 1619 Müller, R. D., Sdrolias, M., Gaina, C., Steinberger, B., and Heine, C., 2008b, Long-Term
 1620 Sea-Level Fluctuations Driven by Ocean Basin Dynamics: *Science*, v. 319, no.
 1621 5868, p. 1357-1362.
 1622 Panasyuk, S. V., and Hager, B. H., 2000, Models of isostatic and dynamic topography,
 1623 geoid anomalies, and their uncertainties: *Journal of Geophysical Research*, v.
 1624 105, no. B12, p. 28199-28209.
 1625 Pari, G., and Peltier, W. R., 1995, The heat flow constraint on mantle tomography-
 1626 based convection models: Towards a geodynamically self-consistent
 1627 inference of mantle viscosity: *Journal of Geophysical Research*, v. 100, no. B7,
 1628 p. 12731-12712,12751.
 1629 Parsons, B., and Daly, S., 1983, The Relationship Between Surface Topography,
 1630 Gravity Anomalies, and Temperature Structure of Convection: *Journal of*
 1631 *Geophysical Research*, v. 88, no. B2, p. 1129-1144.
 1632 Parsons, B., and Sclater, J. G., 1977, An analysis of the variation of ocean floor
 1633 bathymetry and heat flow with age: *Journal of Geophysical Research*, v. 82,
 1634 no. 5, p. 803-827.
 1635 Paulson, A., Zhong, S., and Wahr, J., 2007, Inference of mantle viscosity from GRACE
 1636 and relative sea level data: *Geophysical Journal International*, v. 171, no. 2, p.
 1637 497-508.
 1638 Pekeris, C. L., 1935, Thermal convection in the interior of the Earth: *Geophysical*
 1639 *Journal International*, v. 3, p. 343-367.
 1640 Petersen, K. D., Nielsen, S. B., Clausen, O. R., Stephenson, R., and Gerya, T., 2010,
 1641 Small-Scale Mantle Convection Produces Stratigraphic Sequences in
 1642 Sedimentary Basins: *Science*, v. 329, no. 5993, p. 827-830.
 1643 Pritchard, D., Roberts, G., White, N., and Richardson, C., 2009, Uplift histories from
 1644 river profiles: *Geophysical Research Letters*, v. 36, no. 24, p. L24301.
 1645 Redfield, T. F., 2010, On apatite fission track dating and the Tertiary evolution of
 1646 West Greenland topography: *Journal of the Geological Society*, v. 167, no. 2, p.
 1647 261-271.
 1648 Ricard, Y., Fleitout, L., and Froidevaux, C., 1984, Geoid heights and lithospheric
 1649 stresses for a dynamic Earth: *Annales Geophysicae*, v. 2, p. 267-285.
 1650 Ricard, Y., Richards, M., Lithgow-Bertelloni, C., and Le Stunff, Y., 1993, A Geodynamic
 1651 Model of Mantle Density Heterogeneity: *Journal of Geophysical Research*, v.
 1652 98, no. B12, p. 21895-21821,21909.
 1653 Richards, M. A., and Hager, B. H., 1984, Geoid Anomalies in a Dynamic Earth: *Journal*
 1654 *of Geophysical Research*, v. 89, no. B7, p. 5987-6002.
 1655 Ritsema, J., van Heijst, H. J., and Woodhouse, J. H., 2004, Global transition zone
 1656 tomography: *Journal of Geophysical Research*, v. 109, no. B2, p. B02302.
 1657 Romanowicz, B., 2008, Using seismic waves to image Earth's internal structure:
 1658 *Nature*, v. 447, p. 198-201.
 1659 Rowley, D. B., Forte, A. M., Moucha, R., Mitrovica, J. X., Simmons, N. A., and Grand, S.
 1660 P., 2011, Mantle dynamic impact on passive margin evolution: Implications

1661 for their architecture and derived sea level histories, *Fragile Earth*: Munich,
 1662 Germany.
 1663 Rudge, J. F., Shaw-Champion, M. E., White, N. J., McKenzie, D. P., and Lovell, B., 2008,
 1664 A Plume Model of Transient Diachronous Uplift at the Earth's Surface: *Earth*
 1665 and Planetary Science Letters, v. 267, no. 1-2, p. 146-160.
 1666 Russell, M., and Gurnis, M., 1994, The planform of epeirogeny: Vertical motions of
 1667 Australia during the Cretaceous: *Basin Research*, v. 6, p. 63-76.
 1668 Saleeby, J., Le Pourhiet, L., Saleeby, Z., and Gurnis, M., 2012, Epeirogenic transients
 1669 related to mantle lithosphere removal in the southern Sierra Nevada region,
 1670 California, Part I: Implications of thermo-mechanical modeling: *Geosphere*.
 1671 Sandiford, M., 2007, The tilting continent: A new constraint on the dynamic
 1672 topographic field from Australia: *Earth and Planetary Science Letters*, v. 261,
 1673 no. 1-2, p. 152-163.
 1674 Scholle, P. A., 1977, Geological studies on the COST No. B-2 well, US mid-Atlantic
 1675 outer continental shelf area, US Department of the Interior, Geological
 1676 Survey.
 1677 Schroeder, W., 1984, The empirical age-depth relation and depth anomalies in the
 1678 Pacific Ocean basin: *Journal of Geophysical Research*, v. 89, p. 9873-9884.
 1679 Scotese, C. R., Golonka, J., and Project, P., 1993, Paleogeographic atlas, PALEOMAP
 1680 project, Department of Geology, University of Texas at Arlington.
 1681 Seton, M., Müller, R., Zahirovic, S., Gaina, C., Torsvik, T., Shephard, G., Talsma, A.,
 1682 Gurnis, M., Turner, M., and Chandler, M., 2012, Global continental and ocean
 1683 basin reconstructions since 200 Ma: *Earth-Science Reviews*, v. 113, p. 212-
 1684 270.
 1685 Shaw-Champion, M. E., White, N. J., Jones, S. M., and Lovell, J. P. B., 2008, Quantifying
 1686 transient mantle plume uplift in the Faroe-Shetland basin: *Tectonics*, v. 27.
 1687 Shephard, G. E., Müller, R. D., Liu, L., and Gurnis, M., 2010, Miocene drainage reversal
 1688 of the Amazon River driven by plate-mantle interaction: *Nature Geoscience*,
 1689 v. 3, no. 12, p. 870-875.
 1690 Simmons, N. A., Forte, A. M., and Grand, S. P., 2007, Thermochemical structure and
 1691 dynamics of the African superplume: *Geophysical Research Letters*, v. 34, no.
 1692 2, p. L02301.
 1693 Smith, A. G., Smith, D. G., and Funnell, B. M., 1994, *Atlas of Mesozoic and Cenozoic*
 1694 *Coastlines*, Cambridge, Cambridge University Press.
 1695 Smith, W. H. F., and Sandwell, D. T., 1997, Global sea floor topography from satellite
 1696 altimetry and ship depth soundings: *Science*, v. 277, no. 5334, p. 1956-1962.
 1697 Spasojevic, S., and Gurnis, M., in press, Sea level and vertical motion of continents
 1698 from dynamic earth models since the Late Cretaceous: *AAPG Bulletin*.
 1699 Spasojevic, S., Gurnis, M., and Sutherland, R., 2010, Mantle upwellings above slab
 1700 graveyards linked to the global geoid lows: *Nature Geoscience*, v. 3, no. 6, p.
 1701 435-438.
 1702 Spasojevic, S., Liu, L., and Gurnis, M., 2009, Adjoint models of mantle convection with
 1703 seismic, plate motion, and stratigraphic constraints: North America since the
 1704 Late Cretaceous: *Geochemistry Geophysics Geosystems*, v. 10, no. 5, p.
 1705 Q05W02.

1706 Spasojevic, S., Liu, L., Gurnis, M., and Müller, R. D., 2008, The case for dynamic
 1707 subsidence of the U.S. east coast since the Eocene: *Geophysical Research*
 1708 *Letters*, v. 35, no. 8, p. L08305.
 1709 Stadler, G., Gurnis, M., Burstedde, C., Wilcox, L. C., Alisc, L., and Ghattas, O., 2010, The
 1710 dynamics of plate tectonics and mantle flow: From local to global scales:
 1711 *Science*, v. 329, p. 1033-1038.
 1712 Steinberger, B., 2007, Effects of latent heat release at phase boundaries on flow in
 1713 the Earth's mantle, phase boundary topography and dynamic topography at
 1714 the Earth's surface: *Physics of the Earth and Planetary Interiors*, v. 164, no.
 1715 1-2, p. 2-20.
 1716 Steinberger, B., and Calderwood, A. R., 2006, Models of large-scale viscous flow in
 1717 the Earth's mantle with constraints from mineral physics and surface
 1718 observations: *Geophysical Journal International*, v. 167, no. 3, p. 1461-1481.
 1719 Steinberger, B., and O'Connell, R. J., 1997, Changes of the Earth's rotation axis owing
 1720 to advection of mantle density heterogeneities: *Nature*, v. 387, no. 6629, p.
 1721 169-173.
 1722 Struckmeyer, H. I. M., and Brown, P. J., 1990, Australian Sea Level Curves Part 1:
 1723 Australian Inundation Curves: Bureau of Mineral Resources, 1990/11.
 1724 Sutherland, R., Spasojevic, S., and Gurnis, M., 2010, Mantle upwelling after
 1725 Gondwana subduction death explains anomalous topography and subsidence
 1726 histories of eastern New Zealand and West Antarctica: *Geology*, v. 38, p. 155-
 1727 158.
 1728 Swezey, C. S., 2009, Cenozoic stratigraphy of the Sahara, Northern Africa: *Journal of*
 1729 *African Earth Sciences*, v. 53, no. 3, p. 89-121.
 1730 Sykes, T. J. S., 1996, A correction for sediment load upon the ocean floor: Uniform
 1731 versus varying sediment density estimations—implications for isostatic
 1732 correction: *Marine Geology*, v. 133, no. 1-2, p. 35-49.
 1733 Thoraval, C., and Richards, M. A., 1997, The geoid constraint in global geodynamics:
 1734 viscosity structure, mantle heterogeneity models and boundary conditions:
 1735 *Geophysical Journal International*, v. 131, no. 1, p. 1-8.
 1736 van Wijk, J. W., Baldrige, W. S., van Hunen, J., Goes, S., Aster, R., Coblenz, D. D.,
 1737 Grand, S. P., and Ni, J., 2010, Small-scale convection at the edge of the
 1738 Colorado Plateau: implications for topography, magmatism, and evolution of
 1739 Proterozoic lithosphere: *Geology*, v. 38, no. 7, p. 611-614.
 1740 Veevers, J., and Conaghan, P., 1984, Phanerozoic earth history of Australia, Oxford
 1741 University Press, USA, 418 p.:
 1742 Watts, A., and Thorne, J., 1984, Tectonics, global changes in sea level and their
 1743 relationship to stratigraphical sequences at the US Atlantic continental
 1744 margin: *Marine and Petroleum Geology*, v. 1, no. 4, p. 319-339.
 1745 Weimer, R. J., 1970, Rates of deltaic sedimentation and intrabasin deformation,
 1746 Upper Cretaceous of Rocky Mountain region: *SEPM Special Publication*, v. 15,
 1747 p. 270-292.
 1748 Wernicke, B., 2011, The California River and its role in carving Grand Canyon:
 1749 *Geological Society of America Bulletin*, v. 123, no. 7/8, p. 1288-1326.
 1750 Wessel, P., and Smith, W. H. F., 1998, New, improved version of Generic Mapping
 1751 Tools released: *EOS transactions*, v. 79, p. 579-579.

1752 White, N., 1993, Recovery of strain rate variation from inversion of subsidence data:
 1753 Nature, v. 366, no. 6454, p. 449-452.
 1754 Whittaker, J. M., Müller, R. D., and Gurnis, M., 2010, Development of the Australian-
 1755 Antarctic depth anomaly: *Geochemistry Geophysics Geosystems*, v. 11.
 1756 Winterbourne, J., Crosby, A., and White, N., 2009, Depth, age and dynamic
 1757 topography of oceanic lithosphere beneath heavily sedimented Atlantic
 1758 margins: *Earth and Planetary Science Letters*, v. 287, no. 1–2, p. 137-151.
 1759 Xie, X., Müller, R. D., Li, S., Gong, Z., and Steinberger, B., 2006, Origin of anomalous
 1760 subsidence along the Northern South China Sea margin and its relationship
 1761 to dynamic topography: *Marine and Petroleum Geology*, v. 23, no. 7, p. 745-
 1762 765.
 1763 Zahirovic, S., Müller, R. D., Seton, M., Flament, N., Gurnis, M., and Whittaker, J., 2012,
 1764 Insights on the kinematics of the India-Eurasia collision from global
 1765 geodynamic models: *Geochemistry Geophysics Geosystems*, v. 13, p.
 1766 Q04W11.
 1767 Zandt, G., Gilbert, H., Owens, T. J., Ducea, M., Saleeby, J., and Jones, C. H., 2004, Active
 1768 foundering of a continental arc root beneath the southern Sierra Nevada in
 1769 California: *Nature*, v. 431, p. 41–46.
 1770 Zhang, N., Zhong, S., and Flowers, R. M., 2012, Predicting and testing continental
 1771 vertical motion histories since the Paleozoic: *Earth and Planetary Science*
 1772 *Letters*, v. 317–318, no. 0, p. 426-435.
 1773 Zhong, S., and Davies, G. F., 1999, Effects of plate and slab viscosities on the geoid:
 1774 *Earth and Planetary Science Letters*, v. 170, no. 4, p. 487-496.
 1775 Zhong, S., and Gurnis, M., 1994, Controls on trench topography from dynamic
 1776 models of subducted slabs: *Journal of Geophysical Research*, v. 99, p. 15,683-
 1777 615,695.
 1778 Zhong, S., Zuber, M. T., Moresi, L., and Gurnis, M., 2000, Role of temperature-
 1779 dependent viscosity and surface plates in spherical shell models of mantle
 1780 convection: *Journal of Geophysical Research*, v. 105, no. B5, p. 11,063-
 1781 011,082.
 1782 Zoback, M. L., and Mooney, W. D., 2003, Lithospheric Buoyancy and Continental
 1783 Intraplate Stresses: *International Geology Review*, v. 45, no. 2, p. 95-118.
 1784
 1785

A. This study

B. Steinberger (2007)

C. Panasyuk and Hager (2000)

D. Kaban et al. (2003)

A. This study

B. Steinberger (2007)

C. Ricard et al. (1993)

D. Conrad and Husson (2009)

E. Spasojevic and Gurnis (2012)

(A)

0 Ma

40 Ma

70 Ma

100 Ma

WD (km)

0

1.0

(B)

DT (km)

1.0

0

-1.0

(C)

T (°C)

0

-40

-80

-120

A. 151 Ma

B. 100 Ma

Dynamic topography [m]

C. 49 Ma

D. 0 Ma