

1-1-2018

Relationship of dietary nitrate intake from vegetables with cardiovascular disease mortality: a prospective study in a cohort of older Australians

Alex Liu
University of Western Australia

Catherine Bondonno
University of Western Australia, Edith Cowan University

Joanna Russell
University of Wollongong, jrussell@uow.edu.au

Victoria M. Flood
University Of Sydney, Westmead Hospital, vflood@uow.edu.au

Joshua Lewis
University of Western Australia, Edith Cowan University, Children's Hospital at Westmead, University of Sydney

See next page for additional authors.
Follow this and additional works at: <https://ro.uow.edu.au/sspapers>



Part of the [Education Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Liu, Alex; Bondonno, Catherine; Russell, Joanna; Flood, Victoria M.; Lewis, Joshua; Croft, Kevin D.; Woodman, Richard; Lim, Wai; Kifley, Annette; Wong, Germaine; Mitchell, Paul; Hodgson, Jonathan; and Blekkenhorst, Lauren, "Relationship of dietary nitrate intake from vegetables with cardiovascular disease mortality: a prospective study in a cohort of older Australians" (2018). *Faculty of Social Sciences - Papers*. 4274.

<https://ro.uow.edu.au/sspapers/4274>

Relationship of dietary nitrate intake from vegetables with cardiovascular disease mortality: a prospective study in a cohort of older Australians

Abstract

Purpose: Short-term trials indicate inorganic nitrate and nitrate-rich vegetables may have vascular health benefits. However, few observational studies have explored the relationship between nitrate intake and long-term cardiovascular disease (CVD) outcomes. The primary aim of this study was to investigate the association of nitrate intake from vegetables with CVD mortality in a sample of older Australians.

Methods: A subgroup of participants without diabetes or major CVD at baseline (1992–1994) were included from the Blue Mountains Eye Study, a population-based cohort study of men and women aged ≥ 49 years. Diets were evaluated using a validated food frequency questionnaire at baseline, 5 years and 10 years of follow-up. Vegetable nitrate intake was estimated using a comprehensive vegetable nitrate database. Cox proportional hazard regression was used to explore the association between vegetable nitrate intake and CVD mortality. **Results:** During 14 years of follow-up, 188/2229 (8.4%) participants died from CVD. In multivariable-adjusted analysis, participants in quartile 2 [69.5–99.6 mg/day; HR 0.53 (95% CI 0.35, 0.82)], quartile 3 [99.7–137.8 mg/day; HR 0.51 (95% CI 0.32, 0.80)], and quartile 4 [> 137.8 mg/day; HR 0.63 (95% CI 0.41, 0.95)] of vegetable nitrate intake had lower hazards for CVD mortality compared to participants in quartile 1 (< 69.5 mg/day). **Conclusions:** In older Australian men and women, vegetable nitrate intake was inversely associated with CVD mortality, independent of lifestyle and cardiovascular risk factors. These findings confirm a recent report that intake of vegetable nitrate lowers the risk of CVD mortality in older women and extend these findings to older men.

Disciplines

Education | Social and Behavioral Sciences

Publication Details

Liu, A. H., Bondonno, C. P., Russell, J., Flood, V. M., Lewis, J. R., Croft, K. D., Woodman, R. J., Lim, W. H., Kifley, A., Wong, G., Mitchell, P., Hodgson, J. M. & Blekkenhorst, L. C. (2019). Relationship of dietary nitrate intake from vegetables with cardiovascular disease mortality: a prospective study in a cohort of older Australians. *European Journal of Nutrition*, 58 (7), 2741-2753.

Authors

Alex Liu, Catherine Bondonno, Joanna Russell, Victoria M. Flood, Joshua Lewis, Kevin D. Croft, Richard Woodman, Wai Lim, Annette Kifley, Germaine Wong, Paul Mitchell, Jonathan Hodgson, and Lauren Blekkenhorst

Relationship of dietary nitrate intake from vegetables with cardiovascular disease mortality: a prospective study in a cohort of older Australians

Alex H Liu ¹, Catherine P Bondonno ^{1,2}, Joanna Russell ³, Victoria M Flood ^{4,5}, Joshua R Lewis ^{1,2,6,7}, Kevin D Croft ¹, Richard J Woodman ⁸, Wai H Lim ⁹, Annette Kifley ¹⁰, Germaine Wong ^{6,7}, Paul Mitchell ¹⁰, Jonathan M Hodgson ^{1,2}, Lauren C Blekkenhorst ²

¹ Medical School, Royal Perth Hospital Unit, The University of Western Australia, Perth, WA, Australia

² School of Medical and Health Sciences, Edith Cowan University, Joondalup, WA, Australia

³ School of Health and Society, Faculty of Social Sciences, The University of Wollongong, Australia

⁴ Faculty of Health Sciences, The University of Sydney, Sydney, NSW, Australia

⁵ Western Sydney Local Health District, Westmead Hospital, Westmead, NSW, Australia

⁶ Centre for Kidney Research, Children's Hospital at Westmead, Westmead, NSW, Australia

⁷ School of Public Health, Sydney Medical School, The University of Sydney, NSW, Australia

⁸ Flinders Centre for Epidemiology and Biostatistics, Flinders University, Adelaide, SA, Australia

⁹ Department of Renal Medicine, Sir Charles Gairdner Hospital, Nedlands, WA, Australia

¹⁰ Centre for Vision Research, Department of Ophthalmology, Westmead Institute for Medical Research, The University of Sydney, NSW, Australia

Authors' last names: Liu, Bondonno, Russell, Flood, Lewis, Croft, Woodman, Lim, Wong, Mitchell, Hodgson, Blekkenhorst

Disclaimers: None

Corresponding author:

Lauren Blekkenhorst

School of Medical and Health Sciences

Edith Cowan University

270 Joondalup Drive

JOONDALUP WA 6027 AUSTRALIA

Tel: 61 8 6304 4604

Email: l.blekkenhorst@ecu.edu.au

Acknowledgements: The analysis of this study was supported by a National Health and Medical Research Council (NHMRC) of Australia project grant 1084922. The salaries of JRL and JMH were supported by NHMRC of Australia Fellowships. None of these funding agencies had any role in the conduct of the study;

collection, management, analysis or interpretation of the data; or the preparation, review or approval of the manuscript.

2 **Abstract**

3 **Purpose:** Short-term trials indicate inorganic nitrate and nitrate-rich vegetables may have vascular health benefits.
4 However, few observational studies have explored the relationship between nitrate intake and long-term
5 cardiovascular disease (CVD) outcomes. The primary aim of this study was to investigate the association of nitrate
6 intake from vegetables with CVD mortality in a sample of older Australians.

7 **Methods:** A subgroup of participants without diabetes or major CVD at baseline (1992-1994) were included from
8 the Blue Mountains Eye Study, a population-based cohort study of men and women aged ≥ 49 years. Diets were
9 evaluated using a validated food frequency questionnaire at baseline, 5 years and 10 years of follow-up. Vegetable
10 nitrate intake was estimated using a comprehensive vegetable nitrate database. Cox proportional hazards
11 regression was used to explore the association between vegetable nitrate intake and CVD mortality.

12 **Results:** During 14 years of follow-up, 188/2229 (8.4%) participants died from CVD. In multivariable-adjusted
13 analysis, participants in quartile 2 [69.5-99.6 mg/d; HR: 0.53 (95%CI: 0.35, 0.82)], quartile 3 [99.7-137.8 mg/d;
14 HR: 0.51 (95%CI: 0.32, 0.80)], and quartile 4 [>137.8 mg/d; HR: 0.63 (95%CI: 0.41, 0.95)] of vegetable nitrate
15 intake had lower hazards for CVD mortality compared to participants in quartile 1 (<69.5 mg/d).

16 **Conclusions:** In older Australian men and women, vegetable nitrate intake was inversely associated with CVD
17 mortality, independent of lifestyle and cardiovascular risk factors. These findings confirm a recent report that
18 intake of vegetable nitrate lowers the risk of CVD mortality in older women and extend these findings to older
19 men.

20 **Key words:** cardiovascular disease, mortality, nitrate, vegetables, coronary heart disease, stroke

21 **Introduction**

22 Cardiovascular disease (CVD) is the leading cause of death in both developed and developing countries,
23 accounting for nearly one third of all deaths worldwide [1]. The majority of CVD deaths are caused by stroke or
24 coronary heart disease (CHD) [2,3].

25 Nitric oxide (NO) plays a key role in cardiovascular health. It is continuously produced at low levels in the
26 endothelium, controlling basal vascular tone and integrity, with effects on blood flow, blood pressure, endothelial
27 function and platelet function [4]. Decreased levels are associated with dysfunction to the vasculature with
28 increased arterial stiffness and higher blood pressure observed after inhibition of NO synthesis [5]. Originally
29 thought to be derived solely through the L-arginine-NO synthase pathway, the discovery of an alternate nitrate-
30 nitrite-NO pathway raised the possibility that dietary nitrate could be an important source of NO [6,7]. Vegetables
31 contribute 80-90% to the total nitrate consumed in the diet [7,8], with the remainder coming from other food
32 sources such as cured meats [9]. The addition of nitrate and nitrite to cured meats is controversial due to the
33 potential of nitrate and nitrite to form N-nitrosoamines, which are potential carcinogens [10]. Compounds such as
34 polyphenols, vitamins C and E, and other antioxidants found abundantly in vegetables inhibit the formation of N-
35 nitrosoamines [10]. The evidence of nitrate from specific foods and the risk of certain cancers remains unclear
36 [11]. It is now well-established that after ingestion of nitrate there is an active uptake of plasma nitrate by the
37 salivary glands. In the oral cavity nitrate is converted to nitrite by enzymatic action of bacteria on the dorsal
38 surface of the tongue. After swallowing, this nitrite is absorbed becoming a source of NO [12]. Indeed, data from
39 acute and short-term clinical trials indicate that dietary nitrate may lower blood pressure and improve vascular
40 function via effects on NO [13-20]. The potential impact of these functional changes on long-term CVD risk
41 remains uncertain [9,21].

42 Data from epidemiological studies have indicated a link between diets rich in vegetables [22-24], particularly
43 green leafy vegetables [25,26], and lower risk of CVD. Few epidemiological studies have investigated the
44 associations between nitrate intake from vegetables and CVD mortality. We have recently observed a significantly
45 lower risk of atherosclerotic vascular disease mortality and ischemic cerebrovascular disease events with higher
46 vegetable nitrate intake in a cohort of elderly Australian women [27,28]. The aim of this study was to confirm
47 these findings in another Australian cohort that included both men and women. Nitrate intake from vegetables
48 was our primary focus due to the large proportion of nitrate in the diet coming from vegetables alone [7,8].

49 **Methods**

50 **Study population**

51 Data used in this prospective cohort study were obtained from the Blue Mountains Eye Study (BMES). The BMES
52 was the first population-based study of common eye diseases and other health outcomes in a representative older
53 Australian population. The population was predominantly Caucasian [29], aged 49 years and older, better
54 educated than Australia overall, and fairly stable in terms of follow-up [30,31]. Participants were initially
55 examined in 1992-1994 for baseline characteristics and were followed up at 5-, 10- and 15-year time points.
56 Mortality data were obtained via data linkage with the Australian National Death Index (NDI) [30,31]. Further
57 study details of the BMES have been previously reported [32,33].

58 The BMES was approved by the Sydney West Area Health Service and the University of Sydney Human Research
59 Ethics Committees. The study was conducted according to guidelines from the Declaration of Helsinki. Written
60 informed consents were obtained from all participants.

61 Of the 3654 participants that had completed the main study, 3267 responded to a food frequency questionnaire
62 (FFQ) and 2900 gave FFQ responses of sufficient quality for inclusion. Previous data cleaning processes have
63 been reported elsewhere [32,34]. The eligibility criteria for the present study included available data on all
64 exposure and outcome variables. The exclusion criteria included a report of extreme energy intake (< 500 and <
65 800 kcal/d or > 3500 and > 4000 kcal/d in females and males, respectively), and baseline self-reported history of
66 major CVD (myocardial infarction (MI) and stroke) and/or diabetes. Of the 2900 participants, 163 (5.6%) were
67 excluded for missing data on mortality (n=17), energy intake (n=3), and baseline history of major CVD and
68 diabetes (n=143), 40 (1.4%) were excluded for reporting extreme energy intakes, and data cleaning excluded an
69 additional participant. Participants who reported a baseline history of major CVD (n=264) or diabetes (n=157), or
70 both (n=46) were further excluded (n=467, 16.1%). In total, 2229 out of 2900 (76.9%) participants with FFQ data
71 were eligible for the present study (**Figure 1**).

72 **Dietary assessment**

73 A 145-item self-administered semi-quantitative FFQ, modified from a previous version by Willett et al. [35], was
74 used to assess dietary intake. Participants were required to estimate their usual frequency of consumed food items
75 during the past year. A 9-category frequency scale was used, ranging from never to four or more times per day.

76 Portion sizes were given in household measures (e.g. cup, teaspoon). Twenty-five vegetable items were included
77 in the questionnaire. Reproducibility and validity of the FFQ were verified in a subgroup of the study population
78 (n=79) using three, four-day weighed food records completed four months apart [32]. Nutrient data were analysed
79 using the Australian Tables of Food Composition (NUTTAB90 and NUTTAB95) databases [36,37].

80 *Vegetable nitrate intake*

81 Vegetable nitrate intake was calculated from consumed vegetables, which were assessed using the FFQ, and
82 quantified in grams per day. A recently published comprehensive database, which has nitrate data for 178
83 vegetables from over 250 publications, was applied to assess nitrate intake for each vegetable item [38]. Nitrate
84 intake (mg/d) was calculated by multiplying the amount of each vegetable (g/d) by the median nitrate content
85 (mg/g) for that individual vegetable. The nitrate values from each individual vegetable were then summated to
86 obtain the total daily nitrate intake derived from vegetables.

87 *Total nitrate intake*

88 Nitrate intakes from other food items in the FFQ were also calculated. Nitrate levels from food items were obtained
89 from Inoue-Choi et al [39], the Food Standards Australia New Zealand survey of nitrate and nitrite in food and
90 beverages in Australia [40], and Griesenbeck et al [41]. The value of nitrate was given zero if the food item was
91 not listed in any of the databases aforementioned and if the food item was not similar to any other food item listed
92 with a nitrate value, which was vegemite only. Nitrate intake (mg/d) was calculated by multiplying the daily
93 consumption of that food item (g/d) by the assigned mean nitrate value (mg/g) for that food item. Total nitrate
94 intake (mg/d) was calculated by summing daily nitrate intakes from all other food items together with vegetable
95 items.

96 **Baseline demographic assessments**

97 Demographic information, socio-economic status and lifestyle factors were obtained from the participants through
98 a detailed questionnaire administered by trained interviewers [30,31]. Smoking status was determined using
99 categories of “never smoked”, “past smoker” and “current smoker”, with classification of current smoker
100 including those who stopped smoking within the past 12 months. Weight (kg) and height (m) were assessed using
101 digital scales and a stadiometer, respectively. Body mass index (BMI, kg/m²) was calculated using weight and
102 height measurements. Energy and alcohol intake were estimated using the FFQ described above [30,31]. Physical

103 activity was categorised into three groups according to activity level (not active, light exercise, and vigorous
104 exercise). Assessment of physical activity has been described elsewhere [42].

105 Baseline questionnaires were used to determine values for other potential confounding variables including use of
106 organic nitrate medication, use of antihypertensive medication, use of cholesterol-lowering medication (statins),
107 and use of low-dose aspirin. Fasting blood samples were used to determine total cholesterol levels and glucose
108 concentrations. Renal function was evaluated using Modification of Diet in Renal Disease (MDRD) study
109 equation for estimated glomerular filtration rate (eGFR) [43]. Systolic and diastolic blood pressures were
110 measured using a mercury sphygmomanometer after participants were seated and rested for at least 5 minutes
111 [30,31].

112 The participants provided their previous medical history and current medications, which were verified by their
113 general practitioner. These data were coded using the International Classification of Primary Care-Plus method.
114 This coding methodology allows aggregation of different terms for similar pathologic entities as defined by the
115 International Classification of Disease (ICD-10) coding system. These data were used to determine the presence
116 of pre-existing diabetes mellitus and major CVD (MI and stroke).

117 **Cardiovascular disease mortality assessment**

118 The primary outcome of this study was CVD mortality during 14 years of follow-up, the period available for
119 linkage with the mortality data. CVD mortality data were retrieved by matching the causes of death by CHD or
120 stroke to the NDI. Causes of death were determined by the use of diagnosis codes from the International
121 Classification of Diseases Code, 9th revision (ICD-9) and the International Statistical Classification of Diseases,
122 10th revision (ICD-10). The following codes were used for CHD: (ICD-9: 410.0 to 410.9, 411.0 to 411.8, 412.0,
123 414.0 to 414.9 and ICD-10: I21.0 to I21.9, I22.0 to I22.9, I23.0 to I23.8, I24.0 to I24.9 and I25.0 to I25.9) and
124 stroke: (ICD-9: 430.0 to 438.9 and ICD-10: I60.0 to I69.9). The Australian NDI has been validated and proven to
125 be highly sensitive and specific for CVD mortality (92.5% and 89.6%, respectively) [44].

126 **Statistical analysis**

127 An analytic protocol was developed before formal analyses were performed. Data were analysed using IBM SPSS
128 Statistics for Windows, version 21.0 (IBM) and SAS software, version 9.4 (SAS Institute Inc). Statistical
129 significance was set at a two-sided Type 1 error rate of $P < 0.05$ for all tests. Normally distributed continuous

130 variables were reported as mean \pm standard deviation (SD), non-normally distributed continuous variables were
131 expressed as medians and interquartile ranges (IQRs), and categorical variables were expressed as numbers and
132 proportions (%). Differences in baseline characteristics were tested using one-way ANOVA for normally
133 distributed continuous variables, the Kruskal-Wallis test for non-normally distributed variables, and Chi-squared
134 test for categorical variables.

135 The primary outcome of interest was CVD mortality. Follow-up was calculated in years from each participant's
136 baseline visit until death or 14-year follow-up. Cox proportional hazards (PH) regression was used to investigate
137 the association between exposure variables (vegetable nitrate, total nitrate and non-vegetable nitrate) and CVD
138 mortality. Further analyses exploring the relationship between exposure variables and CHD, stroke, non-CVD,
139 and all-cause mortality were also analysed. We used the mean vegetable nitrate, total nitrate, and non-vegetable
140 nitrate intake, calculated throughout the duration of follow-up (baseline, 5-year and 10-year). We tested for
141 associations between outcomes and exposure variables by comparing a linear relationship with a categorical
142 relationship using quartiles. Because most outcome-exposure relationships had a better fitting model using the
143 exposure variable entered as a categorical variable, we used this approach in all models. We also categorised
144 vegetable nitrate intake using previously reported cut-offs (<52.7 mg/d; 52.7-76.4 mg/d; and >76.4 mg/d) [27,28].
145 These cut-offs were used to compare the results of the present study with reported results from another Australian
146 population [27,28]. Three models of adjustment were used including: (1) unadjusted; (2) age-, gender- and energy-
147 adjusted; and (3) multivariable-adjusted models. The covariates included in the multivariable-adjusted model were
148 age (years), gender (female or male), BMI (kg/m^2), physical activity (not active, light exercise or vigorous
149 exercise), alcohol intake (grams per day), smoking history (never smoked, previous smoker or current smoker),
150 socio-economic status (home/unit owner), use of organic nitrate medications (yes or no), use of antihypertensive
151 medications (yes or no), use of statin medications (yes or no), use of low-dose aspirin (yes or no), renal function
152 (MDRD eGFR, expressed as $\text{ml}\cdot\text{min}^{-1}\cdot 1.73\text{m}^{-2}$) and energy intake (kcal per day). Age, alcohol intake, renal
153 function and energy intake were entered into models as continuous covariates, with the rest entered as categorical
154 covariates. Cox PH assumptions were tested using log-log plots, which were shown to be parallel indicating that
155 PH assumptions were not violated.

156 **Sensitivity analyses**

157 In order to rule out potential bias from reverse causality, we excluded all deaths that occurred in the first 24 months
158 of follow-up, and repeated the multivariable-adjusted Cox PH models for quartiles of vegetable nitrate intake and

159 CVD, CHD and stroke mortality. Pearson product-moment correlation coefficient (r) was used to investigate the
160 relationship between vegetable nitrate intake and total vegetable intake. A forward stepwise Cox PH model for
161 CVD, CHD and stroke mortality including all adjustment variables plus quartiles of vegetable nitrate intake
162 (entered as a categorical variable) and total vegetable intake (entered as a categorical variable) was used to assess
163 multicollinearity between vegetable nitrate intake and total vegetable intake. Total vegetable intake was
164 categorised into servings of vegetables (<2; 2 to <3; 3 to <4; 4 to <5; and ≥ 5). As vegetable nitrate intake may
165 also be a marker for a healthier diet, we further adjusted for diet quality using a modified version of the Healthy
166 Eating Index for Australians. This diet quality index called the Total Diet Score (TDS) has been described
167 elsewhere [31]. The TDS was entered in the multivariable-adjusted model with quartiles of vegetable nitrate intake
168 (entered as a categorical variable) for CVD, CHD and stroke mortality. In addition, greater fish consumption has
169 been shown to be inversely associated with cardiovascular health in the BMES cohort [30]. Therefore, we further
170 adjusted for fish intake in the multivariable-adjusted model. In addition to fish intake, we also individually
171 adjusted for fibre and total flavonoid intakes. Estimation of total flavonoid intake has been described elsewhere
172 [45]. Interaction terms were used to investigate whether there was any evidence of different relationships
173 according to gender and participants aged < and ≥ 70 years.

174 **Results**

175 **Characteristics of the study population**

176 Baseline characteristics and dietary intakes for all participants and by quartiles of vegetable nitrate intakes are
177 shown in **Table 1** and **Table 2**, respectively. Differences were observed between quartiles of vegetable nitrate
178 intake for gender, age, physical activity, and smoking history (Table 1). The mean (SD) vegetable nitrate intake
179 was 109.7 (59.2) mg/d, and total nitrate from all foods was 128.8 (62.0) mg/d. Vegetables contributed 85.2%, fruit
180 5.5%, meat (processed and unprocessed) 3.7%, grains 1.7%, fish 0.2%, and dairy 0.1% to the total nitrate intake.
181 The highest contributing vegetables towards nitrate intake were lettuce (32.2%), spinach and silverbeet (13.2%),
182 and potato (10.7%). Differences in all nutrients were observed between quartiles of vegetable nitrate, except for
183 saturated fat (g/d) (Table 2).

184 **Cardiovascular disease mortality**

185 Over 14 years of follow-up, 188 of 2229 participants (8.4%) died because of CVD. The relationship between
186 quartiles of vegetable nitrate intake and CVD mortality are presented in **Table 3**. Compared with quartile 1,
187 quartiles 2, 3 and 4 were associated with a lower relative hazard of CVD mortality in the multivariable-adjusted
188 model. Multivariable-adjusted cumulative survival curves for CVD by quartiles of vegetable nitrate intake are
189 shown in **Figure 2**. Associations by quartiles of non-vegetable nitrate intake and total nitrate intake with CVD
190 mortality are presented in **Supplemental Table 1** and **Supplemental Table 2**, respectively. Associations for
191 intakes of total nitrate were similar to vegetable nitrate, but non-vegetable nitrate was not associated with CVD
192 mortality.

193 *Coronary heart disease, stroke, non-cardiovascular disease and all-cause mortality*

194 Over 14 years of follow-up, 125 of 2229 participants (5.6%) died of CHD and 63 participants (2.8%) died of
195 stroke. Furthermore, 422 of 2229 participants (18.9%) died of non-CVD causes and 610 of 2229 participants
196 (27.4%) died of any cause. The relationship between quartiles of vegetable nitrate intake and CHD, stroke, non-
197 CVD, and all-cause mortality are presented in Table 3. For CHD, quartile 2 (69.5-99.6 mg/d), compared to the
198 referent, was associated with a lower relative hazard of mortality in the multivariable-adjusted model (P=0.008).
199 Compared with quartile 1, quartile 3 (99.7-137.8 mg/d) was associated with a lower hazard of stroke mortality in
200 the multivariable-adjusted model (P=0.002) (Table 3). Compared with quartile 1, quartile 4 (>137.8 mg/d) was

201 associated with a lower relative hazard of non-CVD ($P=0.009$) and all-cause ($P=0.001$) mortality in multivariable-
202 adjusted models (Table 3). For non-vegetable nitrate intake and total nitrate intake, quartile results are presented
203 in Supplemental Table 1 and Supplemental Table 2, respectively.

204 **Sensitivity analyses**

205 Using previously reported cut-offs in older women [27,28], compared with low intakes of vegetable nitrate (<52.7
206 mg/d), moderate (52.7-76.4 mg/d) and high (>76.4 mg/d) intakes were associated with 50 to 55% lower relative
207 hazard of CVD mortality in multivariable-adjusted models ($P=0.001$) (**Table 4**). Multivariable-adjusted
208 cumulative survival curves for CVD by categories of vegetable nitrate intake using previously reported cut-offs
209 are shown in **Supplemental Figure 1**.

210 Similarly, compared with low intakes of vegetable nitrate (<52.7 mg/d), moderate (52.7-76.4 mg/d) and high
211 (>76.4 mg/d) intakes were associated with 51 to 53% lower relative hazard of CHD mortality ($P=0.017$), and 44
212 to 61% lower relative hazard of stroke mortality ($P=0.036$), in multivariable-adjusted models (Table 4). For non-
213 CVD and all-cause mortality, moderate (52.7-76.4 mg/d) and high (>76.4 mg/d) intakes were associated with 25
214 to 27% ($P=0.111$) and 34 to 37% ($P=0.001$) lower relative hazard, respectively, in comparison with low intakes
215 of vegetable nitrate (<52.7 mg/d) in multivariable-adjusted models (Table 4).

216 To assess the likelihood of reverse causality bias (sick people eat less food/vegetables), we excluded all deaths
217 that occurred in the first 24 months. The observed association between quartiles of vegetable nitrate intake and
218 CVD, CHD, and stroke mortality remained significant in multivariable-adjusted models ($P<0.05$ for all).

219 A strong positive correlation between vegetable nitrate intake and vegetable intake was observed ($r=0.79$,
220 $P<0.001$). In a forward-stepwise Cox PH model for the prediction of CVD mortality, which included all
221 adjustment variables as well as total vegetable intake and vegetable nitrate intake, age ($P<0.001$), gender
222 ($P=0.002$), smoking ($P<0.001$), use of organic nitrate medication ($P=0.001$), and vegetable nitrate intake
223 ($P=0.008$) remained in the final parsimonious model. For CHD mortality, age ($P<0.001$), gender ($P=0.003$),
224 smoking ($P<0.001$), and use of organic nitrate medication ($P=0.004$) remained in the final parsimonious model.
225 For stroke mortality, age ($P<0.001$), BMI ($P=0.023$), use of antihypertensive medication ($P=0.051$), and vegetable
226 nitrate intake ($P=0.009$) remained in the final parsimonious model.

227 After additional adjustment for the TDS, in multivariable-adjusted Cox PH models, results for quartiles of
228 vegetable nitrate intake remained similar for CVD (P=0.006), CHD (P=0.069) and stroke (P=0.013) mortality
229 (**Supplemental Table 3**). After additional adjustment for fish intake, in multivariable-adjusted Cox PH models,
230 results for quartiles of vegetable nitrate intake were similar for CVD (P=0.011) and CHD (P=0.060) mortality,
231 but for stroke mortality, the relationship became borderline significant (P=0.052) (Supplemental Table 3). After
232 individual adjustments for fibre and total flavonoid intakes, in multivariable-adjusted Cox PH models, results for
233 quartiles of vegetable nitrate intake were similar for CVD (P<0.01 for both) and stroke (P<0.05 for both) mortality.
234 For CHD mortality, the relationship was similar after adjusting for total flavonoid intake (P=0.064), and became
235 statistically significant after further adjusting for fibre intake (P=0.035). For CVD, CHD and stroke mortality,
236 there was no evidence that the relationships were different according to gender ($P_{\text{interaction}} > 0.05$ for all) or for ages
237 < 70 and ≥ 70 years ($P_{\text{interaction}} > 0.05$ for all). Quartiles of vegetable nitrate and CVD, CHD and stroke mortality
238 stratified by gender are presented in **Supplemental Table 4**.

239 Discussion

240 This 14-year prospective study demonstrated an inverse association between vegetable nitrate intake and CVD
241 mortality in an older Australian population without diabetes and/or major CVD at baseline. The association
242 remained significant after adjustment for lifestyle and CVD risk factors as well as diet quality and fish, fibre and
243 total flavonoid intakes. To our knowledge, this study is one of the first prospective cohort studies to investigate
244 the association of dietary nitrate intake with CVD in an older general population that includes both men and
245 women.

246 Since Larsen et al. [13] reported a blood pressure lowering effect of dietary nitrate in 2006, numerous small-scale
247 clinical trials have been conducted to investigate dietary nitrate intake with blood pressure, endothelial function
248 and other CVD-related outcomes. Most studies investigating the acute effects of nitrate ingestion have
249 demonstrated positive findings [15,16,46]. However, studies investigating chronic effects have shown
250 contradictory results [14,18,47-50]. Our previous studies have failed to demonstrate blood pressure lowering
251 effects after 7-day consumption of nitrate-rich green leafy vegetables or beetroot juice in pre-hypertensive and
252 treated hypertensive participants, respectively [48,49]. A meta-analysis [51] including 16 acute and short-term
253 studies indicated consumption of inorganic nitrate or nitrate-rich beetroot juice was associated with a reduction in
254 systolic blood pressure. Another meta-analysis [52] including 13 short- and medium-term studies showed positive
255 effects of nitrate ingestion on measures of clinic blood pressure, but not on more accurate measures of ambulatory
256 or home blood pressure. Effects on clinic blood pressure were removed when the authors excluded all studies
257 investigating exercise performance from the analysis. In addition to these meta-analyses on blood pressure, a
258 meta-analysis [53] including 12 acute, short-, and medium-term studies demonstrated beneficial effects of nitrate
259 ingestion on endothelial function. These effects were attenuated in older individuals and individuals with
260 increased CVD risk. Due to these inconsistencies, there is a need to investigate the effects of nitrate ingestion in
261 older and at risk individuals. There is also a need to explore the relationship of habitual intake of dietary nitrate
262 with long-term risk of cardiovascular-related outcomes.

263 To our knowledge, few studies have investigated the long-term associations of nitrate intake with cardiovascular-
264 related outcomes. Bahadoran et al. [54] reported habitual intake of nitrite, rather than nitrate, was inversely
265 associated with the incidence of hypertension over 5.8 years of follow-up. In an Australian population of older
266 women aged ≥ 70 years, we have previously shown an inverse association between vegetable nitrate intake and
267 atherosclerotic vascular disease mortality, which was defined as any death attributed to CHD, ischaemic

268 cerebrovascular disease, peripheral arterial disease and heart failure [27]. We have also shown an inverse
269 relationship between vegetable nitrate intake and ischaemic cerebrovascular disease hospitalisations and deaths
270 [28]. In both prospective studies, the lowest relative hazards were observed for vegetable nitrate intakes between
271 52.7-76.4 mg/d, equivalent to ~1-2 cups of leafy green vegetables per day. Epidemiological studies have also been
272 conducted using biological samples, such as plasma nitrate [55] and urinary nitrate excretion [56]. In a cross-
273 sectional study, Smallwood et al [56] demonstrated urinary nitrate excretion over 24 hours was inversely
274 associated with blood pressure. In a 17-year prospective cohort study, Maas et al [55] found no evidence for an
275 association between plasma nitrate and incident CVD, and demonstrated higher plasma nitrate was associated
276 with an increased risk of death from all causes. This latter relationship is suggested to be in part attributable to
277 higher plasma nitrate being correlated with lower renal function [55]. Using biological samples, such as plasma
278 nitrate and urinary nitrate excretion as markers of nitrate exposure are somewhat controversial. For example,
279 plasma nitrate is likely a representative of both protective and detrimental processes within the body [55].
280 Therefore, caution needs to be taken when interpreting the findings from these studies.

281 In the present study, we demonstrated an inverse relationship between vegetable nitrate intake and CVD mortality.
282 Even though the lowest relative hazard for CVD was observed in participants consuming vegetable nitrate intake
283 ranging from 99.7-137.8 mg/d, the greatest benefit could not be determined as there was overlapping of the
284 confidence intervals of quartiles 2, 3 and 4. However, in agreement with our recent study in older women, the
285 findings suggest that older individuals consuming even relatively low levels of vegetable nitrate (>52.7 mg/day,
286 equivalent to ~1 cup of leafy green vegetables per day), have half the risk of 14-year CVD mortality even after
287 adjusting for cardiovascular risk factors. In sub-group analyses, an inverse relationship between vegetable nitrate
288 intake and stroke was also demonstrated, which aligns with our previous study findings [28]. The lowest relative
289 hazard for stroke mortality was observed in individuals consuming vegetable nitrate intake ranging from 99.7
290 mg/d to 137.8 mg/d, equivalent to ~2-3 cups of leafy green vegetables per day. However, it is important to note
291 that, as mentioned above, the confidence intervals overlap for each quartile and therefore the greatest benefit
292 cannot be determined. It is also important to note that the apparent J-shaped relationship may be an artefact of the
293 cut-points used i.e. quartiles. There is a possibility that by chance there were a lower number of stroke events than
294 expected in the 3rd quartile. The J-shaped relationship in Table 3 disappeared in Table 4 when we collapsed
295 vegetable nitrate intake into three groups based on previously reported cut-points. When collapsing into three
296 groups there is less chance of an artificially small number of events in a single category.

297 As discussed above, clinical trials have demonstrated reductions in blood pressure with nitrate supplementation.
298 In addition, blood pressure is strongly linked with stroke outcomes [57]. This could indicate that moderate habitual
299 intakes of vegetable nitrate could be beneficial for stroke outcomes through blood pressure lowering effects.
300 However, this needs to be explored further. In our study, no difference in blood pressure between quartiles was
301 observed. A single clinic measure may not be sufficient in capturing accurate blood pressure. Home and
302 ambulatory measures of blood pressure are widely accepted to be more accurate [58,59].

303 The intake of vegetable nitrate is likely to vary between populations [60-62]. In the current study, which included
304 both men and women, the mean vegetable nitrate intake was 110 mg/d. In our previously reported study of older
305 women, the mean vegetable nitrate intake was 67 mg/d (difference equivalent to ~0.5-1 cup of leafy green
306 vegetables) [27,28]. Differences between these two cohorts may be due to different ages and/or demographic areas.
307 In the present study, the participants were somewhat younger (mean age 65 versus 75 years) and lived in a semi-
308 urban area, while participants in the previous study were from a metropolitan area. The use of different FFQs in
309 the two studies may also have influenced the estimated nitrate intake. In addition, geographic variability in the
310 nitrate content of specific foods are likely to occur in Australia as it does in the United States and could affect
311 nitrate intake [21].

312 There are several strengths to this study. Firstly, we used a validated dietary assessment tool for dietary evaluation.
313 A comprehensive newly-developed vegetable nitrate database was used to calculate nitrate intake [38]. Secondly,
314 we removed the likelihood of reverse-causality by excluding all deaths in the first 24 months of follow-up. This
315 resulted in a stronger relationship between vegetable nitrate intake and CVD mortality. Thirdly, the long-term
316 nature of the BMES captures an important period in the life course of these older individuals when cardiovascular
317 disease deaths become more common, increasing the power to detect an association between the exposure variable
318 and outcome of interest. Fourthly, using pre-specified cut points, we were able to replicate our previous findings
319 in older women [27,28]. Lastly, the participants of this study were representative of an older Australian population
320 [30,34].

321 Limitations also need to be considered. Due to the observational nature of the study, we are not able to infer
322 causality or rule out residual confounding or the introduction of bias. Vegetable nitrate intake was inversely
323 associated with non-CVD mortality even though the major proposed mechanisms appear specific to CVD. This
324 may support the hypothesis that residual confounding by other dietary or lifestyle factors that may be affecting
325 both CVD and non-CVD mortality. However, evidence suggests nitrate intake may be associated with other

326 health-related outcomes such as increased mitochondrial efficiency [63], improvements in features of metabolic
327 syndrome [64], reductions in pro-inflammatory cytokines [65], and improvements in cognitive function and
328 increased cerebral blood flow [66], all factors that could possibly lead to a reduction in death from other non-
329 CVD causes. In addition, NO has beneficial health effects that go beyond vascular health, including
330 neurotransmission and non-specific host defence [67-69]. The use of vegetable nitrate intake averaged over the
331 period of follow-up could also be considered as a limitation. Compared to baseline vegetable nitrate intake, even
332 though it is more likely to better represent participants' eating habits and minimize the random bias from a single
333 FFQ response, it may also attenuate the strength of the relationship. However, we observed similar results when
334 assessing the relationship using only baseline vegetable nitrate intake in multivariable-adjusted models (data not
335 shown). In addition, vegetable nitrate intake may be a surrogate marker for vegetable consumption and/or diet
336 quality. To address this matter, we conducted a forward-stepwise Cox PH model including both vegetable nitrate
337 intake and total vegetable consumption. In the final parsimonious model, vegetable nitrate intake remained
338 together with age, gender, smoking status, and use of organic nitrate medication. We then conducted further
339 analyses separately adjusting for diet quality (i.e. TDS) and fish [30], fibre and total flavonoid intakes. After the
340 additional separate adjustments for TDS and fish, fibre and total flavonoid intakes, the relationship between
341 vegetable nitrate intake and CVD mortality remained statistically significant in multivariable-adjusted models.
342 However, we still cannot rule out the possibility that vegetable nitrate may be a marker for healthy eating and
343 other beneficial foods/nutrients. In particular, the major source of nitrate intake from vegetables is leafy green
344 vegetables. Leafy green vegetables are inversely associated with cardiovascular disease outcomes [70] and indeed
345 have other beneficial nutrients and phytochemicals that may contribute to these potential benefits [71-73].

346 To conclude, our study demonstrated an inverse relationship between vegetable nitrate intake and CVD and stroke
347 mortality, independent of lifestyle and CVD risk factors in an older Australian population of men and women. We
348 also demonstrated an inverse relationship with non-CVD and all-cause mortality. Further studies are needed in
349 other populations to confirm these results and to evaluate the optimum intake of nitrate from vegetables for the
350 possible prevention of cardiovascular disease.

Declaration of Conflict of Interest: The authors declare that they have no conflict of interest.

Reference list

1. Mendis S, Puska P, Norrving B (2011) Global atlas on cardiovascular disease prevention and control. World Health Organization, Geneva
2. Longstreth W, Bernick C, Fitzpatrick A, Cushman M, Knepper L, Lima J, Furberg C (2001) Frequency and predictors of stroke death in 5,888 participants in the Cardiovascular Health Study. *Neurology* 56 (3):368-375
3. Barnett AG, De Looper M, Fraser JF (2008) The seasonality in heart failure deaths and total cardiovascular deaths. *Australian and New Zealand journal of public health* 32 (5):408-413
4. Hobbs DA, George TW, Lovegrove JA (2013) The effects of dietary nitrate on blood pressure and endothelial function: a review of human intervention studies. *Nutrition Research Reviews* 26 (2):210-222
5. Marti CN, Gheorghide M, Kalogeropoulos AP, Georgiopoulou VV, Quyyumi AA, Butler J (2012) Endothelial dysfunction, arterial stiffness, and heart failure. *Journal of the American College of Cardiology* 60 (16):1455-1469
6. Gkaliagkousi E, Douma S, Zamboulis C, Ferro A (2009) Nitric oxide dysfunction in vascular endothelium and platelets: role in essential hypertension. *J Hypertens* 27 (12):2310-2320
7. Weitzberg E, Lundberg JO (2013) Novel aspects of dietary nitrate and human health. *Annual Review of Nutrition* 33 (1):129-159. doi:10.1146/annurev-nutr-071812-161159
8. Hord NG, Tang Y, Bryan NS (2009) Food sources of nitrates and nitrites: the physiologic context for potential health benefits. *Am J Clin Nutr* 90 (1):1-10
9. Nuñez De González MT, Osburn WN, Hardin MD, Longnecker M, Garg HK, Bryan NS, Keeton JT (2012) Survey of residual nitrite and nitrate in conventional and organic/natural/uncured/indirectly cured meats available at retail in the United States. *J Agric Food Chem* 60 (15):3981-3990. doi:10.1021/jf204611k
10. Blekkenhorst LC, Bondonno NP, Liu AH, Ward NC, Prince RL, Lewis JR, Devine A, Croft KD, Hodgson JM, Bondonno CP (2018) Nitrate, the oral microbiome, and cardiovascular health: a systematic literature review of human and animal studies. *Am J Clin Nutr* 107 (4):504-522. doi:10.1093/ajcn/nqx046
11. Bryan NS, Alexander DD, Coughlin JR, Milkowski AL, Boffetta P (2012) Ingested nitrate and nitrite and stomach cancer risk: an updated review. *Food Chem Toxicol* 50 (10):3646-3665. doi:<https://doi.org/10.1016/j.fct.2012.07.062>
12. Lundberg JO, Weitzberg E, Shiva S, Gladwin MT (2011) The Nitrate–Nitrite–Nitric Oxide Pathway in Mammals. In: *Nitrite and Nitrate in Human Health and Disease*. Springer, pp 21-48
13. Larsen FJ, Ekblom B, Sahlin K, Lundberg JO, Weitzberg E (2006) Effects of dietary nitrate on blood pressure in healthy volunteers. *N Engl J Med* 355 (26):2792-2793
14. Sobko T, Marcus C, Govoni M, Kamiya S (2010) Dietary nitrate in Japanese traditional foods lowers diastolic blood pressure in healthy volunteers. *Nitric Oxide* 22 (2):136-140
15. Bondonno CP, Yang X, Croft KD, Considine MJ, Ward NC, Rich L, Puddey IB, Swinney E, Mubarak A, Hodgson JM (2012) Flavonoid-rich apples and nitrate-rich spinach augment nitric oxide status and improve endothelial function in healthy men and women: a randomized controlled trial. *Free Radical Bio Med* 52 (1):95-102
16. Liu AH, Bondonno CP, Croft KD, Puddey IB, Woodman RJ, Rich L, Ward NC, Vita JA, Hodgson JM (2013) Effects of a nitrate-rich meal on arterial stiffness and blood pressure in healthy volunteers. *Nitric Oxide* 35:123-130
17. Kapil V, Khambata RS, Robertson A, Caulfield MJ, Ahluwalia A (2015) Dietary nitrate provides sustained blood pressure lowering in hypertensive patients: a randomized, phase 2, double-blind, placebo-controlled study. *Hypertension* 65 (2):320-327
18. Rammos C, Hendgen-Cotta UB, Sobierajski J, Bernard A, Kelm M, Rassaf T (2014) Dietary nitrate reverses vascular dysfunction in older adults with moderately increased cardiovascular risk. *J Am Coll Cardiol* 63 (15):1584-1585
19. Webb AJ, Patel N, Loukogeorgakis S, Okorie M, Aboud Z, Misra S, Rashid R, Miall P, Deanfield J, Benjamin N, MacAllister R, Hobbs AJ, Ahluwalia A (2008) Acute blood pressure lowering, vasoprotective, and antiplatelet properties of dietary nitrate via bioconversion to nitrite. *Hypertension* 51 (3):784-790
20. Jackson JK, Patterson AJ, MacDonald-Wicks LK, Oldmeadow C, McEvoy MA (2018) The role of inorganic nitrate and nitrite in cardiovascular disease risk factors: a systematic review and meta-analysis of human evidence. *Nutr Rev* 76 (5):348-371. doi:10.1093/nutrit/nuy005
21. Nuñez de González MT, Osburn WN, Hardin MD, Longnecker M, Garg HK, Bryan NS, Keeton JT (2015) A survey of nitrate and nitrite concentrations in conventional and organic-labeled raw vegetables at retail. *J Food Sci* 80 (5):C942-C949. doi:10.1111/1750-3841.12858
22. Keys A (1995) Mediterranean diet and public health: personal reflections. *American Journal of Clinical Nutrition* 61 (6):1321S-1323S

23. Lundberg JO, Feelisch M, Björne H, Jansson EÅ, Weitzberg E (2006) Cardioprotective effects of vegetables: is nitrate the answer? *Nitric Oxide* 15 (4):359-362
24. Miller ER, Erlinger TP, Appel LJ (2006) The effects of macronutrients on blood pressure and lipids: an overview of the DASH and OmniHeart trials. *Current Atherosclerosis Reports* 8 (6):460-465
25. JSHIPURA KJ, ASCHERIO A, MANSON JE, STAMPFER MJ, RIMM EB, SPEIZER FE, HENNEKENS CH, SPIEGELMAN D, WILLETT WC (1999) Fruit and vegetable intake in relation to risk of ischemic stroke. *Jama* 282 (13):1233-1239
26. JSHIPURA KJ, HU FB, MANSON JE, STAMPFER MJ, RIMM EB, SPEIZER FE, COLDITZ G, ASCHERIO A, ROSNER B, SPIEGELMAN D (2001) The effect of fruit and vegetable intake on risk for coronary heart disease. *Annals of Internal Medicine* 134 (12):1106-1114
27. BLEKKENHORST LC, BONDONNO CP, LEWIS JR, DEVINE A, WOODMAN RJ, CROFT KD, LIM WH, WONG G, BEILIN LJ, PRINCE RL (2017) Association of dietary nitrate with atherosclerotic vascular disease mortality: a prospective cohort study of older adult women. *American Journal of Clinical Nutrition* 106 (1):207-216
28. BONDONNO CP, BLEKKENHORST LC, PRINCE RL, IVEY KL, LEWIS JR, DEVINE A, WOODMAN RJ, LUNDBERG JO, CROFT KD, THOMPSON PL (2017) Association of vegetable nitrate intake with carotid atherosclerosis and ischemic cerebrovascular disease in older women. *Stroke* 49 (6):1-6
29. MITCHELL P, WANG J, HOURIHAN F (1999) The relationship between glaucoma and pseudoexfoliation: The Blue Mountains Eye Study. *Arch Ophthalmol* 117 (10):1319-1324. doi:10.1001/archophth.117.10.1319
30. KAUSHIK S, WANG JJ, FLOOD V, LIEW G, SMITH W, MITCHELL P (2008) Frequency of fish consumption, retinal microvascular signs and vascular mortality. *Microcirculation* 15 (1):27-36
31. RUSSELL J, FLOOD V, ROCHTCHINA E, GOPINATH B, ALLMAN-FARINELLI M, BAUMAN A, MITCHELL P (2013) Adherence to dietary guidelines and 15-year risk of all-cause mortality. *British Journal of Nutrition* 109 (3):547-555
32. SMITH W, MITCHELL P, REAY EM, WEBB K, HARVEY PW (1998) Validity and reproducibility of a self-administered food frequency questionnaire in older people. *Australian and New Zealand journal of public health* 22 (4):456-463
33. MITCHELL P, SMITH W, ATTEBO K, WANG JJ (1995) Prevalence of age-related maculopathy in Australia: the Blue Mountains Eye Study. *Ophthalmology* 102 (10):1450-1460
34. LOUIE JCY, FLOOD VM, BURLUTSKY G, RANGAN AM, GILL TP, MITCHELL P (2013) Dairy consumption and the risk of 15-year cardiovascular disease mortality in a cohort of older Australians. *Nutrients* 5 (2):441-454
35. WILLETT WC, SAMPSON L, BROWNE ML, STAMPFER MJ, ROSNER B, HENNEKENS CH, SPEIZER FE (1988) The use of a self-administered questionnaire to assess diet four years in the past. *American Journal of Epidemiology* 127 (1):188-199
36. LEWIS J, ENGLISH R (1990) NUTTAB 90 - Nutrient Data Table for Use in Australia [floppy disc]. Department of Community Services and Health, Canberra
37. LEWIS J, MILLIGAN GC, HUNT A (1995) NUTTAB95 - Nutrient Data Table for Use in Australia [compact disc]. Australia New Zealand Food Authority, Canberra
38. BLEKKENHORST LC, PRINCE RL, WARD NC, CROFT KD, LEWIS JR, DEVINE A, SHINDE S, WOODMAN RJ, HODGSON JM, BONDONNO CP (2017) Development of a reference database for assessing dietary nitrate in vegetables. *Molecular Nutrition & Food Research* 61 (8):1-13
39. INOUE-CHOI M, VIRK-BAKER MK, ASCHBROOK-KILFOY B, CROSS AJ, SUBAR AF, THOMPSON FE, SINHA R, WARD MH (2016) Development and calibration of a dietary nitrate and nitrite database in the NIH-AARP Diet and Health Study. *Public Health Nutr* 19 (11):1934-1943
40. FOOD STANDARDS AUSTRALIA NEW ZEALAND (2011) Survey of nitrates and nitrites in food and beverages in Australia. FSANZ, Canberra
41. GRIESEBECK JS, STECK MD, HUBER JC, SHARKEY JR, RENE AA, BRENDER JD (2009) Development of estimates of dietary nitrates, nitrites, and nitrosamines for use with the Short Willet Food Frequency Questionnaire. *Nutrition Journal* 8 (1):16
42. GOPINATH B, LIEW G, BURLUTSKY G, MITCHELL P (2014) Physical activity and the 15-year incidence of age-related macular degeneration. *Invest Ophthalmol Vis Sci* 55 (12):7799-7803. doi:10.1167/iovs.14-15575
43. STRIPPOLI GF, CRAIG JC, ROCHTCHINA E, FLOOD VM, WANG JJ, MITCHELL P (2011) Fluid and nutrient intake and risk of chronic kidney disease. *Nephrology* 16 (3):326-334
44. MAGLIANO D, LIEW D, PATER H, KIRBY A, HUNT D, SIMES J, SUNDARARAJAN V, TONKIN A (2003) Accuracy of the Australian National Death Index: comparison with adjudicated fatal outcomes among Australian participants in the Long-term Intervention with Pravastatin in Ischaemic Disease (LIPID) study. *Australian and New Zealand journal of public health* 27 (6):649-653
45. GOPINATH B, LIEW G, KIFLEY A, FLOOD VM, JOACHIM N, LEWIS JR, HODGSON JM, MITCHELL P (2018) Dietary flavonoids and the prevalence and 15-y incidence of age-related macular degeneration. *Am J Clin Nutr* 108 (2):381-387

46. Webb AJ, Patel N, Loukogeorgakis S, Okorie M, Aboud Z, Misra S, Rashid R, Miall P, Deanfield J, Benjamin N (2008) Acute blood pressure lowering, vasoprotective, and antiplatelet properties of dietary nitrate via bioconversion to nitrite. *Hypertension* 51 (3):784-790
47. Gilchrist M, Winyard PG, Aizawa K, Anning C, Shore A, Benjamin N (2013) Effect of dietary nitrate on blood pressure, endothelial function, and insulin sensitivity in type 2 diabetes. *Free Radical Bio Med* 60:89-97. doi:10.1016/j.freeradbiomed.2013.01.024
48. Bondonno CP, Liu AH, Croft KD, Ward NC, Yang X, Considine MJ, Puddey IB, Woodman RJ, Hodgson JM (2014) Short-term effects of nitrate-rich green leafy vegetables on blood pressure and arterial stiffness in individuals with high-normal blood pressure. *Free Radic Biol Med* 77:353-362
49. Bondonno CP, Liu AH, Croft KD, Ward NC, Shinde S, Moodley Y, Lundberg JO, Puddey IB, Woodman RJ, Hodgson JM (2015) Absence of an effect of high nitrate intake from beetroot juice on blood pressure in treated hypertensive individuals: a randomized controlled trial. *American Journal of Clinical Nutrition* 102 (2):368-375
50. Kapil V, Khambata RS, Robertson A, Caulfield MJ, Ahluwalia A (2015) Dietary Nitrate Provides Sustained Blood Pressure Lowering in Hypertensive Patients Novelty and Significance. *Hypertension* 65 (2):320-327
51. Siervo M, Lara J, Ogbonmwan I, Mathers JC (2013) Inorganic nitrate and beetroot juice supplementation reduces blood pressure in adults: a systematic review and meta-analysis. *J Nutr* 143 (6):818-826
52. Ashor AW, Lara J, Siervo M (2017) Medium-term effects of dietary nitrate supplementation on systolic and diastolic blood pressure in adults: a systematic review and meta-analysis. *Journal of Hypertension* 35 (7):1353-1359
53. Lara J, Ashor AW, Oggioni C, Ahluwalia A, Mathers JC, Siervo M (2016) Effects of inorganic nitrate and beetroot supplementation on endothelial function: a systematic review and meta-analysis. *European Journal of Nutrition* 55 (2):451-459
54. Bahadoran Z, Mirmiran P, Ghasemi A, Carlström M, Azizi F, Hadaegh F (2016) Association between dietary intakes of nitrate and nitrite and the risk of hypertension and chronic kidney disease: Tehran Lipid and Glucose Study. *Nutrients* 8 (12):811
55. Maas R, Xanthakis V, Göen T, Müller J, Schwedhelm E, Böger RH, Vasan RS (2017) Plasma nitrate and incidence of cardiovascular disease and all-cause mortality in the community: the Framingham Offspring Study. *Journal of the American Heart Association* 6 (11). doi:10.1161/jaha.117.006224
56. Smallwood MJ, Ble A, Melzer D, Winyard PG, Benjamin N, Shore AC, Gilchrist M (2017) Relationship between urinary nitrate excretion and blood pressure in the InChianti cohort. *Am J Hypertens* 30 (7):707-712. doi:10.1093/ajh/hpx035
57. Lawes CM, Bennett DA, Feigin VL, Rodgers A (2004) Blood pressure and stroke. *Stroke* 35 (3):776-785
58. Banegas JR, Ruilope LM, de la Sierra A, Vinyoles E, Gorostidi M, de la Cruz JJ, Ruiz-Hurtado G, Segura J, Rodríguez-Artalejo F, Williams B (2018) Relationship between clinic and ambulatory blood-pressure measurements and mortality. *N Engl J Med* 378 (16):1509-1520. doi:10.1056/NEJMoal1712231
59. Sharman JE, Howes FS, Head GA, McGrath BP, Stowasser M, Schlaich M, Glasziou P, Nelson MR (2015) Home blood pressure monitoring: Australian Expert Consensus Statement. *J Hypertens* 33 (9):1721
60. Tamme T, Reinik M, Roasto M, Juhkam K, Tenno T, Kiis A (2006) Nitrates and nitrites in vegetables and vegetable-based products and their intakes by the Estonian population. *Food Additives and Contaminants* 23 (4):355-361
61. Siddiqi M, Kumar R, Fazili Z, Spiegelhalter B, Preussmann R (1992) Increased exposure to dietary amines and nitrate in a population at high risk of oesophageal and gastric cancer in Kashmir (India). *Carcinogenesis* 13 (8):1331-1335
62. Dich J, Järvinen R, Knekt P, Penttilä PL (1996) Dietary intakes of nitrate, nitrite and NDMA in the Finnish mobile clinic health examination survey. *Food Additives & Contaminants* 13 (5):541-552
63. Larsen FJ, Schiffer TA, Borniquel S, Sahlin K, Ekblom B, Lundberg JO, Weitzberg E (2011) Dietary inorganic nitrate improves mitochondrial efficiency in humans. *Cell Metabolism* 13 (2):149-159. doi:<https://doi.org/10.1016/j.cmet.2011.01.004>
64. Carlström M, Larsen FJ, Nyström T, Hezel M, Borniquel S, Weitzberg E, Lundberg JO (2010) Dietary inorganic nitrate reverses features of metabolic syndrome in endothelial nitric oxide synthase-deficient mice. *Proceedings of the National Academy of Sciences* 107 (41):17716-17720. doi:10.1073/pnas.1008872107
65. Yang T, Zhang X-M, Tarnawski L, Peleli M, Zhuge Z, Terrando N, Harris RA, Olofsson PS, Larsson E, Persson AEG, Lundberg JO, Weitzberg E, Carlstrom M (2017) Dietary nitrate attenuates renal ischemia-reperfusion injuries by modulation of immune responses and reduction of oxidative stress. *Redox Biology* 13:320-330. doi:<https://doi.org/10.1016/j.redox.2017.06.002>

66. Wightman EL, Haskell-Ramsay CF, Thompson KG, Blackwell JR, Winyard PG, Forster J, Jones AM, Kennedy DO (2015) Dietary nitrate modulates cerebral blood flow parameters and cognitive performance in humans: a double-blind, placebo-controlled, crossover investigation. *Physiology & Behavior* 149:149-158. doi:<https://doi.org/10.1016/j.physbeh.2015.05.035>
67. Bondonno CP, Croft KD, Hodgson JM (2016) Dietary nitrate, nitric oxide, and cardiovascular health. *Critical Reviews in Food Science and Nutrition* 56 (12):2036-2052
68. Hill BG, Dranka BP, Bailey S, Lancaster J, Darley-Usmar V (2010) What part of NO don't you understand? Some answers to the cardinal questions in nitric oxide biology. *Journal of Biological Chemistry*. doi:10.1074/jbc.R110.101618
69. Brecht DS (1999) Endogenous nitric oxide synthesis: biological functions and pathophysiology. *Free Radical Research* 31 (6):577-596
70. Aune D, Giovannucci E, Boffetta P, Fadnes LT, Keum N, Norat T, Greenwood DC, Riboli E, Vatten LJ, Tonstad S (2017) Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality—a systematic review and dose-response meta-analysis of prospective studies. *International Journal of Epidemiology* 46 (3):1029-1056. doi:10.1093/ije/dyw319
71. Threapleton DE, Greenwood DC, Evans CEL, Cleghorn CL, Nykjaer C, Woodhead C, Cade JE, Gale CP, Burley VJ (2013) Dietary fibre intake and risk of cardiovascular disease: systematic review and meta-analysis. *British Medical Journal* 347:1-12. doi:10.1136/bmj.f6879
72. Aburto NJ, Hanson S, Gutierrez H, Hooper L, Elliott P, Cappuccio FP (2013) Effect of increased potassium intake on cardiovascular risk factors and disease: systematic review and meta-analyses. *British Medical Journal* 346. doi:10.1136/bmj.f1378
73. D'Elia L, Barba G, Cappuccio FP, Strazzullo P (2011) Potassium intake, stroke, and cardiovascular disease: a meta-analysis of prospective studies. *J Am Coll Cardiol* 57 (10):1210-1219

Table 1. Baseline characteristics of study participants by categories of vegetable nitrate intake¹

	All participants n=2229	Quartile 1 (<69.5 mg/d) n=557	Quartile 2 (69.5-99.6 mg/d) n=558	Quartile 3 (99.7-137.8 mg/d) n=557	Quartile 4 (>137.8 mg/d) n=557	P-value ²
Participants demographics						
Gender, male	915 (41.0)	268 (48.1)	244 (43.7)	217 (39.0)	186 (33.4)	<0.001
Age, year	64.5 ± 9.1	65.1 ± 9.3	64.4 ± 8.7	63.5 ± 9.0	65.1 ± 9.4	0.010
BMI ³ , kg/m ²	26.0 ± 4.4	26.1 ± 4.5	26.0 ± 4.4	26.4 ± 4.3	25.7 ± 4.2	0.122
SBP ⁴ , mmHg	145.4 ± 20.8	145.6 ± 20.7	145.8 ± 20.3	145.4 ± 20.4	144.9 ± 21.7	0.890
DBP ⁴ , mmHg	83.4 ± 9.8	84.0 ± 10.3	83.3 ± 9.4	83.5 ± 9.5	83.0 ± 9.9	0.319
Physical activity ⁵						<0.001
Not active	476 (21.4)	149 (27.5)	137 (25.1)	104 (18.8)	86 (15.7)	
Light exercise	973 (43.7)	247 (45.6)	231 (42.3)	255 (46.2)	240 (43.8)	
Vigorous exercise	739 (33.2)	146 (26.9)	178 (32.6)	193 (35.0)	222 (40.5)	
Alcohol intake, g/d	2.8 (0.2-14.4)	2.5 (0.2-17.0)	4.5 (0.3-16.8)	3.6 (0.4-15.7)	1.9 (0.2-11.5)	0.058
Smoking history ⁶						0.035
Never smoked	1079 (48.4)	243 (45.3)	262 (48.0)	282 (51.2)	292 (53.3)	
Previous smoker	790 (35.4)	201 (37.5)	196 (35.9)	197 (35.8)	196 (35.8)	
Current smoker	312 (14.0)	92 (17.2)	88 (16.1)	72 (13.1)	60 (10.9)	
Socio-economic status ⁷						
Home/unit owner	1982 (88.9)	481 (88.4)	504 (91.5)	504 (91.5)	493 (90.0)	0.261
Medication use						
Organic nitrate	26 (1.2)	7 (1.3)	6 (1.1)	7 (1.3)	6 (1.1)	0.984

Antihypertensive	632 (28.4)	166 (29.8)	170 (30.5)	154 (27.6)	142 (25.5)	0.243
Statins	66 (3.0)	18 (3.2)	20 (3.6)	15 (2.7)	13 (2.3)	0.615
Low-dose aspirin ⁸						0.766
Less than once per month	1532 (68.7)	374 (73.8)	376 (75.5)	389 (75.8)	392 (77.2)	
More than once per month but less than once per week	124 (5.6)	38 (7.5)	27 (5.4)	32 (6.2)	27 (5.3)	
Once per week or more	371 (16.6)	95 (18.7)	95 (19.1)	92 (17.9)	89 (17.5)	
Biochemical analyses						
MDRD eGFR ⁹ , mL/min per 1.73 m ²	79.1 ± 18.7	78.2 ± 22.2	78.8 ± 17.7	79.3 ± 17.1	80.2 ± 17.1	0.336
Total cholesterol ¹⁰ , mmol/L	6.1 ± 1.1	6.0 ± 1.1	6.0 ± 1.1	6.1 ± 1.0	6.1 ± 1.1	0.583
Fasting glucose ⁹ , mmol/L	4.9 ± 0.6	5.0 ± 0.6	5.0 ± 0.6	5.0 ± 0.6	4.9 ± 0.5	0.135

¹ Vegetable nitrate intake was calculated using the mean intake over the 14 years of follow-up. Values are means ± SDs, medians (IQRs), or n (%). SBP, systolic blood pressure; DBP, diastolic blood pressure; MDRD, Modification of Diet in Renal Disease study equation; eGFR, estimated glomerular filtration rate.

² P-values are a comparison between categories by ANOVA, Kruskal-Wallis test, or chi-square test where appropriate.

³ Measured in 2209 participants.

⁴ Measured in 2225 participants.

⁵ Measured in 2188 participants.

⁶ Measured in 2181 participants.

⁷ Measured in 2194 participants.

⁸ Measured in 2026 participants.

⁹ Measured in 2155 participants.

¹⁰ Measured in 2156 participants.

Table 2. Baseline dietary intakes of all study participants by categories of vegetable nitrate intake¹

	All participants n=2229	Quartile 1 (<69.5 mg/d) n=557	Quartile 2 (69.5-99.6 mg/d) n=558	Quartile 3 (99.7-137.8 mg/d) n=557	Quartile 4 (>137.8 mg/d) n=557	P-value ²
Total vegetable nitrate, mg/d	109.7 ± 59.2	48.3 ± 15.2	85.0 ± 8.8	117.3 ± 10.9	188.7 ± 53.8	<0.001
Total nitrate, mg/d	128.9 ± 62.0	63.9 ± 17.4	103.5 ± 11.1	137.0 ± 12.5	211.1 ± 55.8	<0.001
Baseline dietary intakes						
Energy intake, kcal	2063.9 ± 599.1	1860.7 ± 600.3	2043.8 ± 561.9	2118.5 ± 576.0	2232.8 ± 597.1	<0.001
Total sugar, g/d	124.1 ± 49.8	109.1 ± 48.3	121.0 ± 46.5	126.2 ± 47.6	140.1 ± 51.7	<0.001
Fibre, g/d	27.3 ± 10.4	20.8 ± 8.9	25.5 ± 8.5	28.6 ± 8.7	34.3 ± 10.6	<0.001
Protein, g/d	87.1 ± 27.0	75.1 ± 25.8	86.0 ± 26.0	90.4 ± 23.9	96.9 ± 27.6	<0.001
Fat, g/d	75.1 ± 27.7	69.5 ± 28.1	74.9 ± 26.8	77.0 ± 26.2	79.0 ± 28.8	<0.001
Carbohydrate, g/d	232.7 ± 75.1	206.5 ± 73.3	226.9 ± 68.7	237.6 ± 71.9	259.7 ± 76.5	<0.001
Sodium, mg/d	2034.8 ± 712.4	1847.0 ± 712.1	2005.5 ± 674.1	2090.9 ± 646.9	2196.0 ± 755.6	<0.001
Potassium, mg/d	3721.1 ± 1161.9	2960.5 ± 971.1	3561.5 ± 926.9	3868.8 ± 973.7	4493.9 ± 1197.0	<0.001
Calcium, mg/d	880.8 ± 405.9	767.6 ± 403.7	851.0 ± 394.2	900.9 ± 386.5	1003.5 ± 403.6	<0.001
Phosphorous, mg/d	1513.2 ± 503.8	1304.3 ± 497.5	1478.2 ± 473.3	1569.5 ± 460.8	1700.8 ± 499.6	<0.001
Magnesium, mg/d	336.2 ± 106.1	280.5 ± 97.7	324.9 ± 92.9	349.1 ± 96.8	390.4 ± 105.9	<0.001
Zinc, mg/d	11.5 ± 3.7	9.9 ± 3.6	11.4 ± 3.7	12.0 ± 3.2	12.9 ± 3.8	<0.001
Vitamin C, mg/d	182.4 ± 94.2	131.9 ± 81.3	169.6 ± 78.7	193.3 ± 81.7	234.8 ± 102.3	<0.001
Cholesterol, mg/d	294.0 ± 136.9	268.8 ± 134.1	294.1 ± 128.0	303.3 ± 141.3	309.8 ± 140.7	<0.001
Saturated fat, g/d	29.5 ± 13.0	28.2 ± 13.5	29.7 ± 12.7	30.1 ± 12.7	29.9 ± 13.0	0.073
Monounsaturated fat, g/d	26.6 ± 10.1	24.6 ± 10.1	26.5 ± 9.7	27.3 ± 9.4	28.0 ± 10.8	<0.001

Polyunsaturated fat, g/d	12.5 ± 5.4	11.0 ± 5.2	12.3 ± 5.3	12.9 ± 5.0	13.9 ± 5.7	<0.001
--------------------------	------------	------------	------------	------------	------------	--------

¹Total and vegetable nitrate intakes are the mean calculated using baseline, 5-year, and 10-year dietary data. Values are means ± SDs.

²P-values are a comparison between categories by ANOVA.

Table 3. Cardiovascular disease, coronary heart disease, stroke, non-cardiovascular disease and all-cause mortality for all participants and by quartiles of vegetable nitrate intake¹

	Quartile 1 (<69.5 mg/d) n=557	Quartile 2 (69.5-99.6 mg/d) n=558	Quartile 3 (99.7-137.8 mg/d) n=557	Quartile 4 (>137.8 mg/d) n=557	P-value ²
Cardiovascular disease mortality					
Death, n (%)	61 (11.0)	39 (7.0)	35 (6.3)	53 (9.5)	
Unadjusted	1.00 (Referent)	0.59 (0.40, 0.88)	0.53 (0.35, 0.80)	0.79 (0.55, 1.15)	0.008
Age, gender and energy adjusted	1.00 (Referent)	0.60 (0.40, 0.90)	0.56 (0.37, 0.86)	0.71 (0.48, 1.05)	0.023
Multivariable adjusted ³	1.00 (Referent)	0.53 (0.35, 0.82)	0.51 (0.32, 0.80)	0.63 (0.41, 0.95)	0.006
Coronary heart disease mortality					
Death, n (%)	37 (6.6)	22 (3.9)	30 (5.4)	36 (6.5)	
Unadjusted	1.00 (Referent)	0.55 (0.33, 0.94)	0.75 (0.46, 1.21)	0.89 (0.57, 1.42)	0.143
Age, gender and energy adjusted	1.00 (Referent)	0.56 (0.33, 0.95)	0.78 (0.48, 1.28)	0.80 (0.49, 1.29)	0.201
Multivariable adjusted	1.00 (Referent)	0.46 (0.26, 0.82)	0.73 (0.43, 1.23)	0.68 (0.40, 1.15)	0.066
Stroke mortality					
Death, n (%)	24 (4.3)	17 (3.0)	5 (0.9)	17 (3.1)	
Unadjusted	1.00 (Referent)	0.65 (0.35, 1.21)	0.19 (0.07, 0.50)	0.64 (0.34, 1.19)	0.008
Age, gender and energy adjusted	1.00 (Referent)	0.67 (0.36, 1.26)	0.21 (0.08, 0.56)	0.58 (0.30, 1.12)	0.016
Multivariable adjusted	1.00 (Referent)	0.66 (0.34, 1.26)	0.18 (0.06, 0.52)	0.56 (0.28, 1.11)	0.014
Non-cardiovascular disease mortality					
Death, n (%)	134 (24.1)	105 (18.8)	104 (18.7)	79 (14.2)	
Unadjusted	1.00 (Referent)	0.73 (0.56, 0.94)	0.71 (0.55, 0.92)	0.54 (0.41, 0.72)	<0.001

Age, gender and energy adjusted	1.00 (Referent)	0.77 (0.59, 0.99)	0.81 (0.62, 1.05)	0.56 (0.42, 0.74)	0.001
Multivariable adjusted	1.00 (Referent)	0.89 (0.67, 1.17)	1.00 (0.75, 1.33)	0.65 (0.47, 0.90)	0.034
All-cause mortality					
Death, n (%)	195 (35.0)	144 (25.8)	139 (25.0)	132 (23.7)	
Unadjusted	1.00 (Referent)	0.69 (0.55, 0.85)	0.66 (0.53, 0.81)	0.62 (0.50, 0.78)	<0.001
Age, gender and energy adjusted	1.00 (Referent)	0.72 (0.58, 0.89)	0.73 (0.59, 0.91)	0.61 (0.49, 0.77)	<0.001
Multivariable adjusted	1.00 (Referent)	0.76 (0.60, 0.96)	0.82 (0.65, 1.04)	0.65 (0.51, 0.84)	0.009

¹ Results are presented as HRs (95% CIs).

² P-values are for a comparison between quartiles using the Wald test with Cox PH regression.

³ Multivariable-adjusted models included age, gender, BMI, physical activity, alcohol intake, smoking history, socio-economic status, use of organic nitrate medications, use of antihypertensive medications, use of statin medications, low-dose aspirin use, renal function (MDRD eGFR) and energy intake.

Table 4. Cardiovascular disease, coronary heart disease, stroke, non-cardiovascular disease and all-cause mortality for all participants and by categories of vegetable nitrate intake¹

	Low nitrate (< 52.7 mg/d) n=298	Moderate nitrate (52.7-76.4 mg/d) n=378	High nitrate (> 76.4 mg/d) n=1553	P-value ²
Cardiovascular disease mortality				
Death, n (%)	37 (12.4)	29 (7.7)	122 (7.9)	
Unadjusted	1.00 (referent)	0.58 (0.36, 0.94)	0.57 (0.39, 0.82)	0.009
Age, gender and energy adjusted	1.00 (referent)	0.60 (0.37, 0.99)	0.57 (0.39, 0.84)	0.015
Multivariable adjusted ³	1.00 (referent)	0.50 (0.30, 0.84)	0.45 (0.30, 0.69)	0.001
Coronary heart disease mortality				
Death, n (%)	24 (8.1)	16 (4.2)	85 (5.5)	
Unadjusted	1.00 (referent)	0.50 (0.26, 0.94)	0.62 (0.39, 0.97)	0.053
Age, gender and energy adjusted	1.00 (referent)	0.52 (0.27, 0.98)	0.61 (0.38, 0.97)	0.068
Multivariable adjusted	1.00 (referent)	0.47 (0.24, 0.90)	0.49 (0.29, 0.82)	0.017
Stroke mortality				
Death, n (%)	13 (4.4)	13 (3.4)	37 (2.4)	
Unadjusted	1.00 (referent)	0.73 (0.34, 1.58)	0.48 (0.26, 0.91)	0.061
Age, gender and energy adjusted	1.00 (referent)	0.76 (0.35, 1.65)	0.50 (0.26, 0.97)	0.098
Multivariable adjusted	1.00 (referent)	0.56 (0.24, 1.30)	0.39 (0.19, 0.81)	0.036
Non-cardiovascular disease mortality				
Death, n (%)	79 (26.5)	79 (20.8)	264 (17.0)	
Unadjusted	1.00 (referent)	0.75 (0.55, 1.02)	0.58 (0.45, 0.75)	<0.001

Age, gender and energy adjusted	1.00 (referent)	0.81 (0.59, 1.11)	0.65 (0.50, 0.84)	0.004
Multivariable adjusted	1.00 (referent)	0.75 (0.53, 1.08)	0.73 (0.55, 0.98)	0.111
All-cause mortality				
Death, n (%)	116 (38.9)	108 (28.6)	386 (24.9)	
Unadjusted	1.00 (referent)	0.69 (0.53, 0.90)	0.58 (0.47, 0.71)	<0.001
Age, gender and energy adjusted	1.00 (referent)	0.74 (0.57, 0.97)	0.62 (0.50, 0.77)	<0.001
Multivariable adjusted	1.00 (referent)	0.66 (0.50, 0.89)	0.63 (0.50, 0.80)	0.001

¹ Cut-off values for vegetable nitrate intake categories were determined from previously published data [27,28]. Results are presented as HRs (95% CIs).

² P-values are for a comparison between categories using the Wald test with Cox PH regression.

³ Multivariable-adjusted models included age, gender, BMI, physical activity, alcohol intake, smoking history, socio-economic status, use of organic nitrate medications, use of antihypertensive medications, use of statin medications, low-dose aspirin use, renal function (MDRD eGFR) and energy intake.

Figure legends

Figure 1. Participant flow chart. CVD, cardiovascular disease. DM, diabetes melletus. MI, myocardial infarction.

Figure 2. Multivariable-adjusted cumulative survival curves for cardiovascular disease mortality by quartiles of vegetable nitrate intake. Multivariable adjusted Cox PH model included age, gender, BMI, physical activity, alcohol intake, smoking history, socio-economic status, use of organic nitrate medications, use of antihypertensive medications, use of statin medications, low-dose aspirin use, renal function (MDRD eGFR) and energy intake. CVD, cardiovascular disease.