

January 2020

Does sleep grow on trees? A longitudinal study to investigate potential prevention of insufficient sleep with different types of urban green space

Thomas E. Astell-Burt
University of Wollongong, thomasab@uow.edu.au

Xiaoqi Feng
University of Wollongong, xfeng@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/sspapers>

Recommended Citation

Astell-Burt, Thomas E. and Feng, Xiaoqi, "Does sleep grow on trees? A longitudinal study to investigate potential prevention of insufficient sleep with different types of urban green space" (2020). *Faculty of Social Sciences - Papers*. 4651.
<https://ro.uow.edu.au/sspapers/4651>


Does sleep grow on trees? A longitudinal study to investigate potential prevention of insufficient sleep with different types of urban green space

Abstract

2019 Introduction: To investigate association between urban green space and prevalent and incident cases of insufficient sleep (<6 h sleep per>day). Methods: This longitudinal study examined the odds of prevalent and incident insufficient sleep in relation to indicators of total green space, tree canopy, open grass and other low-lying vegetation in the Sax Institute's 45 and Up Study (baseline 2006-2009; follow-up 2012-2015). Association between green space within 1.6 km road distances and insufficient sleep among 38,982 participants living in Sydney, Wollongong or Newcastle were analysed using multilevel logistic regressions adjusted for confounding. Results: Participants with more total green space had lower odds of prevalent insufficient sleep (e.g. $\geq 30\%$ compared with 0-4% total green space odds ratio (OR) = 0.68, 95% credible interval (95%CI) = 0.53, 0.85). The odds of prevalent insufficient sleep were lower among participants with more tree canopy (e.g. $\geq 30\%$ compared with 0-9% tree canopy OR = 0.78, 95%CI 0.69, 0.88). The odds of incident insufficient sleep were also lower with more tree canopy (e.g. $\geq 30\%$ compared with 0-9% tree canopy OR = 0.87, 95%CI = 0.75, 0.99). There were no statistically significant associations between prevalent or incident insufficient sleep with open grass or other low-lying vegetation, nor incident sufficient sleep with total green space. Conclusions: Prioritising restoration and protection of urban tree canopy may help to promote population-wide prevention of insufficient sleep in middle-to-older aged adults.

Publication Details

Astell-Burt, T. & Feng, X. (2020). Does sleep grow on trees? A longitudinal study to investigate potential prevention of insufficient sleep with different types of urban green space. *SSM - Population Health*, 10



Article

Does sleep grow on trees? A longitudinal study to investigate potential prevention of insufficient sleep with different types of urban green space

Thomas Astell-Burt^{a,c,d,e,*}, Xiaoqi Feng^{a,b,c,e}

^a Population Wellbeing and Environment Research Lab (PowerLab), School of Health and Society, Faculty of Social Sciences, University of Wollongong, Wollongong, NSW, 2522, Australia

^b School of Public Health and Community Medicine, Faculty of Medicine, University of New South Wales, Sydney, NSW, 2052, Australia

^c Menzies Centre for Health Policy, University of Sydney, Sydney, New South Wales Australia

^d School of Public Health, Peking Union Medical College and The Chinese Academy of Medical Sciences, Beijing, China

^e National Institute of Environmental Health, Chinese Center for Disease Control and Prevention, Beijing, China

ARTICLE INFO

Keywords:

Urban green space
Tree canopy
Open grass
Sleep duration
Longitudinal study
Australia
45 and Up Study

ABSTRACT

Introduction: To investigate association between urban green space and prevalent and incident cases of insufficient sleep (<6 h sleep per day).

Methods: This longitudinal study examined the odds of prevalent and incident insufficient sleep in relation to indicators of total green space, tree canopy, open grass and other low-lying vegetation in the Sax Institute's 45 and Up Study (baseline 2006–2009; follow-up 2012–2015). Association between green space within 1.6 km road distances and insufficient sleep among 38,982 participants living in Sydney, Wollongong or Newcastle were analysed using multilevel logistic regressions adjusted for confounding.

Results: Participants with more total green space had lower odds of prevalent insufficient sleep (e.g. $\geq 30\%$ compared with 0–4% total green space odds ratio (OR) = 0.68, 95% credible interval (95%CI) = 0.53, 0.85). The odds of prevalent insufficient sleep were lower among participants with more tree canopy (e.g. $\geq 30\%$ compared with 0–9% tree canopy OR = 0.78, 95%CI 0.69, 0.88). The odds of incident insufficient sleep were also lower with more tree canopy (e.g. $\geq 30\%$ compared with 0–9% tree canopy OR = 0.87, 95%CI = 0.75, 0.99). There were no statistically significant associations between prevalent or incident insufficient sleep with open grass or other low-lying vegetation, nor incident sufficient sleep with total green space.

Conclusions: Prioritising restoration and protection of urban tree canopy may help to promote population-wide prevention of insufficient sleep in middle-to-older aged adults.

Introduction

Insufficient sleep typically defined in the scientific literature as less than 6 h per day is a major public health problem affecting non-trivial percentages of the population (e.g. approximately 12% in Australia according to Adams et al., 2017). Insufficient sleep is associated with elevated risks of workplace injuries (Uehli et al., 2014), road traffic accidents (Connor et al., 2001), noncommunicable diseases (NCDs) (Cappuccio et al., 2010a, 2011), weight gain (Cappuccio et al., 2008; Patel & Hu, 2008) and premature mortality (Cappuccio et al., 2010b). Some of these consequences may be mediated by the effects of insufficient sleep on the immune system (Lange, Dimitrov, & Born, 2010) and glucose metabolism (Spiegel et al., 2009). Person-level characteristics

like older age, lower educational attainment, lower income and unemployment are associated with insufficient sleep (Magee, Iverson, & Caputi, 2009), with the negative role of shift work a major concern in particular (Kecklund & Axelsson, 2016). Studies have also reported associations between sleep duration and the characteristics of neighbourhoods, including air pollution, light, and noise (Hale et al., 2019, pp. 77–84; Muzet, 2007; Sygna et al., 2014). In the search for scalable public health interventions for prevention of insufficient sleep, interest in and research into potential environmental determinants of sleep duration is rising (Hunter & Hayden, 2018; Johnson, Billings, & Hale, 2018).

One modifiable environmental exposure that holds potential for prevention of insufficient sleep is green space. Green space, which

* Corresponding author. Population Wellbeing and Environment Research Lab (PowerLab), School of Health and Society, Faculty of Social Sciences, University of Wollongong, Wollongong, NSW, 2522, Australia.

E-mail addresses: thomasab@uow.edu.au (T. Astell-Burt), xiaoqi.feng@unsw.edu.au (X. Feng).

<https://doi.org/10.1016/j.ssmph.2019.100497>

Received 10 June 2019; Received in revised form 4 October 2019; Accepted 6 October 2019

Available online 7 October 2019

2352-8273/© 2019 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

includes street trees, gardens, grass reserves, woodlands, forests and parks (Frumkin et al., 2017), may influence sleep duration through three domain pathways: (i) mitigation; (ii) restoration; and (iii) instoration (Markevych et al., 2017). On mitigation, studies report proximity to green space may help with abatement of air pollution (Abhijith et al., 2017) and noise annoyance (Dzhambov & Dimitrova, 2015). Evidence from sleep research suggests proximity to green space may help improve air quality and provide quieter environments that are likely to be an important, though not a sufficient, condition for optimal sleep duration (Halperin, 2014; Zanobetti et al., 2010). On restoration, many field experiments and epidemiological studies across several decades have reported contact with green space can lead to reduction of stress, restoration of cognitive capacities and protection of mental health (Hartig et al., 2014). Sleep studies point to the likely bi-directional relationship between insufficient sleep and mental ill-health (Alvaro, Roberts, & Harris, 2013). Finally, green spaces may help to build capacities (instoration) by providing preferential settings that motivate engagement in different types of outdoor physical recreation (Kredlow et al., 2015) and by supplying spaces in which communities come together to foster greater cohesion (Jennings & Bamkole, 2019) and reduce the prevalence of loneliness (e.g. Maas et al., 2009). Numerous studies suggest greater physical activity can support healthier sleep duration (Kredlow et al., 2015). Some research has reported loneliness may lower sleep quality, potentially via a stress pathway (McHugh & Lawlor, 2013), though this association may also be bi-directional (Simon & Walker, 2018).

While these interconnected pathways linking green space provision and sleep duration are plausible, there has been surprisingly little quantitative research so far. Some cross-sectional studies have assessed for potential effects of green space on the odds of insufficient sleep (Astell-Burt, Feng, & Kolt, 2013; Chum, O'Campo, & Matheson, 2015; Grigsby-Toussaint et al., 2015; Johnson et al., 2018b). Three of these studies (Astell-Burt et al., 2013; Grigsby-Toussaint et al., 2015; Johnson et al., 2018b) reported supportive evidence whereas the fourth found no convincing evidence of association (Chum et al., 2015). No longitudinal studies have been conducted. Among the three cross-sectional studies reporting affirmative findings, one found support specifically for benefits from tree canopy (Johnson et al., 2018b). Consideration of different types of green space provision for prevention of insufficient sleep is under-researched. Yet this is a highly policy-relevant issue, as urban and landscape planners seek evidence on which types of green space offer the strongest overall health gains and health economists aim to determine which types of green space are the most cost-effective to invest in for prevention of NCDs.

Accordingly, the purpose of our study was to advance the emerging scientific evidence linking green space and sleep duration in two important ways. First, we investigated the potential benefits of total green space provision on the odds of prevalent and incident cases of insufficient sleep. Second, we explored which types of green space may be key to these associations (if any).

Method

Four samples of participants were extracted from the Sax Institute's 45 and Up Study (45 and Up Study Collaborators, 2008) with intent to analyse odds of prevalent and incident insufficient sleep in January 2019. Participants ($n = 267,153$) in the 45 and Up Study lived in New South Wales (NSW) and had been sampled randomly from the Department of Human Services (formerly Medicare Australia) enrolment database, which provides near complete coverage of the population, and filled in postal surveys at baseline (2006–2009) and follow-up (2012–2015). The baseline response rate was approximately 18%, which is low but relative risks from the 45 and Up Study have been shown to be comparable to those drawn from a representative population health survey (Mealing et al., 2010). Participants provided consent for follow-up, of which 147,000 responded. Participants aged ≥ 80 were

oversampled. The sample was restricted to people living in the cities of Sydney, Wollongong or Newcastle at baseline ($n = 110,223$). This was further restricted to 38,982 participants that completed the follow-up survey and remained in the same local communities over time, as defined by Statistical Area 2s units containing 10,000 residents on average each.

Sleep was measured at baseline and follow-up with the question: "About how many hours in each 24 h day do you usually spend sleeping (including at night and naps)?" Insufficient sleep was defined as < 6 h sleep. The reference group was 6–10 h sleep using complete-outcome baseline sample of 38,982 participants. Participants who reported > 10 h sleep were omitted from the sample, as correlations have been found previously between long sleeps and poor health, which may be attributable to reverse causation (Stamatakis & Punjabi, 2007). Among participants with non-affirmative baseline responses, a second binary outcome was derived based upon incidence of insufficient sleeps at follow-up in a complete-outcome sample of 31,961 participants, respectively.

Each of the two samples were linked to data on percentage green space availability within a 1.6 km (1 mile) network distance buffer around the centroid of each Mesh Block of residence. The Mesh Block is the smallest geographical scale available in Australia at typically 30 to 60 dwellings each. The 1.6 km network distance buffer was selected based upon published guidance (National Prevention Council, 2014). Four green space indicators were derived: (i) total green space; (ii) tree canopy; (iii) open grass without tree canopy; (iv) other low-lying vegetation. Green space data was sourced from Pitney Bowes Ltd in the form of a raster layer ('Geovision') in 2016. This data consisted of a 2-m raster surface classified into descriptive classes captured using machine learning and image classification processes across satellite imagery (8-band multispectral imagery captured by DigitalGlobe's Worldview 3 satellite). Tree canopy included street trees and trees in parks (both deciduous and evergreen). Open grass included herbaceous areas. Open grass and low-lying vegetation were likely under-estimated in areas where they were beneath tree canopy. All four green space indicators were formatted as intervals in order to examine for potentially policy-relevant threshold effects. For total green space and open grass, these intervals were: 0–4%; 5–9%; 10–19%; 20–29%; $\geq 30\%$. For tree canopy and other low-lying vegetation, the intervals were slightly different due to small numbers (tree canopy = 0–9%; 10–19%; 20–29%; $\geq 30\%$, low-lying vegetation = 0–1%; 2–4%; $\geq 5\%$). These categories were based around existing green space standards, for example, the allocation of 10% of subdivisible land to some form of green open space in Perth, Western Australia.

Multilevel logistic regressions estimated with Markov Chain Monte Carlo (MCMC) method (Browne, 2005; Rasbash et al., 2000) took into account clustering of participants within areas. Initial regressions assessed prevalence and incidence of insufficient sleep in relation to each green space indicator, adjusting for age and gender, followed by adjustment for social and economic markers of confounding associated with sleep duration and may also structure access to green space: annual household income; economic status (e.g. employed, retired); educational qualification; and couple status. Each of these markers of potential confounding were self-reported within the 45 and Up Study questionnaire. Weights were not used to address over-sampling of people aged 85 and older.

Results

The prevalence of insufficient sleep at baseline was 18.0% (Table 1). Incidence of insufficient sleep was 13.2%. Comparing participants with $\geq 30\%$ to those with 0–4% total green space, prevalent and incident insufficient sleep were down by 7.4% and 1.6%, respectively. For participants with $\geq 30\%$ tree canopy compared with 0–9% tree canopy, prevalent and incident insufficient sleep were 7.50% and 3.76% lower, respectively. Prevalent and incident insufficient sleep were 2.7% and

Table 1
Descriptive statistics of the study sample.

	Total N (baseline)	% insufficient sleep prevalence	Total N (follow- up)	% insufficient sleep incidence
Total	38982	18.0	31961	13.2
Total green space				
0–4%	631	23.3	484	14.1
5–9%	9350	18.7	7599	13.3
10–19%	10400	18.7	8455	13.8
20–29%	10491	18.0	8602	13.1
≥30%	8110	15.9	6821	12.5
Chi-square (p-value)	43.2	<0.001	6.6	0.161
Tree canopy				
0–9%	4051	22.8	3131	15.4
10–19%	15086	19.3	12172	14.0
20–29%	9827	16.8	8172	12.9
≥30%	10015	15.3	8486	11.6
Chi-square (p-value)	139.7	<0.001	37.9	<0.001
Open grass				
0–4%	4818	16.5	4022	12.0
5–9%	16344	16.8	13600	12.4
10–19%	9996	18.9	8112	13.9
20–29%	6161	20.7	4883	14.8
≥30%	1663	19.2	1344	15.0
Chi-square (p-value)	61.2	<0.001	30.1	<0.001
Low-lying vegetation				
0–4%	25051	17.8	20583	13.0
5–9%	12639	18.4	10317	13.5
≥10%	1292	17.9	1061	14.8
Chi-square (p-value)	1.7	0.438	3.8	0.1
Age group				
45–54y	13695	16.4	11370	14.0
55–64y	14301	17.7	11769	12.6
65–74y	7105	18.8	5770	13.0
≥75y	3881	21.4	3052	13.7
Chi-square (p-value)	43.2	<0.001	10.3	0.016
Gender				
Male	18008	18.2	14732	12.4
Female	20974	17.9	17229	13.9
Chi-square (p-value)	0.7	0.389	16.2	<0.001
Annual household income (\$)				
0–20k	4095	24.6	3086	16.1
21–30k	2458	20.0	1967	13.2
31–40k	2505	17.3	2072	14.5
41–50k	2663	18.0	2183	14.7
51–70k	4541	18.4	3706	12.6
>70k	15693	15.3	13295	11.4
Not determined	7027	19.6	5652	15.3
Chi-square (p-value)	220.4	<0.001	92.5	<0.001
Education				
None	2238	27.5	1622	18.6
School	6394	20.8	5067	15.8
High School	3684	18.3	3010	14.2
Trade	3271	20.5	2602	14.4
Certificate or Diploma	8694	17.1	7206	12.6
University	14363	15.1	12189	11.3
Missing	338	21.6	265	15.5
Chi-square (p-value)	271.2	<0.001	119.3	<0.001

Table 1 (continued)

	Total N (baseline)	% insufficient sleep prevalence	Total N (follow- up)	% insufficient sleep incidence
Total	38982	18.0	31961	13.2
Economic status				
Employed	23410	17.8	19242	12.9
Retired	12467	17.8	10245	13.1
Unemployed	510	24.9	383	18.5
Unpaid work	541	16.1	454	14.1
Disabled	385	30.1	269	21.2
Homemaker	1232	16.6	1028	15.4
Other	437	22.2	340	17.4
Chi-square (p-value)	64.0	<0.001	36.2	<0.001
Couple status				
Not in a couple	8391	23.2	6433	13.9
In a couple	30611	16.6	25528	13.0
Chi-square (p-value)	19.8	<0.001	3.1	0.076

3.1% higher among participants with ≥30% open grass compared with 0–4% open grass nearby, respectively. There was no pattern of prevalent or incident insufficient sleep discernible with respect to low-lying vegetation, so this exposure was not analysed any further.

Fully-adjusted odds ratios for each outcome in relation to total green space, tree canopy and open grass exposures are illustrated in Fig. 1. Participants in total green space intervals higher than 0–4% had lower odds of prevalent short sleep (Fig. 1A). For example, the odds ratio (OR) for participants with 5–9% total green space was 0.76 (95% credible interval (95%CI) = 0.61, 0.95). Although the odds of incident insufficient sleep appeared to be lower at higher intervals of total green space, these associations were not statistically significant (e.g. ≥30% compared with 0–4% OR = 0.90, 95%CI = 0.67, 1.16).

The odds of prevalent insufficient sleep were lower among participants with tree canopy levels at every interval above 0–9% (e.g. ≥30% OR = 0.78, 95%CI 0.69, 0.88, Fig. 1B). The odds of incident insufficient sleep were also lower with more tree canopy, albeit only statistically significantly at the ≥30% level (OR = 0.87, 95%CI = 0.75, 0.99). In contrast to the descriptive analyses, there was no statistically significant evidence of association between prevalent or incident insufficient sleep and the open grass exposure after adjusting for demographic and socioeconomic markers of confounding (Fig. 1C).

Discussion

Our study has enhanced this new area of research in two ways. Lower prevalent insufficient sleep among people with higher total green space available was confirmed. But when examining the data longitudinally, the correlation between total green space and incident insufficient sleep was not convincing. Lower odds of insufficient sleep were more consistently observed among people living near more tree canopy in both the cross-sectional and longitudinal tests. Neither open grass nor other low-lying vegetation were statistically significantly associated with prevalent or incident insufficient sleeps. This suggests urban greening strategies that prioritise increasing the availability of, and reducing inequalities in tree canopy coverage may help to support population-wide improvements in sleep.

Some merits of our study include longitudinal lagged-effects design of the best green space data available and a large residentially stable sample to prevent bias from reverse causation through health-selective migration. Assessment of incident insufficient sleep in this study provided a more robust test of association with green space provision than had previously been achieved, with earlier work relying upon data of cross-sectional design (Astell-Burt et al., 2013; Chum et al., 2015;




Fig. 1. Odds of prevalent and incident short sleep durations in association with total (A) urban green space, (B) tree canopy, and (C) open grass: multilevel logistic regressions adjusted for confounding and estimated using Markov Chain Monte Carlo method. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Grigsby-Toussaint et al., 2015; Johnson et al., 2018b). This longitudinal design combined with the ability to differentiate between different types of green space afforded the identification of tree canopy, rather than any green space, as being associated with lower odds of incident insufficient sleep.

Limitations include self-report sleep duration data and restriction to an ≥ 45 y population. Our results cannot be generalised to people aged ≤ 44 y. However, the prevalence of insufficient sleep increases with age according to a recent nationally representative survey (Adams et al., 2017), which helps to explain why the prevalence of insufficient sleep in our study (18%) was above the national average for adults ≥ 18 y (12%). Potential underestimation in green space land-use, due to loss of provision in some areas over time could not be measured with the data available. Existing evidence suggests 35% of council areas in Australia

have experienced some form of tree canopy loss while only 4% increased tree canopy coverage (Amati et al., 2017). The potential impacts on sleep duration of a change in local tree canopy cover and land-use over time is an important avenue for future research. It is also important to note that our data were limited in respect of open grass cover and other low-lying vegetation being under-estimated in areas where it was located underneath tree canopy.

The identification of tree canopy, rather than any green space *per se*, as potentially important for promoting healthier sleep duration may point towards mitigation and restoration as possible pathways, perhaps via exposure to increased biodiversity (Fuller et al., 2007), natural soundscapes (Annerstedt et al., 2013), amelioration of heat islands (Ng et al., 2012), less psychological distress (Astell-Burt & Feng, 2019) and satisfying human preferences for natural settings with more complex vegetation (Harris et al., 2018). The lack of association between sleep and open grass after adjusting for confounding factors may suggest that pathways involving physical activity are less important in this particular case. However, the association between physical activity and open grass may be contingent upon whether those areas are in private ownership (i. e. back gardens) or are publically accessible (e.g. parks and some sports ovals). Furthermore, research that examines potential pathways using relevant statistical models capable of discerning between serial and parallel mediation would be a prudent next step, as would replication studies and estimates of cost-effectiveness of potential investments in tree canopy. In conclusion, this is the first longitudinal study internationally to link residential green space and sleep duration, with the evidence pointing towards restoration and conservation of urban tree canopy specifically as a scalable public health intervention for prevention of insufficient sleep.

Ethical statements

The University of New South Wales HREC approved the 45 and Up Study and ethics approval for the analyses in this paper was awarded by the University of Wollongong HREC.

Acknowledgements

This study was supported by a National Health and Medical Research Council Boosting Dementia Research Leader Fellowship 1140317 (Dr Astell-Burt) and National Health and Medical Research Council Career Development Fellowship 1148792 (Dr Feng). Drs Astell-Burt and Feng are also supported by grant 1101065 from the National Health and Medical Research Council project and the Green Cities Fund - Hort Innovation Limited, with coinvestment from the University of Wollongong Faculty of Social Sciences, the University of Wollongong Global Challenges initiative, and the Australian Government project GC15005. The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and the decision to submit the manuscript for publication. This research was completed using data collected through the 45 and Up Study (<http://www.saxinstitute.org.au>). The 45 and Up Study is managed by the Sax Institute in collaboration with major partner Cancer Council New South Wales (NSW) and partners the National Heart Foundation of Australia (NSW Division), NSW Ministry of Health, NSW Government Family & Community Services–Ageing, Carers, and the Disability Council NSW, and the Australian Red Cross Blood Service. We thank the many thousands of people participating in the 45 and Up Study.

References

- 45 and Up Study Collaborators. (2008). Cohort profile: The 45 and up study. *International Journal of Epidemiology*, 37(5), 941–947.
- Abhijith, K., Kumar, P., Gallagher, J., et al. (2017). Air pollution abatement performances of green infrastructure in open road and built-up street canyon environments—A review. *Atmospheric Environment*, 162, 71–86.

- Adams, R. J., Appleton, S. L., Taylor, A. W., et al. (2017). Sleep health of Australian adults in 2016: Results of the 2016 sleep health foundation national survey. *Sleep Health*, 3(1), 35–42.
- Alvaro, P. K., Roberts, R. M., & Harris, J. K. (2013). A systematic review assessing bidirectionality between sleep disturbances, anxiety, and depression. *Sleep*, 36(7), 1059–1068.
- Amati, M., Boruff, B., Caccetta, P., et al. (2017). *Where should all the trees go? Investigating the impact of tree canopy cover on socio-economic status and wellbeing*. Melbourne: Innovation Australia Ltd.
- Annerstedt, M., Jönsson, P., Wallergård, M., et al. (2013). Inducing physiological stress recovery with sounds of nature in a virtual reality forest—results from a pilot study. *Physiology & Behavior*, 118, 240–250.
- Astell-Burt, T., & Feng, X. (2019). Association of urban green space with mental health and general health among adults in Australia. *JAMA network open*, 2(7). e198209-e09.
- Astell-Burt, T., Feng, X., & Kolt, G. S. (2013). Does access to neighborhood green space promote a healthy duration of sleep? Novel findings from 259,319 Australians. *BMJ Open*, 3, e003094.
- Browne, W. J. (2005). *MCMC estimation in MLwiN: Version 2.0*. Centre for Multilevel Modelling, University of Bristol.
- Cappuccio, F. P., Cooper, D., Elia, L. D., et al. (2011). Sleep duration predicts cardiovascular outcomes: A systematic review and meta-analysis of prospective studies. *European Heart Journal*, 32, 1484–1492.
- Cappuccio, F. P., D'elia, L., Strazzullo, P., et al. (2010a). Quantity and quality of sleep and incidence of type 2 diabetes: A systematic review and meta-analysis. *Diabetes Care*, 33(2), 414–420.
- Cappuccio, F. P., D'Elia, L., Strazzullo, P., et al. (2010b). Sleep duration and all-cause mortality: A systematic review and meta-analysis of prospective studies. *Sleep*, 33(5), 585–592.
- Cappuccio, F. P., Taggart, F. M., Kandala, N. B., et al. (2008). Meta-analysis of short sleep duration and obesity in children and adults. *Sleep*, 31, 619–626.
- Chum, A., O'Campo, P., & Matheson, F. (2015). The impact of urban land uses on sleep duration and sleep problems. *The Canadian Geographer/Le Géographe canadien*, 59(4), 404–418.
- Connor, J., Whitlock, G., Norton, R., et al. (2001). The role of driver sleepiness in car crashes: A systematic review of epidemiological studies. *Accident Analysis & Prevention*, 33(1), 31–41.
- Dzhambov, A. M., & Dimitrova, D. D. (2015). Green spaces and environmental noise perception. *Urban Forestry and Urban Greening*, 14(4), 1000–1008.
- Frumkin, H., Bratman, G. N., Breslow, S. J., et al. (2017). Nature contact and human health: A research agenda. *Environmental Health Perspectives*, 125(7), 075001-01.
- Fuller, R. A., Irvine, K. N., Devine-Wright, P., et al. (2007). Psychological benefits of greenspace increase with biodiversity. *Biology Letters*, 3(4), 390–394.
- Grigsby-Toussaint, D. S., Turi, K. N., Krupa, M., et al. (2015). Sleep insufficiency and the natural environment: Results from the US behavioral risk factor surveillance system survey. *Preventive Medicine*, 78, 78–84.
- Hale, L., James, S., Xiao, Q., et al. (2019). *Neighborhood factors associated with sleep health. Sleep and Health*. Elsevier.
- Halperin, D. (2014). Environmental noise and sleep disturbances: A threat to health? *Sleep science*, 7(4), 209–212.
- Harris, V., Kendal, D., Hahs, A. K., et al. (2018). Green space context and vegetation complexity shape people's preferences for urban public parks and residential gardens. *Landscape Research*, 43(1), 150–162.
- Hartig, T., Mitchell, R., de Vries, S., et al. (2014). Nature and health. *Annual Review of Public Health*, 35, 207–228.
- Hunter, J. C., & Hayden, K. M. (2018). The association of sleep with neighborhood physical and social environment. *Public Health*, 162, 126–134.
- Jennings, V., & Bamkole, O. (2019). The relationship between social cohesion and urban green space: An avenue for health promotion. *International Journal of Environmental Research and Public Health*, 16(3), 452.
- Johnson, D. A., Billings, M. E., & Hale, L. (2018a). Environmental determinants of insufficient sleep and sleep disorders: Implications for population health. *Current epidemiology reports*, 5(2), 61–69.
- Johnson, B. S., Malecki, K. M., Peppard, P. E., et al. (2018b). Exposure to neighborhood green space and sleep: Evidence from the survey of the health of Wisconsin. *Sleep Health*, 4(5), 413–419.
- Kecklund, G., & Axelsson, J. (2016). Health consequences of shift work and insufficient sleep. *BMJ*, 355, i5210.
- Kredlow, M. A., Capozzoli, M. C., Hearon, B. A., et al. (2015). The effects of physical activity on sleep: A meta-analytic review. *Journal of Behavioral Medicine*, 38(3), 427–449.
- Lange, T., Dimitrov, S., & Born, J. (2010). Effects of sleep and circadian rhythm on the human immune system. *Annals of the New York Academy of Sciences*, 1193(1), 48–59.
- Maas, J., Van Dillen, S. M. E., Verheij, R. A., et al. (2009). Social contacts as a possible mechanism behind the relation between green space and health. *Health & Place*, 15(2), 586–595.
- Magee, C. A., Iverson, D. C., & Caputi, P. (2009). Factors associated with short and long sleep. *Preventive Medicine*, 49(6), 461–467.
- Markevych, I., Schoierer, J., Hartig, T., et al. (2017). Exploring pathways linking greenspace to health: Theoretical and methodological guidance. *Environmental Research*, 158, 301–317.
- McHugh, J. E., & Lawlor, B. A. (2013). Perceived stress mediates the relationship between emotional loneliness and sleep quality over time in older adults. *British Journal of Health Psychology*, 18(3), 546–555.
- Mealing, N. M., Banks, E., Jorm, L. R., et al. (2010). Investigation of relative risk estimates from studies of the same population with contrasting response rates and designs. *BMC Medical Research Methodology*, 10(1), 26.
- Muzet, A. (2007). Environmental noise, sleep and health. *Sleep Medicine Reviews*, 11(2), 135–142.
- National Prevention Council. (2014). *Annual status report*. Washington, DC: Department of Health and Human Services, Office of the Surgeon General.
- Ng, E., Chen, L., Wang, Y., et al. (2012). A study on the cooling effects of greening in a high-density city: An experience from Hong Kong. *Building and Environment*, 47, 256–271.
- Patel, S. R., & Hu, F. B. (2008). Short sleep duration and weight gain: A systematic review. *Obesity*, 16(3), 643–653.
- Rasbash, J., Browne, W., Goldstein, H., et al. (2000). *A user's guide to MLwiN*.
- Simon, E. B., & Walker, M. P. (2018). Sleep loss causes social withdrawal and loneliness. *Nature Communications*, 9.
- Spiegel, K., Tasali, E., Leproult, R., et al. (2009). Effects of poor and short sleep on glucose metabolism and obesity risk. *Nature Reviews Endocrinology*, 5(5), 253.
- Stamatakis, K. A., & Punjabi, N. M. (2007). Long sleep duration: A risk to health or a marker of risk? *Sleep Medicine Reviews*, 11(5), 337.
- Sygn, K., Aasvang, G. M., Aamodt, G., et al. (2014). Road traffic noise, sleep and mental health. *Environmental Research*, 131, 17–24.
- Uehli, K., Mehta, A. J., Miedinger, D., et al. (2014). Sleep problems and work injuries: A systematic review and meta-analysis. *Sleep Medicine Reviews*, 18(1), 61–73.
- Zanobetti, A., Redline, S., Schwartz, J., et al. (2010). Associations of PM10 with sleep and sleep-disordered breathing in adults from seven US urban areas. *American Journal of Respiratory and Critical Care Medicine*, 182(6), 819–825.