

1-1-2020

Loneliness among people with substance use problems: A narrative systematic review

Isabella Ingram

University of Wollongong, ii841@uowmail.edu.au

Peter James Kelly

University of Wollongong, pkelly@uow.edu.au

Frank P. Deane

University of Wollongong, fdeane@uow.edu.au

Amanda Baker

Melvin Goh

University of Wollongong, mcwg178@uowmail.edu.au

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/sspapers>



Part of the [Education Commons](#), and the [Social and Behavioral Sciences Commons](#)

Loneliness among people with substance use problems: A narrative systematic review

Abstract

2020 Australasian Professional Society on Alcohol and other Drugs Issues: Despite the serious implications of loneliness on health and wellbeing, little is understood about this experience across people with substance use problems. This systematic review aimed to examine: (i) correlates and predictors of loneliness; (ii) theories underpinning loneliness; (iii) methods employed to measure loneliness; and (iv) loneliness interventions for people with substance use problems. Approach: Empirical sources were identified from key databases for all publications preceding February 2019. Overall, 41 studies met the eligibility criteria and were included in the review. Key Findings: Findings from this review suggest that loneliness is related to poor physical and mental health, substance use, the quality of relationships, stigma and perception of ill treatment by others. Although cognitive theories have proposed cognitive patterns underlying the onset and maintenance of loneliness, they had not been investigated in relation to measurement or intervention efforts. Just one loneliness measure (UCLA Loneliness Scale) is valid for use with this population. Finally, only a single loneliness intervention had been trialled and was not found to be efficacious in reducing loneliness for people with substance use problems. Implications: Understanding possible links between loneliness and substance use and how to alleviate loneliness is important for this population in terms of their wellbeing and recovery. Conclusion: Loneliness is prevalent and experienced as problematic among people with substance use problems. Future research should focus on employing longitudinal designs, using validated, multidimensional measures of loneliness and on developing and trialling loneliness interventions that meet the specific needs of people with substance use problems.

Disciplines

Education | Social and Behavioral Sciences

Publication Details

Ingram, I., Kelly, P., Deane, F., Baker, A., Goh, M., Raftery, D. & Dingle, G. (2020). Loneliness among people with substance use problems: A narrative systematic review. *Drug and Alcohol Review*,

Authors

Isabella Ingram, Peter James Kelly, Frank P. Deane, Amanda Baker, Melvin Goh, Dayle Raftery, and Genevieve A. Dingle



Loneliness amongst people with substance use problems: A narrative systematic review

Journal:	<i>Drug and Alcohol Review</i>
Manuscript ID	CDAR-2019-0220.R1
Manuscript Type:	Review
Date Submitted by the Author:	10-Feb-2020
Complete List of Authors:	Ingram, Isabella; University of Wollongong School of Psychology, Illawarra Health and Medical Research Institute Kelly, Peter; University of Wollongong School of Psychology, Illawarra Health and Medical Research Institute Deane, Frank; University of Wollongong School of Psychology, Illawarra Health and Medical Research Institute Baker, Amanda; The University of Newcastle School of Medicine and Public Health Goh, Melvin; University of Wollongong School of Psychology, Illawarra Health and Medical Research Institute Raftery, Dayle; University of Wollongong School of Psychology, Illawarra Health and Medical Research Institute Dingle, Genevieve; University of Queensland School of Psychology
Keywords (Please ensure that the Keywords and a short Running Head are also included in the manuscript file):	Loneliness, Systematic review, Addiction, Alcohol and substance dependence

SCHOLARONE™
Manuscripts

Running head: Review: Loneliness and substance dependence

Loneliness amongst people with substance use problems: A narrative systematic review

Isabella Ingram¹, Peter J. Kelly¹, Frank P. Deane¹, Amanda L. Baker², Melvin C. W. Goh¹,
Dayle K. Raftery¹, Genevieve A. Dingle³

¹Illawarra Health and Medical Research Institute and School of Psychology, University of
Wollongong, Wollongong Australia

²School of Medicine and Public Health, University of Newcastle, Newcastle, Australia

³School of Psychology, University of Queensland, Brisbane, Australia

Isabella Ingram GDipProfPsych, PhD Candidate, Peter J. Kelly PhD, Head of Research,
Frank P. Deane PhD, Professor, Amanda L. Baker PhD, Professor, Melvin C. W. Goh
BPsycSc, PhD Candidate, Dayle K. Raftery MProfPsyc, PhD Candidate, Genevieve A.
Dingle PhD, Associate Professor.

Author for correspondence: Ms Isabella Ingram, School of Psychology, University of
Wollongong, Wollongong, NSW 2500, Australia. Tel: 02 4221 4484, Email:

ingram@uow.edu.au

Running head: Review: Loneliness and substance dependence

Abstract

Issues: Despite the serious implications of loneliness on health and wellbeing, little is understood about this experience across people with substance use problems. This systematic review aimed to examine: (i) correlates and predictors of loneliness; (ii) theories underpinning loneliness; (iii) methods employed to measure loneliness; and (iv) loneliness interventions for people with substance use problems.

Approach: Empirical sources were identified from key databases for all publications preceding February 2019. Overall, 41 studies met the eligibility criteria and were included in the review.

Key Findings: Findings from this review suggest that loneliness is related to poor physical and mental health, substance use, the quality of relationships, stigma, and perception of ill treatment by others. Although cognitive theories have proposed cognitive patterns underlying the onset and maintenance of loneliness, they had not been investigated in relation to measurement or intervention efforts. Just one loneliness measure (UCLA Loneliness Scale) is valid for use with this population. Finally, only a single loneliness intervention had been trialled and was not found to be efficacious in reducing loneliness for people with substance use problems.

Implications: Understanding possible links between loneliness and substance use and how to alleviate loneliness is important for this population in terms of their wellbeing and recovery.

Conclusion: Loneliness is prevalent and experienced as problematic among people with substance use problems. Future research should focus on employing longitudinal designs, using validated, multidimensional measures of loneliness, and on developing and trialling loneliness interventions that meet the specific needs of people with substance use problems.

Keywords: loneliness, systematic review, addiction, alcohol and substance dependence

Running head: Review: Loneliness and substance dependence

1
2
3 Loneliness is a global public health issue [1], predicting poor physical and mental
4 health, and morbidity and mortality across the general population [2-5]. Loneliness is a
5 painful emotional state resulting from a discrepancy between the relationships one perceives
6 they have and those they desire [6]. Throughout the literature, loneliness has been considered
7 as both a uni-dimensional construct and also a multi-dimensional construct, encompassing
8 both social and emotional forms of loneliness [7].
9

10
11
12
13
14
15
16
17 Global prevalence rates of loneliness are difficult to ascertain, but it is estimated that
18 40% of older adults [8] and roughly one-third of people across industrialised countries
19 experience loneliness [1]. Recent research has focused on determining those age groups,
20 characteristics and specific populations that may be most vulnerable to experiencing
21 loneliness [e.g. 9-11]. Reviews have concluded that loneliness is highly prevalent amongst
22 elderly populations [12] and people living with serious mental illnesses, such as psychosis
23 [13]. Despite the growth in loneliness research, people with substance use problems are a
24 population that has been largely overlooked, with no reviews having been conducted in this
25 area [14].
26
27
28
29
30
31
32
33
34
35
36

37
38 Illicit drug use has been deemed the most stigmatised health condition in the world,
39 while alcohol dependence is the fourth most stigmatised condition [15, 16]. Research across a
40 range of populations has found that social isolation can result from the effects of stigma [17-
41 19]. While not everyone who is isolated becomes lonely, social isolation and loneliness are
42 closely related [20]. Consequently, people with substance use problems are vulnerable to
43 experiencing loneliness that arises from stigma and social isolation. Additionally, people with
44 substance use problems may make and maintain relationships that meet their needs and
45 support their active substance use but once in recovery, their social needs are likely to have
46 changed (e.g. toward non-using contacts). When abstinent from substance use, there may be a
47 need to avoid those situations and relationships that perpetuate ongoing substance use and
48
49
50
51
52
53
54
55
56
57
58
59
60

Running head: Review: Loneliness and substance dependence

1
2
3 instead attempt to connect with people who support one's recovery [21]. This process is
4
5 likely to increase the risk of loneliness for people recovering from substance use problems.
6
7 Recent research has indicated that 79% of 316 individuals accessing treatment for substance
8
9 use problems reported often feeling lonely. Additionally, this study reported that 69% of
10
11 these participants agreed to the statement "loneliness has been a serious problem for me"
12
13 [22]. While such findings are limited to an Australian population, they suggest loneliness is
14
15 highly prevalent and problematic for people who experience substance use problems.
16
17
18
19

20 Loneliness research is in its infancy amongst those with substance use problems and
21
22 there is a need to better understand the correlates of loneliness and the relationship between
23
24 loneliness and substance use. In general community samples, both younger and older age
25
26 [23], male gender [24] poorer physical and mental health [25-27], poorer quality social
27
28 relationships [28] and poorer quality of life [27] have been associated with loneliness.
29
30 Despite these findings across the broader literature, current research in the context of people
31
32 with substance use disorders has revealed limited and conflicting findings in relation to
33
34 theoretical and empirical correlates and predictors of loneliness.
35
36
37

38 Social and cognitive theories of loneliness have been most prominent in efforts to try
39
40 to better understand predictors and causes of loneliness. For example, attribution theory [6]
41
42 suggests that in attempts to explain the cause of their loneliness, lonely people adopt an
43
44 attribution style that is internal and stable; that is, these individuals believe their loneliness is
45
46 due to some shortcoming of their own (internal) and that this shortcoming is unchangeable
47
48 (stable) [29-31]. Cognitive theories also describe a hypervigilance to social threat in the
49
50 environment, and negative expectations of social interactions, as being central to the onset
51
52 and maintenance of loneliness [26]. While these theories have been used to explain how
53
54 people become lonely, very few studies have referenced these theories in the context of
55
56 addiction. Preliminary research in this field suggests that that cognition may be important in
57
58
59
60

Running head: Review: Loneliness and substance dependence

1
2
3 explaining loneliness [32, 33], yet there is little empirical research available to support this
4
5 assertion. It remains unclear which specific cognitive patterns might be responsible for the
6
7 onset and maintenance of loneliness for people with substance use disorders, and
8
9 consequently, how to best assess and treat this problem.
10
11

12 Loneliness is a difficult construct to measure, with studies across other populations
13
14 using a range of tools that target related social constructs, such as social isolation [34]. The
15
16 University of California, Los Angeles (UCLA) Loneliness Scale [35] is the most widely used
17
18 tool across the broader literature, but there remains ongoing ambiguity surrounding the
19
20 dimensionality of this measure [36]. In addition, most research assessing the validity of
21
22 loneliness scales is focused on college samples or the ageing population [36]. The scarcity of
23
24 research with a focus on substance using populations means that questions remain about how
25
26 loneliness can best be assessed and this in turn impedes research efforts to develop targeted
27
28 interventions for this population.
29
30
31

32
33 Theory may provide guidance in addressing these needs and reviews and meta-
34
35 analyses of intervention studies appear to support cognitive theories of loneliness [14, 37].
36
37 These studies have found that interventions aiming to address maladaptive social cognition
38
39 were most efficacious in reducing loneliness across diverse samples, including children,
40
41 adults and seniors. The impact of interventions that target social cognition for people with
42
43 substance use problems is yet to be examined. In fact, little is known about the efficacy of
44
45 any type of intervention in helping to reduce feelings of loneliness for this population.
46
47
48

49 Given there is little understanding of loneliness across people with substance use
50
51 problems, the purpose of this review is to synthesise the existing literature. Specifically, this
52
53 review aims to examine: (i) correlates and predictors of loneliness (including demographic,
54
55 physical health, mental health, social variables and substance use variables); (ii) prominent
56
57 theories to explain loneliness; (iii) methods to measure loneliness; and (iv) interventions that
58
59
60

Running head: Review: Loneliness and substance dependence

have specifically aimed to target loneliness for people with substance use problems.

Methods

Protocol registration: The review protocol was registered with Prospero International Prospective Register of Systematic Reviews (registration number CRD42018105564) and can be accessed at <https://www.crd.york.ac.uk/prospero/>. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Checklist [see 38] was used to guide reporting of this review.

Information sources: Empirical sources were identified from the databases PsycINFO, PubMed, CINAHL Plus, MEDLINE, Scopus, Web of Science, and The Cochrane Library (Cochrane Database of Systematic Reviews, Cochrane Central Register of Controlled Trials [CENTRAL], Cochrane Methodology Register) for all publications preceding February 2019.

Search strategy: The searches were performed in April 2018 and updated in February 2019 using the search terms: “lonel*” and a range of relevant substance-related key terms (see protocol for a list of specific terms used). These terms were searched for in the *abstract, title, keywords* and *subject* of sources. Reference lists of identified sources were then screened to identify additional relevant studies.

Eligibility criteria: To be included in the review, the sources had to: (i) be published in English language; (ii) report on empirical research; (iii) report on loneliness in their results (loneliness being a research question of the study or specifically measured as part of the study); and (iv) contain a sample that consists of people with substance use problems (i.e. have a diagnosis of substance use disorder, or accessing treatment specifically for substance use problems).

Study selection: Overall, 41 studies met the eligibility criteria and were included in the review. Initial screening of titles and abstracts was undertaken by the first author, and then

Running head: Review: Loneliness and substance dependence

1
2
3 identified full texts were independently screened by the first author and MG. There was a
4
5 high degree of agreement between the first two reviewers, $k=0.84$, $P < 0.001$. Discrepancies in
6
7 decisions to include/exclude full-text sources were resolved through consultation with a third
8
9 reviewer (DR). Figure 1 shows the literature selection process.
10
11
12
13

14
15 INSERT FIGURE 1 ABOUT HERE.
16
17
18

19
20 *Data collection and items:* The first author extracted data from the 41 studies included in the
21
22 review. Data extraction included information related to: authors, the title of the study and
23
24 year of publication, type of study, study setting, participant characteristics and details of:
25
26 tools used to measure loneliness, theoretical discussions, demographics, substance use,
27
28 physical health, mental health, social variables and/or interventions reported in relation to
29
30 loneliness.
31
32

33
34 *Risk of bias in studies:* Two reviewers (II and MG) independently assessed the
35
36 methodological quality and risk of bias of the included quantitative studies against the criteria
37
38 set by the National Institutes of Health Study Quality Assessment Tools [39]. Discrepancies
39
40 in ratings of study quality were resolved through discussion and use of a third reviewer (DR).
41
42 Studies were assigned to one of three categories; 'good', 'fair', or 'poor' quality based on
43
44 their design and conduct. Studies were deemed to be of 'good' quality if validated
45
46 instruments were used (defined as instruments which had been validated for use with people
47
48 who experience substance use problems), the probability of information bias and selection
49
50 bias appeared to be low, follow-up (where relevant) was over a number of years (defined as
51
52 being a timeframe long enough to enable a meaningful analysis to be conducted of the
53
54 relationship between exposures and outcomes), and confounding variables were considered
55
56 and adjusted for. The probability of low selection and information bias was determined by
57
58
59
60

Running head: Review: Loneliness and substance dependence

1
2
3 inspecting the National Institutes of Health items corresponding to each type of bias. As a
4
5 general rule of thumb, the fewer the number of items that were deemed a 'no', then the lower
6
7 the risk of bias and better the overall study quality. The more subjective and qualitative
8
9 method for assessing risk of bias was based on recommendations by Viswanathana *et al.* [40]
10
11 and Wang *et al.* [41]. Qualitative studies that were included in the review were critically
12
13 appraised against the 10-item Critical Appraisal Skills Programme checklist for Qualitative
14
15 studies [42]. Based on the number of items coded 'no' on the Critical Appraisal Skills
16
17 Programme Checklist, indicating potential risk of bias, these qualitative studies were
18
19 categorised into 'good', 'fair' and 'poor' quality, with a greater number of 'no' responses
20
21 indicating poorer quality (See Table S1).
22
23
24
25

26 *Data summary and synthesis:* Data were summarised based on the specified aims of the
27
28 review. While the UCLA measure of loneliness was widely used across studies included in
29
30 the review, a diverse range of correlational and predictor variables were present. Given the
31
32 heterogeneity across these studies in terms of outcomes and methods used, as well as the
33
34 small samples sizes, a narrative synthesis was conducted rather than a meta-analytic
35
36 synthesis. This decision was based on the lack of robust statistical methods available for such
37
38 heterogeneity, and researchers warning against performing underpowered meta-analyses [43,
39
40 44].
41
42
43
44

45 **Results**

46
47 *Study selection:* Of the 1628 records screened, 173 full-text studies were assessed for
48
49 eligibility and 41 were included in the review. Figure 1 shows the study selection process and
50
51 reasons for study exclusions at each stage of the review.
52
53

54 *Study characteristics:* Study characteristics for the 41 studies included in the review
55
56 are presented in Table 1. Overall, nine of the included studies were longitudinal and 32 were
57
58
59
60

Running head: Review: Loneliness and substance dependence

cross-sectional in design, with six of these being qualitative studies. Eleven of the included studies were dissertation theses and the remaining 30 studies were journal articles.

INSERT TABLE 1

Risk of bias of each study: Using the National Institutes of Health quality assessment tool for observational cohort and cross-sectional studies, all studies were rated as either 'good', 'fair' or 'poor' quality. Four studies were rated as 'good', 29 studies as 'fair' and 9 studies as 'poor' (see Table 1). The one quantitative study rated as being of 'good' quality was deemed to have a low risk of information and selection bias and confounding variables were controlled for [64]. The three qualitative studies that were deemed to be 'good' quality [49,57,72] all appeared to present valid results that were clearly described and likely to be informative to relevant practice/policy and/or research literature. Many of the studies appeared to minimise some risk of selection or information bias, yet those that met very few of these criteria were deemed as 'poor' quality. Potential confounding factors were inconsistently assessed across studies, with just two of the cohort studies adjusting for confounding variables [64,65].

Synthesis of results

Across the studies included in this review, the age of participants ranged from 11 to 98 years old, and 65% ($n=9951$) were males. Four studies did not report data related to the proportion of each gender in their sample. Sample sizes ranged from 8 to 652 participants across 40 of the studies included in this review. One study did not specify their sample size [79]. Of the study samples, 49% ($n=20$) were people who used alcohol, 20% ($n=8$) were people who used opiates, 12% ($n=5$) used a mix of drugs and alcohol, 10% ($n=4$) used a mix of drugs only, and 5% ($n=2$) did not report the substance use of their sample. The remaining

Running head: Review: Loneliness and substance dependence

two studies consisted of a sample of people who used methamphetamines, and one sample of people who used heroin. Inpatient substance dependence treatment services were the most common study setting ($n=8$), followed by a combination of inpatient/outpatient settings ($n=7$), methadone maintenance settings ($n=5$), and other outpatient substance dependence treatment services ($n=4$). Fifteen studies used other samples, including Alcoholics Anonymous (AA) populations and inpatient and outpatient mental health services. Two studies did not specify their study setting. Loneliness prevalence was reported in five of the studies [22,32,67,75,81] and ranged from 35% to 79%.

Correlates and predictors of loneliness

Demographics (n=16 studies): Seven studies found no relationship between demographic variables and loneliness. Nine studies ($n=56\%$) reported correlations with some demographic variables. Of those nine, five suggested that younger individuals were lonelier ($n=56\%$ of nine studies) [46,54,65,69,70] and of seven examining gender, four suggested females ($n=57\%$ of seven studies) [46,69,70] may be more likely to be lonely across this population.

Health (n=22 studies): Consistent with findings across other populations, loneliness appears to be related to poor physical and mental health for people with substance use problems [e.g. 22,56,80,81]. Seven studies examined physical health variables and found that poorer sleep quality and quantity [60,67], increased pain intensity [66] and poorer self-rated physical health [22,68,69] was correlated with increased loneliness. In addition, mental health indicators such as depression [22,33,75,78,80,81], self-esteem [32,52,60,68,70,80], suicidality [53,57,83] and poorer wellbeing/quality of life [22,53] were also related to higher loneliness across 15 studies (83% of 18 studies examining mental health variables).

Social and cognitive variables (n=15 studies): For people with alcohol use problems, loneliness was related to dissatisfaction in the quality of their relationships [32,33,46]. In

Running head: Review: Loneliness and substance dependence

1
2
3 addition, loneliness was consistently related to poor social support with all three studies that
4
5 examined this variable finding a correlation [52,56,75]. Three studies also found that their
6
7 samples perceived ill treatment from others [32,56,60] or that they were stigmatised [48] and
8
9 that these factors were related to loneliness for people with substance use problems (i.e. [48]).
10
11 Aligned with the prominent cognitive theories, eight studies (53% of 15 studies examining
12
13 social and cognitive variables) reported factors such as shyness, poor self-esteem and feelings
14
15 of insecurity and inferiority are associated with feelings of loneliness, suggesting that
16
17 negative perceptions of the self in relation to others plays a role in the onset and/or
18
19 maintenance of loneliness [32,33,46,49,57,60,68,70].
20
21
22
23

24 *Substance use variables (n=20 studies):* Of studies that examined substance use
25
26 variables, four (20% of 20 studies) suggest that people use substances to avoid distressing
27
28 feelings such as loneliness [63,74,77,78]. Five studies (25%) that examined signs of
29
30 dependence, such as frequency, severity or duration and symptoms of use, suggest that these
31
32 variables are related to loneliness. When examining loneliness longitudinally, mixed findings
33
34 emerged, with one study suggesting loneliness to be related to signs of substance dependence
35
36 [60], and another study concluding that loneliness was not related to substance use at a two-
37
38 year follow-up [32]. No notable differences emerged in terms of the prevalence or severity of
39
40 loneliness and different types of substance (i.e. alcohol or other substances) [22,81].
41
42 Perceptions of oneself and others noted above appear to ultimately result in feelings of
43
44 loneliness, which in turn may be an antecedent to continued alcohol use [78] or higher
45
46 alcohol consumption [60]. Aligned with these results, some studies found loneliness was a
47
48 risk factor for continued opiate use [62], with opiate use reported to be a means of escaping
49
50 feelings of loneliness [63]. Similar findings emerged in the one study that examined people
51
52 with methamphetamine problems [74], and across people with poly-substance use problems
53
54 [77].
55
56
57
58
59
60

Running head: Review: Loneliness and substance dependence

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Additionally, studies included in the review examined participants across a range of substance use and recovery stages. Two studies (67% of three studies) [47,54] suggest that those in more acute stage of addiction (currently using/detoxification) are lonelier than those in middle and later stages of recovery, while one study found no differences [59]. Across the stages of recovery and treatment settings, common themes emerged in terms of the health, social and substance use variables that were related to loneliness. Specifically, poor sleep, poor self-rated physical health, depression, poorer self-esteem, suicidality and poorer wellbeing/quality of life were related to higher loneliness across samples that were currently using substances and those that were in recovery. Similarly, poor social support, poorer quality and fewer quantities of relationships were consistent social variables that were found to relate to loneliness across different stages of recovery and treatment settings. Finally, reports of substance use as a means of coping with, or escaping loneliness, were reported for samples that were actively using substances, and those that were in recovery.

Few studies ($n=2$, 5% of 41 studies) examined the impact of substance dependence treatment on loneliness. One study [72] suggested that residing in a therapeutic community treatment setting might in fact contribute to feelings of loneliness and social distancing. This qualitative study suggested that within this treatment environment, participants became polarised, whereby alcohol users avoided illicit drug users. While this leaves questions as to the role of treatment services in impacting feelings of loneliness, stigma and social distancing might be a factor that contributes to loneliness in these settings. Targeting stigmatising attitudes of others as well as internalised stigma remains an ongoing target of policy makers and substance use treatment providers (e.g. [84]), and is evidently a necessity in order to aid in reducing feelings of loneliness for this population. Despite the research finding by Neale *et al.* [72] another study found some evidence to suggest that mutual support groups, such as AA groups, might be beneficial in reducing feelings of loneliness [49]. The effect of AA

Running head: Review: Loneliness and substance dependence

might be attributed to the recovery-based social identity gained, and the positive social capital generated through involvement in AA groups [85]. Factors such as social support, feelings of acceptance and shared values that are common in mutual support groups are thought to increase a sense of belonging and reduce feelings of loneliness.

Theories of loneliness (n=2 studies): Studies that describe theories of loneliness across this population are scarce. Just two studies applied theories of loneliness to their design or findings, with both of these studies discussing cognitive theories of loneliness. Akerlind and Hörnquist [32] allude to the social psychological and cognitive perspective (i.e. [7,86-88]). They found loneliness to be related to dissatisfaction with social relationships and to perceived negative treatment from others, and they explained these findings by referring to theory. Notably, these theories were not discussed in depth in relation to their study findings. Johnson [33] discussed her findings, that loneliness decreased with age, in relation to the Attribution Theory of loneliness [88], by suggesting that as one ages, their ability to develop realistic expectations for their relationships improves, as does their ability to overcome social inhibitions. Furthermore, Johnson [33] found that shyness predicted loneliness amongst people who used alcohol. These findings were discussed in the context of Attribution Theory by highlighting that a person's identification as 'shy' is likely to be a way to explain their relational characteristics in a way that is stable, internal and uncontrollable (consistent with Attribution Theory), which in turn decreases the likelihood of developing new relationships.

Measures of loneliness (n=36 studies): A range of measures were used to assess loneliness, yet very few studies used tools that had been specifically validated for use across substance dependent populations. Twelve of the 36 studies used author-developed measures to assess loneliness (33%), while 17 (47%) studies used measures that were developed and validated across a range of populations. Ten (28%) studies used other psychometric tools that included items or subscales that asked about loneliness. The three final studies included in the

Running head: Review: Loneliness and substance dependence

review [49,62,71] did not specifically measure loneliness; rather loneliness was a key theme that emerged as an outcome of these qualitative studies. All measures are summarised in Table 2.

INSERT TABLE 2

While three studies (8% of 36 studies) set out to validate measures of loneliness for substance dependent populations [22,50,73], these studies each used different tools to measure loneliness, with only one study using a multidimensional measure that captured both social and emotional forms of loneliness [22]. The UCLA measure was the most widely-used tool across studies included in this review ($n=11$, 31% of 36 studies), with 'fair' to 'excellent' internal consistency, and 'fair' test-retest reliability reported across study samples (see Table 2). There was limited consistency in the reporting of psychometric properties of loneliness measures. A number of studies included in the review made reference to psychometric properties that were reported for other populations in prior studies (see Table 2 and Appendix 1 for further information). Overall, the mixed reporting of psychometrics, and mixed findings across these studies, continues to create a challenge for future researchers in the field, in terms of selection and application of loneliness measures.

Loneliness interventions (n=1 study): Just one study aimed to evaluate an intervention to alleviate loneliness. Using a non-randomised trial design, Johnson [33] examined a logotherapy intervention. This is an existential form of therapy, which was delivered in a group format for one hour per day over a period of two weeks at an inpatient alcohol treatment facility. This intervention appeared to focus on the identification of personal values, and on goal setting to pursue values-congruent activities. The study involved a treatment and a control group of all males, who were compared on measures of loneliness at post-

Running head: Review: Loneliness and substance dependence

1
2
3 intervention. It was found that there were no differences in loneliness between the
4
5 intervention and control groups following treatment. The non-significant results were
6
7 attributed to the brief nature of the intervention (two weeks) and the broad focus of the
8
9 intervention (not loneliness specific), with some participants having identified values and
10
11 goals that were not social in nature. Findings from this study point to the need for more
12
13 research focusing on lengthier and more targeted interventions for loneliness amongst
14
15 substance dependent populations.
16
17
18
19

20 Discussion

21
22
23 This systematic review aimed to present a comprehensive overview of loneliness
24
25 research conducted across substance dependent populations. Forty-one completed studies met
26
27 the eligibility criteria and were included in this review. Findings from this review provide
28
29 preliminary evidence to suggest that people with substance use problems are lonelier than the
30
31 general population (i.e. [22]) and that females and those younger in age may be lonelier.
32
33 Socioemotional selectivity theory [106] may help to explain some of these findings. This
34
35 theory posits that in later life, individuals tend to cultivate their social networks in order to
36
37 enhance the social and emotional gains they derive from these relationships. This theory
38
39 might explain the reduced loneliness in older age groups found in some studies in this review.
40
41 Findings from this review also suggest that loneliness is consistently related to poor physical
42
43 and mental health for this population. Since data is predominantly correlational in nature, the
44
45 causal sequence of these relationships cannot be determined. It is possible that loneliness
46
47 leads to poorer health, or those experiencing poorer health become lonelier, or both.
48
49
50
51
52
53

54 While it remains unclear whether differences in loneliness exist based on type of
55
56 substance of dependence, a consistent finding was that higher severity/duration of substance
57
58 dependence is related to higher loneliness. Those studies that examined signs of substance
59
60

Running head: Review: Loneliness and substance dependence

1
2
3 dependence and were longitudinal in design ($n=3$) revealed mixed findings in relation to
4
5 substance use variables and loneliness. One of these studies suggested loneliness was cross-
6
7 sectionally but not longitudinally related to substance use variables [32]. However, others
8
9 found loneliness *was* longitudinally related to substance use problems [64], higher alcohol
10
11 consumption, delirium and blackouts [60]. No research has clarified the causal direction or
12
13 dynamics of this relationship, but it is possible that those who use substances to a greater
14
15 extent (i.e. higher severity) are also those who are more likely to have difficulty maintaining
16
17 relationships and/or be stigmatised in society, and ultimately become lonelier as a result of
18
19 social isolation and stigma.
20
21
22
23

24 Overall, just two studies discussed their findings in the context of theories of
25
26 loneliness. Both studies referenced cognitive theories of loneliness and provided some basis
27
28 for the belief that cognition may be important in explaining loneliness amongst substance
29
30 dependent people. However, neither of these studies set out to test specific theories of
31
32 loneliness and the empirical basis to support these theories is lacking. Research that aims to
33
34 test theories of loneliness appears to be in its infancy. Reviews and meta-analyses exploring
35
36 interventions to reduce loneliness amongst a range of populations have suggested cognitive
37
38 interventions as likely to be most effective in alleviating loneliness [14,37,107,108]. Findings
39
40 from these intervention studies suggest that merely increasing social contact has little impact
41
42 on feelings of loneliness, and rather ‘maladaptive social cognition’ or the belief one has about
43
44 themselves and others in relationships, has a greater effect on the subjective feeling of
45
46 loneliness [37]. Conclusions from these intervention studies lend support to the cognitive
47
48 theory of loneliness proposed by Perlman and Peplau [88], yet further research is clearly
49
50 warranted, particularly in field of substance dependence.
51
52
53
54
55

56 The current review revealed that very few of the measures used across the 41 studies
57
58 had been validated for use with people who experience substance use problems. Only three
59
60

Running head: Review: Loneliness and substance dependence

1
2
3 studies [22,50,73] specifically aimed to examine psychometric properties of the loneliness
4
5 measures they used. Of these studies, just one aimed to validate a multi-dimensional measure
6
7 of loneliness in order to further the understanding of this construct amongst this unique
8
9 population [22]. Loneliness is progressively being viewed as a multidimensional construct,
10
11 which cannot be captured using single-item or uni-dimensional measures [109,110]. The
12
13 UCLA Loneliness Scale [35] was the most commonly used tool across the studies included in
14
15 this review. However, this measure does not isolate different types of loneliness that have
16
17 been largely accepted across the broader loneliness literature [see 111]. While this measure
18
19 had been validated for use in methadone maintenance settings, its validity across a broader
20
21 range of substance dependence samples (i.e. inpatient, alcohol, amphetamines) is yet to be
22
23 determined. In addition, this tool potentially poses problems with face validity. The scale
24
25 adopts an indirect approach to measuring loneliness by omitting the word 'lonely', an
26
27 approach that has been found to yield differences in responding compared with a direct
28
29 approach [e.g. 112].
30
31
32
33
34

35
36 One key finding of the current review is the scarcity of studies involving interventions
37
38 specifically aimed at alleviating loneliness for this population. This highlights an important
39
40 future research direction, whereby interventions aimed at targeting loneliness need be
41
42 developed, piloted, and rigorously tested using high quality research designs. Based on
43
44 findings from this review and that of previous research in the field of loneliness [e.g. 14]
45
46 interventions that target one's perceptions of themselves and others, such as how they are
47
48 treated, the support they receive and their self-esteem might be of benefit for this population.
49
50 Research efforts that seek to identify the specific cognitions and social variables to be the
51
52 focus of interventions are needed. Further, clarification about the relationship between
53
54 loneliness and stigma for this population is also warranted, as the effects of stigma on one's
55
56 view of the self and others might be an additional treatment target.
57
58
59
60

Running head: Review: Loneliness and substance dependence

1
2
3 Strengths of this review include the broad scope, inclusion of international samples
4 and studies of all designs. Additionally, the review included unpublished theses. Limitations
5 consist of our inclusion of studies with potential shortcomings in their methodologies, and
6 variability in how loneliness and clinical correlates of loneliness were measured. In addition,
7 our search strategy was refined to studies specifically examining loneliness, rather than
8 including related concepts such as social isolation. It is possible that in doing so, our review
9 may have failed to capture some aspects of the broader social context that are relevant for
10 people with substance use problems. It is also possible that studies examining loneliness have
11 been conducted in treatment service settings, yet the reports of these studies have not been
12 made publicly available. As such, our systematic review is not immune to publication bias, as
13 we were unable to access and include such potential studies due to our search strategy.
14 Finally, rather than examining all dimensions of substance use, including people who use
15 substances occasionally, we narrowed our review to include only people who experience
16 substance use problems. While it is recognised that substance use occurs along a continuum,
17 the rationale for this sample selection was due to the assumption that people with substance
18 use problems are a distinct population from people who occasionally use substances. This is
19 due to the increased stigma, increased mental health difficulties, and transitions in and out of
20 active substance use that people with substance use problems may experience – all of which
21 may enhance their propensity to experiencing loneliness.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 47 **Conclusions**

48
49 Overall, this review of loneliness across substance dependent populations suggests
50 that people with substance use problems are likely to feel lonelier than non-clinical
51 comparator populations. Given the current literature, it is unclear what is most likely to
52 contribute to loneliness for this population, and how best to alleviate loneliness and the
53 associated bearing it has on physical and mental health. There is a lack of research that tests
54
55
56
57
58
59
60

Running head: Review: Loneliness and substance dependence

1
2
3 components of theories of loneliness across this population. Future empirical research should
4
5 focus on testing prominent loneliness theories (e.g. cognitive theory) across people with
6
7 substance use problems in order to determine whether they predict loneliness and
8
9 consequently can inform loneliness interventions. That just one study aimed to examine the
10
11 effect of an intervention for loneliness across an alcohol-dependent population, suggests that
12
13 this is an important gap in the literature. Guided by theoretical work, research exploring
14
15 specific social variables and cognitions that might perpetuate loneliness and be the target of
16
17 loneliness interventions is needed in this field.
18
19
20

21
22 Evidence for sound instruments to capture loneliness and effective interventions to
23
24 alleviate loneliness for substance dependent populations was limited. At present, the UCLA
25
26 measure of loneliness appears to be the most widely used. Advantages of using this tool are
27
28 the potential for replicability and comparisons with other populations, as well as the potential
29
30 of ruling out measurement variance in understanding research in this area. However, of note
31
32 is that this tool has only been validated for a methadone maintenance sample and that it is an
33
34 indirect approach to measurement, potentially posing challenges for face validity. Future
35
36 research needs to focus on replicating validation studies that have already been conducted
37
38 across this population, and/or qualitative work to determine the acceptability of current
39
40 loneliness measures and potential necessity of developing tools that are more appropriate.
41
42
43

44
45 The majority of the studies reviewed did not utilise sound methodologies in that
46
47 confounders were rarely adjusted for, and validated measures of loneliness for this specific
48
49 population were rarely used. Future research should focus on employing longitudinal designs,
50
51 with use of comparator groups and use of validated, multidimensional measures of loneliness.
52
53 Furthering the theoretical understanding of loneliness and its related constructs will help to
54
55 inform the development of targeted interventions, and ultimately overcome loneliness for this
56
57 vulnerable population.
58
59
60

Running head: Review: Loneliness and substance dependence

1
2
3
4
5
6
7 **Acknowledgements**
8

9 This research has been conducted with the support of the Australian Government Research
10 Training Program Scholarship. This research did not receive any specific grant from funding
11 agencies in the public, commercial, or not-for-profit sectors. All authors have contributed to
12 the article preparation and have approved the final article.
13
14
15
16
17

18
19 **Conflict of interest**
20

21 The authors have no conflicts of interest to declare.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Running head: Review: Loneliness and substance dependence

References

1. Cacioppo JT, Cacioppo S. The growing problem of loneliness. *Lancet* 2018;391:426.
2. Holt-Lunstad J. The potential public health relevance of social isolation and loneliness: Prevalence, epidemiology, and risk factors. *Public Policy Aging Rep* 2017;27:127-30.
3. Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: A meta-analytic review. *PLoS Med* 2010;7:e1000316.
4. Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and social isolation as risk factors for mortality: A meta-analytic review. *Perspect Psychol Sci* 2015;10:227-37.
5. Rico-Urbe LA, Caballero FF, Martín-María N, Cabello M, Ayuso-Mateos JL, Miret M. Association of loneliness with all-cause mortality: A meta-analysis. *PLoS One* 2018;13:e0190033.
6. Peplau LA, Perlman D. Loneliness: a sourcebook of current theory, research, and therapy. USA: Wiley Interscience; 1982.
7. Weiss RS. Loneliness: The experience of emotional and social isolation. Cambridge, MA, US: The MIT Press; 1973.
8. Hawkey LC. Loneliness and social embeddedness in old age. In: Pachana N, editor. *Encyclopedia of Geropsychology*. Singapore: Springer; 2015.
9. Böger A, Huxhold O. Age-related changes in emotional qualities of the social network from middle adulthood into old age: How do they relate to the experience of loneliness? *Psychol Aging* 2018;33:482-96.
10. Hawkins-Elder H, Milfont TL, Hammond MD, Sibley CG. Who are the lonely? A typology of loneliness in New Zealand. *Aust N Z J Psychiatry* 2018;52:357-64.

Running head: Review: Loneliness and substance dependence

11. Lasgaard M, Friis K, Shevlin M. "Where are all the lonely people?" A population-based study of high-risk groups across the life span. *Soc Psychiatry Psychiatr Epidemiol* 2016;51:1373-84.
12. Landeiro F, Barrows P, Musson EN, Gray AM, Leal J. Reducing social isolation and loneliness in older people: a systematic review protocol. *BMJ Open* 2017;7:e013778.
13. Lim MH, Gleeson JFM, Alvarez-Jimenez M, Penn DL. Loneliness in psychosis: a systematic review. *Soc Psychiatry Psychiatr Epidemiol* 2018;53:221-38.
14. Mann F, Bone JK, Lloyd-Evans B, Frerichs J, Pinfold V, Ma R, et al. A life less lonely: the state of the art in interventions to reduce loneliness in people with mental health problems. *Soc Psychiatry Psychiatr Epidemiol* 2017;52:627-38.
15. Crisp A, Gelder M, Goddard E, Meltzer H. Stigmatization of people with mental illnesses: a follow-up study within the Changing Minds campaign of the Royal College of Psychiatrists. *World Psychiatry* 2005;4:106-13.
16. Room R, Rehm J, Trotter RT, Paglia A, UÜstün TB. Cross-cultural views on stigma valuation parity and societal attitudes towards disability. In: Rehm J, editor. *Disability and culture: Universalism and diversity*. Seattle, WA: Hofgrebe & Huber; 2001. p. 247–91.
17. Lichtenstein B, Laska MK, Clair JM. Chronic sorrow in the HIV-positive patient: Issues of race, gender, and social support. *AIDS Patient Care STDs* 2002;16:27-38.
18. Ware NC, Wyatt MA, Tugenberg T. Social relationships, stigma and adherence to antiretroviral therapy for HIV/AIDS. *AIDS Care* 2006;18:904-10.
19. Linz SJ, Sturm BA. The phenomenon of social isolation in the severely mentally ill. *Perspect Psychiatr Care* 2013;49:243-54.
20. Petersen J, Kaye J, Jacobs PG, Quinones A, Dodge H, Arnold A, et al. Longitudinal relationship between loneliness and social isolation in older adults: Results from the Cardiovascular Health Study. *J Aging Health* 2016;28:775-95.

Running head: Review: Loneliness and substance dependence

21. Best D, Gow J, Taylor A, Knox A, White W. Recovery from heroin or alcohol dependence: A qualitative account of the recovery experience in Glasgow. *J Drug Issues* 2011;41:359-77.
22. Ingram I, Kelly PJ, Deane FP, Baker AL, Raftery DK. Loneliness in treatment-seeking substance-dependent populations: Validation of the Social and Emotional Loneliness Scale for Adults-Short Version. *J Dual Diagn* 2018;14:211-9.
23. Luhmann M, Hawkley LC. Age differences in loneliness from late adolescence to oldest old age. *Devl Psychol* 2016;52:943.
24. Baker D. All the lonely people: loneliness in Australia, 2001–2009. Canberra: The Australia Institute; 2012.
25. Leigh-Hunt N, Bagguley D, Bash K, Turner V, Turnbull S, Valtorta N, et al. An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public Health* 2017;152:157-71.
26. Hawkley LC, Cacioppo JT. Loneliness matters: a theoretical and empirical review of consequences and mechanisms. *Ann Behav Med* 2010;40:218-27.
27. Australian Psychology Society. Australian Loneliness Report: A survey exploring the loneliness levels of Australians and the impact on their health and wellbeing Melbourne, Australia: The Australian Psychological Society Ltd; 2018.
28. Cohen-Mansfield J, Hazan H, Lerman Y, Shalom V. Correlates and predictors of loneliness in older-adults: a review of quantitative results informed by qualitative insights. *Int Psychogeriatr* 2016;28:557-76.
29. Anderson CA. Attributional style, depression, and loneliness: A cross-cultural comparison of american and chinese students. *Pers Soc Psychol Bull* 1999;25:482-99.
30. Solano CH. Loneliness and perceptions of control: General traits versus specific attributions. *J Soc Behav Pers* 1987;2:201.

Running head: Review: Loneliness and substance dependence

31. Vanhalst J, Soenens B, Luyckx K, Van Petegem S, Weeks MS, Asher SR. Why do the lonely stay lonely? Chronically lonely adolescents' attributions and emotions in situations of social inclusion and exclusion. *J Pers Soc Psychol*. 2015;109:932-48.
32. Akerlind I, Hörnquist JO. Stability and change in feelings of loneliness: a two-year prospective longitudinal study of advanced alcohol abuse. *Scand J Psychol* 1989;30:102-12.
33. Johnson RA. Loneliness among hospitalized alcohol abusers: Exploration and treatment. US: ProQuest Information & Learning; 1985.
34. Valtorta NK, Kanaan M, Gilbody S, Hanratty B. Loneliness, social isolation and social relationships: what are we measuring? A novel framework for classifying and comparing tools. *BMJ Open* 2016;6:e010799.
35. Russell D, Peplau LA, Ferguson ML. Developing a measure of loneliness. *J Pers Assess* 1978;42:290-4.
36. Penning MJ, Liu G, Chou PHB. Measuring loneliness among middle-aged and older adults: The UCLA and de Jong Gierveld Loneliness Scales. *Social Indicators Research* 2014;118:1147-66.
37. Masi CM, Chen HY, Hawkey LC, Cacioppo JT. A meta-analysis of interventions to reduce loneliness. *Pers Soc Psychol Rev* 2011;15:219-66.
38. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 2009;6:e1000097.
39. National Heart Lung and Blood Institute NIH. Study Quality Assessment Tools: U.S. Department of Health and Human Services; 2019 [Available from: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>].

Running head: Review: Loneliness and substance dependence

- 1
- 2
- 3
- 4 40. Viswanathana M, Patnodeb CD, Berkmana ND, Bassc EB, Changd S, Hartlinge L, et
- 5
- 6 al. Recommendations for assessing the risk of bias in systematic reviews of health-care
- 7
- 8 interventions. *J Clin Epidemiol* 2018;97:26-34.
- 9
- 10
- 11 41. Wang Z, Taylor K, Allman-Farinelli M, Armstrong B, Askie L, Ghersi D, et al. A
- 12
- 13 systematic review: Tools for assessing methodological quality of human observational
- 14
- 15 studies. NHMRC; 2019.
- 16
- 17 42. Critical Appraisal Skills Programme. CASP Qualitative Checklist 2018 [Available
- 18
- 19 from: <https://casp-uk.net/casp-tools-checklists/>.
- 20
- 21
- 22 43. Jackson D, Turney R. Power analysis for random-effects meta-analysis. *Res Syn Meth*
- 23
- 24 2017;8:290-302.
- 25
- 26
- 27 44. van Assen MALM, van Aert RCM, Wicherts JM. Meta-Analysis Using Effect Size
- 28
- 29 Distributions of Only Statistically Significant Studies. *Psychol Methods* 2014;20:293-309.
- 30
- 31 45. Akerlind I, Hörnquist JO, Bjurulf P. Prognosis in alcoholic rehabilitation: the relative
- 32
- 33 significance of social, psychological, and medical factors. *Int J Addict* 1988;23:1171-95.
- 34
- 35 46. Akerlind I, Hörnquist JO, Hansson B. Loneliness correlates in advanced alcohol
- 36
- 37 abusers. I. Social factors and needs. *Scand J Soc Med.* 1987;15:175-83.
- 38
- 39
- 40 47. Allen HA, Peterson JS, Whipple S. Loneliness and alcoholism: a study of three
- 41
- 42 groups of male alcoholics. *Int J Addict* 1981;16:1255-8.
- 43
- 44
- 45 48. Armstrong JB. Loneliness and perceived stigmatization among older adults enrolled
- 46
- 47 in opiate substitution treatment programs and the utilization of mental health services. US:
- 48
- 49 ProQuest Information & Learning; 2016.
- 50
- 51
- 52 49. Boyles BR. How does Alcoholics Anonymous affect drinking outcomes? A grounded
- 53
- 54 theory perspective. US: ProQuest Information & Learning; 2018.
- 55
- 56 50. Britton PC, Conner KR. Reliability of the UCLA Loneliness Scale in opiate
- 57
- 58 dependent individuals. *J Pers Assess* 2007;88:368-71.
- 59
- 60

Running head: Review: Loneliness and substance dependence

- 1
2
3 51. Russell DW. UCLA Loneliness Scale (Version 3): Reliability, validity, and factor
4 structure. *J Pers Assess* 1996;66:20-40.
5
6
- 7 52. Cao Q, Liang Y. Perceived social support and life satisfaction in drug addicts: Self-
8 esteem and loneliness as mediators. *J Health Psychol* 2017 [Epub ahead of print].
9
- 10 53. Conner KR, Britton PC, Sworts LM, Joiner Jr TE. Suicide attempts among
11 individuals with opiate dependence: The critical role of belonging. *Addict Behav*
12 2007;32:1395-404.
13
14
- 15 54. Elton HL, Hörnquist JO. Abusers of alcohol granted disability pension: Prospective
16 longitudinal and multidisciplinary studies: Linköping University; 1983.
17
18
- 19 55. Hörnquist JO, Elton HF. A prospective longitudinal study of abusers of alcohol
20 granted disability pension. *Scand J Soc Med* 1983;11:91-6.
21
22
- 23 56. Essex EL, Petras D, Massat CR. Predictors of loneliness among court-involved and
24 substance abusing mothers. *Women Crim Justice* 2007;17:63-74.
25
26
- 27 57. Evans TJ. Sober and alone: A phenomenological exploration of the loneliness
28 experienced by recovering alcoholics. US: ProQuest Information & Learning; 2010.
29
30
- 31 58. Funk PE. A descriptive analysis of selected intrapersonal characteristics of drug
32 abusers. ProQuest Information & Learning, US; 1973. Retrieved from
33 [http://ezproxy.uow.edu.au/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=](http://ezproxy.uow.edu.au/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=psych&AN=1976-00695-001&site=ehost-live)
34 [psych&AN=1976-00695-001&site=ehost-live](http://ezproxy.uow.edu.au/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=psych&AN=1976-00695-001&site=ehost-live) Available from EBSCOhost psych database.
35 Accessed 16 April 2018.
36
37
- 38 59. Harris KS. The developmental effects of alcoholism in the adolescent. US: ProQuest
39 Information & Learning; 1983.
40
41
- 42 60. Hörnquist JO, Akerlind I. Loneliness correlates in advanced alcohol abusers. II.
43 Clinical and psychological factors. *Scand J Soc Med*, 1987;15:225-32.
44
45
- 46 61. Hörnquist JO, Hansson B, Akerlind I. The working capacity of the alcohol abuser.
47 Prognostic multiple regression analyses. *Scand J Soc Med* 1988;16:27-33.
48
49
50
51
52
53
54
55
56
57
58
59
60

Running head: Review: Loneliness and substance dependence

62. Hosseinbor M, Yassini Ardekani SM, Bakhshani S, Bakhshani S. Emotional and social loneliness in individuals with and without substance dependence disorder. *Int J High Risk Behav Addict* 2014;3:e22688-e.
63. Itzick M, Segal JN, Possick C. Relationships in the lives of Israeli women coping with drug addiction: An ecosystemic perspective. *J Soc Pers Relat* 2019;36:741-60.
64. Kuerbis A, Mereish EH, Hayes M, Davis CM, Sijing S, Morgenstern J, et al. Testing cross-sectional and prospective mediators of internalized heterosexism on heavy drinking, alcohol problems, and psychological distress among heavy drinking men who have sex with men. *J Stud Alcohol Drugs* 2017;78:113-23.
65. Kuerbis A, Padovano HT, Shao SJ, Houser J, Muench FJ, Morgenstern J. Comparing daily drivers of problem drinking among older and younger adults: An electronic daily diary study using smartphones. *Drug Alcohol Depend* 2018;183:240-6.
66. Li F, Xu Y-M, Zhu J-H, Lu J, Zhong B-L. Pain of methadone-maintained heroin addicts: lonelier individuals feel more intense pain. *Oncotarget* 2017;8:79948-52.
67. Li H-J, Zhong B-L, Xu Y-M, Zhu J-H, Lu J. Sleep in lonely heroin-dependent patients receiving methadone maintenance treatment: longer sleep latency, shorter sleep duration, lower sleep efficiency, and poorer sleep quality. *Oncotarget* 2017;8:89278-83.
68. Medora NP. Variables affecting loneliness among individuals undergoing treatment in alcohol rehabilitation centers. US: ProQuest Information & Learning; 1983.
69. Medora NP, Woodward JC. Loneliness and alcoholism. *Wellness Perspectives* 1990;6:42.
70. Medora NP, Woodward JC. Factors associated with loneliness among alcoholic in rehabilitation centers. *J Soc Psychol* 1991;131:769-79.
71. Michaels AW. The prevalence of loneliness in alcoholic versus non-alcoholic treatment populations. ProQuest Information & Learning, US; 1982. Retrieved from <http://ezproxy.uow.edu.au/login?url=http://search.ebscohost.com/login.aspx?direct=true&db>

Running head: Review: Loneliness and substance dependence

[=psych&AN=1983-51365-001&site=ehost-live](#) Available from EBSCOhost psych database.

72. Neale J, Tompkins CNE, Strang J. Qualitative exploration of relationships between peers in residential addiction treatment. *Health Soc Care Community* 2018;26:e39-e46.
73. Nerviano VJ, Gross WF. Loneliness and locus of control for alcoholic males: Validity against Murray need and Cattell trait dimensions. *J Clin Psychol* 1976;32:479-84.
74. Newton TF, De La Garza R, Kalechstein AD, Tziortzis D, Jacobsen CA. Theories of addiction: methamphetamine users' explanations for continuing drug use and relapse. *Am J Addict* 2009;18:294-300.
75. Perodeau GM, du Fort GG. Psychotropic drug use and the relation between social support, life events, and mental health in the elderly. *J Appl Gerontol* 2000;19:23-41.
76. Price RH, Curlee-Salisbury J. Patient-treatment interactions among alcoholics. *J Stud Alcohol* 1975;36:659-69.
77. Schmidt DR. An ethnographic process evaluation of a faith-based therapeutic community for chemically addicted men: Trinity Evangelical Divinity School; 2002.
78. Schonfeld L, Dupree LW, Rohrer GE. Age-specific differences between younger and older alcohol abusers. *J Clin Geropsychol* 1995;1:219-27.
79. Siddique F, Ahmad Mann A, Ali T. Influence of social factors on drug use behavior in Punjab, Pakistan. *Pak J Nutr* 2012;11:1099-100.
80. Van Hasselt VB, Null JA, Kempton T, Bukstein OG. Social skills and depression in adolescent substance abusers. *Addict Behav* 1993;18:9-18.
81. Yang YJ, Xu YM, Chen WC, Zhu JH, Lu J, Zhong BL. Loneliness and its impact on quality of life in Chinese heroin-dependent patients receiving methadone maintenance treatment. *Oncotarget* 2017;8:79803-8.
82. Yeh M-C. Loneliness, alcohol and marijuana use among male college students. US: ProQuest Information & Learning; 2002.

Running head: Review: Loneliness and substance dependence

- 1
2
3 83. Zhong B-L, Xu Y-M, Zhu J-H, Liu X-J. Non-suicidal self-injury in Chinese heroin-
4
5 dependent patients receiving methadone maintenance treatment: Prevalence and associated
6
7 factors. *Drug Alcohol Depend* 2018;189:161-5.
8
9
10 84. Lancaster K, Seear K, Ritter A. Monograph 26: Reducing stigma and discrimination
11
12 for people experiencing problematic alcohol and other drug use. National Drug and Alcohol
13
14 Research Centre Sydney: National Drug and Alcohol Research Centre; 2018.
15
16
17 85. Best D, Beckwith M, Haslam C, Haslam SA, Jetten J, Mawson E, et al. Overcoming
18
19 alcohol and other drug addiction as a process of social identity transition: the social identity
20
21 model of recovery (SIMOR). *Addict Res Theory* 2016;24:111-23.
22
23
24 86. Levin I, Stokes JP. An examination of the relation of individual difference variables
25
26 to loneliness. *J Pers* 1986;54:717-33.
27
28
29 87. de Jong-Gierveld J. Developing and testing a model of loneliness. *J Pers Soc Psychol*
30
31 1987;53:119-28.
32
33
34 88. Perlman D, Peplau LA. Toward a Social Psychology of Loneliness. In: R. Gilmour,
35
36 Duck S, editors. *Personal Relationships: 3 Relationships in Disorder*. London: Academic
37
38 Press; 1981. p. 31-56.
39
40
41 89. Russell D, Peplau LA, Cutrona CE. The revised UCLA loneliness scale: Concurrent
42
43 and discriminant validity evidence. *J Pers Soc Psychol* 1980;39:472-80.
44
45
46 90. Hughes ME, Waite LJ, Hawkley LC, Cacioppo JT. A short scale for measuring
47
48 loneliness in large surveys: Results from two population-based studies. *Res Aging*
49
50 2004;26:655-72.
51
52
53 91. Ditommaso E, Brannen C, Best LA. Measurement and validity characteristics of the
54
55 short version of the social and emotional loneliness scale for adults. *Educ Psychol Meas*
56
57 2004;64:99-119.
58
59
60

Running head: Review: Loneliness and substance dependence

92. Jowker B. Psychometric properties of the short form of social and emotional loneliness scale for adults (SELSA-S). *J Behav Sci* 2012;5:311-7.
93. Bradley R. Measuring loneliness. Unpublished doctoral dissertation: Washington State University; 1969.
94. Belcher MJ. Loneliness: A review, theoretical framework, and its therapeutic implications. 1971.
95. Belcher MJ. The measurement of loneliness: A validation of the Belcher Extended Loneliness Scale (BELS). *Dissertation Abstracts International: Illinois Institute of Technology*; 1973.
96. Sisenwein RJ. Loneliness and the individual as viewed by himself and others. Unpublished doctoral dissertation: Columbia University, New York; 1964.
97. Elton M, Hörnquist JO. Grounds for disability pension: Younger abusers of alcohol as compared with older ones. *Scand J Soc Med* 1983;11:53-8.
98. Woodward JC. Loneliness and Solitude: Phenomena, Incidence and Factorial Relationships.: University of Nebraska-Lincoln; 1967 - Delete?.
99. Woodward J. The solitude of loneliness. Lexington, MA: Lexington Books; 1988.
100. Fillenbaum C, Dellinger D, Maddox G, Pfeiffer E. Assessment of individual functional status in a program evaluation and resource allocation model. In: *Development CftSoAaH*, editor. *Multidimensional functional assessment: The OARS Methodology*. Durham, NC: Duke University; 1978. p. 3-12.
101. Harel Z, Deimling GT. Social resources and mental health: An empirical refinement. *J Gerontol* 1984;39:747-52.
102. Marlatt GA, Gordon JR. Determinants of relapse: Implications for the maintenance of behavior change. In: Davidson PO, Davidson SM, editors. *Behavioral medicine: Changing health lifestyles*. New York: Brunner/Mazel; 1980. p. 1410-52.

Running head: Review: Loneliness and substance dependence

103. Marlatt GA, Gordon JR. Relapse prevention: Maintenance strategies in the treatment of addictive behaviors. New York: Guilford Press; 1985.
104. Asher SR, Hymel S, Renshaw PD. Loneliness in Children. *Child Dev* 1984;55:1456-64.
105. Cicchetti DV. Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychol Assess* 1994;6:284.
106. Carstensen LL. Social and emotional patterns in adulthood: support for socioemotional selectivity theory. *Psychol Aging* 1992;7:331-8.
107. Cacioppo S, Grippo AJ, London S, Goossens L, Cacioppo JT. Loneliness: Clinical import and interventions. *Perspect Psychol Sci* 2015;10:238-49.
108. Cohen-Mansfield J, Perach R. Interventions for alleviating loneliness among older persons: a critical review. *Am J Health Promot* 2015;29:e109-e25.
109. Maes M, Klimstra T, Van den Noortgate W, Goossens L. Factor structure and measurement invariance of a multidimensional loneliness scale: Comparisons across gender and age. *J Child Fam Stud* 2014;24:1829-37.
110. Heinrich LM, Gullone E. The clinical significance of loneliness: A literature review. *Clin Psychol Rev* 2006;26:695-718.
111. de Jong Gierveld J, van Tilburg T, Dykstra P. Loneliness and social isolation: New ways of theorizing and conducting research. In: Vangelisti AL, Perlman D, editors. *The Cambridge Handbook of Personal Relationships*. 2 ed. Cambridge, England: Cambridge University Press; 2018. p. 391-404.
112. Shiovitz-Ezra S, Ayalon L. Use of direct versus indirect approaches to measure loneliness in later life. *Res Aging*. 2012;34:572-91.

Running head: Review: Loneliness and substance dependence

- 1
2
3 113. Russell D. The Measurement of Loneliness. In Peplau L, Perlman D, editors.
4 Loneliness: A Sourcebook of Current Theory, Research and Therapy. United States: Wiley;
5
6 1982. p. 81-104.
7
8 114. Solano CH. Two measures of loneliness: A comparison. Psychol Rep 1980;46:23-8.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only




Figure 1. PRISMA Flow diagram. SUD, substance use disorder.

Table 1. *Study characteristics*

Article	Objectives	Sample size and sociodemographic	Methods	Outcome	Quality rating
Akerlind, Hornquist, Bjurulf, 1988 [45]	To determine the importance of social, psychological, and medical factors in prediction of post-treatment functioning. The outcome measure, longitudinal working capacity, was assumed to reflect overall functional capacity.	N: 34 Response rate: Not reported Country: Sweden Gender: Male Age: 24-60 ($M=41.5$, $SD=10.5$) Substance use: Alcohol	Design: Longitudinal Setting: Community/ Inpatient alcoholism treatment service/ Sociomedical outpatient clinic/ Psychiatric outpatient clinic/ Vocational rehabilitation service. Year: 1978 and 1979 Procedure: The participants were classified into three groups. The subjects followed the regular treatment programs at the different settings and completed the assessment measures.	<ul style="list-style-type: none"> Loneliness appeared as the most important factor for working capacity in the stepwise multiple regression analysis of the data. Rehabilitation of advanced alcoholics, normal psychological mechanisms related to perceived well-being and quality of life play an important part. 	Fair
Akerlind I, Hornquist JO, Hansson B, 1987 [46]	To determine whether loneliness correlates with quality of life (comprised of six domains: structural, material, social, activity, psychological, physical).	N: 95 (Group 1: 54 males, 7 females; Group 2: 34 males) Response rate: Not reported Country: Sweden Gender: Males and Females Age: 24-60 ($M=44.5$, $SD=10.2$) Substance use: Alcohol	Design: Longitudinal Setting: Applicants for disability pension in Ostergotland, Sweden Year: 1978-1981 Procedure: Participants were contacted and examined on two occasions. Participants self-rated their loneliness. Correlations between loneliness scores and participant's social background (12 parameters), external social network (20 parameters), work and activities (27 parameters), societal position (9 parameters), life priorities (29 parameters) and life satisfaction (25 parameters) were calculated.	<ul style="list-style-type: none"> No results presented or comment made about whether loneliness changes over time. Quantitative aspects of social network was not found to be related to loneliness. Dissatisfaction with quality of relationships found to relate to loneliness. Loneliness related to quality of life. Age was negatively related to loneliness; Females were lonelier; Income, education & occupation were not related to loneliness; Residence in an urban area was related to loneliness; Difficulties in the current administration of one's economy were related to loneliness; insecurity and inferiority was related to loneliness. 	Poor
Akerlind I, Hornquist JO, 1989 [32]	To analyse covariations between longitudinal changes in loneliness and changes in social network, psychological wellbeing, life satisfaction, activities, adaption to work and non-work situation,	N: 95 (88 males, 7 females) Response rate: Not reported	Design: Cross-sectional and Longitudinal Setting: Community/inpatient alcoholism treatment service, sociomedical outpatient clinic, psychiatric outpatient clinic, vocational rehabilitation service.	<ul style="list-style-type: none"> <i>Cross-sectional:</i> Loneliness correlated with wellbeing, life satisfaction, psychiatry (anxiety, autonomic disturbance, muscular tension, reduced sleep, passivity, global rating of illness), activity (sports and recreation, shopping), adaptation (meaningfully occupied in free time), social network (number and availability of friends and acquaintances) and 	Fair

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

	psychiatric symptoms, and alcohol consumption.	<p>Country: Sweden</p> <p>Gender: Males and Females</p> <p>Age: 24-60 years ($M=44.5, SD=10.2$)</p> <p>Substance use: Alcohol</p>	<p>Year: 1978 and 1979</p> <p>Procedure: Participants were examined on two occasions initially and re-examined after two years. Participant's self-rated loneliness. Other variables assessed via structured interview by a social worker and a semi-structured interview conducted by a psychiatrist.</p>	<p>alcohol consumption at second assessment only.</p> <ul style="list-style-type: none"> • <i>Longitudinal:</i> Change in loneliness over two years was accompanied by changes in wellbeing (indolence, inferiority, perceived negative treatment from others), psychiatry (sadness, inability to feel, pessimistic thoughts, suicidal thoughts, autonomic disturbance), life satisfaction, activity (time spent in sports and rec, resting and relaxation), adaptation (meaningfully occupied in free time) social network (availability of friends and acquaintances, availability of close friends). • Loneliness related to perceived quality and satisfaction with relationships, not to quantity of relationships. • Loneliness not longitudinally related to alcohol consumption. • Psychological wellbeing (particularly indolence and self-evaluation in relation to others) most salient correlate with loneliness and changes to loneliness. • At both examinations a great minority of individuals (45-35%) scored at the upper half of the scale, indicating salient feelings of loneliness. 	
Allen HA, Peterson JS, Whipple S, 1981 [47]	To investigate the relationship between alcoholism and loneliness at various stages of recovery. It was hypothesised that people who use alcohol in the acute group would be lonelier than those in the chronic group, who would in turn be lonelier than the recovering group.	<p>N: 45 (all males)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male</p> <p>Age: Acute Group (detoxification) $M=45.1$; Chronic Group (outpatient treatment) $M=32.3$; Recovering Group (members of AA) $M=41$</p> <p>Substance use: Alcohol</p>	<p>Design: Cross-sectional</p> <p>Setting: Outpatient mental health treatment service.</p> <p>Year: Not reported</p> <p>Procedure: Participants were volunteers who had had contact with the mental health centre. The loneliness scale was given within 3 days of entry to the program.</p>	<ul style="list-style-type: none"> • Differences exist within this population, with those currently drinking (acute group) most lonely, those with 1 year's sobriety (members of AA) the least lonely, and those in treatment (chronic group) falling in the middle. 	Poor

Armstrong JB, 2016 [48]	To examine the role that loneliness and perceived stigmatisation play in the decision to seek mental health services among older adults enrolled in opiate substitution treatment.	<p>N: 94</p> <p>Response rate: 98%</p> <p>Country: America</p> <p>Gender: Not reported</p> <p>Age: 50-71 ($M=57.22$, $SD=5.13$)</p> <p>Substance use: Opioids</p>	<p>Design: Cross-sectional</p> <p>Setting: Outpatient substance dependence treatment service</p> <p>Year: Not reported</p> <p>Procedure: Participants were actively enrolled in the opiate substitution treatment program at the service. Opportunity to volunteer to participate was offered by their regular chemical dependency counsellors. Clients were offered a \$3 McDonald's gift card as an incentive. The counsellors administered the surveys at the beginning of the monthly session.</p>	<ul style="list-style-type: none"> Participants who reported utilising more available mental health services also tended to indicate higher degrees of both loneliness and perceived stigmatisation. 	Fair
Boyles BR, 2018 [49]	<p><i>Study One:</i> To inform an emic understanding of change processes embedded in AA's literature that may explain continuous abstinence experienced by community AA members.</p> <p><i>Study Two:</i> To identify AA change processes and their functioning within AA's fellowship in order to inform a guiding theory that details how continuous abstinence occurs for AA members.</p>	<p>N: <i>Study One:</i> $N=3$ (3 males); <i>Study Two:</i> $N=5$ (4 males, 1 female)</p> <p>Response rate: <i>Study One:</i> 33% (9 invited to participate, of the 7 that accepted, 2 withdrew, 1 did not submit data, 1 non-adherent); <i>Study Two:</i> 50% (10 invited, 7 accepted, 2 of these withdrew)</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: <i>Study One:</i> 30-59 years; <i>Study Two:</i> 48-69 years</p> <p>Substance use: Alcohol and opioids</p>	<p>Design: Cross-sectional, qualitative</p> <p>Setting: Members of AA</p> <p>Year: 2016</p> <p>Procedure: <i>Study One:</i> This study uses qualitative content analysis of the AA literature and a grounded theory approach to AA's fellowship. <i>Study Two:</i> The qualitative data were collected through focus groups to inform an understanding of how change occurs within AA. Six focus groups were conducted 1-2 weeks apart.</p>	<p><i>Study One:</i></p> <ul style="list-style-type: none"> Only one of the passages identified by the content analysis suggests a change in loneliness (e.g. "we can be alone at perfect peace and ease", p.75). A keyword search of the AA text indicates loneliness is discussed at least 12 times. <p><i>Study Two:</i></p> <ul style="list-style-type: none"> Grounded theory analysis of focus group data suggests that AA involvement produced changes in insecurity, loneliness (marked by a sense of belonging), life meaning, anxiety and shame. An early stage of the AA experience, reported by the participants, included a reduction in loneliness. In middle stage of AA, the participants reported becoming aware of an internal or intrapersonal loneliness. The focus group participants indicated problems with loneliness were not fully resolved until late AA. 	Good
Britton PC, Conner KR, 2007	To examine the internal consistency and test-retest	<p>N: 121 (61 females, 60</p>	<p>Design: Longitudinal</p>	<ul style="list-style-type: none"> The internal consistency (0.87) and test-retest reliability ($r=0.77$) of the UCLA-LS in a 	Fair

[50]	reliability of the self-report UCLA-LS (Russell 1996 [51]) in methadone maintenance patients at an urban university hospital.	<p>males)</p> <p>Response rate: 96% baseline ($n=117$); 57% follow-up ($n=67$)</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: 21-59 years ($M=41.9$, $SD=9.7$)</p> <p>Substance use: Opioids</p>	<p>Setting: Outpatient substance dependence treatment service</p> <p>Year: 2005</p> <p>Procedure: Participants recruited through poster advertisements at methadone clinic. Interviewer met with participants for 1 hour to complete battery of self-report and interviewer-based instruments. Participants invited to return after 14 days for a follow-up session. Participants paid a \$30 grocery gift card following each assessment.</p>	<p>clinical sample of individuals with opiate dependence in MMT were comparable to those obtained from the UCLA-LS in other samples (Russell 1996 [51]).</p> <ul style="list-style-type: none"> No differences in mean loneliness scores (i.e. male vs. female, African American vs. White, levels of education). No differences in loneliness scores from first to second administration of the UCLA-LS. 	
Cao Q, Liang Y, 2017 [52]	To test the mediating effect of self-esteem and loneliness on the relationship between social support and life satisfaction in people who use drugs, and to investigate the contribution of each specific mediator variable.	<p>N: 110 (91 males, 19 females)</p> <p>Response rate: 84.6% (110 of 130 distributed)</p> <p>Country: China</p> <p>Gender: Male and female</p> <p>Age: 18-54 years ($M=38.47$, $SD=7.31$)</p> <p>Substance use: Heroin: 80.7% ($n = 88$); Methamphetamine: 8.2% ($n = 9$); Cocaine: 4.5% ($n = 5$); Marijuana: 3.3% ($n = 4$); Other: 3.3% ($n = 4$)</p>	<p>Design: Cross-sectional</p> <p>Setting: Guangdong Fangcun Brain Hospital in China.</p> <p>Year: Not reported</p> <p>Procedure: Participants completed the questionnaires in a waiting room of the hospital. Instruments took approximately 30 minutes to complete. Confirmatory Factor analysis conducted.</p>	<ul style="list-style-type: none"> Perceived social support was positively related to self-esteem and life satisfaction, and had a negative correlation with loneliness Although higher perceived social support could increase self-esteem, it could not reduce loneliness, which might weaken life satisfaction. Self-esteem and loneliness played a mediating role in the relationship between social support and life satisfaction. 	Fair
Conner KR, Britton PC, Sworts LM, Joiner TE, 2007 [53]	<p>To determine if low belonging, high burdensomeness, and high loneliness are associated with attempted suicide among individuals with opiate dependence.</p> <p>To explore whether or not</p>	<p>N: 131 (69 females, 62 males)</p> <p>Response rate: Not reported</p> <p>Country: America</p>	<p>Design: Cross-sectional</p> <p>Setting: Outpatient substance dependence treatment service</p> <p>Year: 2005</p> <p>Procedure: Participants recruited through poster</p>	<ul style="list-style-type: none"> Moderate difference in loneliness between suicide attempters and non-attempters. Higher scores on loneliness scale were associated with a higher probability of an attempt. However, after adjusting for covariates, higher scores on loneliness were not statistically associated with a higher probability of an attempt. Small differences in loneliness between 	Fair

	unintentional overdose is also associated with perceived belonging, burdensomeness, and loneliness.	<p>Gender: Male and female</p> <p>Age: 21-59 years ($M=41.8$, $SD=9.6$)</p> <p>Substance use: Opioids</p>	<p>advertisements at methadone clinic. Interviewer met with participants for 1 hour to complete battery of self-report and interviewer-based instruments. Participants invited to return after 14 days for a follow-up session. Participants paid a \$30 gift card following each assessment.</p>	<p>overdose and non-overdose subjects.</p> <ul style="list-style-type: none"> With the exception of the unadjusted analysis, the results did not support an association of loneliness and suicidal behaviour. 	
<p>Elton HL, Hornquist JO, 1983 [54]</p> <p>NB. Thesis</p>	<p>Aimed to compare alcohol users with non-users on variables (such as loneliness) prior to and after receiving their disability pension. These groups were also compared to a group of users (in rehabilitation) who were not receiving a pension.</p> <p><i>*NB. Thesis contained 6 studies – all with same sample – only studies 2, 3, and 6 of this thesis were included in the review.</i></p> <p><i>Study 5 was included in the review as a separate published journal article (Hornquist & Elton, 1983 [55])</i></p> <p><i>Studies 1 and 4 did not meet inclusion criteria for the review.</i></p>	<p>N: 78 (pensioned users; 54 males, 7 females)</p> <p><i>Matched reference groups:</i> 27 (pensioned non-users); 30 (non-pensioned users undergoing rehabilitation)</p> <p>Response rate: 82% ($n=61$) at baseline; 92% ($n=56$) at 2-years (Study 6 only)</p> <p>Country: Sweden</p> <p>Gender: Male and female</p> <p>Age: $M=46$ years</p> <p>Substance use: Alcohol</p>	<p>Design: Cross-sectional and longitudinal</p> <p>Setting: People who use alcohol in the community accessing disability pension; and a matched sample of people who use alcohol accessing rehabilitation (inpatient or outpatient unknown)</p> <p>Year: 1 January 1978-1 July 1979</p> <p>Procedure:</p> <p><i>Study 2:</i> Cross-sectional interviews.</p> <p><i>Study 3:</i> Cross-sectional interviews same as Study 2. Sample divided into older and younger subgroups to make comparisons.</p> <p><i>Study 6:</i> Longitudinal. Interviews conducted prior to receiving a pension and again after two-years.</p>	<ul style="list-style-type: none"> <i>Study 2:</i> Mean loneliness scores were found to be different amongst pensioned people who use alcohol compared to the matched reference groups (pensioned people who do not use alcohol / non-pensioned rehabilitators). Pensioned group was found to have higher levels of loneliness compared to the matched samples. <i>Study 3:</i> Younger people who use alcohol found to be lonelier than older people who use alcohol. <i>Study 6:</i> Loneliness amongst pensioned people who use alcohol contributed to changes (from baseline to two-year follow-up) to wellbeing, psychiatric status, need satisfaction, intellectual performance, and alcohol consumption. 	<p>Study 2: Poor</p> <p>Study 3: Poor</p> <p>Study 6: Fair</p>

Essex EL, Petras, D, Massat CR, 2007 [56]	To determine what predicts loneliness for substance using, court-involved mothers.	<p>N: 94 (all females)</p> <p>Response rate: 94%</p> <p>Country: America</p> <p>Gender: Female</p> <p>Age: 19-50 ($M=35.4$, $SD=6.3$)</p> <p>Substance use: Heroin (66%)</p> <p>Cocaine (22%)</p> <p>Other (12%)</p>	<p>Design: Cross-sectional</p> <p>Setting: Adults who were substance involved and convicted of a criminal offense and ordered to TASC.</p> <p>Year: Pooled samples from 2000 and 2004-2005</p> <p>Procedure: Potential subjects were referred to the researchers by the Chicago metropolitan area offices of TASC. Flyers and letters describing the study were distributed to women who met eligibility criteria. Subjects completed 1.5-hour structured interviews. Data was pooled from two cross-sectional exploratory studies.</p>	<ul style="list-style-type: none"> • Found support for viewing loneliness in this population as stemming from: <ul style="list-style-type: none"> (i) Characteristics of the women themselves (the presence or absence of co-occurring conditions); (ii) Child characteristics (number of minors in the home); (iii) Partner relations (degree of domestic violence experienced in their relations with partners); and (iv) Informal and formal social supports. • Mothers' satisfaction with substance use services was found to be a negative predictor of loneliness. • Level of informal social support had strongest relationship to loneliness. 	Fair
Evans TJ, 2010 [57]	<p>To understand loneliness, as experienced by recovering alcoholics.</p> <p>To explore what factors recovering alcoholics identified as contributing to their experience of loneliness.</p>	<p>N: 8 (5 males, 3 females)</p> <p>Response rate: N/A</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: Not reported</p> <p>Substance use: Alcohol</p>	<p>Design: Cross-sectional/ Qualitative</p> <p>Setting: AA meetings</p> <p>Year: 2009</p> <p>Procedure: Participants were recruited from local AA meetings. Potential participants were asked to take a short self-report survey. Face to face in-depth interviews then completed and lasted 1.5-2 hours each. Interviews were tape recorded and transcribed verbatim.</p>	<ul style="list-style-type: none"> • Loneliness described to include feelings of: fearful, empty, hopeless, overwhelmed, misunderstood, suicidal, isolated from others, and alone in a crowd. • Participants attributed their experience with loneliness during recovery to a variety of factors, including severed or strained relationships, the inability to trust, a history of insecure or inept parental attachment, and the re-occurrence of negative thoughts. 	Good
Funk PE, 1973 [58]	To describe six intrapersonal characteristics and their possible relationship to drug use. The six intrapersonal characteristics were: anxiety, loneliness, affection, guilt, punishment and frustration.	<p>N: 35 (23 females, 12 males)</p> <p>Response rate: 81% (8 removed by researchers)</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: 15-55 years</p>	<p>Design: Cross-sectional/qualitative</p> <p>Setting: Volunteer who used drugs residing in the community</p> <p>Year: Not reported</p> <p>Procedure: Participants were invited to the study through counsellors at the Gallatin County Help Centre, teachers, and friends, as well as newspaper classified ads. Data collected via audio tapes and client self-perceptions. Clients</p>	<ul style="list-style-type: none"> • Loneliness, anxiety, affection, frustration, punishment, and guilt were found to be present in some form and to some degree in nearly every individual studied. • A lack of strong attachments and feelings of alienation seemed to characterise the lives of some of the individuals studied. • Loneliness and anxiety seemed most pronounced and were most obvious to the investigator. • Researcher concludes that loneliness could very likely be a cause of drug abuse. 	Poor

		Substance use: Not reported ("drug users")	completed the interviews at the counselling laboratory in Montana State University.		
Harris KS, 1983 [59]	To examine how alcoholism related to development in the areas of egocentrism, ego identity and intimacy.	N: 72; <i>Alcoholic group:</i> 26 (13 male, 13 female); <i>Non-alcoholic group:</i> 26 (13 male, 13 female); <i>Recovered group:</i> 20 (9 male, 11 female) Response rate: Not reported Country: America Gender: Male and female Age: 18-27; <i>Alcoholic group</i> (M=19.81); <i>Non-alcoholic group</i> (M=20); <i>Recovered group</i> (M=22.55) Substance use: Alcohol	Design: Cross-sectional Setting: University classes ('Alcoholic ' and 'Non-alcoholic' groups); AA meetings (Recovered group) Year: Not reported Procedure: Participants invited at university class and questionnaire packs distributed. Recovered group approached at AA meetings and completed same questionnaire pack (2 participants from recovered group were recruited at the university)	<ul style="list-style-type: none"> No difference in loneliness across groups (non-alcoholic, alcoholic, or recovered alcoholic). No differences in loneliness across gender. Concluded that in general, loneliness did not seem to be a factor in the developmental delay predicted for the alcoholic group. 	Fair
Hornquist JO, Akerlind I, 1987 [60]	To extend a preceding correlational analysis of loneliness to clinical and psychological parameters in a sample of 95 alcohol users.	N: 95 (88 males, 7 females) Sample 1: 61 (54 males, 7 females) Sample 2: 34 (all males) Response rate: 82% Country: Sweden Gender: Male and female Age: 24-60 (M=44.5, SD=10.2) Substance use: Alcohol	Design: Longitudinal Setting: Community/Inpatient alcoholism treatment service, Sociomedical outpatient clinic, Psychiatric outpatient clinic, Vocational rehabilitation service. Year: 1978 and 1979 cross sectional data collected (then again at two-year interval) Procedure: Participants were first-time applicants for disability pension. Participants were examined twice with an approximate interval of two years. Self-rating scales were used as well as broader medical-psychological examination, including clinical methods such as mental test, interview, judgment by psychiatrist and biochemical test.	<p>Intellectual ability:</p> <ul style="list-style-type: none"> Achievement on mental tests was not related to feelings of loneliness. <p>Wellbeing:</p> <ul style="list-style-type: none"> Loneliness found to be related to indolence, life dissatisfaction, inferiority, self-esteem and negative ego-concept. A perception of ill treatment from others was one of the strongest correlates of loneliness in the study. <p>Psychiatric symptoms:</p> <ul style="list-style-type: none"> Suicidal thoughts, sadness or depression, autonomic disturbances, anxiety or tension, emotional inhibition, lassitude, indecision, sexual and sleep disturbances were stable correlates with loneliness. <p>Characteristics of alcohol use:</p> <ul style="list-style-type: none"> Loneliness was related to higher alcohol consumption, delirium, and blackouts. 	Fair

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

<p>Hornquist JO, Elton HF, 1983 [55]</p> <p>NB. Also in thesis</p>	<p>To determine what changes in quality of life occur when people who use alcohol are granted disability pension.</p>	<p>N: Pensioned people who use alcohol=78; Pensioned people who do not use alcohol=27; Rehabilitators=30</p> <p>Response rate: <i>Pensioned alcohol group:</i> 78% at baseline, 92% at follow-up <i>Pensioned non-alcohol group:</i> 85% at follow-up <i>Rehabilitators:</i> 97% at 2-year follow-up.</p> <p>Country: Sweden</p> <p>Gender: Male and female</p> <p>Age: 20-60 years</p> <p>Substance use: Alcohol</p>	<p>Design: Longitudinal</p> <p>Setting: Community/ Inpatient alcoholism treatment service/ Sociomedical outpatient clinic/ Psychiatric outpatient clinic/ Vocational rehabilitation service.</p> <p>Not clearly stated where “pensioned users” group was recruited.</p> <p>Year: 1978-1982</p> <p>Procedure: Participants were people who had applied for disability pension for the first time. Three groups determined: (i) pensioned alcohol use group; (ii) pensioned non-alcohol use group; (iii) rehabilitators recruited from either: inpatient treatment, accessing treatment at socio-medical clinic, psychiatric clinic, vocational rehabilitation). Groups were matched for age, sex, and type of pension granted (e.g. back issues, heart disease). Participants' background, social network, daily activities, material situation and structured situation, abuse of alcohol, psychiatric status, wellbeing, mental tests and laboratory studies were investigated at both time points.</p>	<ul style="list-style-type: none"> Loneliness was related to having a relative with an alcohol use problem. Feelings of loneliness in pensioned alcohol using group diminished over two-year period. Not reported whether feelings of loneliness also reduced for the “pensioned non-users” and “rehabilitators” groups. This study does not report where “pensioned users” were recruited from/ if they were accessing treatment across this two-year interval. As such, comments about the potential effects of substance dependence treatment on reduced loneliness cannot be determined. Functional disability of pensioned alcohol using group does not change, yet general wellbeing and psychiatric status does change over two-year period. 	<p>Fair</p>
<p>Hörnquist JO, Hansson B, Akerlind I, 1988 [61]</p>	<p>To determine variables predictive of a regained working capacity in alcohol users undergoing rehabilitation. Variables included: psychological, attitudinal, behavioural, use characteristics, psychiatric and biochemical variables.</p>	<p>N: 34 (all male)</p> <p>Response rate: Not reported</p> <p>Country: Sweden</p> <p>Gender: Male</p> <p>Age: 24-60 (<i>M</i>=41.5, <i>SD</i>=10.5)</p> <p>Substance use: Alcohol</p>	<p>Design: Longitudinal</p> <p>Setting: Inpatient alcoholism treatment service/ Sociomedical outpatient clinic/ Psychiatric outpatient clinic/ Vocational rehabilitation service.</p> <p>Year: Baseline data collected 1978 and 1979 then followed up after two-years</p> <p>Procedure: The people who use alcohol were not being given any special method of treatment within the frame of this study but followed the ordinary routines at their different institutes of</p>	<ul style="list-style-type: none"> Loneliness was the most significant predictor of the subsequent vocational situation of people who use alcohol. Possibility of a better vocational outcome was strengthened when the individual had no drinking buddies and did not feel lonely. The user who actually is alone but does not feel lonely seems to be best equipped for rehabilitation. 	<p>Fair</p>

			treatment. Self-rating scales were completed at both time points.		
Hosseinbor M, Yassini ASM, Bakhshani S, Bakhshani S, 2014 [62]	To assess emotional, social, romantic, and familial dimensions of loneliness in people who use drugs and people who do not use drugs.	<p>N: 228 (152 males, 76 females)</p> <p>Response rate: Not reported</p> <p>Country: Iran</p> <p>Gender: Male and female</p> <p>Age: <i>Drug use group:</i> 16-55 ($M = 29.43$, $SD = 7.83$); <i>Non-drug use group:</i> 17-50 ($M = 27.99$, $SD = 8.54$)</p> <p>Substance use: Narcotics</p>	<p>Design: Cross-sectional</p> <p>Setting: Inpatient mental health treatment service</p> <p>Year: Not reported</p> <p>Procedure: Drug use group were enrolled through random sampling method, and individuals without history of using narcotic drugs were selected through available sampling method from drug use groups companions, students or staff of the Baharan Psychiatric Hospital as non-drug use group. All participants completed a clinical interview and the loneliness scale.</p>	<ul style="list-style-type: none"> Individuals diagnosed with substance dependency scored higher on the romantic, family, social, and emotional subscales of SELSA than those of individuals without substance dependency. No significant difference between substance dependent men and women on loneliness scores Significant difference between scores of non-dependent men and women on romantic subscale. Social and emotional feelings of loneliness deemed to be a high-risk factor for initiation of drug use and its maintenance 	Fair
Ingram I, Kelly PJ, Deane FP, Baker AL, Raftery DK, 2018 [22]	To validate the SELSA-S measure of loneliness for use in substance dependent treatment populations. The study also aimed to determine which demographic and physical and mental health variables were related to loneliness.	<p>N: 346</p> <p>Response rate: 91%</p> <p>Country: Australia</p> <p>Gender: Male and female</p> <p>Age: $M=37.71$, $SD=9.85$</p> <p>Substance use: Alcohol and other drugs</p>	<p>Design: Cross-sectional</p> <p>Setting: Inpatient substance dependence treatment service</p> <p>Year: 2017</p> <p>Procedure: At each service, all residents were invited to attend a meeting where the study was explained. Completion of the anonymous surveys indicated tacit consent was given. Surveys took 30 minutes to complete.</p>	<ul style="list-style-type: none"> Further research needed to determine the validity of the SELSA-S measure for use with substance-dependent populations. Substance-dependent populations experience higher rates of loneliness compared with the general population. Frequency of loneliness was 79%, 69% reported loneliness to be problematic. Loneliness appeared to be primarily experienced in the form of romantic loneliness. Higher psychological distress and lower quality of life were associated with higher loneliness scores. Higher levels of loneliness were related to poorer mental health and poorer physical health. 	Fair
Itzick M, Segal JN, Possick C, 2019 [63]	To use a contextual, ecosystemic framework to explore the experience and meaning of relationships,	<p>N: 12</p> <p>Response rate: Not</p>	<p>Design: Cross-sectional/ Qualitative</p> <p>Setting: Narcotics Anonymous Groups</p>	<ul style="list-style-type: none"> Loneliness was the dominant feeling expressed by most of the women ($n= 9$). Drugs were often used as a way to escape overwhelming feelings of loneliness, but 	Fair

	amongst women experiencing problems with substances.	reported Country: Israel Gender: Female Age: 20-33 Substance use: Narcotics	Year: Not reported Procedure: Participants recruited through outreach programs. The interview began with a general invitation for the women to tell their life stories. Participants then questioned about their relationships with significant others in the past and the present. Interviews were 1–1.5 hours in duration. The interviews were recorded and transcribed verbatim.	ultimately did not help. <ul style="list-style-type: none"> It seems that the loneliness expressed by the women stemmed from a feeling that no one really sees them. It may also develop as the women retreat inward, distancing themselves from others as a result of the trauma they experienced in close relationships. 	
Johnson RA, 1985 [33]	To conduct an exploratory study to: (i) investigate the experience of loneliness among people who are dependent on alcohol; and (ii) to evaluate the effectiveness of an existential form of group therapy (Logoanalysis).	N: <i>Study One:</i> 56; <i>Study Two:</i> 20 (10 in treatment and 10 in control) Response rate: Study Two treatment group 71% (not reported for Study one or control condition of Study Two). Country: America Gender: Male Age: <i>Baseline:</i> 27-62 (<i>M</i> =39.91) <i>Experimental:</i> Treatment condition 29-58 (<i>M</i> =41.10); Control condition, 29-63 (<i>M</i> =42.10) Substance use: Alcohol	Design: Cross-sectional Setting: Inpatient substance dependence treatment service Year: Not reported Procedure: Participants invited to participate during routine intake interviews conducted at the time of admission to the six-week inpatient alcohol treatment program at a large Veterans Administration neuropsychiatric hospital. The study was conducted in two phases, baseline and experimental. <i>Baseline</i> - self-report measures to compare alcoholic subjects and other identified groups (e.g. college students). <i>Experimental</i> - compared treatment group with control subjects. Participants enrolled on a voluntary basis in the two-week, daily group based on the principles of Logoanalysis. Baseline questionnaires completed and a second administration of the scales took place two weeks later at the end of the group program. Logoanalysis group ran for one hour a day.	<i>Study One:</i> <ul style="list-style-type: none"> Loneliness was slightly higher than reported for college students, but equal to or lower than those reported for various groups of "high risk" adults. Loneliness unrelated to age Loneliness related to shyness, depression and purpose in life Loneliness related to number of friendships and satisfaction of friendships Loneliness not related to frequency of contact with friendships Loneliness was also negatively related to the degree of intimacy felt in romantic relationships, frequency of contact with romantic partners, and the amount of satisfaction experienced None of the characteristics associated with family relationships was significantly related to loneliness <i>Study Two</i> <ul style="list-style-type: none"> Logoanalysis was not found to be effective in the alleviation of loneliness, nor did it affect any of the other variables of interest in the present study. 	Fair

<p>Kuerbis A, Mereish EH, Hayes M, Davis CM, Sijing S, Morgenstern J, Shao S, 2017 [64]</p>	<p>To explore how coping and social factors (i.e. loneliness) mediate the relationships between internalised heterosexism and health outcomes.</p>	<p>N: 198</p> <p>Response rate: 49% at baseline, 86.4% at 3 month follow-up, 96% 9 months follow-up</p> <p>Country: America</p> <p>Gender: Male</p> <p>Age: "participants were about 36 years old"</p> <p>Substance use: Alcohol</p>	<p>Design: Longitudinal</p> <p>Setting: Community (88% Alcohol Use Disorder diagnosis, 11% met alcohol abuse criteria)</p> <p>Year: Not reported</p> <p>Procedure: Recruitment strategies included online and community-based advertising. Participants were randomly assigned to two possible 12-week treatments: (a) 4 sessions of motivational interviewing or (b) 12 sessions of motivational interviewing plus behavioural self-control therapy. All groups were followed at equivalent time points. Follow-up interviews were implemented at 3 and 9 months after baseline.</p>	<ul style="list-style-type: none"> Loneliness was found to mediate the relationship between internalised heterosexism and alcohol problems. Concluded that addressing loneliness as a potential risk factor for alcohol problems may be crucial in preventing alcohol problems and psychological distress 	<p>Good</p>
<p>Kuerbis A, Padovano HT, Shao SJ, Houser J, Muench FJ, Morgenstern J, 2018 [65]</p>	<p>This study used secondary data analysis to test whether age moderated relationships between variables (i.e. loneliness) and drinking among problem drinkers.</p>	<p>N: 139 (56.8% female, 43.2% male)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: 20–73 ($M=43.2$, $SD=12.5$)</p> <p>Substance use: Alcohol</p>	<p>Design: Cross-sectional cohort</p> <p>Setting: Community</p> <p>Year: Not reported</p> <p>Procedure: Participants recruited using advertising online and in local media. Participants had to have a current Alcohol Use Disorder. Participants completed a series of standard, global self-report assessments. Ecological momentary assessment (EMA) online surveys completed daily over 7 days prior to randomisation. Participants were then assessed again at baseline, the point of randomisation.</p>	<ul style="list-style-type: none"> Loneliness affected daily drinking across all ages equally Older participants reported being less lonely. 	<p>Poor</p>
<p>Li F, Xu Y, Zhu J, Lu J, Zhong B, 2017 [66]</p>	<p>This study examined the association between loneliness and pain intensity in people who use heroin receiving MMT</p>	<p>N: 603 (68.3% males, 31.7% females)</p> <p>Response rate: Not reported</p> <p>Country: China</p> <p>Gender: Male and female</p>	<p>Design: Cross-sectional</p> <p>Setting: Outpatient substance dependence treatment services</p> <p>Year: 2009-2010</p> <p>Procedure: The study consecutively enrolled adults who use heroin who met DSM-IV criteria for a lifetime diagnosis of heroin dependence.</p>	<ul style="list-style-type: none"> Loneliness was significantly associated with an increase in pain intensity After controlling for the potential confounding effects of other covariates, loneliness remained significantly associated with pain, suggesting an independent effect of loneliness on pain in people who use heroin. Concluded loneliness is a significant contributor to increased pain intensity. 	<p>Fair</p>

		Age: $M=38.1, SD=7.0$	Self-administered questionnaires were completed.		
		Substance use: Opioids			
Li H, Zhong B, Xu Y, Zhu J, Lu J, 2017 [67]	To examine the impact of loneliness on sleep patterns in a sample of Heroin Dependent Patients receiving MMT	N: 603 (68.3% males, 31.7% females) Response rate: Not reported Country: China Gender: Male and female Age: 21-59 ($M=38.1, SD=7.0$) Substance use: Opioids	Design: Cross-sectional Setting: Outpatient substance dependence treatment services Year: 2009-2010 Procedure: The study consecutively enrolled adults who use heroin who met DSM-IV criteria for a lifetime diagnosis of heroin dependence and were taking methadone for drug rehabilitation. Self-administered questionnaires were completed.	<ul style="list-style-type: none"> Prevalence of loneliness was 55.9% among people with heroin dependence accessing MMT clinics. There was a significant association between loneliness and poor sleep in terms of quality and quantity, including longer sleep latency, shorter sleep duration, and lower sleep efficiency, indicating that loneliness may exacerbates sleep disturbance. 	Fair
Medora NP, 1983 [68]	(i) To determine the extent of loneliness among alcohol dependent individuals. (ii) To assess loneliness in relation to a range of demographic health and social variables. (iii) To determine whether differences in loneliness exist amongst different populations.	N: 152 (92 Males, 60 Females) Response rate: 61% Country: America Gender: Male and female Age: 19-75 Substance use: Alcohol	Design: Cross-sectional Setting: Participants were undergoing treatment for alcoholism in seven alcohol rehabilitation centres (not reported if inpatient or outpatient settings). Year: Not reported Procedure: Questionnaires distributed at services by counsellors at the service. Investigator provided 'de-briefing' sessions following completion of the anonymous questionnaire whereby the objectives of the study were explained for interested participants.	<ul style="list-style-type: none"> Mean loneliness score was higher than all other samples, except 'low-income single adolescent mother' group. Best predictors of loneliness were: self-esteem, age, self-rated marital satisfaction. Age: Individuals aged 56-65 years were found to be less lonely than those 15-45 years. Gender: Females had higher loneliness scores than males. Marital status: Individuals who were divorced or remarried had highest loneliness scores, while married or de facto had lowest loneliness scores and satisfaction with marital status was related to loneliness. Health: self-rated good/excellent health was related to lower loneliness scores, and poor health was related to higher loneliness scores Mental health: Loneliness was related to self-esteem; Difference in loneliness scores between people who experience happiness in past year and those who did not. Social: Loneliness related to difficulty making friends. People who went out "with relatives" were found to be more lonely than those who went out with a date, or with friends Alcohol history: Difference found for 	Fair

				loneliness between people who had past history of alcoholism in the family, and those who did not. Relationship found between number of years alcohol consumed and loneliness.	
				<ul style="list-style-type: none"> • <i>Other:</i> Loneliness related to job satisfaction. 	
Medora NP, Woodward JC, 1990 [69]	The extent of loneliness was investigated in relationship to gender, religiosity, age, education, adequacy of income, social class, number of close friends, self-rated health, ease in making friends, and frequency of participating in social activities.	<p>N: 152 (92 males, 60 females)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: 19 - 55 years</p> <p>Substance use: Alcohol</p>	<p>Design: Cross-sectional</p> <p>Setting: Participants undergoing treatment in seven alcohol rehabilitation centres (not reported if inpatient or outpatient settings)</p> <p>Year: Not reported</p> <p>Procedure: Questionnaires distributed at services by counsellors at the service.</p>	<ul style="list-style-type: none"> • Younger persons were found to be significantly lonelier than older persons • Women were significantly lonelier than men. • There was a relationship between loneliness and self-rated state of health and ease in making friends. • The following variables did not affect loneliness--education, socioeconomic status, adequacy of income, religiosity, number of close friends, and frequency of going out for social reasons. 	Fair
Medora NP, Woodward JC, 1991 [70]	Objective was to examine the extent of loneliness in relation to demographic and social variables of participants undergoing treatment at alcohol rehabilitation centres.	<p>N: 152 (92 male, 60 female)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: 19 - 55 years</p> <p>Substance use: Alcohol</p>	<p>Design: Cross-sectional</p> <p>Setting: Participants undergoing treatment in seven alcohol rehabilitation centres (not reported if inpatient or outpatient settings)</p> <p>Year: Not reported</p> <p>Procedure: Self-report questionnaires were administered to participants at alcohol rehabilitation centres.</p>	<ul style="list-style-type: none"> • Females lonelier than males. • Negative relationship between loneliness and self-rated marital satisfaction. • Awareness of history of alcoholism had higher loneliness scores than people who were unaware of a family history of alcoholism. • Negative relationship between loneliness and self-esteem. • Negative relationship between loneliness and number of year's alcohol has been consumed. • Higher ratings of happiness of the past year ("very happy") were lonelier than people who rated they were "happy". 	Fair
Michaels AW, 1982 [71]	To compare alcoholic and non-alcoholic treatment populations in terms of denial, and to determine whether denial is related to loneliness and/or to alcoholism	<p>N: 60 (36 males, 24 females)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p>	<p>Design: Cross-sectional</p> <p>Setting: Outpatient substance dependence treatment service; Outpatient mental health treatment service</p> <p>Year: Not reported</p> <p>Procedure: Handout distributed in waiting rooms</p>	<ul style="list-style-type: none"> • Mean loneliness score were not significantly different between groups • Among alcoholics: loneliness scores were correlated with education and income, but not correlated with age. • Among outpatients loneliness not correlated with any demographic variables. • The higher the level of denial, the lower the loneliness score • The lower the level of denial, the higher the 	Fair

		Age: 21-69 ($M=34$) Substance use: Alcohol	at the clinics to recruit participants. Participants completed a survey pack in the waiting room.	<ul style="list-style-type: none"> subject is likely to score on loneliness Loneliness was found to correlate with need for inpatient care. 	
Neale J, Tompkins CNE, Strang J, 2018 [72]	The qualitative study aimed to provide further insights into relationships between peers in residential substance use treatment services	N: 21 (13 males, 8 females) Response rate: Not reported Country: England Gender: Male and female Age: 23-57 Substance use: Not reported	Design: Cross-sectional/qualitative Setting: Inpatient substance dependence treatment service Year: 2015-2016 Procedure: Member of the research team visited treatment services to recruit current residents. Staff at the services contacted former residents for recruitment. Researcher then approached a subgroup of those expressing interest. Semi-structured interviews completed which asked about backgrounds, substance use and experiences of residential treatment, including relationships with their peers. All interviews were audio-recorded and transcribed verbatim.	<ul style="list-style-type: none"> Contrary to expectations, few residents described bonding with their peers. Interpersonal differences polarised residents. Residents more often reported isolation, loneliness, wariness, bullying, manipulation, intimidation, social distancing, tensions and conflict. Overall, findings undermine the notion of the therapeutic community as a method of positive behaviour change. 	Good
Nerviano VJ, Gross WF, 1976 [73]	Aim was a revision and/or regrouping of Bradley's 38 loneliness items for a population of chronic alcoholics. Evidence for the discriminant and convergent validity of newly derived subscales was to be sought.	N: 349 (all males) Response rate: Not reported Country: America Gender: Male Age: $M = 44$ Substance use: Alcohol	Design: Cross-sectional Setting: Inpatient substance dependence treatment service Year: Not reported Procedure: All participants were detoxified and completed the assessment battery at time of admission	<ul style="list-style-type: none"> The Bradley scale, revised factor-analytically into two correlated dimensions termed Interpersonal Anxiety (LS1) and Sense of Rejection and Abandonment (LS2), showed many moderately high correlations. These correlates were seen as highly supportive of the construct validity of the new scales. Overall, the experience of loneliness seems more related to the degree of general success in interpersonal relations than to single needs or traits. Loneliness correlated with personality variables of: immature, interpersonally inhibited, low impulse organisation, poor self-presentation; high loneliness scores related to high subjective distress and emotional instability. Consistent relationship of high loneliness scores to the factor markers for PF Anxiety, an indication of high subjective distress and emotional instability. 	Poor
Newton TF, De La Garza R,	To investigate perceptions of the reasons for	N: 73 (12 females, 61	Design: Cross-sectional	<ul style="list-style-type: none"> 23% of the sample reported using drugs 'very much' to make bad feelings like boredom, 	Fair

Kalechstein AD, Tziortzis D, Jacobsen CA, 2009 [74]	methamphetamine use.	<p>males)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: $M=36.3, SD=9.2$</p> <p>Substance use: Methamphetamines</p>	<p>Setting: Community (all participants met DSM-IV criteria for methamphetamine-dependence)</p> <p>Year: Not reported</p> <p>Procedure: Participants completed an initial battery of questionnaires and were then administered a variety of assessments. At the time of assessment, a toxicology screen was performed.</p>	<p>loneliness, or apathy go away; while 30.1% of the sample reported they did 'not at all'.</p> <ul style="list-style-type: none"> • 26% of the sample reported relapsing mostly to make bad feelings like boredom, loneliness, or apathy go away 	
Perodeau GM, du Fort GG, 2000 [75]	To compare elderly people who use or do not use, psychotropic drugs on mental health and psychosocial characteristics.	<p>N: 199 (82.1% female, 17.9% male)</p> <p>Response rate: 88%</p> <p>Country: Canada</p> <p>Gender: Male and female</p> <p>Age: 62-98</p> <p><i>Drug use group: $M=77.8, SD=7.8$;</i></p> <p><i>Non-drug use group: $M=80.9, SD=7.4$</i></p> <p>Substance use: Psychotropic drugs (not for medical purposes)</p>	<p>Design: Cross-sectional</p> <p>Setting: Community (People who use psychotropic drugs were defined as individuals who reported using one or more psychotropic drugs in the preceding 3 months)</p> <p>Year: Not reported</p> <p>Procedure: Two 90-minute face-to-face interviews were conducted in French or English language, by experienced female interviewers in the elder's home with a 1-week interval between interviews. The first interview concerned health patterns, and the second focused on psychosocial issues.</p>	<ul style="list-style-type: none"> • Feelings of loneliness reported by a higher percentage of people who used psychotropic drugs (40%) than people who did not use drugs (16%). • Anxiety related to loneliness in people who used drugs. • Depression related to loneliness in people who use drugs. • People who use drugs appear to have a greater sensitivity to perceived weaknesses in the social support system than nonusers. 	Fair
Price RH, Curlee-Salisbury J, 1975 [76]	Examined different reactions to treatment settings based on responses of the patient group. Additionally, this study aimed to examine subsets of patients who showed different patterns of response to various	<p>N: 51 (all males)</p> <p>Response rate: Not reported</p> <p>Country: America</p>	<p>Design: Cross-sectional</p> <p>Setting: Inpatient substance dependence treatment service</p> <p>Year: Not reported</p> <p>Procedure: Patients tested individually within 3-4</p>	<ul style="list-style-type: none"> • Conclusions do not specifically refer to loneliness • Conclusions here have been inferred from Table 2 of the source: it appears that loneliness had one of highest ranks when people were in hospital, and one of the lowest ranks (compared to other domains such as pleasantness, vigour, affiliation etc.) for other aspects of alcoholism treatment. 	Poor

	treatment modalities.	<p>Gender: Male</p> <p>Age: 21-56 ($M=44$)</p> <p>Substance use: Alcohol</p>	<p>days discharge. Patients ranked 8 aspects of treatment program on 8 response dimensions. Patients also completed Minnesota Multiphasic Personality Inventory (MMPI-II) and 46 the Shipley Full Scale IQ and 5 the Quick test. Loneliness was one response dimension that was rated across 8 different treatment modalities (e.g. group therapy, lecture, free time).</p>		
Schmidt DR, 2002 [77]	To investigate the impact of a recovery program for recovering adult male addicts. The research was divided into etiological issues and intervention strategy. Within these two main sections three primary areas were explored: (i) loneliness and social relationships; (ii) identity and self-esteem issues; and (iii) self-control issues.	<p>N: 5 staff and 13 clients</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: Not reported</p> <p>Substance use: All substances</p>	<p>Design: Cross-sectional/ Qualitative</p> <p>Setting: Inpatient substance dependence treatment service</p> <p>Year: 2001</p> <p>Procedure: Qualitative cross-sectional interviews were conducted in an ethnographic style asking for description and probing within the answers received. Compared the client interview findings with staff interviews.</p>	<ul style="list-style-type: none"> A lack of quality trusting relationships and isolation when using substances were linked to loneliness according to the staff interviewed. Because of the detached and abusive way that most were treated, they felt often like "survivors" and "on their own" which included feelings of loneliness. Participants identified loneliness as being linked to an urge to escape into drugs and alcohol. The staff and residents were in agreement that there are problems of loneliness and social relationships amongst almost all of the residents. 	Poor
Schonfeld L, Dupree LW, Rohrer GE, 1995 [78]	To determine pre-treatment drinking behaviours of younger and older people who use alcohol, and to identify any differences in antecedents to drinking.	<p>N: <i>Older sample:</i> 109 (69 males, 40 females) <i>Younger Sample:</i> 47 (37 males, 10 females)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: <i>Older sample:</i> 55-84 ($M=64.9$, $SD=6.3$); <i>Younger sample:</i> 20-53 ($M=34.9$, $SD=7.7$)</p> <p>Substance use: Alcohol</p>	<p>Design: Cross-sectional</p> <p>Setting: Inpatient & Outpatient substance dependence treatment service</p> <p>Year: Not reported</p> <p>Procedure: Assessments administered by staff via interviews. For the older sample, the interview was conducted within first two-weeks of program entry. In the younger sample, the interview was conducted at any time during 6-week stay.</p>	<ul style="list-style-type: none"> Older people who use alcohol were found to drink in response to feelings of depression, loneliness, and related interpersonal and emotional states. Younger people who use alcohol tended to drink with other people, away from home, and in response to a wider variety of antecedents. Concluded that differences appear to exist in terms of antecedent to alcohol abuse between younger and older people who use alcohol. Category including loneliness – "depressed, lonely, and bored" – was determined to be an antecedent to drinking alcohol for 45% of the older sample, and 21% of the younger sample. 	Fair

Siddique F, Ahmad Mann A, Ali T, 2012 [79]	To investigate drug use behaviour and its relationship with social characteristics.	<p>N: Specific sample size not reported ("a proportion of 500 drug users")</p> <p>Response rate: Not reported</p> <p>Country: Pakistan</p> <p>Gender: Male</p> <p>Age: Not reported</p> <p>Substance use: Not reported</p>	<p>Design: Cross-sectional</p> <p>Setting: 5 government model drug abuse and rehabilitation centres (not specified if inpatient or outpatient)</p> <p>Year: Not reported</p> <p>Procedure: Participants were interviewed randomly from a list of 500 potential participants available at these centres.</p>	<ul style="list-style-type: none"> • Concluded that loneliness influenced the behaviour of people who experience drug dependence. • Conclusions made about loneliness and drug use don't appear to be supported by the data presented throughout the article 	Poor
Van Hasselt VB, Null JA, Kempton T, Bukstein OG, 1993 [80]	To conduct an evaluation of social skills and depression in adolescents who use substances.	<p>N: 104 (53 females, 51 males)</p> <p>Response rate: Not reported</p> <p>Country: America</p> <p>Gender: Male and female</p> <p>Age: 11.4 - 18.8 years ($M=15.3$, $SD=1.7$)</p> <p>Substance use: Not reported</p>	<p>Design: Cross-sectional</p> <p>Setting: Inpatient mental health treatment service</p> <p>Year: Not reported</p> <p>Procedure: A self-report battery was administered within the first week of admission that assessed level of assertion, social satisfaction, loneliness, depression, hopelessness, and self-esteem. In addition, the relationship between social skills and depression was examined.</p>	<ul style="list-style-type: none"> • Higher levels of depression were related to less assertion skill and increased loneliness. • Loneliness found to be correlated with depression, hopelessness and self-esteem 	Fair
Yang Y, Xu Y, Chen W, Zhu J, Lu J, Zhong B, 2017 [81]	This study determined the prevalence and socio-demographic and clinical correlates of loneliness and its impact on quality of life in Chinese heroin-dependent patients receiving MMT.	<p>N: 603 (68.3% female)</p> <p>Response rate: 92.5%</p> <p>Country: China</p> <p>Gender: Male and female</p> <p>Age: 21-59 ($M=38.1$, $SD=7.0$)</p>	<p>Design: Cross-sectional</p> <p>Setting: Outpatient substance dependence treatment services</p> <p>Year: 2015</p> <p>Procedure: The cross-sectional survey was conducted in three city-owned MMT clinics.</p>	<ul style="list-style-type: none"> • Found a high prevalence of loneliness (55.9%) in Chinese heroin dependent patients receiving MMT. • Efforts to reduce loneliness may be useful to target on those who are unmarried, unemployed, and depressed, and have religious beliefs, get along with others poorly, and have a history of injecting heroin. • Loneliness related to being unmarried, unemployed, having religious beliefs. • Loneliness related to having a history of injecting heroin. • Loneliness related to depression. 	Fair

		Substance use: Opioids		<ul style="list-style-type: none"> Those who got along poorly with others were found to be lonelier. 	
Yeh M, 2002 [82]	To examine the relationships among and between global loneliness, social loneliness, and emotional loneliness, and the degree/severity of alcohol or marijuana use related problems among referred alcohol and marijuana users and a random sample drawn from the university's general male student body.	N: 116 (all males) Experimental, N=49 Control, N=67 Response rate: 91% Country: America Gender: Male Age: Not reported Substance use: Alcohol and marijuana	Design: Cross-sectional Setting: University/outpatient substance dependence treatment service Year: 2000 Procedure: Participants were selected in two different ways: (i) from a referral group sample (experimental group); and (ii) from a random sample (control or comparison group). A purposive sampling design was used for recruiting participants from the referral group. Participants completed a survey including demographics, UCLA-LS and substance use measure.	<ul style="list-style-type: none"> No significant relationship found between drug and alcohol score and the global loneliness score, the emotional loneliness score, and the social loneliness score, for both groups. Significant relationships found among global, emotional, and social loneliness scores for the substance use group and the control group. Significant difference between the substance use group and the control group with respect to the emotional loneliness score and the social loneliness. The control group had a higher social loneliness score. Substance use group had a higher emotional loneliness score. Within the substance use group, the marijuana users were emotionally lonelier and had a higher degree/severity of alcohol or marijuana use related problems than alcohol users. No difference in the global loneliness score between the substance use group and the control group. 	Fair
Zhong B, Xu Y, Zhu J, Liu X, 2018 [83]	To examine the prevalence of NSSI, its subtypes, and factors significantly associated with NSSI.	N: 652 Response rate: 92.5% Country: China Gender: Male and female Age: $M=38.1$, $SD=7$ Substance use: Heroin	Design: Cross-sectional Setting: Outpatient substance dependence treatment services Year: 2009-2010 Procedure: Investigators reviewed medical charts and interviewed patients for eligibility. All patients independently and anonymously completed the questionnaires.	<ul style="list-style-type: none"> Relative to the no NSSI group, patients in the NSSI group were more likely to feel lonely. Loneliness was one factor found to be significantly associated with non-suicidal self-injury amongst this sample. 	Fair

AA, Alcoholics Anonymous; MMT, methadone maintenance treatment; NSSI, non-suicidal self-injury; SELSA, Social and Emotional Loneliness Scale for Adults; SELSA-S, Social and Emotional Loneliness Scale for Adults, short version; TASC, Treatment Alternatives for Safe Communities; UCLA-LS, University of California, Los Angeles Loneliness Scale.

Table 2. *Measures of loneliness*

Measure used	Psychometric properties
<i>UCLA Loneliness Scale (Version 3)</i> [51]	
Armstrong JA, 2016 [48]	Not reported for study sample. Cited psychometric properties reported by Russell (1996) [51] (elderly population, aged >65, n=301) <i>Test-retest reliability</i> : ICC =0.73 (fair)
Britton PC, Conner KR, 2007 [50]	<i>Internal consistency</i> : $\alpha = 0.87$ (good) <i>Test-retest reliability</i> : ICC = 0.76 (fair) Cited psychometric properties reported by Russell (1996) [51]: <i>Internal consistency</i> : $\alpha = 0.89 - 0.94$ (excellent) <i>Test-retest reliability</i> : ICC = 0.73 (fair) <i>Criterion validity</i> : negatively correlated with measure of belonging, $r = -0.67$, $P < 0.001$
Conner KR, Britton PC, Sworts LM, Joiner TE, 2007 [53]	<i>Internal consistency</i> : $\alpha = 0.87$ (good) <i>Test-retest reliability</i> : ICC = 0.76 (fair) <i>Criterion validity</i> : negatively correlated with measure of belonging, $r = -0.67$, $P < 0.001$, and burdensomeness and loneliness was $r = -0.67$, $P < 0.001$.
Kuerbis A, Mereish EH, Hayes M, Davis CM, Sijing S, Morgenstern J, Shao S, 2017 [64]	<i>Internal consistency</i> : $\alpha = 0.91 - 0.93$ across three-month time frame (excellent)
<i>Revised UCLA Loneliness Scale</i> [89]	
Cao Q, Liang Y, 2017 [52]	<i>Internal Consistency</i> : $\alpha = 0.79$ (fair) Cited psychometric properties reported by Russell <i>et al.</i> (1980) [89]: <i>Internal consistency</i> : $\alpha = 0.94$ (excellent)
Essex EL, Petras D, Massat CR, 2007 [56]	<i>Internal consistency</i> : $\alpha = 0.89$ (good)
Evans TJ, 2010 [57]	Measure used to identify eligible participants for the study. Not reported.
Harris KS, 1983 [59]	<i>Internal consistency</i> : $\alpha = 0.94$ (excellent) Cited psychometric properties reported by Russell <i>et al.</i> (1980) [89]: <i>Concurrent validity</i> : UCLA correlated with Beck Depression Inventory, $r = 0.62$, and the Costello-Comrey Anxiety measure, $r = 0.82$
Johnson RA, 1985 [33]	Measure used to identify eligible participants for the study. Cited psychometrics reported by Russell <i>et al.</i> [89]: <i>Internal consistency</i> : $\alpha = 0.94$ (excellent) <i>Test-retest reliability</i> : coefficient reported to be 0.70 over two months (fair) <i>Convergent validity</i> : Revised UCLA correlated with original UCLA scale, $r = 0.91$
Yeh M, 2002 [82]	Cited psychometrics reported elsewhere [47,84,85]: <i>Internal Consistency</i> : $\alpha = 0.84 - 0.96$ (good – excellent) <i>Convergent validity</i> : Russell [89] provided correlations with the NYU Loneliness Scale, $r = 0.65$ and The Differential Loneliness Scale, $r = -0.72$ <i>Concurrent validity</i> : Russell <i>et al.</i> [89] found loneliness scores were related to the experience of affects that have been linked to loneliness (not specified what these are) <i>Discriminant validity</i> : Russell <i>et al.</i> [89] demonstrated loneliness scores to be distinct from social desirability, social support, depression, lack of

	affiliative motivation, and low social risk taking
3-item Revised UCLA Loneliness Scale [90]	
Ingram I, Kelly PJ, Deane FP, Baker AL, Raftery DK, 2018 [22]	<i>Internal consistency:</i> $\alpha = 0.82$ for study sample (good)
<i>Social and Emotional Loneliness Scale for Adults (SELSA-S; [91])</i>	
Ingram I, Kelly PJ, Deane FP, Baker AL, Raftery DK, 2018 [22]	<i>Internal consistency:</i> Total of the 15 items: $\alpha = 0.81$ (good); social loneliness subscale $\alpha = 0.80$ (good), family loneliness $\alpha = 0.83$ (good), and romantic loneliness $\alpha = 0.82$ (good). <i>Concurrent validity:</i> SELSA-S total score correlated with UCLA 3-item score, $r = 0.49$, $P < 0.001$. Satisfaction with one's marital status was related to scores on the romantic subscale of the SELSA-S, $r(305) = 0.23$, $P < 0.001$. <i>Discriminant validity:</i> SELSA-S total score correlated with psychological distress, $r = 0.33$, $P < 0.001$. SELSA-S total score inversely correlated with quality of life, $r = -0.39$, $P < 0.001$. <i>Validity:</i> Three-factor model was the best fit, yet this model still fit the data poorly: ($\chi^2/df = 4.33$, CFI = 0.86, Tucker-Lewis Index (TLI) = 0.83, and RMSEA = 0.10). Factor loadings were high, with all loadings exceeding the .30 cutoff criteria.
Hosseini M, Yassini ASM, Bakhshani S, Bakhshani S, 2014 [62]	Cited psychometric properties reported by Jowker [92]: <i>Internal consistency:</i> $\alpha = 0.92$ (romantic subscale) (excellent), $\alpha = 0.84$ (social subscale) (good) and $\alpha = 0.78$ for (family subscale) (fair). Cited psychometric properties reported by Ditommaso and Brannen [91] <i>Internal consistency:</i> $\alpha = 0.87 - 0.89$ (good)
<i>Bradley Loneliness Scale [93]</i>	
Michaels AW, 1982 [71]	Cited psychometrics reported by Belcher [94,95]: <i>Test-retest reliability:</i> College student sample over two-week time frame and 8-weeks $r = 0.89$, $P < 0.001$. High face validity stated (psychometrics not reported) <i>Discriminant validity:</i> established with inmate population against MMPI Depression score (psychometrics not reported).
Nerviano VJ, Gross WF, 1976 [73]	<i>Construct validity:</i> $r = 0.18 - 0.55$ (the authors deemed this moderate-high)
<i>Sisenwein Loneliness Scale [96]</i>	
Allen HA, Peterson JS, Whipple S, 1981 [47]	Not reported
<i>Wellbeing Questionnaire developed by Elton and Hornquist [97]</i>	
Akerlind, Hornquist, Bjurulf, 1988 [45]	<i>Internal consistency:</i> $\alpha = 0.64 - 0.89$ (unacceptable - good) <i>Validity:</i> $r = 0.44$
Akerlind I, Hörnquist JO, Hansson B, 1987 [46]	<i>Internal consistency:</i> $\alpha = 0.86$ (good)
Akerlind I, Hörnquist JO, 1989 [32]	<i>Internal consistency:</i> $\alpha = 0.86$ (good)
Elton HL, Hörnquist JO, 1983 [54]	<i>Internal consistency:</i> $\alpha = 0.85 - 0.89$ (good)
Hörnquist JO, Elton HF, 1983 [55]	Not reported
Hörnquist JO, Akerlind I, 1987 [60]	<i>Internal consistency:</i> $\alpha = 0.86$ and $\alpha = 0.85$ at two-year re-examination (good)
Hörnquist JO, Hansson B, Akerlind I, 1988 [61]	<i>Internal consistency:</i> $\alpha = 0.85 - 0.89$ (good)
<i>Loneliness Inventory developed by Woodward [98]</i>	

Medora NP, 1983 [68]	<i>Internal consistency:</i> $\alpha = 0.96$ (excellent) <i>Validity:</i> not reported for this study, yet author alluded to other studies that have established this measure's validity against the Sisenwein Loneliness Self-Rating Scale [96]
Medora NP, Woodward JC, 1990 [69]	<i>Internal Consistency:</i> $\alpha = 0.96$ (excellent) <i>Test-retest reliability:</i> ICC = 0.97 (excellent) Cited psychometrics reported by Woodward [99]: <i>Criterion validity:</i> correlation with a single-item self-report measure of loneliness: $r = 0.93$, $P < 0.001$. Correlation with UCLA: $r = 0.87$, $P < 0.001$
Medora NP, Woodward JC, 1991 [70]	<i>Internal Consistency:</i> $\alpha = 0.96$ (excellent) <i>Test-retest reliability:</i> ICC = 0.97 (excellent) Cited psychometrics reported by Woodward [99]: <i>Criterion validity:</i> correlation with a single-item self-report measure of loneliness: $r = 0.93$, $P < 0.001$ Correlation with UCLA: $r = 0.87$, $P < 0.001$
<i>Single Item Measures</i>	
Kuerbis A, Padovano HT, Shao SJ, Houser J, Muench FJ, Morgenstern J, 2018 [65]	Not reported. One item measured loneliness, "In the past hour, how lonely do you feel?" The response set on these items ranged from 0 (not at all) to 8 (extremely).
Li F, Xu Y, Zhu J, Lu J, Zhong B, 2017 [66]	Not reported. Loneliness was measured with a single-item self-report question "How often do you feel lonely?" with a five-point scale: 1 (never), 2 (seldom), 3 (sometimes), 4 (often), 5 (always).
Li H, Zhong B, Xu Y, Zhu J, Lu J, 2017 [67]	Not reported. Loneliness was measured with a single-item self-report question "How often do you feel lonely?" with a five-point scale: 5 (never), 4 (seldom), 3 (sometimes), 2 (often), 1 (always). The authors classified participants as being 'lonely' if they indicated feeling lonely at least 'sometimes'.
Price RH, Curlee-Salisbury J, 1975 [76]	Not reported for items related to loneliness. Loneliness was assessed through a single item that was part of an 8-item scale. The item read: "I felt lonely".
Yang Y, Xu Y, Chen W, Zhu J, Lu J, Zhong B, 2017 [81]	Not reported for items related to loneliness. Loneliness was assessed with a single question asking how often the respondent feels lonely on a 5-point Likert scale: 1 (always), 2 (often), 3 (sometimes), 4 (seldom), 5 (never). Participants were classified as lonely if they indicated their loneliness was "sometimes", "often", or "always", while participants who reported "never" or "seldom" were classified as not lonely.
Zhong B, Xu Y, Zhu J, Liu X, 2018 [83]	Not reported for items related to loneliness. Loneliness was assessed with a single question asking how often the respondent feels lonely on a 5-point Likert scale: 1 (always), 2 (often), 3 (sometimes), 4 (seldom), 5 (never). The five category loneliness variable was transformed into a binary variable: lonely (≥ 3) and not lonely (≤ 2).
<i>Qualitative studies</i>	
Boyles BR, 2018 [49]	Not applicable
Funk PE, 1973 [58]	Not applicable. Interview question: "Within the framework of the phenomenon of loneliness are there similarities in descriptions of that phenomenon by some members of the population?"
Itzick M, Segal JN, Possick C, 2019 [63]	Not applicable
Neale J, Tompkins CNE, Strang J, 2018 [72]	Not applicable
Schmidt DR, 2002 [77]	Not reported for items related to loneliness. Interview question for staff: "Talk to me about the philosophy and approach that Faith Recovery Center has in addressing the issues of loneliness and social relationships." Loneliness was not explicitly asked of clients.

<i>Other measures</i>	
Newton TF, De La Garza R, Kalechstein AD, Tziortzis D, Jacobsen CA, 2009 [74]	Not reported for items related to loneliness Loneliness was a category on a questionnaire used to identify self-perceived reasons for taking drugs or for relapsing. The item read: Do you use drugs mostly to make bad feelings like boredom, loneliness, or apathy go away? Responses ranged from 1 (not at all) to 7 (very much).
Perodeau GM, du Fort GG, 2000 [75]	Not reported for items related to loneliness Older Americans Resources and Services [100]. One of the subjective items on this scale was "feeling lonely" and the scoring procedure described by Harel and Deimling [101] was used.
Schonfeld L, Dupree, LW, Rohrer GE, 1995 [78]	Not reported for items related to loneliness Loneliness described as a category of 'intrapersonal determinants of drinking' which was devised by the researchers based on classifications developed previously by Marlatt and Gordon [102,103].
Siddique F, Ahmad Mann A, Ali T, 2012 [79]	Not reported for items related to loneliness A questionnaire was used which asked about loneliness. Table 3 in the study indicates that participants rated loneliness "To a great extent", "To some extent", or "Not at all". No information about how the item was framed.
Van Hasselt VB, Null JA, Kempton T, Bukstein OG, 1993 [80]	The Loneliness Scale [104] is a 24-item questionnaire developed to evaluate children's feelings of isolation and social dissatisfaction. Children indicate on a five-point scale the extent to which each statement is a true description of them. Cited psychometrics reported by Asher <i>et al.</i> [104]: Split-half reliability = 0.91 (excellent)

Note. Internal consistency cutoffs based on Cicchetti (1994) [105]. For information about the validity and reliability of the UCLA Loneliness Scale (Version 3; [51]); The Bradley Loneliness Scale [93], The Sisenwein Loneliness Scale [96] and the Social and Emotional Loneliness Scale for Adults – Short Version [91] across other populations, see Appendix 1. ICC, intraclass correlation coefficient.