University of Wollongong

Research Online

Faculty of Engineering and Information
Sciences - Papers: Part A

Faculty of Engineering and Information
Sciences

1-1-2017

Degradation of pharmaceuticals and personal care products by white-rot fungi-a critical review

Muhammad Bilal Asif

University of Wollongong, mba409@uowmail.edu.au

Faisal I. Hai

University of Wollongong, faisal@uow.edu.au

Lakhveer Singh

Universiti Malaysia Pahang

William E. Price

University of Wollongong, wprice@uow.edu.au

Long D. Nghiem

University of Wollongong, longn@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/eispapers>

Part of the [Engineering Commons](#), and the [Science and Technology Studies Commons](#)

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Degradation of pharmaceuticals and personal care products by white-rot fungi-a critical review

Abstract

White-rot fungi (WRF) mediated treatment can offer an environmentally friendly platform for the removal of pharmaceuticals and personal care products (PPCPs) from wastewater. These PPCPs may have adverse impacts on aquatic organisms and even human and thus their removal during wastewater treatment is of significant interest to the water industry. Whole-cell WRF or their extracellular lignin modifying enzymes (LMEs) have been reported to efficiently degrade PPCPs that are persistent to conventional activated sludge process. WRF mediated treatment of PPCPs depends on a number of factors including physicochemical properties of PPCPs (e.g., hydrophobicity and chemical structure) and wastewater matrix (e.g., pH, temperature, and dissolved constituents), type of WRF species and their specific extracellular enzymes. This review critically analyzes the performance of whole-cell WRF and their LMEs for the removal of PPCPs; particularly, it offers insights into PPCP removal mechanisms (e.g., biosorption vs. biodegradation) and degradation pathways as well as the formation of intermediate byproducts.

Keywords

pharmaceuticals, personal, care, products, white-rot, fungi-a, critical, degradation, review

Disciplines

Engineering | Science and Technology Studies

Publication Details

Asif, M. B., Hai, F. I., Singh, L., Price, W. E. & Nghiem, L. D. (2017). Degradation of pharmaceuticals and personal care products by white-rot fungi—a critical review. *Current Pollution Reports*, 3 (2), 88-103.

Degradation of pharmaceuticals and personal care products by white-rot fungi – A critical review

Post-peer reviewed version

Current Pollution Reports

<https://dx.doi.org/10.1007/s40726-017-0049-5>

(February, 2017)

Muhammad B. Asif^a, Faisal I. Hai^{a*}, Lakhveer Singh^b, William E. Price^c, Long D. Nghiem^a

^a Strategic Water Infrastructure Lab, School of Civil, Mining and Environmental Engineering, University of Wollongong (UOW), Wollongong, NSW 2522, Australia. (E-mail: mba409@uowmail.edu.au; faisal@uow.edu.au; longn@uow.edu.au)

^b Faculty of Engineering Technology, Universiti Malaysia Pahang, Lebuhraya Tun Razak, 26300 Gambang, Kuantan, Pahang, Malaysia. (Email: lakhveer@ump.edu.my)

^c Strategic Water Infrastructure Lab, School of Chemistry, University of Wollongong (UOW), Wollongong, NSW 2522, Australia. (E-mail: wprice@uow.edu.au)

*Corresponding author: Faisal I. Hai (E-mail: faisal@uow.edu.au)

Citing information:

Asif, M.B., Hai, F.I., Singh, L., Price, W.E., Nghiem, L.D. 2017. Degradation of Pharmaceuticals and Personal Care Products by White-Rot Fungi—a Critical Review. *Curr Pollution Rep*, 1-16 (DOI: 10.1007/s40726-017-0049-5).

Abstract:

White-rot fungi (WRF) mediated treatment can offer an environmentally friendly platform for the removal of pharmaceuticals and personal care products (PPCPs) from wastewater. These PPCPs may have adverse impacts on aquatic organisms and even human and thus their removal during wastewater treatment is of significant interest to the water industry. Whole-cell WRF or their extracellular lignin modifying enzymes (LMEs) have been reported to efficiently degrade PPCPs that are persistent to conventional activated sludge process. WRF mediated treatment of PPCPs depends on a number of factors including physicochemical properties of PPCPs (e.g., hydrophobicity and chemical structure) and wastewater matrix (e.g., pH, temperature and dissolved constituents), type of WRF species and their specific extracellular enzymes. This review critically analyzes the performance of whole-cell WRF and their LMEs for the removal of PPCPs: particularly it offers insights into PPCP removal mechanisms (*e.g.*, biosorption *vs.* biodegradation) and degradation pathways as well as the formation of intermediate byproducts.

Keywords: White-rot fungi; lignin modifying enzymes (LMEs); Pharmaceuticals and personal care products (PPCPs); Biodegradation; Redox-mediators; metabolites.

1. Introduction

Trace organic contaminants (TrOCs) include diverse groups of chemicals such as pharmaceuticals and personal care products (PPCPs), pesticides, surfactants and industrial chemicals. Due to their ineffective removal by the conventional wastewater treatment processes, these TrOCs are commonly detected in the aquatic environment including surface water and groundwater, and even seawater [1, 2, 3]. Most TrOCs are biologically active even at trace concentrations (in the range of few ng/L) and can impose detrimental impacts on aquatic environment as well as on human health [4, 5]. For example, PPCPs such as synthetic hormones can induce endocrine disrupting effects in aquatic lives [6, 7]. Some of these TrOCs can also accumulate in aquatic biota. Exposure to a non-lethal dose of antibiotics may result in the development of antibiotic resistant genes in bacteria which is an emerging concern for human health according to the World Health Organization [8, 9, 10, 11]. Biological treatment processes are environmentally friendly and cost effective [12, 13, 14]. The conventional activated sludge and membrane bioreactor processes can efficiently remove bulk organics, nutrients and pathogens. However, certain groups of PPCPs such as significantly hydrophilic compounds and/or compounds with strong electron withdrawing functional groups are poorly removed by the conventional biological treatment processes [2, 15, 16, 17]. Thus, effective treatment processes to remove PPCPs from wastewater is urgently needed.

Extracellular enzymes of white-rot fungi (WRF) are characterized by their capacity to degrade the complex structure of lignin, which also enables them to degrade a wide range of resistant pollutants including synthetic dyes, chlorophenols, polychlorinated biphenyls and polycyclic aromatic hydrocarbons [18, 19, 20, 21, 22, 23]. WRF and their lignin modifying enzymes (LMEs) have been investigated recently for the degradation of a broad spectrum of PPCPs [24, 25, 26, 27]. The potential of WRF for the removal of PPCPs has been investigated mostly in batch mode. There are only a few studies on continuous flow reactor configurations [24, 26, 28, 29, 30]. Despite a significant research effort, efficient removal of PPCPs by WRF mediated treatment, their removal mechanisms and degradation pathways remain largely to be elucidated. There have been a few excellent reviews on bioremediation of

recalcitrant compounds by WRF and their LMEs [31, 32, 33, 34, 35]. However, aspects such as comparative performance of different whole-cell WRF species, the impacts of different reactor configurations as well as the performance under non-sterile environment deserve further attention. Moreover, factors controlling the performance of the crude and/or purified LMEs for the degradation of PPCPs, particularly the redox-mediator enhanced laccase systems remain largely unaddressed. This critical review fills these gaps. Degradation mechanisms and pathways for the removal of PPCPs are also critically discussed and the major metabolites or intermediate products following WRF treatment are identified. In addition, the technological challenges associated with the continuous fungal bioreactors are discussed and possible solutions are outlined. Directions for future research are also provided.

2. Enzymatic Systems of WRF

Several features of WRF make their application for PPCP removal attractive: (i) the non-specificity of their enzyme systems, facilitating degradation of a broad spectrum of pollutants [36, 37]; (ii) the hyphal growth, a defining property of filamentous fungi including WRF, enabling them to access pollutants by colonizing fast; (iii) the secretion of extracellular enzymes enabling them to degrade pollutants with low water solubility; and (iv) the ability of some WRF species to degrade pollutants in nutrient deficient environment over a wide range of pH. Notable, however, is that resistant pollutant degradation by WRF is a co-metabolic process, *i.e.*, occurs in presence of an easily degradable substrate. Since, PPCP degradation by WRF depends mainly on their enzyme systems, these are described in this section.

Three different types of extracellular LME, namely, lignin peroxidases (LiPs), laccase and manganese-dependent peroxidases (MnPs), are secreted by WRF. The main difference between laccase and peroxidases is the electron acceptor where oxygen and hydrogen peroxide are the respective electron acceptors. [38, 39]. Not every WRF species produces all three enzymes, and combination of major LME varies from one WRF species to another. Even the secretion pattern of enzymes varies within a WRF species. For instance, different strains of *T. versicolor* has been reported to secrete all three enzymes but laccase is the main enzyme secreted by the strain ATCC 7731 [40, 41, 42]. In addition, composition of

growth medium and culture conditions can influence the secretion of a specific enzyme. Degradation of some pollutants such as phenolic compounds, peptides, and organic acids by WRF may result in the formation of low molecular weight mediators which can enhance the spectrum of compounds degraded by WRF [37, 39, 43]. Based on the secretion patterns of enzymes, WRF can be categorized as [44]: (i) MnP-laccase group such as *T. versicolor*, *P. ostreatus*, *D. squalens* and *P. tigrinus*; (ii) LiP-laccase group such as *P. ochraceofulva*; and (iii) LiP-MnP group such as *P. chrysosporium*.

In addition to the extracellular enzymes, intercellular enzymes may play an important role in the degradation of xenobiotics. Intracellular cytochrome P450 enzyme system has been observed to play a vital role in the degradation of some PPCPs such as and chlorinated hydrocarbons and polycyclic aromatic hydrocarbons [45, 46, 47]. Cytochrome P450 is a group of monooxygenases which can degrade PPCPs by catalyzing a number of reactions such as heteroatom oxygenation, dehalogenation and hydroxylation [48].

3. Modes of PPCP Degradation by WRF

Different WRF and their LME have been investigated under different experimental conditions for the removal of PPCPs - either by whole-cell WRF culture or by using the crude and/or purified enzyme extracts. In this section, different modes of PPCP degradation via WRF and their LMEs are critically reviewed.

3.1. Removal by Whole-cell WRF

3.1.1. Degradation capacity of different WRF

Although whole-cell WRF cultures, either in submerged or solid media, have been used for the removal of PPCPs, submerged whole-cell WRF cultures have been more commonly reported. Since WRF species harbor different enzyme systems, the extent of PPCP removal achieved by different WRF species also varies. PPCP removal performance by different WRF species is presented in Table 1. Notably, temperature and pH for these studies were in the range of 25-30°C and 4.5-5, respectively, with glucose used as the common electron donor.

T. versicolor, also known as *Coriolus versicolor* [49, 50], has been investigated in several studies for the removal of PPCPs both in batch and continuous reactors [24, 26, 27, 43, 51, 52, 53]. Depending on the strain, *T. versicolor* may contain laccase, LiP and MnP, with laccase being the predominant enzyme in some strains. It can be observed from Table 1 that *T. versicolor* achieved significant removal (>70%) for most of the tested PPCPs. PPCPs such as steroid hormones, nonylphenol and octocrylene can induce endocrine disrupting effects [54, 55]. Their removal by *T. versicolor* in literature ranges from 80-99% regardless of the operating conditions. Conversely, poor and unstable removal was reported for particular pharmaceuticals namely ciprofloxacin (35%), salicylic acid (0-5%), azithromycin (26%), tetracycline (0-5%) and carbamazepine (negligible to 90%). Low and/or unstable removal of these compounds can be attributed to the presence of strong electron withdrawing functional groups in their chemical structure [31]. Compared to the pharmaceuticals, the removal of personal care products has been communicated in only a limited number of studies [27, 52]. However, *T. versicolor* has been shown to achieve high removal of ingredients of personal care products such as triclosan, oxybenzone, octocrylene and nonylphenol (Table 1).

[TABLE 1]

Other WRF species such as *P. chrysosporium* [56, 57], *B. adusta*, *D. squalens* and *P. ostreatus* containing different combinations of LMEs have also been investigated for the removal of PPCPs (Table 1). However, these studies reported the removal of only a few PPCPs such as ibuprofen, diclofenac and triclosan. Despite the difference in enzyme secretion pattern of these WRF, efficient removal (in the range of 70-99%) was achieved for most tested PPCPs, with uncertain/unstable removal reported for carbamazepine, which is a known persistent pharmaceutical. It is important to note that a direct comparison of PPCP removal data from different studies may not be valid due to the differences in operating conditions and reactor configurations as well as difference in their enzymatic systems. However, Table 1 serves the purpose of providing a general overview.

3.1.2. WRF Reactor configurations

Different reactor configurations have been explored for the continuous treatment of PPCPs in WRF-based systems (Table 1). Since whole-cell WRF based treatment systems are still in their development phase, the reactor design and configurations are of significant importance. Hence, the salient features of the reactor configurations studied to date for the removal of PPCPs, namely, stirred tank bioreactors [56, 57], bubble column bioreactors, fluidized or packed bed bioreactors [59, 60, 61] and membrane bioreactors [62, 63, 64, 65] are briefly discussed here to provide a general overview.

Among all the reactor configurations, stirred tank bioreactor has been the most common type of reactor used for the treatment of PPCPs in WRF based treatment systems. This reactor type has been explored mostly in batch and sterile modes [56, 57]. In this reactor aeration is provided usually at the bottom of the reactor which is dispersed via mechanical agitation. Mechanical mixing also ensures the uniform mixing of the growth medium and wastewater in the reactor [59]. Enhanced production of extracellular enzymes could be achieved in stirred tank bioreactors compared to other reactor configurations. For instance, Babič and Pavko [66] investigated the production of laccase and MnP from *D. squalens* in stirred tank bioreactor and bubble column reactor under different operating conditions such as incubation time and agitation speed: they observed that laccase production was as much as 70% higher in the former, although the production of MnP was comparable [66]. Agitation speed and high shear rate may influence the morphology of the fungal biomass. In a study by Cao et al. [67], impact of two reactor configurations, namely stirred tank bioreactor and airlift bioreactor, on the morphology of *P. sanguineus* was investigated. They found that the morphology of the fungal biomass was adversely impacted in the stirred tank bioreactor which uses strong mechanical mixing. Consistent with the finding of Cao et al. [67], it has been reported that excessive agitation may lead to the rupturing of fungal hyphae [59, 68, 69]. Therefore, agitation/mixing speed in bioreactors is an important parameter governing fungal morphology and enzymatic activity.

Fluidized bed bioreactor is another type that has been used for the removal of PPCPs from municipal and synthetic wastewater [24, 58]. In this type of reactor, fungal biomass rapidly moves around the solid carrier ('bed'), allowing uniform mixing of the reaction mixture [70]. However, aggregation of fungal biomass may cause poor fluidization, resulting in spouting of bed. Biomass aggregation can be avoided by intermittent and partial purging of the biomass [71].

A bubble column reactor coupled with a microfiltration membrane has been recently explored for the removal of PPCPs [26, 27]. Such fungal membrane bioreactor was expected to offer some additional advantages over bubble column reactors [14, 72, 73] such as: (i) formation of biofilm on the membrane surface that may enhance the removal of recalcitrant PPCPs; (ii) maintenance of high fungal biomass concentration improving biodegradation rate; and (iii) effective prevention of enzyme washout. Nevertheless, bacterial contamination may hamper the growth and enzymatic activity of whole-cell WRF in any reactor configuration. Impacts of bacterial contamination on WRF are discussed in the next section.

3.1.3. Performance under non-sterile environment

The capacity of WRF for PPCP removal has been commonly investigated under sterile conditions to avoid bacterial contamination. Fungal bioreactors have also been operated under non-sterile conditions but for a short period of time [24, 26, 58]. Bacterial contamination in the bioreactor adversely affects PPCP removal efficiency by: (i) creating competition between bacteria and fungi for substrate; (ii) damaging the fungal mycelium; and (iii) disrupting the growth of fungi [26, 70]. Therefore, it is important to develop techniques for an uninterrupted growth of fungal strains and to enhance their competitiveness over bacteria. Some of the strategies to avoid bacterial contamination are outlined below:

- (i) Operation under acidic pH: Optimum pH for the growth and reproduction of fungi ranges from 4.5-5. Conversely, bacteria grow at or near neutral pH. Bacterial growth can be suppressed by maintaining the pH in the range of 4.5-5. However, this is a temporary solution to the problem of bacterial contamination because some bacteria can eventually adapt to acidic environment [74].

- (ii) Immobilization or attached growth of fungi: Immobilization of fungal strains onto different carriers under non-sterile conditions shows promising results. For instance, by immobilizing *C. versicolor* onto a plastic support, Hai et al. [72] was able to prevent bacterial contamination for an extended period of time while operating the reactor under non-sterile conditions for the treatment of an azo dye.
- (iii) Nitrogen limited feed: Bacterial contamination can be avoided by using a media deficient of nitrogen. However, this strategy can only help during the startup of the bioreactor. Bacterial contamination would occur with the passage of operating time as bacteria would start consuming carbon and nitrogen available in fungal mycelium [74].
- (iv) Coupling of bioreactor with micro-screen: Bioreactors can be coupled with a micro-screen which would retain fungal biomass but allow the washout of bacteria with effluent. Moreover, this strategy will benefit from using shorter hydraulic retention times, which will enhance the washout of bacteria from the reactor [75].
- (v) Use of disinfecting agents: Inactivation of bacteria without imposing any harmful effects on fungal biomass can be a promising strategy. Depending on the wastewater characteristics, it is important to carefully select the type, dose and contact time of the disinfectant. Disinfection of wastewater using ozone has been used successfully to selectively inactivate bacteria [76, 77].
- (vi) Biomass replacement: Periodic biomass replacement and purging strategy can be used to carry out long term operation of fungal bioreactors. In this strategy, biomass in the bioreactor is purged and renewed in different fractions (*e.g.*, $\frac{1}{2}$ or $\frac{1}{4}$ th of the initial biomass volume) at different frequencies [78, 79].
- (vii) Pretreatment of wastewater: Coagulation-flocculation pretreatment of non-sterile wastewater can help to reduce the initial bacterial count which would allow an extended operation of the fungal bioreactor [79].

3.2. Removal by Extracellular LME

Individual extracellular LME have been tested for the removal of a wide range of pollutants. Use of the harvested enzyme instead of a whole-cell preparation allows decoupling of fungal growth and pollutant degradation steps and this can be a suitable strategy to avoid bacterial contamination issues. The capacity of both crude and purified/commercially available extracellular LME for PPCP removal has been reported previously [27, 41, 42, 80, 81, 82, 83, 84, 85, 86]. However, whole-cell fungi may achieve relatively better removal of a broader spectrum of pollutants than extracellular LME due to the availability of extracellular, intercellular and/or mycelium bound enzymes in addition to sorption of pollutants onto fungal biomass. In this section the performance of crude, purified or commercially available extracellular enzymes are discussed.

3.2.1. Performance of Crude LMEs

Crude extracellular enzymes have been investigated for the degradation of PPCPs in both batch and continuous mode. For instance, Wen et al. [87] achieved significant degradation of two pharmaceuticals, oxytetracycline (84%) and tetracycline (72%), by using crude MnP (40 U/L) extracted from *P. chrysosporium*. Similarly, crude enzyme solution extracted from *T. versicolor* containing MnP (30 U/L) and laccase (1500 U/L) was tested for the degradation of 10 pharmaceuticals at an initial concentration of 10 µg/L each [80]. They achieved complete degradation of five pharmaceuticals diclofenac, ibuprofen, naproxen, indomethacin and fenopufen, while the rest were partially removed. LiP extracted from *P. chrysosporium* was tested for the degradation of diclofenac and carbamazepine at different pH [88]. It was observed that degradation of carbamazepine was mostly <10%, whereas a complete degradation was achieved for diclofenac at pH 3-4.5. Similarly, crude extract from *T. versicolor* was used for the removal of 30 TrOCs including PPCPs [81]. The results revealed that all steroid hormones were almost completely removed (>95%), while removal of two PPCPs namely diclofenac and triclosan ranged from 50-60% with poor removal of the remaining PPCPs (<20%). In another study [89], crude enzyme extract from *P. ostreatus* achieved low removal (<20 %) for two PPCPs, namely, oxybenzone and naproxen. Since WRF

also secretes natural mediators along with LMEs, crude enzyme may achieve better PPCP removal efficiency compared to purified and/or commercially available LME. For instance, Tran et al. [80] highlighted that complete removal (>99%) of some pharmaceuticals such as diclofenac, indomethacin and ibuprofen was due to the presence of some natural mediators in crude laccase solution.

Removal performance of selected PPCPs by individual crude enzymes has been systematically presented in Table 2. It can be seen that the extent of PPCP removal is different for each type of LME. Moreover, crude enzyme extract from different fungi may perform differently. For instance, Weng et al. [90] collected crude LiP from two WRF species, namely, *P. sordida* and *P. chrysosporium* for the treatment of EDCs and found that LiP from *P. sordida* was more effective than LiP from *P. chrysosporium*. Similarly, removal of DEET, an insect repellent, by *T. versicolor* was 55% in real wastewater compared to its 20% removal in acetate buffer. High removal of DEET in real wastewater was attributed to the presence of other compounds (such as phenolic compounds) which may act as redox-mediators [91], as further discussed in a following section.

[TABLE 2]

3.2.2. Performance of Purified LME

Purified LME, mostly laccase from different WRF, has been used for the treatment of PPCPs in both batch and continuous reactors. Average removal of PPCPs by purified laccase reported in recent studies is presented in Figure 1. Purified laccases are more effective for the removal of phenolic compounds such as oxybenzone, triclosan and steroid hormones. Removal of non-phenolic compounds such as carbamazepine, naproxen and ketoprofen is generally poor/unstable. Their degradation depends on their physicochemical properties such as hydrophobicity and chemical structure as well as the oxidation reduction potential (ORP) of the enzyme [31, 102]. Indeed, based on the results of the recent studies incorporated in Figure 1, removal of phenolic compounds ranges from 70-99%, while the removal of non-phenolics is generally <20%. However, relatively higher removal of some non-phenolic compounds, namely, diclofenac (40-50%), octocrylene (>80%) and ibuprofen (30-45%) has been reported because

these compounds contain both electron donating and electron withdrawing functional groups. Although both crude and purified LMEs demonstrated degradation of a range of pollutants, crude LMEs achieved better removal of some PPCPs such as diclofenac and naproxen compared to purified LMEs [27, 41, 81, 82, 83, 85], possibly due to the presence of natural mediators in crude LMEs secreted by WRF. For instance, Tran et al. [80] achieved almost complete removal of three pharmaceuticals namely diclofenac, ibuprofen and naproxen after treatment with crude enzyme extracted from *T. versicolor*, whereas purified laccase from *T. versicolor* and *A. oryzae* achieved 20-50% removal of these compounds [82, 83, 84, 103]. Additionally, utilization of crude LMEs for the treatment of PPCPs may considerably reduce the cost of the treatment process. However, extracellular extract *i.e.*, the crude enzyme solution also contains significant amount of the unspent growth media, and dosing crude enzyme means additional organic loading from this.

[FIGURE 1]

3.3. Performance of Redox-mediator Enhanced Laccase Catalyzed System

Laccase catalyzes mono-electronic oxidation of PPCPs. However, the extent of removal depends on the ORP of the enzyme and individual PPCP. Poor removal of non-phenolic PPCPs can be attributed to: (i) presence of a strong EWG in the structure of non-phenolic compounds causing steric hindrance; and (ii) higher ORP of non-phenolic compounds than laccase [102]. Removal of non-phenolic compounds can be enhanced by introducing a low-molecular weight redox-mediator. In a redox-mediator catalyzed system, highly reactive radicals are formed due to the oxidation of a mediator by laccase and these radicals then serve as an electron transfer shuttle between PPCPs and laccase, facilitating enhanced removal of recalcitrant compounds. Three oxidation mechanisms, namely hydrogen atom transfer (HAT), ionic mechanisms and electron transfer have been reported for mediators. For instance, 1-hydroxybenzotriazole (HBT) and syringaldehyde (SA) follow HAT mechanism, while 2,2-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid (ABTS) and 2,2,6,6-tetramethylpiperidinyloxy (TEMPO) follow electron transfer and ionic mechanisms, respectively [89, 109].

Properties of different redox-mediators used to enhance the removal of different PPCPs are summarized in Table 3. Although the mediators perform differently, two mediators namely HBT and SA have been commonly used to broaden the spectrum of compounds significantly degraded by laccase. Studies involving these mediators confirmed that they not only improved the ORP of reaction mixture but also the extent of removal. Only a few studies elucidated the performance of different mediators based on the type of substrate (i.e., phenolic vs. non-phenolic). For instance, N-OH type mediators (VA and HBT) were found to achieve the best removal of non-phenolic PPCPs such as clofibric acid, naproxen and carbamazepine, whereas SA and ABTS performed better for phenolic compounds such as salicylic acid and steroid hormones [82, 83, 85].

Mediator type, concentration and compound properties influence the performance of a mediator catalyzed system. For instance, removal of diclofenac improved from 40 to 80% by increasing the concentration of SA from 0.1 to 0.5 mM [103]. Similarly, Nguyen et al. [85] achieved an improvement of 35% in the removal of diclofenac in an enzymatic membrane reactor by increasing the dose of SA from 0.01 mM to 0.1 mM. However, beyond a threshold concentration increasing mediator dose may not improve the removal of PPCPs. For instance, Ashe et al. [89] observed that increasing the concentration of VA and ABTS from 0.5 to 1 mM could not enhance the removal of oxybenzone and naproxen. They also observed that removal of atrazine was reduced by 15-25 % when the concentration of VA and HBT was increased from 0.1 to 0.25 mM. This may be attributed to the complex interactions between laccase and the radicals generated due to degradation of the mediator by laccase, as discussed below.

[TABLE 3]

It has been observed in almost all the studies that enzymatic activity significantly drops with the addition of mediators. For instance, Hata et al. [111] observed 90% decline in enzymatic activity within first 8 h of incubation in the presence of HBT. Similarly, rapid decline in enzymatic activity was also observed with the addition of VA, HBT or ABTS [89], although rate of enzyme inactivation depends on the relative stability of the radicals generated by mediators. In the absence of any known enzyme inhibitor, rapid

enzyme inactivation in mediator catalyzed system can be attributed to: (i) the blockage of active sites of the enzyme by charged radicals and metabolites; and (ii) the reaction of metabolites with enzyme-active sites to convert them into non-productive complexes [112, 113].

Despite rapid inactivation of enzymes by different mediators, mediators can compensate by enhancing the rate of reaction, eventually achieving rapid and enhanced removal of PPCPs. However, periodic replenishment of enzyme is needed to maintain the removal efficiency of PPCPs, constraining the long term operation of mediator-enzyme based wastewater treatment processes. Therefore, selection of mediator type and its concentration is vital for effective and long term operation of such systems.

4. Impacts of Physicochemical Characteristics of Wastewater

Performance of WRF in wastewater treatment depends on a number of factors such as environmental conditions and physiochemical properties of the wastewater as well as the properties of PPCPs. Influence of PPCP properties on their overall removal in a WRF mediated treatment process has been reviewed by Yang et al. [31]. Briefly, physicochemical properties of water/wastewater such as pH, temperature and the presence of dissolved organic and/or inorganic compounds may influence WRF performance. Because information about these properties are vital to design and optimize WRF mediated treatment systems, these are discussed in this section.

The operating temperature not only affects the stability of WRF enzymatic systems but also the rate of reaction. It is believed that the rate of reaction increases with increase in temperature [88]. However, depending on the WRF strain, rapid thermal inactivation of LMEs has been observed at temperature $>40^{\circ}\text{C}$ [114, 115]. Only a few studies have investigated the impact of temperature on LME activity and removal efficiency in a reaction mixture. For instance, increase in temperature of the reaction mixture from 20 to 25°C enhanced the removal efficiency of chlorophenols in laccase mediated treatment [116]. Similarly, Ullah et al. [117] investigated the removal of pentachlorophenol by varying the temperature from $10\text{--}45^{\circ}\text{C}$ and found that the optimal temperature was 25°C to achieve maximum laccase activity and

pentachlorophenol removal. Temperature range of 37-40 °C has been reported to achieve optimal activity of MnP and LiP [87, 118].

Enzyme performance and substrate properties can be influenced by pH of the reaction mixture, thereby affecting the extent of PPCP removal. For instance, optimum pH for the removal of diclofenac (60-100%) by purified laccase (*T. versicolor* and *M. thermophila*) and LiP (*P. chrysosporium*) was in the range of 3.0 – 4.5 [83, 88, 103, 105]. Similarly, Wen et al. [87] observed that the removal of two antibiotics, namely tetracycline and oxytetracycline, by crude LiP and MnP was pH dependent and the best removal was achieved at pH 4.5-5.0. However, optimum pH for the removal of a personal care product triclosan was in the range of 5-5.5 during the treatment with laccase from *T. versicolor* [93, 116]. Media pH may not affect the removal of compounds highly amenable to laccase-catalyzed degradation. For instance, laccase from *T. versicolor* achieved significant removal (>70%) of steroid hormones over a pH range of 4 - 7 [103]. Similarly, changing the pH from 2.5- 6 could not improve the removal of carbamazepine by LiP [88]. In light of the above discussion, pH ranging from 4-5.5 is recommended for WRF mediated treatment processes.

The impact of dissolved organic and inorganic constituents on laccase activity and PPCP removal has been communicated in a limited number of studies. Many compounds such as sulphides, halides [119, 120], natural/synthetic organics [121, 122, 123, 124, 125, 126] and heavy metals [92, 127] can inhibit the catalytic activity of laccases [128]. Moreover, each compound may have different mode of laccase inhibition. For instance, fatty acids inhibit laccase catalytic potential by blocking the binding sites for phenolic substrates [129, 130, 131]. On the other hand, spectrophotometric assays, electron spin resonance spectroscopy and catalytic voltammetry analysis confirmed that anionic inhibitors such as halides and sulphides could block the access to the active copper sites in laccase [132, 133, 134]. Among anionic inhibitors, fluoride and azide are the most effective and can rapidly reduce the catalytic activity of laccase by 50% even at μM concentrations [135]. Although inhibition of laccases by halides can proceed in the following order: fluoride > chloride > bromide, the concentration of halides required to inhibit

laccases varies, with no fixed correlation with their inhibition potential. For instance, chloride concentration ranging from 100 μ M – 100 mM may cause a 50% drop in activity of laccase from different species [35, 136].

5. Removal Mechanisms

Removal mechanisms during treatment with WRF whole-culture include (i) sorption onto the fungal biomass; (ii) degradation by extracellular enzymes; and (iii) degradation by mycelium bound or intercellular enzymes. A schematic of fungal mediated treatment process with possible removal mechanisms is presented as Figure 2. Hydrophobicity (log D) of PPCPs is a key property that governs biosorption onto fungal biomass and could facilitate enhanced removal of some compounds. For instance, a batch study to investigate the contribution of biosorption and degradation by extracellular enzymes confirmed that hydrophobic PPCPs (log D>4) were highly removed by both mechanisms [41]. Moreover this study also confirmed that biosorption of significantly hydrophobic compounds facilitated the biodegradation of these compounds. On the other hand, a few studies have reported that removal of some PPCPs such as 17 β -Estradiol, 17 α -Ethinylestradiol, triclosan and nonylphenol by whole-cell WRF and extracellular enzymes is comparable (see Table 1 and 2), indicating a negligible impact of biosorption on their removal.

Biodegradation by whole-cell can be due to intracellular, extracellular and mycelium-associated enzymes. This can lead to significant differences in removal by whole-cell WRF and harvested enzyme. For instance, carbamazepine, containing a strong EWG, was significantly removed by some WRF species namely *P. ostreatus* (100%) [45] and *T. versicolor* ATCC 42530 (76%) [80], whereas crude [45, 80, 81] and purified laccase [82, 85] could only achieve 5-15% carbamazepine removal. Similarly, ibuprofen and naproxen were almost completely removed by whole-cell WRF [24, 43, 58]. In contrast, their removal by crude and purified laccase was in the range of 10-40% [27, 82, 85, 103]. Since both naproxen and ibuprofen are hydrophilic compounds (log D <3), role of biosorption in their removal would be limited. However, almost complete removal of these compounds in whole-cell WRF substantiates the role of

mycelium bound and/or or intercellular enzymes. Indeed, the role of intercellular enzyme (*i.e.* cytochrome P450) in the degradation of naproxen, diclofenac and carbamazepine has been demonstrated [43, 45, 137]. These studies showed that naproxen, diclofenac and carbamazepine were partially removal (15-50%) in the presence of cytochrome P450 inhibitor during whole-cell WRF treatment. Therefore, it can be concluded that high removal of some PPCPs in whole-cell WRF treatment is due to the synergetic effects of extracellular, intercellular and/or mycelium bound enzymes. Moreover, secretion of natural mediators may also help in enhancing the removal of these compounds.

[FIGURE 2]

6. Degradation Pathways and Identification of Intermediates

PPCP degradation pathways and their intermediate products have been identified for some compounds such as carbamazepine, diclofenac, triclosan and ibuprofen. However, each WRF species may follow a different degradation pathway. Degradation pathways and intermediate products of some PPCPs are discussed in this section.

Fungal mediated treatment of diclofenac starts with the conversion of the hydroxyl group in its structure into an intermediate product, namely, hydroxy diclofenac. Hydroxy diclofenac can be amenable to further fungal biodegradation [137, 138]. In *in vivo* and *in vitro* experiments for the degradation of diclofenac showed that laccase (*T. versicolor*) catalyzed degradation leads to the formation of biodegradable compounds such as: (i) hydroxylated metabolites (appeared in both *in vivo* and *in vitro* experiments); and (ii) 4-(2,6 dichlorophenylamino)-1,3-benzenedimethanol metabolite (appeared only in *in vivo* experiments) [138]. Diclofenac and its metabolites both disappeared after 24 h of incubation, reducing the ecotoxicity of the treated effluent. Degradation of ketoprofen was reported to initiate by the intercellular cytochrome P450 that converts ketoprofen into (2-[3-(4-hydroxybenzoyl)phenyl]-propanoic acid) and (2-[(3-hydroxy(phenyl)-methyl)phenyl]-propanoic acid) via hydroxylation and reduction of the ketone

group, respectively [138]. Moreover, it was also observed that the role of extracellular enzyme (laccase) in the degradation of ketoprofen was insignificant.

Both laccase and cytochrome P450 can catalyze the degradation of naproxen in WRF based treatment. Formation of two intermediates, namely 6-desmethylnaproxen and 1-(6-methoxynaphthalen-2-yl)ethanone), was observed possibly via P450-mediated desmethylation and laccase catalysis, respectively [43, 57]. Moreover, naproxen and associated intermediates were completely removed from the reaction mixture and the treated effluent was non-toxic [43].

Ibuprofen conversion predominantly starts with the formation of hydroxy-ibuprofen via hydroxylation. Marco-Urrea et al. [139] investigated the degradation pathways of ibuprofen by WRF *T. versicolor*. They observed that oxidation of isopropyl chain resulted in the formation of 1-hydroxy ibuprofen and 2-hydroxy ibuprofen. These intermediates were then degraded by *T. versicolor* to 1,2-dihydroxy ibuprofen, which was not further degraded. Hence, the ecotoxicity of treated effluent was higher than that of the initial solution [139].

Degradation pathways for carbamazepine by different whole-cell WRF including *T. versicolor* and *P. ostreatus* have also been studied. These have identified a number of stable intermediate products namely, 10,11-dihydro-10,11-dihydroxy-carbamazepine, acridine, acridone and 10,11-epoxy-carbamazepine. Interestingly, the treated effluent showed less toxicity [45, 53].

Intermediates or metabolites of triclosan during whole-cell treatment depends on WRF species. Major intermediates of triclosan during the treatment with *C. polyzona* WRF were dimer, trimer and tetramer [140], while 2-O-(2,4,4-trichlorodiphenyl ether)-b-D-xylopyranoside, 2-O-(2,4,4-trichlorodiphenyl ether)-b-D-glucopyranoside and 2,4-dichlorophenol were identified as major intermediates following its treatment with *T. versicolor* [141].

7. Future Scope of Research

Performance of whole-cell WRF has mostly been investigated in sterile batch tests. Only a few studies have been conducted in continuous flow reactors under non-sterile conditions [24, 26, 27, 58], because bacterial contamination can seriously affect WRF performance. Therefore, more efforts are required to solve this limitation.

Another limitation of WRF mediated treatment process is the washout of the enzyme and mediators with treated effluent. This is particularly problematic for enzymatic reactors where enzymes instead of whole-cell WRFs are used. Enzyme immobilization or coupling of enzymatic reactor with a membrane with suitable pore size can be used to avoid enzyme washout [27, 82, 83, 85, 142, 143]. Coupling of membrane separation with an enzymatic reactor offers several advantages over immobilized enzymatic systems such as: (i) more effective enzyme retention; (ii) reduced mass transfer limitations; (iii) ease of enzyme replenishment and (iv) ease of operation and maintenance [29, 144]. Moreover, enhanced TrOC removal by enzymatic membrane bioreactors (EMBR) has been demonstrated in a few recent studies [82, 83]. While investigating the removal of 30 TrOCs in an EMBR, Nguyen et al. [82] observed retention of PPCPs such as gemfibrozil, naproxen and ketoprofen on the enzyme gel layer formed over the membrane. It was also confirmed that the compounds retained by the enzyme gel layer were subsequently biodegraded [82]. Despite enhanced removal of some PPCPs, EMBR may not be effective to protect enzymes from thermal and chemical denaturation. In such situations, immobilized enzymatic systems have been reported to be more effective than free enzymes [145, 146, 147]. In enzyme-mediator systems, redox-mediators may still pass through the membrane with the treated effluent, thus increasing the operating costs. Washout of redox-mediators can be prevented by integrating a high retention membrane such as nanofiltration, membrane distillation or forward osmosis membrane with the enzymatic reactor, or formulating strategies to recover mediators from effluent.

Degradation pathways of only a few PPCPs such as carbamazepine, diclofenac, ibuprofen and naproxen have been reported in literature. The intermediate products in some cases are more toxic than the parent

compound. Therefore, more efforts are required to identify and elucidate the formation of intermediate products during WRF mediated treatment of PPCPs which would help to further understand the degradation mechanisms of PPCPs.

8. Conclusion

WRF mediated treatment of PPCPs is a promising and environmentally friendly technology. Varieties of PPCPs are efficiently removed by both whole-cell WRF and crude/purified LMEs. Hydrophilic and persistent PPCPs such as naproxen, ketoprofen and carbamazepine are significantly removed only in whole-cell WRF treatment because of the synergistic effects of extracellular and intercellular enzymes as well as sorption onto fungal biomass. Different redox-mediators have been introduced to enhance the removal of persistent PPCPs but continuous addition of mediators is an expensive practice. Therefore, mediator type and concentration must be selected carefully and techniques to recover mediators be optimized. Operating conditions of the enzymatic reactor including pH, temperature and interfering dissolved contaminants could influence the performance of WRF mediated treatment process. Despite the better performance of whole-cell WRF treatment than crude and purified enzymes, long term continuous operation of fungal bioreactor could not be possible due to bacterial contamination. This problems needs to be solved before their full scale applications.

Acknowledgments: This work has been partially funded by the GeoQuEST research centre of the University of Wollongong. A PhD scholarship to Muhammad B. Asif from the University of Wollongong is greatly appreciated.

Conflict of Interest: The authors declare no conflict of interest.

References

1. Combi T, Pintado-Herrera MG, Lara-Martin PA, Miserocchi S, Langone L, Guerra R. Distribution and fate of legacy and emerging contaminants along the Adriatic Sea: A comparative study. *Environ Pollut.* 2016;218:1055-64.
2. Luo Y, Guo W, Ngo HH, Nghiem LD, Hai FI, Zhang J, Liang S, Wang XC. A review on the occurrence of micropollutants in the aquatic environment and their fate and removal during wastewater treatment. *Sci Total Environ.* 2014;473:619-41.
3. Sui Q, Cao X, Lu S, Zhao W, Qiu Z, Yu G. Occurrence, sources and fate of pharmaceuticals and personal care products in the groundwater: a review. *Emerging Contaminants.* 2015;1(1):14-24.

4. Gavrilescu M, Demnerová K, Aamand J, Agathos S, Fava F. Emerging pollutants in the environment: present and future challenges in biomonitoring, ecological risks and bioremediation. *New Biotechnology*. 2015;32(1):147-56.
5. Pal A, He Y, Jekel M, Reinhard M, Gin KY-H. Emerging contaminants of public health significance as water quality indicator compounds in the urban water cycle. *Environ Int*. 2014;71:46-62.
6. Vandenberg LN, Colborn T, Hayes TB, Heindel JJ, Jacobs Jr DR, Lee D-H, Shioda T, Soto AM, vom Saal FS, Welshons WV. Hormones and endocrine-disrupting chemicals: low-dose effects and nonmonotonic dose responses. *Endocr Rev*. 2012;33(3):378-455.
7. Chen Z-F, Ying G-G. Occurrence, fate and ecological risk of five typical azole fungicides as therapeutic and personal care products in the environment: A review. *Environ Int*. 2015;84:142-53.
8. Ortiz de García S, Pinto GP, García-Encina PA, Mata RI. Ranking of concern, based on environmental indexes, for pharmaceutical and personal care products: An application to the Spanish case. *J Environ Manage*. 2013;129:384-97.
9. Camargo MC, García A, Riquelme A, Otero W, Camargo CA, Hernandez-García T, Candia R, Bruce MG, Rabkin CS. The problem of *Helicobacter pylori* resistance to antibiotics: a systematic review in Latin America. *The American Journal of Gastroenterology*. 2014;109(4):485-95.
10. Rizzo L, Manaia C, Merlin C, Schwartz T, Dagot C, Ploy M, Michael I, Fatta-Kassinos D. Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the environment: a review. *Sci Total Environ*. 2013;447:345-60.
11. Nazaret S, Aminov R. Role and prevalence of antibiosis and the related resistance genes in the environment. *Frontiers in Microbiology*. 2014;5.
12. Zhang Y, Arends JB, Van de Wiele T, Boon N. Bioreactor technology in marine microbiology: from design to future application. *Biotechnol Adv*. 2011;29(3):312-21.
13. Benner J, Helbling DE, Kohler H-PE, Wittebol J, Kaiser E, Prasse C, Ternes TA, Albers CN, Aamand J, Horemans B, Springael D, Walravens E, Boon N. Is biological treatment a viable alternative for micropollutant removal in drinking water treatment processes? *Water Res*. 2013;47(16):5955-76.
14. Hai FI, Yamamoto K, Fukushi K. Hybrid treatment systems for dye wastewater. *Crit Rev Environ Sci Technol*. 2007;37(4):315-77.
15. Hai FI, Alturki A, Nguyen LN, Price WE, Nghiem LD. Removal of Trace Organic Contaminants by Integrated Membrane Processes for Water Reuse Applications. In: Ngo HH, Guo W, Surampalli RY, Zhang TC, editors. *Green Technologies for Sustainable Water Management*. Reston: American Society of Civil Engineers; 2016. p. 533-78. (ISBN: 9780784414422).
16. Hai FI, Nghiem LD, Khan SJ, Price WE, Yamamoto K. Wastewater reuse: removal of emerging trace organic contaminants (TrOC). In: Hai FI, Yamamoto K, Lee C, editors. *Membrane Biological Reactors: Theory, Modeling, Design, Management and Applications to Wastewater Reuse*. United Kingdom: IWA Publishing; 2014. p. 165-205. (ISBN: 9781780400655).
17. Hai FI, Nguyen LN, Nghiem LD, Liao B-Q, Koyuncu I, Price WE. Trace Organic Contaminants Removal by Combined Processes for Wastewater Reuse. In: Fatta-Kassinos D, Dionysiou DD, Kümmerer K, editors. *Advanced Treatment Technologies for Urban Wastewater Reuse*. 45. New York: Springer; 2014. p. 39-77. (ISBN: 9783319238869).
18. Hai FI, Yamamoto K, Nakajima F, Fukushi K. Factors governing performance of continuous fungal reactor during non-sterile operation—the case of a membrane bioreactor treating textile wastewater. *Chemosphere*. 2009;74(6):810-7.
19. Hai FI, Yamamoto K, Nakajima F, Fukushi K. Removal of structurally different dyes in submerged membrane fungi reactor—Biosorption/PAC-adsorption, membrane retention and biodegradation. *J Membr Sci*. 2008;325(1):395-403.

20. Chen M, Xu P, Zeng G, Yang C, Huang D, Zhang J. Bioremediation of soils contaminated with polycyclic aromatic hydrocarbons, petroleum, pesticides, chlorophenols and heavy metals by composting: Applications, microbes and future research needs. *Biotechnol Adv.* 2015;33(6):745-55.
21. Anasonye F, Winkvist E, Kluczek-Turpeinen B, Räsänen M, Salonen K, Steffen KT, Tuomela M. Fungal enzyme production and biodegradation of polychlorinated dibenzo-p-dioxins and dibenzofurans in contaminated sawmill soil. *Chemosphere.* 2014;110:85-90.
22. Hai FI, Yamamoto K. Suitability of Membrane Bioreactor for treatment of recalcitrant textile dye wastewater utilising white-rot fungi. *International Journal of Environmental Engineering.* 2009;2(1-3):43-55.
23. Hai FI, Yamamoto K, Fukushi K. Development of a submerged membrane fungi reactor for textile wastewater treatment. *Desalination.* 2006;192(1):315-22.
24. Cruz-Morató C, Ferrando-Climent L, Rodríguez-Mozaz S, Barceló D, Marco-Urrea E, Vicent T, Sarrà M. Degradation of pharmaceuticals in non-sterile urban wastewater by *Trametes versicolor* in a fluidized bed bioreactor. *Water Res.* 2013;47(14):5200-10.
25. Rodríguez-Rodríguez CE, García-Galán MJ, Blánquez P, Díaz-Cruz MS, Barceló D, Caminal G, Vicent T. Continuous degradation of a mixture of sulfonamides by *Trametes versicolor* and identification of metabolites from sulfapyridine and sulfathiazole. *J Hazard Mater.* 2012;213:347-54.
26. Yang S, Hai FI, Nghiem LD, Nguyen LN, Roddick F, Price WE. Removal of bisphenol A and diclofenac by a novel fungal membrane bioreactor operated under non-sterile conditions. *Int Biodeterior Biodegrad.* 2013;85:483-90.
27. Nguyen LN, Hai FI, Yang S, Kang J, Leusch FD, Roddick F, Price WE, Nghiem LD. Removal of trace organic contaminants by an MBR comprising a mixed culture of bacteria and white-rot fungi. *Bioresour Technol.* 2013;148:234-41.
28. Rodríguez Porcel E, Casas López J, Sanchez Perez J, Chisti Y. Enhanced production of lovastatin in a bubble column by *Aspergillus terreus* using a two-stage feeding strategy. *J Chem Technol Biotechnol.* 2007;82(1):58-64.
29. Modin O, Hai FI, Nghiem LD, Basile A, Fukushi K. Gas-diffusion, extractive, biocatalytic and electrochemical membrane biological reactors. In: Hai FI, Yamamoto K, Lee C, editors. *Membrane Biological Reactors: Theory, Modeling, Design, Management and Applications to Wastewater Reuse.* United Kingdom IWA Publishing; 2014. p. 299-333. (ISBN: 9781780400655).
30. Hai FI, Nghiem LD, Modin O. Biocatalytic membrane reactors for the removal of recalcitrant and emerging pollutants from wastewater. In: Basile A, editor. *Handbook of Membrane Reactors: Reactor Types and Industrial Applications.* 2. United Kingdom: Woodhead Publishing Limited; 2013. p. 763-807. (ISBN: 9780857094155).
31. Yang S, Hai FI, Nghiem LD, Price WE, Roddick F, Moreira MT, Magram SF. Understanding the factors controlling the removal of trace organic contaminants by white-rot fungi and their lignin modifying enzymes: a critical review. *Bioresour Technol.* 2013;141:97-108.
32. Gao D, Du L, Yang J, Wu W-M, Liang H. A critical review of the application of white rot fungus to environmental pollution control. *Crit Rev Biotechnol.* 2010;30(1):70-7.
33. Tortella G, Durán N, Rubilar O, Parada M, Diez M. Are white-rot fungi a real biotechnological option for the improvement of environmental health? *Crit Rev Biotechnol.* 2015;35(2):165-72.
34. Kües U. Fungal enzymes for environmental management. *Curr Opin Biotechnol.* 2015;33:268-78.
35. Rodgers CJ, Blanford CF, Giddens SR, Skamnioti P, Armstrong FA, Gurr SJ. Designer laccases: a vogue for high-potential fungal enzymes? *Trends Biotechnol.* 2010;28(2):63-72.
36. Rouches E, Herpoël-Gimbert I, Steyer J, Carrere H. Improvement of anaerobic degradation by white-rot fungi pretreatment of lignocellulosic biomass: A review. *Renewable and Sustainable Energy Reviews.* 2016;59:179-98.

37. Pointing S. Feasibility of bioremediation by white-rot fungi. *Appl Microbiol Biotechnol*. 2001;57(1-2):20-33.
38. Lundell TK, Mäkelä MR, Hildén K. Lignin-modifying enzymes in filamentous basidiomycetes—ecological, functional and phylogenetic review. *J Basic Microbiol*. 2010;50(1):5-20.
39. Guillén F, Gómez-Toribio Vc, Martínez MaJ, Martínez AT. Production of hydroxyl radical by the synergistic action of fungal laccase and aryl alcohol oxidase. *Archives of Biochemistry and Biophysics*. 2000;383(1):142-7.
40. Bending GD, Friloux M, Walker A. Degradation of contrasting pesticides by white rot fungi and its relationship with ligninolytic potential. *FEMS Microbiol Lett*. 2002;212(1):59-63.
41. Yang S, Hai FI, Nghiem LD, Roddick F, Price WE. Removal of trace organic contaminants by nitrifying activated sludge and whole-cell and crude enzyme extract of *Trametes versicolor*. *Water Science and Technology*. 2013;67(6):1216-23.
42. Nguyen LN, Hai FI, Yang S, Kang J, Leusch FDL, Roddick F, Price WE, Nghiem LD. Removal of pharmaceuticals, steroid hormones, phytoestrogens, UV-filters, industrial chemicals and pesticides by *Trametes versicolor*: Role of biosorption and biodegradation. *Int Biodeterior Biodegrad*. 2014;88:169-75.
43. Marco-Urrea E, Pérez-Trujillo M, Blánquez P, Vicent T, Caminal G. Biodegradation of the analgesic naproxen by *Trametes versicolor* and identification of intermediates using HPLC-DAD-MS and NMR. *Bioresour Technol*. 2010;101(7):2159-66.
44. Hatakka A. Lignin-modifying enzymes from selected white-rot fungi: production and role from in lignin degradation. *FEMS Microbiol Rev*. 1994;13(2-3):125-35.
45. Golan-Rozen N, Chefetz B, Ben-Ari J, Geva J, Hadar Y. Transformation of the recalcitrant pharmaceutical compound carbamazepine by *Pleurotus ostreatus*: role of cytochrome P450 monooxygenase and manganese peroxidase. *Environ Sci Technol*. 2011;45(16):6800-5.
46. Marco-Urrea E, Gabarrell X, Sarrà M, Caminal G, Vicent T, Reddy CA. Novel aerobic perchloroethylene degradation by the white-rot fungus *Trametes versicolor*. *Environ Sci Technol*. 2006;40(24):7796-802.
47. Marco-Urrea E, Parella T, Gabarrell X, Caminal G, Vicent T, Reddy CA. Mechanistics of trichloroethylene mineralization by the white-rot fungus *Trametes versicolor*. *Chemosphere*. 2008;70(3):404-10.
48. Bernhardt R. Cytochromes P450 as versatile biocatalysts. *J Biotechnol*. 2006;124(1):128-45.
49. Paice MG, Jurasek L, Bourbonnais R, Archibald F. CORIOLUS (TRAMETES) VERSICOLOR. In: Kirk TK, Chang H, editors. *Biotechnology in Pulp and Paper Manufacture: Applications and Fundamental Investigations*. Reed Publishing USA; 2013. p. 123-30 (ISBN: 9780409901924).
50. Durán N, Cuevas R, Cordi L, Rubilar O, Diez MC. Biogenic silver nanoparticles associated with silver chloride nanoparticles (Ag@ AgCl) produced by laccase from *Trametes versicolor*. *SpringerPlus*. 2014;3(1):1.
51. Blánquez P, Guieysse B. Continuous biodegradation of 17 β -estradiol and 17 α -ethynylestradiol by *Trametes versicolor*. *J Hazard Mater*. 2008;150(2):459-62.
52. Cajthaml T, Křesinová Z, Svobodová K, Möder M. Biodegradation of endocrine-disrupting compounds and suppression of estrogenic activity by ligninolytic fungi. *Chemosphere*. 2009;75(6):745-50.
53. Jelic A, Cruz-Morató C, Marco-Urrea E, Sarrà M, Perez S, Petrović M, Barcelo D. Degradation of carbamazepine by *Trametes versicolor* in an air pulsed fluidized bed bioreactor and identification of intermediates. *Water Res*. 2012;46(4):955-64.
54. Bergman Å, Heindel JJ, Jobling S, Kidd KA, Zoeller RT, Jobling SK. State of the science of endocrine disrupting chemicals 2012: an assessment of the state of the science of endocrine disruptors prepared by a group of experts for the United Nations Environment Programme and World Health Organization. Switzerland: World Health Organization; 2013.

55. Rebuli ME, Patisaul HB. Assessment of sex specific endocrine disrupting effects in the prenatal and pre-pubertal rodent brain. *The Journal of Steroid Biochemistry and Molecular Biology*. 2016;160:148-59.
56. Rodarte-Morales A, Feijoo G, Moreira M, Lema J. Biotransformation of three pharmaceutical active compounds by the fungus *Phanerochaete chrysosporium* in a fed batch stirred reactor under air and oxygen supply. *Biodegradation*. 2012;23(1):145-56.
57. Rodarte-Morales A, Feijoo G, Moreira M, Lema J. Operation of stirred tank reactors (STRs) and fixed-bed reactors (FBRs) with free and immobilized *Phanerochaete chrysosporium* for the continuous removal of pharmaceutical compounds. *Biochem Eng J*. 2012;66:38-45.
58. Cruz-Morató C, Lucas D, Llorca M, Rodríguez-Mozaz S, Gorga M, Petrovic M, Barceló D, Vicent T, Sarrà M, Marco-Urrea E. Hospital wastewater treatment by fungal bioreactor: removal efficiency for pharmaceuticals and endocrine disruptor compounds. *Sci Total Environ*. 2014;493:365-76.
59. Moreira M, Feijoo G, Lema J. Fungal bioreactors: applications to white-rot fungi. *Reviews in Environmental Science and Biotechnology*. 2003;2(2-4):247-59.
60. Andleeb S, Atiq N, Robson GD, Ahmed S. An investigation of anthraquinone dye biodegradation by immobilized *Aspergillus flavus* in fluidized bed bioreactor. *Environmental Science and Pollution Research*. 2012;19(5):1728-37.
61. Kunii D, Levenspiel O. Bubbling bed model for kinetic processes in fluidized beds. Gas-solid mass and heat transfer and catalytic reactions. *Industrial & Engineering Chemistry Process Design and Development*. 1968;7(4):481-92.
62. Ozgun H, Dereli RK, Ersahin ME, Kinaci C, Spanjers H, van Lier JB. A review of anaerobic membrane bioreactors for municipal wastewater treatment: integration options, limitations and expectations. *Separation and Purification Technology*. 2013;118:89-104.
63. Mutamim NSA, Noor ZZ, Hassan MAA, Yuniarto A, Olsson G. Membrane bioreactor: applications and limitations in treating high strength industrial wastewater. *Chem Eng J*. 2013;225:109-19.
64. Hai FI, Yamamoto K, Lee C-H. *Membrane Biological Reactors: Theory, Modeling, Design, Management and Applications to Wastewater Reuse*: IWA Publishing; 2013. (ISBN:9781780400655).
65. Hai FI, Yamamoto K, Fukushi K, Nakajima F. Fouling resistant compact hollow-fiber module with spacer for submerged membrane bioreactor treating high strength industrial wastewater. *J Membr Sci*. 2008;317(1):34-42.
66. Babič J, Pavko A. Enhanced enzyme production with the pelleted form of *D. squalens* in laboratory bioreactors using added natural lignin inducer. *J Ind Microbiol Biotechnol*. 2012;39(3):449-57.
67. Cao J, Zhang H-J, Xu C-P. Culture characterization of exopolysaccharides with antioxidant activity produced by *Pycnoporus sanguineus* in stirred-tank and airlift reactors. *Journal of the Taiwan Institute of Chemical Engineers*. 2014;45(5):2075-80.
68. Ünyayar A, Demirbilek M, Turkoglu M, Celik A, Mazmanci MA, Erkurt EA, Ünyayar S, Cekic Ö, Atacag H. Evaluation of cytotoxic and mutagenic effects of *Coriolus versicolor* and *Funalia trogii* extracts on mammalian cells. *Drug Chem Toxicol*. 2006;29(1):69-83.
69. Tepwong P, Giri A, Ohshima T. Effect of mycelial morphology on ergothioneine production during liquid fermentation of *Lentinula edodes*. *Mycoscience*. 2012;53(2):102-12.
70. Espinosa-Ortiz EJ, Rene ER, Pakshirajan K, van Hullebusch ED, Lens PN. Fungal pelleted reactors in wastewater treatment: applications and perspectives. *Chem Eng J*. 2016;283:553-71.
71. Rodríguez-Rodríguez CE, Jelić A, Llorca M, Farré M, Caminal G, Petrović M, Barceló D, Vicent T. Solid-phase treatment with the fungus *Trametes versicolor* substantially reduces pharmaceutical concentrations and toxicity from sewage sludge. *Bioresour Technol*. 2011;102(10):5602-8.
72. Hai FI, Yamamoto K, Nakajima F, Fukushi K, Nghiem LD, Price WE, Jin B. Degradation of azo dye acid orange 7 in a membrane bioreactor by pellets and attached growth of *Coriolus versicolour*. *Bioresour Technol*. 2013;141:29-34.

73. Hai FI, Yamamoto K, Nakajima F, Fukushi K. Application of a GAC-coated hollow fiber module to couple enzymatic degradation of dye on membrane to whole cell biodegradation within a membrane bioreactor. *J Membr Sci.* 2012;389:67-75.
74. Libra JA, Borchert M, Banit S. Competition strategies for the decolorization of a textile-reactive dye with the white-rot fungi *Trametes versicolor* under non-sterile conditions. *Biotechnol Bioeng.* 2003;82(6):736-44.
75. Van Leeuwen J, Hu Z, Yi T, Pometto III A, Jin B. Kinetic model for selective cultivation of microfungi in a microscreen process for food processing wastewater treatment and biomass production. *Acta Biotechnol.* 2003;23(2-3):289-300.
76. Cheng Z, Xiang-Hua W, Ping N. Continuous Acid Blue 45 decolorization by using a novel open fungal reactor system with ozone as the bactericide. *Biochem Eng J.* 2013;79:246-52.
77. Sankaran S, Khanal SK, Pometto AL, van Leeuwen JH. Ozone as a selective disinfectant for nonaseptic fungal cultivation on corn-processing wastewater. *Bioresour Technol.* 2008;99(17):8265-72.
78. Blázquez P, Sarrà M, Vicent MT. Study of the cellular retention time and the partial biomass renovation in a fungal decolourisation continuous process. *Water Res.* 2006;40(8):1650-6.
79. Mir-Tutusa JA, Sarrà M, Caminal G. Continuous treatment of non-sterile hospital wastewater by *Trametes versicolor*: How to increase fungal viability by means of operational strategies and pretreatments. *J Hazard Mater.* 2016;318:561-70.
80. Tran NH, Urase T, Kusakabe O. Biodegradation characteristics of pharmaceutical substances by whole fungal culture *Trametes versicolor* and its laccase. *Journal of Water and Environment Technology.* 2010;8(2):125-40.
81. Nguyen LN, Hai FI, Kang J, Leusch FD, Roddick F, Magram SF, Price WE, Nghiem LD. Enhancement of trace organic contaminant degradation by crude enzyme extract from *Trametes versicolor* culture: Effect of mediator type and concentration. *Journal of the Taiwan Institute of Chemical Engineers.* 2014;45(4):1855-62.
82. Nguyen LN, Hai FI, Price WE, Kang J, Leusch FD, Roddick F, van de Merwe JP, Magram SF, Nghiem LD. Degradation of a broad spectrum of trace organic contaminants by an enzymatic membrane reactor: complementary role of membrane retention and enzymatic degradation. *Int Biodeterior Biodegrad.* 2015;99:115-22.
83. Nguyen LN, Hai FI, Price WE, Leusch FD, Roddick F, McAdam EJ, Magram SF, Nghiem LD. Continuous biotransformation of bisphenol A and diclofenac by laccase in an enzymatic membrane reactor. *Int Biodeterior Biodegrad.* 2014;95:25-32.
84. Nguyen LN, Hai FI, Price WE, Leusch FD, Roddick F, Ngo HH, Guo W, Magram SF, Nghiem LD. The effects of mediator and granular activated carbon addition on degradation of trace organic contaminants by an enzymatic membrane reactor. *Bioresour Technol.* 2014;167:169-77.
85. Nguyen LN, van de Merwe JP, Hai FI, Leusch FD, Kang J, Price WE, Roddick F, Magram SF, Nghiem LD. Laccase-syringaldehyde-mediated degradation of trace organic contaminants in an enzymatic membrane reactor: Removal efficiency and effluent toxicity. *Bioresour Technol.* 2016;200:477-84.
86. Spina F, Varese GC. Fungal Bioremediation of Emerging Micropollutants in Municipal Wastewaters. In: Purchase D, editor. *Fungal Applications in Sustainable Environmental Biotechnology.* Berlin, Germany: Springer; 2016. p. 115-41 (ISBN: 9783319428529).
87. Wen X, Jia Y, Li J. Enzymatic degradation of tetracycline and oxytetracycline by crude manganese peroxidase prepared from *Phanerochaete chrysosporium*. *J Hazard Mater.* 2010;177(1):924-8.
88. Zhang Y, Geißen S-U. In vitro degradation of carbamazepine and diclofenac by crude lignin peroxidase. *J Hazard Mater.* 2010;176(1):1089-92.

89. Ashe B, Nguyen LN, Hai FI, Lee D-J, van de Merwe JP, Leusch FD, Price WE, Nghiem LD. Impacts of redox-mediator type on trace organic contaminants degradation by laccase: Degradation efficiency, laccase stability and effluent toxicity. *Int Biodeterior Biodegrad*. 2016.
90. Wang J, Majima N, Hirai H, Kawagishi H. Effective removal of endocrine-disrupting compounds by lignin peroxidase from the white-rot fungus *Phanerochaete sordida* YK-624. *Curr Microbiol*. 2012;64(3):300-3.
91. Tran NH, Hu J, Urase T. Removal of the insect repellent N, N-diethyl-m-toluamide (DEET) by laccase-mediated systems. *Bioresour Technol*. 2013;147:667-71.
92. Kim Y-J, Nicell JA. Impact of reaction conditions on the laccase-catalyzed conversion of bisphenol A. *Bioresour Technol*. 2006;97(12):1431-42.
93. Kim YJ, Nicell JA. Laccase-catalysed oxidation of aqueous triclosan. *J Chem Technol Biotechnol*. 2006;81(8):1344-52.
94. Saito T, Kato K, Yokogawa Y, Nishida M, Yamashita N. Detoxification of bisphenol A and nonylphenol by purified extracellular laccase from a fungus isolated from soil. *J Biosci Bioeng*. 2004;98(1):64-6.
95. Tamagawa Y, Hirai H, Kawai S, Nishida T. Removal of estrogenic activity of endocrine-disrupting genistein by ligninolytic enzymes from white rot fungi. *FEMS Microbiol Lett*. 2005;244(1):93-8.
96. Suzuki K, Hirai H, Murata H, Nishida T. Removal of estrogenic activities of 17 β -estradiol and ethinylestradiol by ligninolytic enzymes from white rot fungi. *Water Res*. 2003;37(8):1972-5.
97. Lloret L, Hollmann F, Eibes G, Feijoo G, Moreira M, Lema J. Immobilisation of laccase on Eupergit supports and its application for the removal of endocrine disrupting chemicals in a packed-bed reactor. *Biodegradation*. 2012;23(3):373-86.
98. Auriol M, Filali-Meknassi Y, Tyagi RD, Adams CD. Laccase-catalyzed conversion of natural and synthetic hormones from a municipal wastewater. *Water Res*. 2007;41(15):3281-8.
99. Auriol M, Filali-Meknassi Y, Adams CD, Tyagi RD, Noguerol T-N, Pina B. Removal of estrogenic activity of natural and synthetic hormones from a municipal wastewater: efficiency of horseradish peroxidase and laccase from *Trametes versicolor*. *Chemosphere*. 2008;70(3):445-52.
100. Hirai H, Nakanishi S, Nishida T. Oxidative dechlorination of methoxychlor by ligninolytic enzymes from white-rot fungi. *Chemosphere*. 2004;55(4):641-5.
101. Hirano T, Honda Y, Watanabe T, Kuwahara M. Degradation of bisphenol A by the lignin-degrading enzyme, manganese peroxidase, produced by the white-rot basidiomycete, *Pleurotus ostreatus*. *Biosci, Biotechnol, Biochem*. 2000;64(9):1958-62.
102. d'Acunzo F, Galli C, Gentili P, Sergi F. Mechanistic and steric issues in the oxidation of phenolic and non-phenolic compounds by laccase or laccase-mediator systems. The case of bifunctional substrates. *New Journal of Chemistry*. 2006;30(4):583-91.
103. Lloret L, Eibes G, Lú-Chau TA, Moreira MT, Feijoo G, Lema JM. Laccase-catalyzed degradation of anti-inflammatories and estrogens. *Biochem Eng J*. 2010;51(3):124-31.
104. Margot J, Copin P-J, von Gunten U, Barry DA, Holliger C. Sulfamethoxazole and isoproturon degradation and detoxification by a laccase-mediator system: Influence of treatment conditions and mechanistic aspects. *Biochem Eng J*. 2015;103:47-59.
105. Margot J, Maillard J, Rossi L, Barry DA, Holliger C. Influence of treatment conditions on the oxidation of micropollutants by *Trametes versicolor* laccase. *New Biotechnology*. 2013;30(6):803-13.
106. Lloret L, Eibes G, Feijoo G, Moreira MT, Lema JM. Application of response surface methodology to study the removal of estrogens in a laccase-mediated continuous membrane reactor. *Biocatalysis Biotransformation*. 2013;31(4):197-207.
107. Becker D, Varela Della Giustina S, Rodriguez-Mozaz S, Schoevaart R, Barceló D, de Cazes M, Belleville M-P, Sanchez-Marcano J, de Gunzburg J, Couillerot O, Völker J, Oehlmann J, Wagner M.

Removal of antibiotics in wastewater by enzymatic treatment with fungal laccase – Degradation of compounds does not always eliminate toxicity. *Bioresour Technol.* 2016;219:500-9.

108. Lloret L, Eibes G, Moreira MT, Feijoo G, Lema JM. Removal of Estrogenic Compounds from Filtered Secondary Wastewater Effluent in a Continuous Enzymatic Membrane Reactor. Identification of Biotransformation Products. *Environ Sci Technol.* 2013;47(9):4536-43.

109. Astolfi P, Brandi P, Galli C, Gentili P, Gerini MF, Greci L, Lanzalunga O. New mediators for the enzyme laccase: mechanistic features and selectivity in the oxidation of non-phenolic substrates. *New Journal of Chemistry.* 2005;29(10):1308-17.

110. Murugesan K, Chang Y-Y, Kim Y-M, Jeon J-R, Kim E-J, Chang Y-S. Enhanced transformation of triclosan by laccase in the presence of redox mediators. *Water Res.* 2010;44(1):298-308.

111. Hata T, Shintate H, Kawai S, Okamura H, Nishida T. Elimination of carbamazepine by repeated treatment with laccase in the presence of 1-hydroxybenzotriazole. *J Hazard Mater.* 2010;181(1):1175-8.

112. Purich DL. Enzyme kinetics: catalysis & control: a reference of theory and best-practice methods. Netherlands: Elsevier; 2010. (ISBN: 9780123809247).

113. Khelifi R, Belbahri L, Woodward S, Ellouz M, Dhouib A, Sayadi S, Mechichi T. Decolourization and detoxification of textile industry wastewater by the laccase-mediator system. *J Hazard Mater.* 2010;175(1):802-8.

114. Mukhopadhyay A, Dasgupta AK, Chakrabarti K. Enhanced functionality and stabilization of a cold active laccase using nanotechnology based activation-immobilization. *Bioresour Technol.* 2015;179:573-84.

115. Sampaio LM, Padrão J, Faria J, Silva JP, Silva CJ, Dourado F, Zille A. Laccase immobilization on bacterial nanocellulose membranes: Antimicrobial, kinetic and stability properties. *Carbohydrate Polymers.* 2016;145:1-12.

116. Zhang J, Liu X, Xu Z, Chen H, Yang Y. Degradation of chlorophenols catalyzed by laccase. *Int Biodeterior Biodegrad.* 2008;61(4):351-6.

117. Ullah MA, Bedford CT, Evans CS. Reactions of pentachlorophenol with laccase from *Coriolus versicolor*. *Appl Microbiol Biotechnol.* 2000;53(2):230-4.

118. Bosco F, Capolongo A, Ruggeri B. Effect of temperature, pH, ionic strength, and sodium nitrate on activity of LiPs: Implications for bioremediation. *Bioremediation J.* 2002;6(1):65-76.

119. Sondhi S, Sharma P, Saini S, Puri N, Gupta N. Purification and characterization of an extracellular, thermo-alkali-stable, metal tolerant laccase from *Bacillus tequilensis* SN4. *PloS One.* 2014;9(5):e96951.

120. Hu X, Wang C, Wang L, Zhang R, Chen H. Influence of temperature, pH and metal ions on guaiacol oxidation of purified laccase from *Leptographium qinlingensis*. *World Journal of Microbiology and Biotechnology.* 2014;30(4):1285-90.

121. Bhattacharya S, Das A, Prashanthi K, Palaniswamy M, Angayarkanni J. Mycoremediation of Benzo [a] pyrene by *Pleurotus ostreatus* in the presence of heavy metals and mediators. *3 Biotech.* 2014;4(2):205-11.

122. Sun K, Luo Q, Gao Y, Huang Q. Laccase-catalyzed reactions of 17 β -estradiol in the presence of humic acid: Resolved by high-resolution mass spectrometry in combination with ¹³ C labeling. *Chemosphere.* 2016;145:394-401.

123. Sun J, Bostick BC, Mailloux BJ, Ross JM, Chillrud SN. Effect of oxalic acid treatment on sediment arsenic concentrations and lability under reducing conditions. *J Hazard Mater.* 2016;311:125-33.

124. Schmidt G, Krings U, Nimtz M, Berger RG. A surfactant tolerant laccase of *Meripilus giganteus*. *World Journal of Microbiology and Biotechnology.* 2012;28(4):1623-32.

125. Wan J, Zeng G, Huang D, Huang C, Lai C, Li N, Wei Z, Xu P, He X, Lai M. The oxidative stress of *Phanerochaete chrysosporium* against lead toxicity. *Appl Biochem Biotechnol.* 2015;175(4):1981-91.

126. Zeng G, Li N, Huang D, Lai C, Zhao M, Huang C, Wei Z, Xu P, Zhang C, Cheng M. The stability of Pb species during the Pb removal process by growing cells of *Phanerochaete chrysosporium*. *Appl Microbiol Biotechnol*. 2015;99(8):3685-93.
127. Murugesan K, Kim Y-M, Jeon J-R, Chang Y-S. Effect of metal ions on reactive dye decolorization by laccase from *Ganoderma lucidum*. *J Hazard Mater*. 2009;168(1):523-9.
128. D'Souza-Ticlo D, Sharma D, Raghukumar C. A thermostable metal-tolerant laccase with bioremediation potential from a marine-derived fungus. *Mar Biotechnol*. 2009;11(6):725-37.
129. Zhang J, Kjonaas R, Flurkey WH. Does N-hydroxyglycine inhibit plant and fungal laccases? *Phytochemistry*. 1999;52(5):775-83.
130. Gianfreda L, Sannino F, Filazzola M, Leonowicz A. Catalytic behavior and detoxifying ability of a laccase from the fungal strain *Cerrena unicolor*. *Journal of Molecular Catalysis B: Enzymatic*. 1998;4(1):13-23.
131. Kreuter T, Steudel A, Pickert H. On the inhibition of laccase by lower fatty acids. *Acta Biotechnol*. 1991;11(1):81-3.
132. Xu F. Dioxygen reactivity of laccase. *Appl Biochem Biotechnol*. 2001;95(2):125-33.
133. Naqui A, Varfolomeev S. Inhibition mechanism of *Polyporus* laccase by fluoride ion. *FEBS Lett*. 1980;113(2):157-60.
134. Blanford CF, Foster CE, Heath RS, Armstrong FA. Efficient electrocatalytic oxygen reduction by the 'blue' copper oxidase, laccase, directly attached to chemically modified carbons. *Faraday Discussions*. 2009;140:319-35.
135. Bento I, Martins LO, Lopes GG, Carrondo MA, Lindley PF. Dioxygen reduction by multi-copper oxidases; a structural perspective. *Dalton Transactions*. 2005(21):3507-13.
136. Xu F. Oxidation of phenols, anilines, and benzenethiols by fungal laccases: correlation between activity and redox potentials as well as halide inhibition. *Biochemistry*. 1996;35(23):7608-14.
137. Hata T, Kawai S, Okamura H, Nishida T. Removal of diclofenac and mefenamic acid by the white rot fungus *Phanerochaete sordida* YK-624 and identification of their metabolites after fungal transformation. *Biodegradation*. 2010;21(5):681-9.
138. Marco-Urrea E, Pérez-Trujillo M, Cruz-Morató C, Caminal G, Vicent T. Degradation of the drug sodium diclofenac by *Trametes versicolor* pellets and identification of some intermediates by NMR. *J Hazard Mater*. 2010;176(1):836-42.
139. Marco-Urrea E, Pérez-Trujillo M, Vicent T, Caminal G. Ability of white-rot fungi to remove selected pharmaceuticals and identification of degradation products of ibuprofen by *Trametes versicolor*. *Chemosphere*. 2009;74(6):765-72.
140. Cabana H, Jones J, Agathos SN. Elimination of endocrine disrupting chemicals using white rot fungi and their lignin modifying enzymes: a review. *Eng Life Sci*. 2007;7(5):429-56.
141. Hundt K, Martin D, Hammer E, Jonas U, Kindermann MK, Schauer F. Transformation of triclosan by *Trametes versicolor* and *Pycnoporus cinnabarinus*. *Appl Environ Microbiol*. 2000;66(9):4157-60.
142. Nair RR, Demarche P, Agathos SN. Formulation and characterization of an immobilized laccase biocatalyst and its application to eliminate organic micropollutants in wastewater. *New Biotechnology*. 2013;30(6):814-23.
143. Ansari SA, Husain Q. Potential applications of enzymes immobilized on/in nano materials: A review. *Biotechnol Adv*. 2012;30(3):512-23.
144. Lloret L, Eibes G, Feijoo G, Moreira M, Lema J. Degradation of estrogens by laccase from *Myceliophthora thermophila* in fed-batch and enzymatic membrane reactors. *J Hazard Mater*. 2012;213:175-83.
145. Hou J, Dong G, Luu B, Sengpiel RG, Ye Y, Wessling M, Chen V. Hybrid membrane with TiO₂ based bio-catalytic nanoparticle suspension system for the degradation of bisphenol-A. *Bioresour Technol*. 2014;169:475-83.

146. Hou J, Dong G, Ye Y, Chen V. Laccase immobilization on titania nanoparticles and titania-functionalized membranes. *J Membr Sci.* 2014;452:229-40.
147. Nguyen LN, Hai FI, Dosseto A, Richardson C, Price WE, Nghiem LD. Continuous adsorption and biotransformation of micropollutants by granular activated carbon-bound laccase in a packed-bed enzyme reactor. *Bioresour Technol.* 2016;210:108-16.

List of Tables

Table 1: Removal (%) of PPCPs by different species of WRF (whole-cell) under different operating conditions

WRF Specie	Reactor Type	Operating conditions	PPCPs	Initial concentration (mg/L)	Removal efficiency (%)	References
<i>P. chrysosporium</i> (LiP, MnP)	Stirred tank (Batch-fed)	Inoculum= 1.2 g/L	Diclofenac	0.8	>99	[56]
		Reactor volume = 2 L, HRT= 24 h	Ibuprofen	0.8	75-99	
		Mixing speed = 200 rpm	Naproxen	0.8	>99	
		pH = 4.5; Temperature= 30°C				
	Stirred tank (Continuous)	Electron donor= glucose				[57]
		Operating time= 30 days				
		Reactor volume = 1.5 L	Diclofenac	1	92	
		HRT= 24 h	Ibuprofen	1	95	
		pH = 4.5; Temperature= 30°C	Naproxen	1	95	
		Electron donor= glucose	Carbamazepine	0.5	25-60	
		Operating time= 50 days	Diazepam	0.25-0.5	0	
<i>T. versicolor</i> (Laccase, LiP, MnP)	Fixed bed (Continuous)	Inoculum= 3.2 g/L				[51]
		Reactor volume = 0.13 L	17β-estradiol (E1)	3-18.8	>95	
		HRT= 120 days	17α-ethynylestradio (EE2)	7.3	>95	
		pH = 4.5; Temperature= 22°C				
	Fluidized bed (Batch-fed)	Electron donor= glucose				[53]
		Operating time= 26 days				
		Inoculum= 3.8 g	Carbamazepine	0.05-9	61-94	
		Reactor volume = 1.5 L				
		pH = 4.5; Temperature= 25°C				
		Electron donor= glucose				
	Fluidized bed (Batch-fed)	Operating time= 15 days				[24]
		Inoculum= 1.5 g/L	Ibuprofen	2.34	100	
		Reactor volume = 10 L	Acetaminophen	1.56	100	
		pH = 4.5; Temperature= 25°C	Ketoprofen	0.08	100	
		Electron donor= glucose	Ciprofloxacin	84.71	35	
		Operating time= 8 days	Azithromycin	4.31	100	
		Real hospital wastewater	Propranolol	0.06	100	
	Fluidized bed	Non sterile conditions	Carbamazepine	0	-50	[58]
		Inoculum= 1.4 g/L	Acetaminophen	109	100	

(Continuous)	Reactor volume = 10 L pH = 4.5; Temperature= 25°C Electron donor= glucose Operating time= 8 days Real hospital wastewater Non sterile conditions	Naproxen	1.62	100	
		Ibuprofen	35.5	100	
		Ketoprofen	2.17	95	
		Diclofenac	0.477	100	
		Codeine	0.606	100	
		Phenazone	0.497	96	
		Salicylic Acid	0.606	0	
		Ofloxacin	3.34	98	
		Ciprofloxacin	13.0	99	
		Sulfamethoxazole	1.41	100	
		Trimethoprim	0.853	100	
		Metronidazole	0.912	85	
		Azithromycin	1.37	26	
		Clarithromycin	2.20	80	
		Erithromycin	0.008	100	
		Tetracyclin	0.011	0	
		Caffeine	149	39	
		Carbamazepine	0.056	0	
		Atenolol	2.99	75	
		Metoprolol	0.019	95	
Membrane bioreactor (Continuous)	Inoculum= 3.5 g/L	Diclofenac	0.3-1.5	55	[26]
	Reactor volume = 5.5 L, HRT= 1 day				
	pH = 5.4; Temperature= 27°C				
	Operating time= 90 days				
	Non sterile conditions				[27]
Membrane bioreactor (Continuous)	Inoculum= 3 g/L	Ibuprofen	0.005	95	
	Reactor volume = 5.5 L, HRT= 2 day	Naproxen	0.005	95	
	pH = 4.5; Temperature= 27°C	Ketoprofen	0.005	90	
	Operating time= 110 days	Diclofenac	0.005	50	
	Non sterile conditions	Carbamazepine	0.005	20	
		Metronidazole	0.005	40	
		Gemfibrozil	0.005	95	
		Amitriptyline	0.005	85	
		Estriol (E3)	0.005	>95	
		17-β-estradiol	0.005	>95	

			17- β -estradiol – acetate	0.005	>95	
			17- α ethinylestradiol	0.005	>95	
			Triclosan	0.005	95	
			Benzophenone	0.005	80	
			Oxybenzone	0.005	92	
			Octocrylene	0.005	94	
	Erlenmeyer flask (Batch-fed)	Inoculum= 2-3 mg/test	Triclosan	10	98	
		pH = 4.5; Temperature= 28°C	17- α ethinylestradiol	10	94	[52]
		Operating time= 14 days Electron donor: glucose	Nonylphenol	10	90	
	Erlenmeyer flask (Batch-fed)	Inoculum= 2-3 mg	Triclosan	10	98	
<i>P. ostreatus</i> (Laccase, MnP)		pH = 4.5; Temperature= 28°C	17- α ethinylestradiol	10	62	[52]
		Operating time= 14 days Electron donor: glucose	Nonylphenol	10	93	
	Erlenmeyer flask (Batch-fed)	Inoculum= 2-3 mg	Triclosan	10	98	
<i>D. squalens</i> (Laccase, MnP)		pH = 4.5; Temperature= 28°C	17- α ethinylestradiol	10	78	[52]
		Operating time= 14 days	Nonylphenol	10	85	
	Erlenmeyer flask (Batch-fed)	Inoculum= 2-3 mg	Triclosan	10	98	
<i>B. adusta</i> (Laccase, LiP, MnP)		pH = 4.5; Temperature= 28°C	17- α ethinylestradiol	10	78	[52]
		Operating time= 14 days	Nonylphenol	10	85	

Table 2: Performance of crude LME for the removal (%) of PPCPs

PPCPs	Laccase ^[80, 81, 89, 91, 92, 93, 94, 95, 96, 97, 98, 99]			LiP ^[88, 90, 100]			MnP ^[80, 95, 96, 101]		
Non-phenolic compounds	Removal (%)	Initial concentration (mg/L)	Incubation time (h)	Removal (%)	Initial concentration (mg/L)	Incubation time (h)	Removal (%)	Initial concentration (mg/L)	Incubation time (h)
Carbamazepine	5 – 37	0.01-0.1	24-48	<10	5	2	14-20	0.01-4.7	24-48
Ibuprofen	<5 – 38	0.01-0.1	24-48	-	-	-	20	0.01	48
Naproxen	20-100	0.01-0.5	24-48	-	-	-	95	0.01	48
Diclofenac	60-100	0.01-0.1	3-48	>99	5	2	100	0.01	48
Gemfibrozil	20-25	0.01-0.1	24-48	-	-	-	30	0.01	48
ketoprofen	<5 -12	0.01-0.1	24-48	-	-	-	22	0.01	48
Clofebric acid	10-20	0.01-0.1	24-48	-	-	-	<10	0.01	48
Benzophenone	<5	0.1	24	-	-	-	-	-	-
DEET	20-55	0.01	4	-	-	-	-	-	-
Octocrylene	20	0.1	24	-	-	-	-	-	-
Phenolic compounds									
Estrone	70-100	0.1-27	1-24	60	27	24	>99	5	8
17 β -Estradiol	>99	0.01-5	1-24	40, 85	0.6, 27	1, 24	>99	2.96	1
17 α -Ethinylestradiol	>99	0.1-2.96	1-24	20, 82	6.6, 30	8, 24	>99	2.96	1
Oxybenzone	10-25	0.1-0.5	24	-	-	-	-	-	-
Nonylphenol	100	22	1	-	-	-	-	-	-
Triclosan	70-90	0.1-144	24	-	-	-	-	-	-
“-”: not reported									

Table 3: Properties of redox-mediators used to improve the performance of laccase based treatment of PPCPs

Redox-mediator	Type of mediator	Free radicals	Oxidation mechanism	Application for TrOC removal	References
1-hydroxibenzotriazole (HBT)	N-OH/ synthetic	aminoxyl	Hydrogen atom transfer	PPCPs, EDCs, pesticides and industrial chemicals	[43, 81, 89, 104, 105, 110]
Violuric acid (VA)	N-OH/ natural	aminoxyl	Hydrogen atom transfer	PPCPs, pesticides and industrial chemicals	[89, 90]
N-hydroxyphthalimide (HPI)	N-OH/ synthetic	aminoxyl	Hydrogen atom transfer	PPCPs and pesticides	[89]
Syringaldehyde (SA)	C ₆ H ₄ (OH)(OCH ₃)/ natural	phenoxyl	Hydrogen atom transfer	PPCPs, EDCs, pesticides and industrial chemicals	[82, 83, 84, 85, 89]
Vanillin (VAN)	C ₆ H ₄ (OH)(OCH ₃)/ synthetic	phenoxyl	Hydrogen atom transfer	PPCPs, EDCs, pesticides and industrial chemicals	[89, 90, 110]
2,2,6,6-tetramethylpiperidinyloxyl (TEMPO)	N – O / synthetic	Oxoammonium	Ionic	PPCPs, EDCs, pesticides and industrial chemicals	[89, 90, 110]
2,2-azino-nis(3-ethylbenzothiazoline-6-sulfonic acid (ABTS)	ABTS/ synthetic	ABTS ⁺ and ABTS ⁺⁺	Electron transfer	PPCPs, EDCs, pesticides and industrial chemicals	[89, 90]

List of Figures

Figure 1. Average removal of phenolic and non-phenolic PPCPs after treatment with purified/commercially available laccase. Error bar indicates average $\pm$ standard deviation. Numbers within parenthesis indicates number of data points. Data was collected from the following studies: [82, 83, 84, 85, 103, 104, 105, 106, 107, 108].

Figure 2. A schematic representation of whole-cell WRF mediated treatment process