



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science - Papers (Archive)

Faculty of Science, Medicine and Health

2009

Responses of plants in polar regions to UVB exposure : a meta-analysis

K. K. Newsham

British Antarctic Survey, Natural Environment Research Council, UK

Sharon A. Robinson

University of Wollongong, sharonr@uow.edu.au

Publication Details

This article was originally published as Newsham, KK and Robinson, SA, Responses of plants in polar regions to UV-B radiation : a meta-analysis, *Global Change Biology*, 15, 2009, 2574-2589. The definitive version is available at www3.interscience.wiley.com

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Responses of plants in polar regions to UVB exposure : a meta-analysis

Abstract

We report a meta-analysis of data from 34 field studies into the effects of UV-B radiation on Arctic and Antarctic bryophytes and angiosperms. The studies measured plant responses to decreases in UV-B radiation under screens, natural fluctuations in UV-B irradiance, or increases in UV-B radiation applied from fluorescent UV lamps. Exposure to UV-B radiation was found to increase the concentrations of UV-B absorbing compounds in leaves or thalli by 7% and 25% (expressed on a mass or area basis, respectively). UV-B exposure also reduced aboveground biomass and plant height by 15% and 10%, respectively, and increased DNA damage by 90%. No effects of UV-B exposure were found on carotenoid or chlorophyll concentrations, net photosynthesis, F_v/F_m or $\Phi PSII$, belowground or total biomass, leaf mass, leaf area or specific leaf area (SLA). The methodology adopted influenced the concentration of UV-B absorbing compounds, with screens and natural fluctuations promoting significant changes in the concentrations of these pigments, but lamps failing to elicit a response. Greater reductions in leaf area and SLA, and greater increases in concentrations of carotenoids, were found in experiments based in Antarctica than in those in the Arctic. Bryophytes typically responded in the same way as angiosperms to UV-B exposure. Regression analyses indicated that the percentage difference in UV-B dose between treatment and control plots was positively associated with concentrations of UV-B absorbing compounds and carotenoids, and negatively so with aboveground biomass and leaf area. We conclude that, despite being dominated by bryophytes, the vegetation of polar regions responds to UV-B exposure in a similar way to higher plant-dominated vegetation at lower latitudes. In broad terms, the exposure of plants in these regions to UV-B radiation elicits the synthesis of UV-B absorbing compounds, reduces aboveground biomass and height, and increases DNA damage.

Keywords

aboveground biomass, angiosperms, Antarctic, Arctic, bryophytes, DNA damage, flavonoids, height, methodology, UVB absorbing compounds

Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

This article was originally published as Newsham, KK and Robinson, SA, Responses of plants in polar regions to UV-B radiation : a meta-analysis, *Global Change Biology*, 15, 2009, 2574-2589. The definitive version is available at www3.interscience.wiley.com

1 Responses of plants in polar regions to UV-B exposure: a 2 meta-analysis

3

4 Kevin K. Newsham* and Sharon A. Robinson†

5

6 *British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road,
7 Cambridge, CB3 0ET, UK,

8 †Institute for Conservation Biology, University of Wollongong, NSW 2522, Australia

9

10 Correspondence: K.K. Newsham, tel. +44/ 1223 221400, fax +44/ 1223 362616, e-mail:

11 kne@bas.ac.uk

12

13 *Keywords:* aboveground biomass, angiosperms, Antarctic, Arctic, bryophytes, DNA damage,
14 flavonoids, height, methodology, UV-B absorbing compounds

15

16 *Running title:* meta-analysis of polar plant response to UV-B

17

18 Abstract

19 We report a meta-analysis of data from 34 field studies into the effects of UV-B radiation on Arctic and
20 Antarctic bryophytes and angiosperms. The studies measured plant responses to decreases in UV-B
21 radiation under screens, natural fluctuations in UV-B irradiance, or increases in UV-B radiation applied
22 from fluorescent UV lamps. Exposure to UV-B radiation was found to increase the concentrations of
23 UV-B absorbing compounds in leaves or thalli by 7% and 25% (expressed on a mass or area basis,
24 respectively). UV-B exposure also reduced aboveground biomass and plant height by 15% and 10%,
25 respectively, and increased DNA damage by 90%. No effects of UV-B exposure were found on
26 carotenoid or chlorophyll concentrations, net photosynthesis, F_v/F_m or Φ_{PSII} , belowground or total
27 biomass, leaf mass, leaf area or specific leaf area (SLA). The methodology adopted influenced the
28 concentration of UV-B absorbing compounds, with screens and natural fluctuations promoting
29 significant changes in the concentrations of these pigments, but lamps failing to elicit a response.
30 Greater reductions in leaf area and SLA, and greater increases in concentrations of carotenoids, were
31 found in experiments based in Antarctica than in those in the Arctic. Bryophytes typically responded in
32 the same way as angiosperms to UV-B exposure. Regression analyses indicated that the percentage
33 difference in UV-B dose between treatment and control plots was positively associated with
34 concentrations of UV-B absorbing compounds and carotenoids, and negatively so with aboveground
35 biomass and leaf area. We conclude that, despite being dominated by bryophytes, the vegetation of
36 polar regions responds to UV-B exposure in a similar way to higher plant-dominated vegetation at

1 lower latitudes. In broad terms, the exposure of plants in these regions to UV-B radiation elicits the
2 synthesis of UV-B absorbing compounds, reduces aboveground biomass and height, and increases
3 DNA damage.

5 **Introduction**

6 The depletion of ozone in the stratosphere over polar regions is particularly intense (Farman *et al.*,
7 1985; Newman *et al.*, 1997). Chemical reactions on the surfaces of polar stratospheric clouds, which
8 form each winter in cold (< -78 °C) air masses over the Arctic and Antarctic, convert chlorine, derived
9 from chlorofluorocarbons, into chlorine monoxide, which photocatalyses ozone to oxygen (World
10 Meteorological Organization, 2007). Because ozone is the principal gas in the atmosphere that absorbs
11 solar ultraviolet-B radiation (UV-B; 280-315 nm), its depletion exposes plants in polar regions to
12 elevated irradiances of UV-B during boreal or austral spring, often as the plants emerge from melting
13 snow and ice. Since UV-B is absorbed by DNA and biological membranes, these increased irradiances
14 of UV-B have possible deleterious effects on plant physiology and growth, and consequences for the
15 functioning of these ecosystems (Caldwell *et al.*, 1995; Rozema *et al.*, 1997; Paul & Gwynn-Jones,
16 2003).

17 In recent years, considerable effort has been invested in determining the responses of polar
18 vegetation, which is dominated by bryophytes, to UV-B exposure. However, the UV-B responses of
19 plants in polar regions are at present poorly defined, with wide variation in responses found between
20 different studies. For example, negative effects on biomass accumulation have been recorded in some
21 experiments (e.g. Xiong & Day, 2001; Xiong *et al.*, 2002; Robinson *et al.*, 2005) but not in others (e.g.
22 Lappalainen *et al.*, 2008). The responses of photosynthetic pigments to UV-B exposure also often
23 differ between studies, with different experiments showing either increases (e.g. Niemi *et al.*, 2002a),
24 no change (e.g. Niemi *et al.*, 2002b; Lappalainen *et al.*, 2008) or decreases (e.g. Gehrke, 1998; 1999;
25 Robinson *et al.*, 2005) in the concentration of chlorophylls in leaves or thalli. Damage to DNA in
26 aboveground plant parts also varies widely, with increases in damage in some studies (e.g. Lud *et al.*,
27 2001a; Turnbull & Robinson, 2009) but no apparent effects in others (e.g. Lud *et al.*, 2003; Boelen
28 *et al.*, 2006; Rozema *et al.*, 2006). One of the most consistent responses of plants at lower latitudes to
29 elevated UV-B exposure is the synthesis of UV-B absorbing compounds in foliage (Searles *et al.*,
30 2001), but, even for this response, polar UV-B experiments have yielded conflicting results, with
31 consistent positive effects on the concentrations of these pigments in some studies (e.g. Xiong & Day,
32 2001; Newsham *et al.*, 2002; Newsham, 2003), but no effects in others (e.g. Gehrke, 1998; 1999;
33 Boelen *et al.*, 2006; Rozema *et al.*, 2006).

34 The reasons for the disparate results from these studies are presently unclear. One potential
35 factor is the differences between studies in the methodologies used to alter the dose of UV-B radiation
36 received by plants. Three main approaches have been used in polar UV-B experiments. The first of
37 these is to cover plants with screens made from materials that either absorb or transmit UV-B radiation,
38 with Mylar polyester (cut off $\lambda = 314$ nm) typically being used to remove UV-B from solar radiation.
39 The responses of plants under this material are compared with those of plants under materials such as
40 Aclar, Teflon, or polymethylmethacrylate (PMMA), each of which transmit most or all of the

1 wavelengths of solar UV-B radiation that reach the Earth's surface (e.g. Xiong & Day, 2001; Newsham
2 *et al.*, 2005; Albert *et al.*, 2008). The second approach is to use fluorescent UV lamps, suspended over
3 plants from frames, to increase the flux of UV-B in solar radiation, typically simulating between 15%
4 and 30% loss of ozone from the atmosphere, assuming cloudless skies over the study site (e.g. Gehrke,
5 1998; 1999; Björn *et al.*, 1999; Rozema *et al.*, 2006). In order to apply UV-B radiation to treatment
6 plots and to remove wavelengths of UV-C (200-280 nm), which are not present in solar radiation
7 reaching the Earth's surface, energised fluorescent UV lamps are wrapped in cellulose acetate (cut off λ
8 = 292 nm). The responses of plants under cellulose acetate-filtered lamps are then compared with those
9 in control plots, under energised lamps wrapped in Mylar polyester or encased in glass (cut off λ = 318
10 nm; Gehrke, 1999). The third, and much less commonly used approach, is to examine plant response to
11 natural fluctuations in ambient UV-B irradiance. This non-manipulative approach samples plants
12 repeatedly from the same location in the natural environment under varying ozone column depths,
13 exposing plants to wide fluctuations in UV-B irradiance (e.g. Newsham *et al.*, 2002; Dunn & Robinson,
14 2006).

15 A further explanation for the disparities that exist between studies is the wide variation in the
16 doses of UV-B radiation received by plants in treatment and control plots. Such variation is often
17 associated with the methodologies used in different experiments. For example, UV-B attenuating
18 screens reduce incident UV-B dose by up to 86% (Xiong *et al.*, 2002), whereas lamps, depending on
19 which action spectrum the spectral irradiances are weighted with and the level of ozone depletion
20 simulated, are typically used to increase the dose of UV-B radiation received by plants by *c.* 30%
21 above that present in solar radiation (e.g. Gehrke, 1998; 1999; Björn *et al.*, 1999; Niemi *et al.*, 2002a).
22 The substantial effect of UV-B attenuating screens on the dose of UV-B radiation received by plants
23 has been put forward as the reason for the consistent, but as yet unquantified, effects of screens on plant
24 performance, compared with the apparently less consistent effects of fluorescent UV lamps (Rozema *et*
25 *al.*, 2005). This is a difficult issue to address, since studies often do not report the absolute UV-B doses
26 received by plants in treatment and control plots, or use different biological action spectra to weight the
27 calculated UV-B doses, hampering the direct comparison of data from different studies.

28 Another potential factor that could account for the differences between studies in plant
29 response to UV-B exposure is the taxa that are studied. Different plant forms are known to transmit
30 different amounts of UV-B radiation to the mesophyll (Day *et al.*, 1992; Day & Vogelmann, 1993),
31 possibly because of the presence of hairs, cuticles and surface waxes on leaves (Day, 1993), which are
32 more frequent in angiosperms and gymnosperms than in bryophytes (Gehrke, 1999). It has also been
33 suggested that higher plants and bryophytes could differ in their abilities to synthesize UV-B absorbing
34 compounds (Gwynn-Jones *et al.*, 1999), although recent data do not support this argument (Newsham
35 *et al.*, 2002; Newsham, 2003; Dunn & Robinson, 2006; Clarke & Robinson, 2008). In addition,
36 previous experiments may have underestimated the UV-B screening potential of bryophytes, since a
37 recent study has demonstrated that UV-B absorbing compounds located in the cell walls of Antarctic
38 mosses are not efficiently extracted with acidified methanol (Clarke & Robinson, 2008).

39 In order to resolve why these differences exist in plant response to UV-B exposure between
40 studies, and to identify common effects of UV-B radiation on polar vegetation, a meta-analysis of data

1 from the literature on the UV-B responses of plants in polar regions was performed. Meta-analyses
2 have been used in similar ways to clarify the responses of plants to warming (Arft *et al.*, 1999),
3 elevated concentrations of carbon dioxide and ozone in air (Curtis & Wang, 1998; Stiling and
4 Cornelissen, 2007; Feng *et al.*, 2008) and simulated ozone depletion (Searles *et al.*, 2001). The latter
5 study analysed data on higher plant responses to UV-B radiation applied from fluorescent UV lamps,
6 typically at mid-latitudes, with data from only three polar field experiments included in the analyses. In
7 the current study, we augment the analyses of Searles *et al.* (2001) by including data from screen and
8 non-manipulative studies, the latter of which did not exist in the literature until 2002. We aimed to
9 determine whether differences in plant response to UV-B exposure can be ascribed to variation in
10 methodology, plant form, UV-B dosage or the duration of exposure to UV-B treatments. Given that a
11 previous narrative review suggests that plant response to UV-B exposure varies between the Arctic and
12 Antarctic (Rozema *et al.*, 2005), we also tested for differences in plant response between polar regions.

14 **Materials and methods**

15 Data were extracted from peer-reviewed publications in primary research journals on angiosperm and
16 bryophyte response to UV-B exposure in the Arctic, sub-Arctic, maritime Antarctic and continental
17 Antarctic. Although lichens are a key component of the polar flora, data on the responses of the
18 symbiosis to UV-B radiation were not included in the analysis, since it was centred on higher and lower
19 plant response to UV-B exposure. It was deemed necessary to impose several selection criteria on the
20 data before they were included in the meta-analysis. These limited the numbers of observations for
21 certain responses in subsequent analyses, but increased the robustness of the conclusions that could be
22 drawn from them. Data were included from studies that met the following criteria: (i) *Experimental*
23 *location*. Data were only included from outdoor experiments. Those from laboratory and glasshouse
24 studies, in which UV-B radiation is known to have anomalously large effects on plants (Rozema *et al.*,
25 1997), were disregarded. (ii) *Screen methodology*. Screens affect both the temperature and humidity of
26 vegetation (Kennedy, 1995), and hence have significant effects on plant growth and photosynthetic
27 parameters (Huiskes *et al.*, 2001). Data from studies using screens were hence only included when both
28 control and treatment plots had been covered with screens. Those from studies comparing the responses
29 of plants under screens with those of plants outside of screens were disregarded. It should be noted,
30 however, that screens, because they warm vegetation and increase humidity, can alter plant response to
31 UV-B exposure, and hence do not provide an accurate simulation of ozone depletion (Rozema *et al.*,
32 2005). (iii) *UV-A exposure*. Data from screen studies that attenuated only UV-B radiation were
33 included in the analyses. Those that attenuated both UV-A (315 - 400 nm) and UV-B radiation with
34 screens were excluded, since UV-A radiation is known to have significant effects on plant growth (e.g.
35 Flint & Caldwell, 2003). (iv) *Lamp methodology*. Energised Mylar-filtered fluorescent UV lamps are
36 known to have effects on plant growth relative to unenergised lamps (Newsham *et al.*, 1996), and so
37 data from lamp studies were only included when comparisons had been made between cellulose
38 acetate-filtered and glass- or Mylar-filtered lamps. Those from studies comparing the responses of
39 plants under cellulose acetate-filtered lamps with those under unenergised lamps, or those in
40 unirradiated plots, were disregarded. One study that did not use Mylar-filtered lamps as a control

1 (Niemi *et al.*, 2002a) was included in the analysis since the authors explicitly state that no effects of
2 Mylar-filtered fluorescent UV lamps were found on plant response in a previous study using the same
3 irradiation facility.

4 Of the 46 publications that were considered for inclusion, 12 did not meet the above criteria
5 and were hence excluded from the analyses. The 34 studies that met the criteria, and which were hence
6 included, are listed in Appendix 1. Thirty two species of bryophyte and angiosperm were represented in
7 the analysis (Appendix 1). In order to enable comparisons between studies, only commonly-measured
8 parameters were analysed. These included acclimation responses, such as the synthesis of UV-B
9 absorbing compounds (putative flavonoids) in leaves or thalli, measures of biomass and growth (above-
10 , belowground and total biomass, height, individual leaf mass, total leaf area and specific leaf area) and
11 indications of DNA damage, measured as the accumulation of cyclobutane pyrimidine dimers (CPDs).
12 In addition to changes in the concentrations of photosynthetic pigments such as chlorophylls and
13 carotenoids, three measures of photosynthetic physiology were included, *viz.*, net photosynthetic gas
14 exchange (P_n) and the two chlorophyll fluorescence parameters that measure photosynthetic yield,
15 Φ_{PSII} and F_v/F_m (Appendix 1).

16 In order to avoid bias towards studies that report multiple measurements of the same
17 parameter, a mean value of each parameter was calculated for each plant species in each publication.
18 The exceptions to this were when measurements had been made at more than one location in the same
19 study or in different years. In cases where two levels of elevated UV-B radiation were applied, mean
20 treatment plot response values were calculated and entered into the analyses. If the response to a factor
21 other than UV-B radiation was reported (e.g. warming; Day *et al.*, 2008), then only the data from the
22 UV-B treatment plots were included. When specific leaf area (SLA) was reported (Xiong & Day, 2001;
23 Xiong *et al.*, 2002), then data for concentrations of UV-B absorbing compounds were expressed per
24 unit of leaf mass and leaf area. Only chlorophyll and carotenoid concentrations that were expressed on
25 a leaf mass basis were included, since few publications express the concentrations of these pigments
26 per unit of leaf area. The length of the longest leaf of a grass and a forb species was used a proxy for
27 height in three studies (Day *et al.*, 2001; Ruhland & Day, 2000; Xiong *et al.*, 2002).

28 The mean value of each parameter in treatment plots and control plots was determined, and
29 the response ratio ($\ln R$; Hedges *et al.*, 1999) calculated:

30
31
$$\ln R = \ln (\text{treatment mean} / \text{control mean})$$

32
33 This enabled the expression of data as relative values, correcting for size differences between studies
34 and plant species. Values for treatment means were derived from plots exposed to high doses of UV-B
35 radiation, i.e., those from under cellulose acetate-filtered UV fluorescent lamps or from under Aclar,
36 Teflon or PMMA screens. Those for control means were from plots exposed to low doses of UV-B
37 radiation, under glass- or Mylar-filtered lamps or Mylar-filtered screens. For non-manipulative studies,
38 data were regressed against daily UV-B dose and treatment and control means were respectively
39 entered as the highest and lowest values on the y-axis, within the range of the data, along the line of
40 best fit.

1 We then used a combination of weighted and unweighted meta-analyses in order to determine
2 responses of $\ln R$ to UV-B exposure. For the weighted analyses, which used the reciprocal of the
3 sampling variance to weight $\ln R$ (Rosenberg *et al.*, 2000), n and standard deviations (s.d.) or standard
4 errors (s.e.) values were extracted from the literature. Mean weighted effect sizes were calculated, and,
5 because of the small sample size for most parameters, 95% bootstrap confidence intervals were
6 calculated, using resampling tests generated from 999 iterations (Rosenberg *et al.*, 2000). An effect size
7 was considered to be significantly different at $P < 0.05$ when its confidence intervals did not bracket
8 zero (Curtis & Wang, 1998; Feng *et al.*, 2008). The data from two reports using fluorescent UV lamps,
9 in which the stated level of n is several times higher than that reported in previous studies using the
10 same experimental facility, were excluded from the weighted analyses.

11 As in previous meta-analyses (e.g. Searles *et al.*, 2001; Dormann & Woodin, 2002), we
12 encountered several obstacles to extracting s.d. or s.e. values from the literature. For example, some of
13 the selected publications report errors but do not stipulate whether they represent s.d. or s.e., whilst one
14 publication does not report errors, reporting only mean values for treatment and control plots. In
15 several publications, error bars associated with treatment means are obscured by those associated with
16 control means, and *vice versa*. Therefore, as in previous studies (e.g. Dormann & Woodin, 2002), we
17 also used unweighted meta-analyses in order to maximise the number of observations in our study. By
18 using unweighted analyses, we were able include data that are not reported in the literature but which
19 can be derived from the published data, such as total biomass when above- and belowground biomass
20 are reported (e.g. Xiong & Day, 2001). Unweighted analyses also enabled us to determine the
21 magnitude of response to UV-B exposure in non-manipulative experiments, which would not have
22 been possible using weighted analyses. One-tailed *t*-tests were used in the unweighted analyses to
23 determine significant differences between $\ln R$ and zero for each parameter, and general linear models,
24 along with weighted analyses, were used to determine the effects of categorical variables
25 (methodology, plant form or polar region) on $\ln R$. Statistical analyses were made in MINITAB 15
26 (MINITAB inc., State College, PA, USA) and MetaWin 2.0 (Rosenberg *et al.*, 2000).

27 28 *Regression analyses*

29 We used the percentage difference between treatment and control plots in UV-B dose weighted with
30 the generalised plant action spectrum (Caldwell, 1971; $UV-B_{PAS}$) as a predictor variable for
31 untransformed response ratios in regression analyses. The data from several studies (e.g. Niemi *et al.*,
32 2002a, b; Lud *et al.*, 2003) that report UV-B doses weighted with the DNA damage or CIE action
33 spectra (Setlow, 1974; McKinlay & Diffey, 1987) were excluded from the regression analyses, since it
34 is likely, owing to the different shapes of these action spectra from that of the generalised plant action
35 spectrum (Holmes, 1997), that the percentage difference in UV-B exposure between treatment and
36 control plots would have differed substantially from studies that used $UV-B_{PAS}$ to weight irradiances.
37 We did, however, include data from reports that used both formulations of the generalised plant action
38 spectrum, either that of Green *et al.* (1974) or Thimijan *et al.* (1978), because the shapes of these two
39 formulations are similar (McLeod, 1997) and, as in previous studies (e.g. Björn *et al.*, 1997), we hence
40 anticipated only minor effects of these differences on the analyses. For experiments based at Abisko in

1 northern Sweden, UV-B_{PAS} data were derived from Johanson *et al.* (1995b). For the non-manipulative
2 studies, we calculated the difference between the maximum and minimum daily UV-B_{PAS} doses on the
3 *x*-axes in the regression analyses referred to above, treating these as supplemental studies. To aid
4 visualisation of the data, the inverses of the untransformed response ratios for screen studies were
5 entered into these analyses. Data for two parameters (height and specific leaf area) were removed from
6 the regression analyses, since, in each case, there were less than four levels of the predictor variable
7 against which to regress the response ratios. In addition, the duration of the UV-B treatment (in days)
8 was used as a predictor for untransformed response ratios in regression analyses. Linear and quadratic
9 regressions were used to determine the associations between response ratios and UV-B_{PAS} exposure.

11 Results

12 *Effects of UV-B exposure*

13 Weighted and unweighted meta-analyses indicated a significant influence of UV-B exposure on *ln R*
14 for concentrations of UV-B absorbing compounds, expressed on a mass basis, with a mean increase of
15 7.4% in plant tissues sampled from treatment plots, relative to those from control plots (Table 1; Fig.
16 1). Both methods of analysis also showed there to be a significant effect of UV-B exposure on *ln R* for
17 concentrations of UV-B absorbing compounds expressed on an area basis (Table 1), with a mean
18 increase of 24.6% in the concentrations of these pigments in plant tissues sampled from treatment plots,
19 compared with those sampled from control plots (Fig. 1). Response ratios for photosynthetic
20 parameters (F_v/F_m , Φ_{PSII} and P_n) and concentrations of associated pigments (total carotenoids and
21 chlorophylls) were unaffected by UV-B exposure (Table 1; Fig. 1).

22 Both methods of analysis showed that UV-B exposure significantly decreased *ln R* for
23 aboveground biomass (Table 1), with a 14.7% reduction in this parameter in treatment plots, relative to
24 control plots (Fig. 1). Plant height was also significantly affected by UV-B exposure, with unweighted
25 and weighted analyses both indicating a mean 10.0% reduction in height in treatment plots compared
26 with control plots (Table 1; Fig. 1). Neither method of analysis showed an effect of UV-B exposure on
27 *ln R* for belowground biomass, nor total biomass (Table 1; Fig. 1). The response ratio for total leaf area
28 was unaffected by UV-B exposure in the unweighted analyses, but was significantly reduced by 6.1%
29 in the weighted analyses (Table 1; Fig. 1). Neither method of analysis showed individual leaf mass or
30 SLA to be affected by UV-B exposure (Table 1; Fig. 1), but both indicated that UV-B exposure
31 significantly influenced *ln R* for DNA damage, with the formation of CPDs increasing by 90.2% in
32 DNA from treatment plots, compared with that from control plots (Table 1; Fig. 1).

34 *Effects of methodology*

35 Methodology had a significant effect on the response ratio for concentrations of UV-B absorbing
36 compounds expressed per unit of mass (Table 1). In unweighted analyses, studies that used screens to
37 attenuate UV-B radiation, or recorded plant response to natural fluctuations in UV-B irradiance in non-
38 manipulative experiments, showed significant 12.2% and 17.3% increases in the concentrations of UV-
39 B absorbing compounds in response to UV-B exposure (both $n = 8$, $P < 0.01$ and $P < 0.05$, respectively;
40 Fig. 2a). In contrast, *ln R* for concentrations of these compounds from studies that applied UV-B

1 radiation from fluorescent UV lamps was not significantly different from zero in unweighted analyses
2 ($n = 26$, $P > 0.05$; Fig. 2a). Weighted analyses similarly indicated no change in the concentrations of
3 these pigments under fluorescent UV lamps ($P > 0.05$), but a significant positive response to UV-B
4 radiation under screens ($P < 0.05$). Belowground biomass was also influenced by methodology (Table
5 1): unweighted analyses showed that the exposure of plants to UV-B radiation under screens led to a
6 16.9% reduction in this parameter ($n = 5$; Fig. 2b). In contrast, no effect was recorded on belowground
7 biomass when plants had been exposed to UV-B radiation under fluorescent UV lamps (Fig. 2b). It
8 should be noted, however, that the number of observations for this response was low ($n = 2$).

9 10 *Effects of plant form*

11 Unweighted and weighted analyses both indicated that the response ratios for UV-B absorbing
12 compounds expressed per unit of leaf area differed between bryophytes and angiosperms (Table 1): $\ln R$
13 for concentrations of UV-B absorbing compounds expressed in this way was higher for angiosperms
14 than for bryophytes, with both methods of analysis indicating a significant 34.5% increase in the
15 concentrations of these pigments, relative to controls, in angiosperms ($n = 8$, Fig. 2c). A significant
16 effect of UV-B exposure on $\ln R$ for concentrations of these pigments, relative to controls, was not
17 found for bryophytes using either method of analysis ($n = 6$; Fig. 2c). It is important to note, however,
18 that the latter data were all derived from one species of bryophyte sampled from a mixture of lamp and
19 non-manipulative experiments in different years (Appendix 1). There was also a marginally significant
20 effect of plant form on $\ln R$ for F_v/F_m in unweighted analyses (Table 1), with a reduction of 2.2% in
21 this parameter in angiosperms, but an increase of 4.5% in bryophytes in response to UV-B exposure.
22 However, neither response ratio was significantly different from zero (data not shown).

23 24 *Effects of region*

25 Unweighted analyses indicated different response ratios for carotenoid concentration in studies based in
26 the Arctic and Antarctic (Table 1). There was no effect of UV-B treatment on $\ln R$ for carotenoid
27 concentrations in plant tissues in the Arctic ($n = 9$, $P > 0.05$), but $\ln R$ for concentrations of these
28 pigments increased by 17.1% in Antarctic studies ($n = 6$; Fig. 2d). Unweighted and weighted analyses
29 showed there to be an effect of region on $\ln R$ for SLA, with the former analyses also indicating an
30 effect on total leaf area (Table 1). The response ratios for these parameters did not differ from zero in
31 Arctic studies ($n = 10$ and 14 , respectively, both $P > 0.05$), but were respectively reduced by 25.3% and
32 21.4% in Antarctic experiments ($n = 4$ and 3 , respectively; Fig. 2e, f). It should be noted, however, that
33 the latter data were all derived from work by one research group on two plant species (Appendix 1).
34 Unweighted analyses also showed there to be a marginally significant effect of region on $\ln R$ for total
35 biomass (Table 1), with a mean increase of 8.5% in this parameter in Arctic studies, but a 10.4%
36 decrease in Antarctic studies. However, neither response ratio was significantly different from zero
37 (data not shown). Although there was no main effect of region on plant height (Table 1), separate
38 analyses indicated that there was a significant 8.9% reduction in height when plants had been exposed
39 to UV-B radiation applied from lamps simulating 15% ozone depletion in Arctic studies ($n = 10$;
40 unweighted analyses $P < 0.01$, weighted $P < 0.05$; data not shown).

1 *Associations between response ratios and predictor variables*

2 The percentage difference in UV-B_{PAS} dose between treatment and control plots was a significant
3 predictor for four untransformed response ratios (Table 2). The response ratio for concentrations of
4 UV-B absorbing compounds, expressed per unit of mass, was positively associated with the difference
5 in UV-B_{PAS} dose between control and treatment plots (Table 2). There were consistent increases and
6 decreases in the response ratios for these pigments in non-manipulative and screen studies, respectively
7 (Fig. 3a). The association between the two variables was best described by a quadratic function (Fig.
8 3a). In lamp experiments, the range in response ratios for UV-B absorbing compounds was
9 considerable, varying between 0.7 and 1.2, compared with the ranges for screen and non-manipulative
10 studies of 0.8 to 1.0 and 1.1 to 1.4, respectively (Fig. 3a). A linear association was found for
11 concentrations of carotenoids: there was a positive association between the response ratios for the
12 concentrations of these pigments and difference in UV-B_{PAS} dose, with unweighted analyses indicating
13 significant 22.7% increases in the concentrations of these pigments in non-manipulative studies, but
14 lamps and screens having no effect on carotenoid concentrations (Fig. 3b). Aboveground biomass was
15 negatively and linearly associated with the difference in UV-B_{PAS} dose between treatment and control
16 plots (Fig. 3c). Both unweighted and weighted analyses indicated that shielding plants from UV-B
17 radiation under screens significantly increased aboveground biomass by 27.5%, while exposure to UV-
18 B from lamps had no significant effect on this parameter (Fig. 3c). Leaf area was also negatively
19 associated with difference in UV-B_{PAS} dose, with unweighted and weighted analyses both indicating a
20 significant 34.5% increase in the response ratio for this parameter when UV-B was removed from solar
21 radiation with screens, but no effect on the ratio when supplemental UV-B radiation had been applied
22 from lamps (Fig. 3d).

23 One significant association between the duration of exposure to UV-B radiation and
24 untransformed response ratios was recorded: the ratio for carotenoid concentration decreased linearly
25 with the length of exposure to UV-B radiation (Fig. 3b, *inset*). The duration of exposure to UV-B was
26 otherwise not associated with untransformed response ratios (r^2 adj. = 0.228 – 0.258; all $P > 0.05$; data
27 not shown).

28

29 **Discussion**

30 Here we show, using a combination of weighted and unweighted meta-analyses, that the exposure of
31 polar bryophytes and angiosperms to UV-B radiation increases the concentration of UV-B absorbing
32 compounds in leaves and thalli by 7% and 25% (expressed per unit of mass and area, respectively), and
33 decreases the height and aboveground biomass of plants by 10% and 15%, respectively. These results
34 are strikingly similar to those from a previous meta-analysis of plant response to simulated
35 stratospheric ozone depletion (Searles *et al.*, 2001). In the previous analysis, which used data from 62
36 studies, predominantly on temperate angiosperms and gymnosperms exposed to supplemental UV-B
37 radiation from fluorescent UV lamps, UV-B absorbing compounds were found to be increased in
38 concentration by 10% following UV-B exposure, height was diminished by 3% and shoot biomass was
39 reduced by 16%, but only when studies simulated > 20% loss of ozone from the atmosphere (Searles *et*
40 *al.*, 2001). Furthermore, total leaf area, which the weighted analyses in the current study indicated was

1 reduced by 6% following UV-B exposure, was also reduced by 13% in the previous analysis (Searles *et al.*, 2001). Other parameters common to the meta-analysis here and that of Searles *et al.* (2001), *viz.*,
2 chlorophyll and carotenoid concentrations, P_n and F_v/F_m , were each unaffected by UV-B exposure,
3 supporting the view that photosynthesis and the concentrations of associated pigments are unaffected
4 by UV-B radiation in field experiments (e.g. Fiscus & Booker, 1995; Allen *et al.*, 1998; Sullivan &
5 Rozema, 1999).
6

7 Although Searles *et al.* (2001) did not analyse data on DNA damage in plant foliage, the
8 current study also recorded a significant 90% increase in the accumulation of CPDs in DNA following
9 exposure to UV-B radiation. This substantial increase in DNA damage in the foliage of plants exposed
10 to elevated UV-B radiation in treatment plots relative to that in control plots is apparently owing to
11 unrealistically high doses of UV-B radiation having been applied to plants: two of the highest response
12 ratios for this parameter were derived from lamp studies that applied Setlow (1974)-weighted doses of
13 UV-B radiation to treatment plots at an order of magnitude higher than were received in control plots
14 (Lud *et al.*, 2002; 2003). However, even though the frequency of CPDs in DNA was tripled by
15 exposure to elevated UV-B radiation in these studies, repair took place rapidly, with DNA damage in
16 plants in treatment plots falling to the same level as in control plots within 12 h post irradiation (Lud *et al.*,
17 2002; 2003). The recent finding that fewer DNA photoproducts accumulate in desiccated,
18 compared with hydrated, mosses also suggests that screening and passive defence mechanisms are well
19 developed in Antarctic bryophytes (Turnbull *et al.*, 2009).
20

21 Bryophytes are thought to respond in the same way as vascular plants to UV-B exposure
(Rozema *et al.*, 2005). The analyses here support this view: although 60% of the plant species included
22 in our analyses were bryophytes, similar results were found to those of Searles *et al.* (2001), who only
23 analysed data from angiosperms and gymnosperms. Only one response ratio differed between these two
24 plant forms in the current study. The higher response ratio of UV-B absorbing compounds expressed
25 per unit of area for angiosperms than for bryophytes corroborates earlier suggestions that screening
26 pigments in bryophytes are less responsive to UV-B exposure than those in angiosperms (Gwynn-Jones
27 *et al.*, 1999). However, this probably reflects a species-specific response, since data from only one
28 study (Lappalainen *et al.*, 2008) on a single bryophyte species were included in the analysis of these
29 pigments. We did not anticipate finding this effect: since higher plants are able to attenuate UV-B
30 radiation through the presence of epidermal hairs, cuticles and surface waxes on leaves (Day, 1993),
31 they should not need to synthesize screening pigments to the same extent as bryophytes, which tend to
32 lack these protective features (Gehrke, 1999).
33

34 Our study demonstrates that the method which is used to alter the dose of UV-B radiation
35 received by plants has an effect on the synthesis of UV-B absorbing compounds in foliage. The
36 analyses show that exposure to UV-B radiation in screen and non-manipulative studies increases the
37 concentrations of UV-B absorbing compounds in leaves and thalli by 12% and 17%, respectively, but
38 that UV-B applied from fluorescent UV lamps fails to elicit a change in the concentration of these
39 compounds. These findings suggest that the use of fluorescent UV lamps in polar environments does
40 not elicit the same response as at lower latitudes, since the exposure of plants to UV-B applied from
fluorescent lamps, typically at mid-latitudes, significantly increases the concentrations of UV-B

1 absorbing compounds in foliage by 10% (Searles *et al.*, 2001). It is unclear as to why this difference
2 might exist. It is possible that it can be accounted for by higher doses of UV-B radiation having been
3 applied to plants in the experiments analysed by Searles *et al.* (2001). Although the mean level of
4 ozone depletion simulated in the lamp studies analysed here (18%) fell within the range of ozone
5 depletion simulated in the majority of studies analysed in the previous meta-analysis (10-20%), the
6 absolute doses of UV-B radiation applied to plants for a given level of ozone depletion in polar habitats
7 will be less than at mid-latitudes because of lower solar elevation angles at higher latitudes. Another
8 issue that might have contributed to the lack of effect of UV-B exposure on the concentrations of UV-B
9 absorbing compounds in plant tissues could have been the application of inappropriate biological
10 weighting functions. Recently, it has been suggested that the generalised plant action spectrum
11 (Caldwell, 1971), which is often used to weight supplemental UV-B radiation, results in unrealistically
12 low UV-B doses being applied from fluorescent UV lamps to vegetation (Flint & Caldwell, 2003;
13 Caldwell *et al.*, 2006). Our analyses also indicate that lamps introduce considerable heterogeneity into
14 plant response to UV-B exposure in polar habitats, with a wider range in response ratios for UV-B
15 absorbing compounds in lamp studies than in screen or non-manipulative studies, indicating positive
16 responses in some lamp experiments and null, or negative, responses in others. We suspect that the
17 difficulties of maintaining stable outputs from fluorescent UV lamps at low temperatures may be
18 responsible for some of the heterogeneity in screening pigment response to UV-B exposure identified
19 here (Johanson & Zeuthen, 1998; Rozema *et al.*, 2001).

20 In their narrative review of UV-B effects on polar vegetation, Rozema *et al.* (2005) conclude
21 that plant response to UV-B exposure may differ between the Arctic and Antarctic. Our analyses
22 suggest the same effect, showing that concentrations of carotenoids increase, and that leaf area and
23 SLA decrease, in Antarctic experiments, but that no effects occur on these parameters in Arctic studies.
24 However, these apparent differences in plant response between polar regions can largely be explained
25 by the different methodological approaches used in the two regions: in the current study, the data for
26 leaf area and SLA from Antarctic experiments were all derived from studies using screens (Ruhland &
27 Day, 2001; Xiong & Day, 2001; Xiong *et al.*, 2002), which have consistent effects on these leaf
28 parameters. Most, but not all, of the Arctic studies from which leaf area and SLA were derived applied
29 UV-B radiation from fluorescent UV lamps, with only two studies (Albert *et al.*, 2005b; Albert *et al.*,
30 2008) using screens. Our analyses thus suggest that the main reason plant response differs between the
31 two polar regions is because screens, which have consistent effects on plant performance, are typically
32 used in the Antarctic, and fluorescent UV lamps, which have less consistent effects, are more
33 frequently used in the Arctic.

34 Rozema *et al.* (2005) surmise that, with the exception of studies close to Anvers Island in the
35 maritime Antarctic (e.g. Day *et al.*, 1999; 2001), most field-based lamp or screen studies indicate
36 negligible effects of UV-B exposure on plants. Whilst our data support this view for lamp studies, they
37 do not corroborate this view for screens, with analyses for screen studies showing effects on plant
38 response irrespective of geographical location. For example, for UV-B absorbing compounds, in
39 addition to data from screen studies from close to Anvers Island (Ruhland & Day, 2000; Ruhland &
40 Day, 2001; Xiong & Day, 2001; Xiong *et al.*, 2002), data from one Antarctic and three Arctic screen

1 studies (Newsham *et al.*, 2005; Albert *et al.*, 2005b; Albert *et al.*, 2008; Kotilainen *et al.*, 2008), were
2 also included in the meta-analysis, all of which showed positive response ratios for concentrations of
3 UV-B absorbing compounds expressed on a mass basis. It thus seems likely that the attenuation of UV-
4 B radiation by screens does have consistent and generic effects on plant performance in polar field
5 experiments.

6 Similar results were found in the current study when data were analysed using weighted or
7 unweighted meta-analyses. This was most probably owing to the similar level of replication used in
8 most of the lamp and screen experiments analysed here, which usually employed a modest level of
9 replication, with typically three or four control or treatment plots used in each study. As in previous
10 meta-analyses (e.g. Searles *et al.*, 2001), we noted effects of data from apparently pseudoreplicated
11 (Hurlbert, 1984) studies on the weighted analyses, with these data having anomalously large effects on
12 the analyses because of the extra weight given to them, as a result of the large reciprocals of the
13 sampling variances associated with their response ratios. These data were thus removed from the
14 weighted analyses, constraining our ability to detect UV-B treatment effects on response ratios.
15 Another factor that constrained our ability to detect effects on the response ratios was the low number
16 of observations for certain parameters, which arose partly from the limited number of studies in the
17 literature on the UV-B responses of polar vegetation, and partly from the selection criteria that we
18 imposed on the data before they were included in the analysis. Caution is needed when interpreting the
19 data for certain parameters because of the low number of observations, particularly those for
20 photosynthetic parameters, belowground biomass, individual leaf mass and DNA damage. Our analyses
21 do identify, however, that additional data are needed in the literature on the responses of these
22 parameters to UV-B exposure in polar environments.

23 Although we found the percentage difference in UV-B_{PAS} dose between treatment and control
24 plots to be a predictor for four response ratios in the current study, the duration of exposure to UV-B
25 treatment had little apparent influence on the response ratios. Only that for carotenoid concentration
26 was associated with the duration of exposure, with the data indicating a reduction in the magnitude of
27 response to UV-B exposure with increasing time. These data broadly indicate that the magnitude of
28 effect of UV-B radiation on plant response does not increase or decrease with longer periods of
29 exposure, corroborating the view that exposure to UV-B radiation does not have cumulative long-term
30 effects on plant growth in polar regions (Rozema *et al.*, 2006). For pigments, these data also broadly
31 support the view that plants can rapidly acclimate to UV-B exposure, even in polar habitats at
32 temperatures close to freezing point (Newsham *et al.*, 2002; Newsham *et al.*, 2005; Dunn & Robinson,
33 2006; Snell *et al.*, 2009).

34 Given the responses shown here, then in broad terms, what will have been the likely
35 consequences of increased UV-B exposure for polar vegetation in recent years? Assuming cloudless
36 skies, then the mean 15% springtime loss of ozone from the stratosphere over the Arctic in the last two
37 decades (World Meteorological Organization, 2007) will have led to a 30% increase at solar noon in
38 UV-B_{PAS} irradiance in late spring at 68° N (Johanson *et al.*, 1995b), which, as reported here, reduces
39 plant height by 9%. Data in Fig. 3 suggest that the 30% springtime loss of ozone from the stratosphere
40 over the Antarctic in the last two decades (World Meteorological Organization, 2007), which has led to

1 an approximate doubling in UV-B_{PAS} exposure at solar noon in late spring at 64°-67° S, latitudes at
2 which the majority of Antarctic UV-B studies have been made (Xiong *et al.*, 2002; Newsham *et al.*,
3 2002; Dunn & Robinson, 2006), might have reduced aboveground biomass by some 20%. The data in
4 Fig. 3 also suggest that although there are unlikely to have been effects on the concentrations of UV-B
5 absorbing compounds in plant tissues in the Arctic in recent years, the doubling in UV-B_{PAS} exposure in
6 the Antarctic would have led to an approximate 5% increase in the concentrations of these compounds
7 in leaves and thalli. Recent studies corroborate this view, with larger increases over the last three
8 decades in the concentrations of UV-B absorbing compounds in clubmoss tissues sampled from South
9 Georgia in the Scotia Sea than in those from Greenland (Lomax *et al.*, 2008). Such increases will most
10 probably have had effects on decomposition and herbivory (Caldwell *et al.*, 1995; Paul & Gwynn-
11 Jones, 2003), as well as subtle negative effects on plant growth, owing to the carbon drain associated
12 with synthesizing flavonoids in leaves and thalli (Snell *et al.*, 2009).

13

14 *Conclusions*

15 In broad terms, this meta-analysis indicates that vegetation in polar regions responds to UV-B radiation
16 exposure in a similar way to that at lower latitudes, often by increasing the synthesis of UV-B
17 absorbing compounds in leaves and thalli. Photosynthetic parameters show little consistent response to
18 UV-B exposure, but there is evidence of negative effects on aboveground biomass, plant height and
19 DNA. Given that a recovery of the ozone layer is not expected for several decades (World
20 Meteorological Organization, 2007), we advocate further studies, preferably using the same
21 methodologies in the Arctic and Antarctic, to assess the broader influence of UV-B radiation on overall
22 ecosystem functioning in polar regions.

23

24 **Acknowledgements**

25 This work was funded by the Natural Environment Research Council through BAS's Long Term
26 Monitoring and Survey programme (KKN) and by the Australian Antarctic Division (SAR). Satu
27 Huttunen, Kristian Albert, Tad Day, Peter Boelen and Jelte Rozema supplied unpublished data, Peter
28 Rothery gave valuable statistical advice and Jon Shanklin, Andy McLeod, Pete Convey and three
29 anonymous reviewers supplied helpful comments. All are gratefully acknowledged. Data were
30 accessed in late 2008.

1 **References**

- 2 Albert KR, Ro-Poulsen H, Mikkelsen TN, Bredahl L, Haakansson KB (2005a) Effects of reducing the
3 ambient UV-B radiation in the high arctic on *Salix arctica* and *Vaccinium uliginosum*. *Phyton-*
4 *Annales Rei Botanicae*, **45**, 41-49.
- 5 Albert KR, Mikkelsen TN, Ro-Poulsen H (2005b) Effects of ambient versus reduced UV-B radiation
6 on high arctic *Salix arctica* assessed by measurements and calculations of chlorophyll a
7 fluorescence parameters from fluorescence transients. *Physiologia Plantarum*, **124**, 208-226.
- 8 Albert KR, Mikkelsen TN, Ro-Poulsen H (2008) Ambient UV-B radiation decreases photosynthesis in
9 high arctic *Vaccinium uliginosum*. *Physiologia Plantarum*, **133**, 199-210.
- 10 Allen DJ, Nogués S, Baker NR (1998) Ozone depletion and increased UV-B radiation: is there a real
11 threat to photosynthesis? *Journal of Experimental Biology*, **49**, 1775-1788.
- 12 Arft AM, Walker MD, Gurevitch J *et al.* (1999) Responses of tundra plants to experimental warming:
13 meta-analysis of the International Tundra Experiment. *Ecological Monographs*, **69**, 491-511.
- 14 Björn L-O, Callaghan TV, Gehrke C, Gwynn-Jones D, Lee JA, Johanson U, Sonesson M, Buck ND
15 (1999) Effects of ozone depletion and increased ultraviolet-B radiation on northern vegetation.
16 *Polar Research*, **18**, 331-337.
- 17 Björn L-O, Callaghan TV, Johnsen I *et al.* (1997) The effects of UV-B radiation on European
18 heathland species. *Plant Ecology*, **128**, 252-264.
- 19 Boelen P, de Boer MK, de Bakker NVJ, Rozema J (2006) Outdoor studies on the effects of solar UV-B
20 on bryophytes: Overview and methodology. *Plant Ecology*, **182**, 137-152.
- 21 Bredahl L, Ro-Poulsen H, Mikkelsen TN (2004) Reduction of the ambient UV-B radiation in the high-
22 arctic increases F_v/F_m in *Salix arctica* and *Vaccinium uliginosum* and reduces stomatal conductance
23 and internal CO₂ concentration in *Salix arctica*. *Arctic, Antarctic and Alpine Research*, **36**, 364-369.
- 24 Caldwell MM (1971) Solar UV radiation and the growth and development of higher plants. In:
25 *Photophysiology* (ed Giese AC), pp. 131-177. Academic Press, New York.
- 26 Caldwell MM, Bornman JF, Ballaré CL, Flint SD, Kulandaivelu G (2006) Terrestrial ecosystems,
27 increased solar ultraviolet radiation, and interactions with other climate change factors. In:
28 *Environmental effects of ozone depletion and its interactions with climate change: 2006 assessment*.
29 pp. 65-94. United Nations Environment Programme, Nairobi, Kenya.
- 30 Caldwell MM, Teramura AH, Tevini M, Bornman JF, Björn L-O, Kulandaivelu G (1995) Effects of
31 increased solar ultraviolet radiation on terrestrial plants. *Ambio*, **24**, 166-173.
- 32 Clarke LJ, Robinson SA (2008) Cell wall-bound ultraviolet-screening compounds explain the high
33 ultraviolet tolerance of the Antarctic moss, *Ceratodon purpureus*. *New Phytologist*, **179**, 776-783.
- 34 Curtis PS, Wang X (1998) A meta-analysis of elevated CO₂ effects on woody plant mass, form, and
35 physiology. *Oecologia*, **113**, 299-313.
- 36 Day TA (1993) Relating UV-B radiation screening effectiveness of foliage to absorbing-compound
37 concentration and anatomical characteristics in a diverse group of plants. *Oecologia*, **95**, 542-550.
- 38 Day TA, Ruhland CT, Grobe CW, Xiong F (1999) Growth and reproduction of Antarctic vascular
39 plants in response to warming and UV radiation reductions in the field. *Oecologia*, **119**, 24-35.

- 1 Day TA, Ruhland CT, Xiong F (2001) Influence of solar ultraviolet-B radiation on Antarctic terrestrial
2 plants: results from a 4-year field study. *Journal of Photochemistry and Photobiology B: Biology*,
3 **62**, 78-87.
- 4 Day TA, Ruhland CT, Xiong FS (2008) Warming increases aboveground plant biomass and C stocks in
5 vascular-plant dominated Antarctic tundra. *Global Change Biology*, **14**, 1827-1843.
- 6 Day TA, Vogelmann TC (1993) Penetration of UV-B radiation in foliage: evidence that the epidermis
7 behaves as a non-uniform filter. *Plant, Cell and Environment*, **16**, 735-741.
- 8 Day TA, Vogelmann TC, Delucia EH (1992) Are some plant life forms more effective than others in
9 screening out ultraviolet-B radiation? *Oecologia*, **92**, 513-519.
- 10 Dormann CF, Woodin SJ (2002) Climate change in the Arctic: using plant functional types in a meta-
11 analysis of field experiments. *Functional Ecology*, **16**, 4-17.
- 12 Dunn JL, Robinson SA (2006) Ultraviolet B screening potential is higher in two cosmopolitan moss
13 species than in a co-occurring Antarctic endemic moss: implications of continuing ozone depletion
14 *Global Change Biology*, **12**, 2282-2296.
- 15 Farman JC, Gardiner BG, Shanklin JD (1985) Large losses of total ozone in Antarctica reveal seasonal
16 ClO_x/NO_x interaction. *Nature*, **315**, 207-210.
- 17 Feng Z, Kobayashi K, Ainsworth EA (2008) Impact of elevated ozone concentration on growth,
18 physiology, and yield of wheat (*Triticum aestivum* L.): a meta analysis. *Global Change Biology*, **14**,
19 2696-2708.
- 20 Fiscus EL, Booker FL (1995) Is increased UV-B a threat to crop photosynthesis and productivity?
21 *Photosynthesis Research*, **43**, 81-92.
- 22 Flint SD, Caldwell MM (2003) Field testing of UV biological spectral weighting functions for higher
23 plants. *Physiologia Plantarum*, **117**, 145-153.
- 24 Gehrke C (1998) Effects of enhanced UV-B radiation on production-related properties of a *Sphagnum*
25 *fuscum* dominated subarctic bog. *Functional Ecology*, **12**, 940-947.
- 26 Gehrke C (1999) Impacts of enhanced ultraviolet-B radiation on mosses in a subarctic heath ecosystem.
27 *Ecology*, **80**, 1844-1851.
- 28 Green AES, Sawada T, Shettle EP (1974) The middle ultraviolet reaching the ground. *Photochemistry*
29 *and Photobiology*, **19**, 251-259.
- 30 Gwynn-Jones D (2001) Short-term impacts of enhanced UV-B radiation on photo-assimilate allocation
31 and metabolism: a possible interpretation for time-dependent inhibition of growth. *Plant Ecology*,
32 **154**, 67-73.
- 33 Gwynn-Jones D, Lee JA, Johanson U, Phoenix GK, Callaghan TV, Sonesson M (1999) The responses
34 of plant functional types to enhanced UV-B radiation. In: *Stratospheric ozone depletion: the effects*
35 *of enhanced UV-B radiation on terrestrial ecosystems* (ed Rozema J), pp. 173-185. Backhuys
36 Publishers, Leiden, The Netherlands.
- 37 Hedges LV, Gurevitch J, Curtis PS (1999) The meta-analysis of response ratios in experimental
38 ecology. *Ecology*, **80**, 1150-1156.

- 1 Holmes MG (1997) Action spectra for UV-B effects on plants: monochromatic and polychromatic
2 approaches for analysing plant responses. In: *Plants and UV-B: Responses to Environmental*
3 *Change* (ed. Lumsden PJ), pp. 31-50. Cambridge University Press, Cambridge.
- 4 Huiskes A, Lud D, Moerdijk-Poorvliet T (2001) Field research on the effects of UV-B filters on
5 terrestrial Antarctic vegetation. *Plant Ecology*, **154**, 77-86.
- 6 Hurlbert SH (1984) Pseudoreplication and the design of ecological field experiments. *Ecological*
7 *Monographs*, **54**, 187-211.
- 8 Johanson U (1997) *Future stratospheric ozone depletion will affect a subarctic dwarf shrub ecosystem*.
9 Dissertation. Lund University, Sweden. ISBN 91-628-2402-X.
- 10 Johanson U, Gehrke C, Bjorn LO, Callaghan TV (1995a) The effects of enhanced UV-B radiation on
11 the growth of dwarf shrubs in a sub-arctic heathland. *Functional Ecology*, **9**, 713-719.
- 12 Johanson U, Gehrke C, Björn LO, Callaghan TV, Sonesson M (1995b) The effects of enhanced UV-B
13 radiation on a subarctic heath ecosystem. *Ambio*, **24**, 106-111.
- 14 Johanson U, Zeuthen J (1998) The ecological relevance of UV-B enhancement studies is dependent on
15 tube type, tube age, temperature and voltage. *Journal of Photochemistry and Photobiology. B.*
16 *Biology*, **44**, 169-174.
- 17 Kennedy AD (1995) Simulated climate change: are passive greenhouses a valid microcosm for testing
18 the biological effects of environmental perturbations? *Global Change Biology* **1**, 29-42.
- 19 Kotilainen T, Tegelberg R, Julkunen-Tiitto R, Lindfors A, Aphalo PJ (2008) Metabolite specific effects
20 of solar UV-A and UV-B on alder and birch leaf phenolics. *Global Change Biology*, **14**, 1294-1304.
- 21 Lappalainen NM, Huttunen S, Suokanerva H (2008) Acclimation of the pleurocarpous moss
22 *Pleurozium schreberi* (Britt.) Mitt. to enhanced ultraviolet radiation *in situ*. *Global Change Biology*,
23 **14**, 321-333.
- 24 Lomax BH, Fraser WT, Sephton MA *et al.* (2008) Plant spore walls as a record of long-term changes in
25 ultraviolet-B radiation. *Nature Geoscience*, **1**, 592-596.
- 26 Lud D, Buma AGJ, van de Poll WH, Moerdijk TCW, Huiskes AHL (2001a) DNA damage and
27 photosynthetic performance in the Antarctic terrestrial alga *Prasiola crispa* ssp. *antarctica*
28 (Chlorophyta) under manipulated UV-B radiation. *Journal of Phycology*, **37**, 459-467.
- 29 Lud D, Huiskes A, Moerdijk T, Rozema J (2001b) The effects of altered levels of UV-B radiation on
30 an Antarctic grass and lichen. *Plant Ecology*, **154**, 89-99.
- 31 Lud D, Moerdijk TCW, van de Poll WH, Buma AGJ, Huiskes AHL (2002) DNA damage and
32 photosynthesis in Antarctic and Arctic *Sanionia uncinata* (Hedw.) Loeske under ambient and
33 enhanced levels of UV-B radiation. *Plant, Cell and Environment*, **25**, 1579-1589.
- 34 Lud D, Schlenso M, Schroeter B, Huiskes AHL (2003) The influence of UV-B radiation on light-
35 dependent photosynthetic performance in *Sanionia uncinata* (Hedw.) Loeske in Antarctica. *Polar*
36 *Biology*, **26**, 225-232.
- 37 McKinlay AF, Diffey BL (1987) A reference action spectrum for ultra-violet induced erythema in
38 human skin. In: *Human Exposure to Ultra-Violet Radiation: Risks and Regulations* (eds Passchier
39 WF, Bosnjakovic BFM), pp. 83-87. Elsevier, Amsterdam.

- 1 McLeod AR (1997) Outdoor supplementation systems for studies of the effects of increased UV-B
2 radiation. *Plant Ecology*, **128**, 78-92.
- 3 Mendez M, Gwynn-Jones D, Manetas Y (1999) Enhanced UV-B radiation under field conditions
4 increases anthocyanin and reduces the risk of photoinhibition but does not affect growth in the
5 carnivorous plant *Pinguicula vulgaris*. *New Phytologist*, **144**, 275-282.
- 6 Newman PA, Gleason JF, McPeters RD, Stolarski RS (1997) Anomalously low ozone over the Arctic.
7 *Geophysical Research Letters*, **24**, 2689-2692.
- 8 Newsham KK (2003) UV-B radiation arising from stratospheric ozone depletion influences the
9 pigmentation of the moss *Andreaea regularis*. *Oecologia*, **135**, 327-331.
- 10 Newsham KK, Geissler P, Nicolson M, Peat HJ, Lewis Smith RI (2005) Sequential reduction of UV-B
11 radiation in the field alters the pigmentation of an Antarctic leafy liverwort. *Environmental and*
12 *Experimental Botany*, **54**, 22-32.
- 13 Newsham KK, Hodgson DA, Murray AWA, Peat HJ, Lewis Smith RI (2002) Response of two
14 Antarctic bryophytes to stratospheric ozone depletion. *Global Change Biology*, **8**, 972-983.
- 15 Newsham KK, McLeod AR, Greenslade PD, Emmett BA (1996) Appropriate controls in outdoor UV-
16 B supplementation experiments. *Global Change Biology*, **2**, 319-324.
- 17 Niemi R, Martikainen P, Silvola J, Sonninen E, Wulff A, Holopainen T (2002a) Responses of two
18 *Sphagnum* moss species and *Eriophorum vaginatum* to enhanced UV-B in a summer of low UV
19 intensity. *New Phytologist*, **156**, 509-515.
- 20 Niemi R, Martikainen P, Silvola J, Wulff A, Turtola S, Holopainen T (2002b) Elevated UV-B radiation
21 alters fluxes of methane and carbon dioxide in peatland microcosms. *Global Change Biology*, **8**,
22 361-371.
- 23 Paul ND, Gwynn-Jones D (2003) Ecological roles of solar UV radiation: towards an integrated
24 approach. *Trends in Ecology and Evolution*, **18**, 48-55.
- 25 Phoenix GK, Gwynn-Jones D, Lee JA, Callaghan TV (2000) The impacts of UV-B radiation on the
26 regeneration of a sub-arctic heath community. *Plant Ecology*, **146**, 67-75.
- 27 Rinnan R, Keinänen MM, Kasurinen A *et al.* (2005) Ambient ultraviolet radiation in the Arctic reduces
28 root biomass and alters microbial community composition but has no effects on microbial biomass.
29 *Global Change Biology*, **11**, 564-574.
- 30 Robinson SA, Turnbull JD, Lovelock CE (2005) Impact of changes in natural ultraviolet radiation on
31 pigment composition, physiological and morphological characteristics of the Antarctic moss,
32 *Grimmia antarctici*. *Global Change Biology*, **11**, 476-489.
- 33 Rosenberg MS, Adams DC, Gurevitch J (2000) *MetaWin: statistical software for meta-analysis*.
34 Version 2. Sinauer Associates, Sunderland, Massachusetts.
- 35 Rozema J, Boelen P, Blokker P (2005) Depletion of stratospheric ozone over the Antarctic and Arctic:
36 Responses of plants of polar terrestrial ecosystems to enhanced UV-B, an overview. *Environmental*
37 *Pollution*, **137**, 428-442.
- 38 Rozema J, Boelen P, Solheim B *et al.* (2006) Stratospheric ozone depletion: High arctic tundra plant
39 growth on Svalbard is not affected by enhanced UV-B after 7 years of UV-B supplementation in the
40 field. *Plant Ecology*, **182**, 121-135.

- 1 Rozema J, Broekman R, Lud D *et al.* (2001) Consequences of depletion of stratospheric ozone for
2 terrestrial Antarctic ecosystems: the response of *Deschampsia antarctica* to enhanced UV-B
3 radiation in a controlled environment. *Plant Ecology*, **154**, 103-115.
- 4 Rozema J, van de Staaij, Björn L-O, Caldwell MM (1997) UV-B as an environmental factor in plant
5 life: stress and regulation. *Trends in Ecology and Evolution*, **12**, 22-28.
- 6 Ruhland CT, Day TA (2000) Effects of ultraviolet-B radiation on leaf elongation, production and
7 phenylpropanoid concentrations of *Deschampsia antarctica* and *Colobanthus quitensis* in
8 Antarctica. *Physiologia Plantarum*, **109**, 244-251.
- 9 Ruhland CT, Day TA (2001) Size and longevity of seed banks in Antarctica and the influence of
10 ultraviolet-B radiation on survivorship, growth and pigment concentrations of *Colobanthus*
11 *quitensis* seedlings. *Environmental and Experimental Botany*, **45**, 143-154.
- 12 Searles PS, Flint SD, Caldwell MM (2001) A meta-analysis of plant field studies simulating
13 stratospheric ozone depletion. *Oecologia*, **127**, 1-10.
- 14 Semerdjieva SI, Sheffield E, Phoenix GK, Gwynn-Jones D, Callaghan TV, Johnson GN (2003)
15 Contrasting strategies for UV-B screening in sub-Arctic dwarf shrubs. *Plant, Cell and Environment*,
16 **26**, 957-964.
- 17 Setlow RB (1974) The wavelengths in sunlight effective in producing skin cancer. *Proceedings of the*
18 *National Academy of Sciences, USA*, **71**, 3363-3366.
- 19 Smith RIL (1999) Biological and environmental characteristics of three cosmopolitan mosses dominant
20 in continental Antarctica. *Journal of Vegetation Science*, **10**, 231-242.
- 21 Snell KRS, Kokubun T, Griffiths H, Convey P, Hodgson DA, Newsham KK (2009). Quantifying the
22 metabolic cost to an Antarctic liverwort of responding to UV-B radiation exposure. *Global Change*
23 *Biology*, in press.
- 24 Stiling P, Cornelissen T (2007) How does elevated carbon dioxide (CO₂) affect plant-herbivore
25 interactions? A field experiment and meta-analysis of CO₂-mediated changes on plant chemistry
26 and herbivore performance. *Global Change Biology*, **13**, 1823-1842.
- 27 Sullivan JH, Rozema J (1999) UV-B effects on terrestrial plant growth and photosynthesis. In:
28 *Stratospheric ozone depletion: the effects of enhanced UV-B radiation on terrestrial ecosystems* (ed
29 Rozema J), pp. 39-57. Backhuys Publishers, Leiden, The Netherlands.
- 30 Thimijan RW, Carns HR, Campbell LE (1978) *Final report (EPA-IAG-D6-0168): Radiation sources*
31 *and relative environmental control for biological and climatic effects of UV research (BACER)*.
32 Environmental Protection Agency, Washington, DC.
- 33 Turnbull JD, Leslie S, Robinson SA (2009) Desiccation protects two Antarctic mosses from ultraviolet-
34 B induced DNA damage. *Functional Plant Biology*, **36**, 214-221.
- 35 Turnbull JD, Robinson SA (2009) Accumulation of DNA damage in Antarctic mosses: correlations
36 with ultraviolet-B radiation, temperature and turf water content vary amongst species. *Global*
37 *Change Biology*, **15**, 319-329.
- 38 World Meteorological Organization (2007) *Scientific Assessment of Ozone Depletion: 2006*. Global
39 Ozone Research and Monitoring Project - report No. 50. Geneva, Switzerland.

- 1 Xiong F, Day T (2001) Effect of solar ultraviolet-B radiation during springtime ozone depletion on
- 2 photosynthesis and biomass production of Antarctic vascular plants. *Plant Physiology*, **125**, 738-
- 3 751.
- 4 Xiong F, Ruhland CT, Day TA (2002) Effect of springtime solar ultraviolet-B radiation on growth of
- 5 *Colobanthus quitensis* at Palmer Station, Antarctica. *Global Change Biology*, **8**, 1146-1155.

1 *Figure legends*

2 **Fig. 1** Untransformed response ratios for the effects of UV-B exposure on measures of pigmentation,
 3 photosynthesis, growth, biomass and DNA damage in polar vegetation. Where 95% confidence interval bars do
 4 not cross the horizontal dotted line, natural logarithm-transformed response ratios were significantly different
 5 from zero in unweighted analyses (* $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$). All response ratios indicated as
 6 significantly different from zero were also significant at $P < 0.05$ in weighted analyses, including that for leaf area.
 7 Numbers of observations for each parameter are shown in Table 1. *Abbreviations:* UV-B_{ac} mass and UV-B_{ac} area,
 8 UV-B absorbing compounds expressed per unit of mass and area, respectively; carots, carotenoids; chl,
 9 chlorophyll; F_v/F_m , maximum quantum yield of PSII; Φ_{PSII} , actual quantum yield of PSII; P_n , net photosynthesis;
 10 AGB, BGB and TB; above-, belowground and total biomass, respectively; SLA, specific leaf area; CPDs,
 11 cyclobutane pyrimidine dimers.

12

13 **Fig. 2** The influence of categorical variables on untransformed response ratios for (a) UV-B absorbing
 14 compounds expressed per unit of mass, (b) belowground biomass, (c) UV-B absorbing compounds
 15 expressed per unit of area, (d) carotenoids, (e) leaf area and (f) specific leaf area. Where 95%
 16 confidence interval bars do not cross the horizontal dotted lines, natural logarithm-transformed
 17 response ratios were significantly different from zero in unweighted analyses (* $P < 0.05$, ** $P < 0.01$
 18 and *** $P < 0.001$). All response ratios indicated as significantly different from zero were also
 19 significant at $P < 0.05$ in weighted analyses. See Table 1 for P values from unweighted analyses,
 20 indicating effects of categorical variables on response ratios. *Abbreviation:* non-manip, non-
 21 manipulative studies.

22

23 **Fig. 3** Untransformed response ratios for (a) UV-B absorbing compounds expressed per unit of mass,
 24 (b) carotenoid concentration, (c) aboveground biomass and (d) leaf area as a function of percentage
 25 difference in UV-B_{PAS} dose between treatment and control plots in (○) screen, (●) lamp and (●) non-
 26 manipulative studies. Note that the inverses of response ratios are shown for screen studies. Dashed
 27 lines indicate lines of best fit. Line in (a) is a quadratic function. See Table 2 for details of regression
 28 analyses. Significant differences, from unweighted analyses, between the natural logarithm-
 29 transformed response ratios for each method and zero are indicated by * $P < 0.05$, ** $P < 0.01$ and ***
 30 $P < 0.001$ in (b)-(d). All significant effects were also significant at $P < 0.05$ in weighted analyses, except
 31 that in (b), which could not be analysed in this way. See Fig. 2a for the significance of effects of UV-B
 32 exposure on the response ratios shown in (a). *Inset* in (b) is association between untransformed
 33 response ratios for carotenoid concentration and duration of exposure to UV-B radiation.

Table 1. Results from unweighted and weighted meta-analyses testing for the effects of UV-B exposure and categorical variables (methodology, plant form and polar region) on the pigmentation, photosynthesis, biomass, growth and DNA damage of polar vegetation. *P* values from unweighted analyses are shown, with significant values in bold. Those significant at $P < 0.05$ and below in weighted analyses are indicated by (*). All data were natural logarithm-transformed prior to analysis. Number of observations (*n*) for unweighted analyses are shown, with those for weighted analyses in parentheses.

| | pigmentation | | | | photosynthesis | | | biomass, growth and DNA damage | | | | | | | |
|--------------------------|----------------------------|----------------------------|--------------------|-----------------------------|----------------|---------------|-------|--------------------------------|-----------------------------|------------------|----------------------|--------------|--------------|----------------------|--------------------|
| | UV-B _{ac} mass | UV-B _{ac} area | carotenoids | chlorophyll <i>a + b</i> | F_v/F_m | Φ_{PSII} | P_n | above- ground biomass | below- ground biomass | total biomass | height | leaf area | leaf mass | SLA | CPDs |
| <i>n</i> | 42 (30) | 14 (10) | 15 (12) | 17 (14) | 14 (14) | 7 (7) | 9 (9) | 18 (15) | 7 (7) | 14 (9) | 19 (19) | 14 (13) | 10 (9) | 17 (12) | 12 (10) |
| UV-B exposure | 0.015 (*) | <0.001 (*) | 0.713 | 0.396 | 0.543 | 0.956 | 0.125 | 0.003 (*) | 0.440 | 0.817 | <0.001 (*) | 0.165 (*) | 0.914 | 0.503 | 0.030 (*) |
| methodology ¹ | 0.043 [†] | 0.377 [†] | 0.220 [†] | 0.536 | 0.735 | 0.879 | 0.126 | 0.893 | 0.043 | 0.117 | 0.719 | 0.128 | - | 0.359 | 0.174 [†] |
| plant form ² | 0.420 | 0.015 (*) | 0.626 | 0.621 | 0.054 | 0.196 | 0.082 | 0.709 | - | 0.934 | 0.719 | - | 0.887 | 0.462 | 0.220 |
| region ³ | 0.847 | 0.234 | 0.039 | 0.664 | 0.438 | 0.709 | 0.989 | 0.534 | 0.515 | 0.051 | 0.719 | 0.020 | - | <0.001 (*) | 0.400 |

¹lamp, screen or non-manipulative studies; ²bryophyte or angiosperm; ³Arctic or Antarctic. Dashes indicate that analyses were not possible owing to insufficient replication, [†]indicate that weighted analyses were not possible since data from non-manipulative studies could not be included. See Fig. 1 for abbreviations.

Table 2. Number of observations (n), adjusted r^2 and P values from regressions between untransformed response ratios and percentage difference in UV-B_{PAS} between treatment and control plots. Significant P values are marked in bold. Analyses were not made on photosynthetic parameters, height, leaf mass or SLA because of insufficient replication.

| | response | n | adj. r^2 (%) | P |
|--------------------------------|-------------------------|-----|----------------|--------------|
| pigmentation | UV-B _{ac} mass | 29 | 30.7* | 0.001 |
| | UV-B _{ac} area | 10 | 21.4 | 0.100 |
| | carotenoids | 8 | 55.4 | 0.021 |
| | chlorophyll $a + b$ | 9 | 0.0 | 0.368 |
| biomass, growth and DNA damage | aboveground biomass | 13 | 43.9 | 0.008 |
| | belowground biomass | 5 | 0.0 | 0.887 |
| | total biomass | 10 | 0.0 | 0.399 |
| | leaf area | 13 | 45.8 | 0.007 |
| | CPDs | 7 | 0.0 | 0.679 |

*best fit provided by a quadratic function. Abbreviations as in Fig. 1.

Appendix 1. Literature from which the data for the meta-analysis were taken

| reference | method | species (plant form) | region | pigmentation | | | | photosynthesis | | | biomass, growth and DNA damage | | | | | | | |
|--|--------|-----------------------------------|--------|--------------------|--------------------|--------|---------|----------------|---------------|-------|--------------------------------|-----|----|------|------|------|-----|------|
| | | | | UV-B _{ac} | UV-B _{ac} | carots | chl | F_v/F_m | Φ_{PSII} | P_n | AGB | BGB | TB | ht | leaf | leaf | SLA | CPDs |
| | | | | mass | area | | $a + b$ | | | | | | | area | mass | | | |
| Albert <i>et al.</i> (2005a) | S | <i>Salix arctica</i> (A) | Arc | | | | | x | | | | | | | | | | |
| Albert <i>et al.</i> (2005b) | S | <i>Salix arctica</i> (A) | Arc | x | | | | x | | | | | | | | x | x | |
| Albert <i>et al.</i> (2008) [†] | S | <i>Vaccinium uliginosum</i> (A) | Arc | x | | | | x | | x | | x | x | x | x | x | x | |
| Bredahl <i>et al.</i> (2004) | S | <i>Salix arctica</i> (A) | Arc | | | | | x | | x | | | | | | | | |
| | | <i>Vaccinium uliginosum</i> (A) | | | | | | x | | | | | | | | | | |
| Kotilainen <i>et al.</i> (2008) | S | <i>Alnus incana</i> (A) | Arc | x | | | | | | | | | | | | | | |
| | | <i>Betula pubescens</i> (A) | | x | | | | | | | | | | | | | | |
| Rinnan <i>et al.</i> (2005) | S | <i>Vaccinium uliginosum</i> (A) | Arc | | | | | | | | | | x | | | | | |
| Day <i>et al.</i> (2001) | S | <i>Colobanthus quitensis</i> (A) | Ant | | | | | | | | | x | | | x | | | |
| | | <i>Deschampsia antarctica</i> (A) | | | | | | | | | | x | | | x | | | |
| Day <i>et al.</i> (2008) | S | <i>Colobanthus quitensis</i> (A) | Ant | | | | | | | | | x | | | | | | |
| | | <i>Deschampsia antarctica</i> (A) | | | | | | | | | | x | | | | | | |
| Smith (1999) [†] | S | <i>Bryum argenteum</i> (B) | Ant | | | | | | | | | | | | x | | | |
| | | <i>B. pseudotriquetrum</i> (B) | | | | | | | | | | | | | x | | | |
| | | <i>Ceratodon purpureus</i> (B) | | | | | | | | | | | | | x | | | |
| Newsham <i>et al.</i> (2005) | S | <i>Cephaloziella varians</i> (B) | Ant | x | | | | x | | | | | | | | | | |
| Ruhland & Day (2000) | S | <i>Colobanthus quitensis</i> (A) | Ant | | x | | | | | | | | | | | x | | |
| | | <i>Deschampsia antarctica</i> (A) | | | x | | | | | | | | | | | x | | |
| Ruhland & Day (2001) | S | <i>Colobanthus quitensis</i> (A) | Ant | | x | | | x | | | | | | | | | x | |
| Xiong & Day (2001) | S | <i>Colobanthus quitensis</i> (A) | Ant | x | x | x | x | x | x | x | x | x | x | x | x | x | | x |
| | | <i>Deschampsia antarctica</i> (A) | | x | x | x | x | x | x | x | x | x | x | x | x | x | | x |
| Xiong <i>et al.</i> (2002) [†] | S | <i>Colobanthus quitensis</i> (A) | Ant | x | x | | | x | | | | x | x | x | x | x | | |
| Newsham <i>et al.</i> (2002) | NM | <i>Cephaloziella varians</i> (B) | Ant | x | | | | | | | | | | | | | | |
| | | <i>Sanionia uncinata</i> (B) | | x | | | | | | | | | | | | | | |
| Newsham (2003) | NM | <i>Andreaea regularis</i> (B) | Ant | x | | | | | | | | | | | | | | |
| Dunn & Robinson (2006) | NM | <i>Bryum pseudotriquetrum</i> (B) | Ant | x | | | | | | | | | | | | | | |
| | | <i>Ceratodon purpureus</i> (B) | | x | | | | | | | | | | | | | | |




Fig. 1 Untransformed response ratios for the effects of UV-B exposure on measures of pigmentation, photosynthesis, growth, biomass and DNA damage in polar vegetation. Where 95% confidence interval bars do not cross the horizontal dotted line, natural logarithm-transformed response ratios were significantly different from zero in unweighted analyses (* $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$). All response ratios indicated as significantly different from zero were also significant at $P < 0.05$ in weighted analyses, including that for leaf area. Numbers of observations for each parameter are shown in Table 1. Abbreviations: UV-Bac mass and UV-Bac area, UV-B absorbing compounds expressed per unit of mass and area, respectively; carots, carotenoids; chl, chlorophyll; F_v/F_m, maximum quantum yield of PSII; Φ_{PSII}, actual quantum yield of PSII; P_n, net photosynthesis; AGB, BGB and TB; above-, belowground and total biomass, respectively; SLA, specific leaf area; CPDs, cyclobutane pyrimidine dimers.

209x296mm (500 x 500 DPI)




Fig. 2 The influence of categorical variables on untransformed response ratios for (a) UV-B absorbing compounds expressed per unit of mass, (b) belowground biomass, (c) UV-B absorbing compounds expressed per unit of area, (d) carotenoids, (e) leaf area and (f) specific leaf area. Where 95% confidence interval bars do not cross the horizontal dotted lines, natural logarithm-transformed response ratios were significantly different from zero in unweighted analyses (* $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$). All response ratios indicated as significantly different from zero were also significant at $P < 0.05$ in weighted analyses. See Table 1 for P values from unweighted analyses, indicating effects of categorical variables on response ratios. Abbreviation: non-manip, non-manipulative studies.

209x296mm (500 x 500 DPI)




Fig. 3 Untransformed response ratios for (a) UV-B absorbing compounds expressed per unit of mass, (b) carotenoid concentration, (c) aboveground biomass and (d) leaf area as a function of percentage difference in UV-BPAS dose between treatment and control plots in (□) screen, (□) lamp and (□) non-manipulative studies. Note that the inverses of response ratios are shown for screen studies. Dashed lines indicate lines of best fit. Line in (a) is a quadratic function. See Table 2 for details of regression analyses. Significant differences, from unweighted analyses, between the natural logarithm-transformed response ratios for each method and zero are indicated by * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$ in (b)-(d). All significant effects were also significant at $P < 0.05$ in weighted analyses, except that in (b), which could not be analysed in this way. See Fig. 2a for the significance of effects of UV-B exposure on the response ratios shown in (a). Inset in (b) is association between untransformed response ratios for carotenoid concentration and duration of exposure to UV-B radiation.

209x296mm (500 x 500 DPI)

Table 1. Results from unweighted and weighted meta-analyses testing for the effects of UV-B exposure and categorical variables (methodology, plant form and polar region) on the pigmentation, photosynthesis, biomass, growth and DNA damage of polar vegetation. *P* values from unweighted analyses are shown, with significant values in bold. Those significant at $P < 0.05$ and below in weighted analyses are indicated by (*). All data were natural logarithm-transformed prior to analysis. Number of observations (*n*) for unweighted analyses are shown, with those for weighted analyses in parentheses.

| | pigmentation | | | | photosynthesis | | | biomass, growth and DNA damage | | | | | | | |
|--------------------------|----------------------------|----------------------------|--------------------|-----------------------------|----------------|---------------|-------|--------------------------------|-----------------------------|------------------|----------------------|--------------|--------------|----------------------|--------------------|
| | UV-B _{ac} mass | UV-B _{ac} area | carotenoids | chlorophyll <i>a + b</i> | F_v/F_m | Φ_{PSII} | P_n | above- ground biomass | below- ground biomass | total biomass | height | leaf area | leaf mass | SLA | CPDs |
| <i>n</i> | 42 (30) | 14 (10) | 15 (12) | 17 (14) | 14 (14) | 7 (7) | 9 (9) | 18 (15) | 7 (7) | 14 (9) | 19 (19) | 14 (13) | 10 (9) | 17 (12) | 12 (10) |
| UV-B exposure | 0.015 (*) | <0.001 (*) | 0.713 | 0.396 | 0.543 | 0.956 | 0.125 | 0.003 (*) | 0.440 | 0.817 | <0.001 (*) | 0.165 (*) | 0.914 | 0.503 | 0.030 (*) |
| methodology ¹ | 0.043 [†] | 0.377 [†] | 0.220 [†] | 0.536 | 0.735 | 0.879 | 0.126 | 0.893 | 0.043 | 0.117 | 0.719 | 0.128 | - | 0.359 | 0.174 [†] |
| plant form ² | 0.420 | 0.015 (*) | 0.626 | 0.621 | 0.054 | 0.196 | 0.082 | 0.709 | - | 0.934 | 0.719 | - | 0.887 | 0.462 | 0.220 |
| region ³ | 0.847 | 0.234 | 0.039 | 0.664 | 0.438 | 0.709 | 0.989 | 0.534 | 0.515 | 0.051 | 0.719 | 0.020 | - | <0.001 (*) | 0.400 |

¹lamp, screen or non-manipulative studies; ²bryophyte or angiosperm; ³Arctic or Antarctic. Dashes indicate that analyses were not possible owing to insufficient replication, [†] indicate that weighted analyses were not possible since data from non-manipulative studies could not be included. See Fig. 1 for abbreviations.

Table 2. Number of observations (n), adjusted r^2 and P values from regressions between untransformed response ratios and percentage difference in UV-B_{PAS} between treatment and control plots. Significant P values are marked in bold. Analyses were not made on photosynthetic parameters, height, leaf mass or SLA because of insufficient replication.

| | response | n | adj. r^2 (%) | P |
|--------------------------------|-------------------------|-----|----------------|--------------|
| pigmentation | UV-B _{ac} mass | 29 | 30.7* | 0.001 |
| | UV-B _{ac} area | 10 | 21.4 | 0.100 |
| | carotenoids | 8 | 55.4 | 0.021 |
| | chlorophyll $a + b$ | 9 | 0.0 | 0.368 |
| biomass, growth and DNA damage | aboveground biomass | 13 | 43.9 | 0.008 |
| | belowground biomass | 5 | 0.0 | 0.887 |
| | total biomass | 10 | 0.0 | 0.399 |
| | leaf area | 13 | 45.8 | 0.007 |
| | CPDs | 7 | 0.0 | 0.679 |

*best fit provided by a quadratic function. Abbreviations as in Fig. 1.

Appendix 1. Literature from which the data for the meta-analysis were taken

| reference | method | species (plant form) | region | pigmentation | | | | photosynthesis | | | biomass, growth and DNA damage | | | | | | | |
|--|--------|-----------------------------------|--------|--------------------|--------------------|--------|---------|----------------|---------------|-------|--------------------------------|-----|----|------|------|------|-----|------|
| | | | | UV-B _{ac} | UV-B _{ac} | carots | chl | F_v/F_m | Φ_{PSII} | P_n | AGB | BGB | TB | ht | leaf | leaf | SLA | CPDs |
| | | | | mass | area | | $a + b$ | | | | | | | area | mass | | | |
| Albert <i>et al.</i> (2005a) | S | <i>Salix arctica</i> (A) | Arc | | | | | x | | | | | | | | | | |
| Albert <i>et al.</i> (2005b) | S | <i>Salix arctica</i> (A) | Arc | x | | | | x | | | | | | | | x | x | |
| Albert <i>et al.</i> (2008) [†] | S | <i>Vaccinium uliginosum</i> (A) | Arc | x | | | | x | | x | | x | x | x | | x | x | |
| Bredahl <i>et al.</i> (2004) | S | <i>Salix arctica</i> (A) | Arc | | | | | x | | x | | | | | | | | |
| | | <i>Vaccinium uliginosum</i> (A) | | | | | | x | | | | | | | | | | |
| Kotilainen <i>et al.</i> (2008) | S | <i>Alnus incana</i> (A) | Arc | x | | | | | | | | | | | | | | |
| | | <i>Betula pubescens</i> (A) | | x | | | | | | | | | | | | | | |
| Rinnan <i>et al.</i> (2005) | S | <i>Vaccinium uliginosum</i> (A) | Arc | | | | | | | | | | x | | | | | |
| Day <i>et al.</i> (2001) | S | <i>Colobanthus quitensis</i> (A) | Ant | | | | | | | | | x | | | x | | | |
| | | <i>Deschampsia antarctica</i> (A) | | | | | | | | | | x | | | x | | | |
| Day <i>et al.</i> (2008) | S | <i>Colobanthus quitensis</i> (A) | Ant | | | | | | | | | x | | | | | | |
| | | <i>Deschampsia antarctica</i> (A) | | | | | | | | | | x | | | | | | |
| Smith (1999) [†] | S | <i>Bryum argenteum</i> (B) | Ant | | | | | | | | | | | | | x | | |
| | | <i>B. pseudotriquetrum</i> (B) | | | | | | | | | | | | | | x | | |
| | | <i>Ceratodon purpureus</i> (B) | | | | | | | | | | | | | | x | | |
| Newsham <i>et al.</i> (2005) | S | <i>Cephaloziella varians</i> (B) | Ant | x | | | | x | | | | | | | | | | |
| Ruhland & Day (2000) | S | <i>Colobanthus quitensis</i> (A) | Ant | | x | | | | | | | | | | | x | | |
| | | <i>Deschampsia antarctica</i> (A) | | | x | | | | | | | | | | | x | | |
| Ruhland & Day (2001) | S | <i>Colobanthus quitensis</i> (A) | Ant | | x | | | x | | | | | | | | | x | |
| Xiong & Day (2001) | S | <i>Colobanthus quitensis</i> (A) | Ant | x | x | x | x | x | x | x | x | x | x | x | x | | x | |
| | | <i>Deschampsia antarctica</i> (A) | | x | x | x | x | x | x | x | x | x | x | x | x | | x | |
| Xiong <i>et al.</i> (2002) [†] | S | <i>Colobanthus quitensis</i> (A) | Ant | x | x | | | x | | | | x | x | x | x | | | |
| Newsham <i>et al.</i> (2002) | NM | <i>Cephaloziella varians</i> (B) | Ant | x | | | | | | | | | | | | | | |
| | | <i>Sanionia uncinata</i> (B) | | x | | | | | | | | | | | | | | |
| Newsham (2003) | NM | <i>Andreaea regularis</i> (B) | Ant | x | | | | | | | | | | | | | | |
| Dunn & Robinson (2006) | NM | <i>Bryum pseudotriquetrum</i> (B) | Ant | x | | | | | | | | | | | | | | |
| | | <i>Ceratodon purpureus</i> (B) | | x | | | | | | | | | | | | | | |

