

University of Wollongong

Research Online

Faculty of Science, Medicine and Health -
Papers: part A

Faculty of Science, Medicine and Health

1-1-2013

Abundance of *Tesseropora rosea* at the margins of its biogeographic range is closely linked to recruitment, but not fecundity

Eszter Z. Hidas

University of Wollongong, ezh998@uow.edu.au

Kenneth G. Russell

Charles Sturt University, kerussell@csu.edu.au

David J. Ayre

University of Wollongong, david_ayre@uow.edu.au

Todd E. Minchinton

University of Wollongong, tminch@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>



Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Hidas, Eszter Z.; Russell, Kenneth G.; Ayre, David J.; and Minchinton, Todd E., "Abundance of *Tesseropora rosea* at the margins of its biogeographic range is closely linked to recruitment, but not fecundity" (2013).

Faculty of Science, Medicine and Health - Papers: part A. 781.

<https://ro.uow.edu.au/smhpapers/781>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Abundance of *Tesseropora rosea* at the margins of its biogeographic range is closely linked to recruitment, but not fecundity

Abstract

For benthic marine invertebrates with planktonic larvae, adult abundance at the species' range limit may depend on local fecundity, and recruitment from central, source populations. We tested the importance of these processes using a hierarchical design at multiple spatial scales (from tens to hundreds of kilometres) to assess relationships between patterns of adult abundance, fecundity and recruitment in the rocky intertidal barnacle *Tesseropora rosea*, approaching its geographical range limit on the southeast coast of Australia. Across the 600 km region, adult abundance, recruitment and fecundity all declined towards the range limit, but only adult abundance and recruitment were positively correlated, suggesting that recruitment is negatively affected at the range limit and that recruitment plays a significant role in setting the geographical patterns of abundance of *T. rosea*. In contrast, at smaller spatial scales within this region, adult abundance and recruitment varied by up to 15- and 100-fold, respectively, among some locations only tens of kilometres apart, and no obvious patterns in adult-recruit relationships were revealed towards the range limit. Local patterns of abundance and life-cycle relationships do not, therefore, translate to biogeographic scales, but investigation at this multitude of spatial scales helps to identify the defining processes that account for spatial variation and determine population structure towards the range limits of benthic marine invertebrates.

Keywords

tesseropora, rosea, not, margins, its, biogeographic, range, closely, abundance, recruitment, but, linked, fecundity

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Hidas, E. Z., Russell, K. G., Ayre, D. J. & Minchinton, T. E. (2013). Abundance of *Tesseropora rosea* at the margins of its biogeographic range is closely linked to recruitment, but not fecundity. *Marine Ecology: Progress Series*, 483 199-208.

ARTICLE TYPE: RESEARCH ARTICLE

**TITLE: LINKS BETWEEN ADULTS, FECUNDITY AND RECRUITS: DO
LOCAL RELATIONSHIPS TRANSLATE TO BIOGEOGRAPHIC SCALES?**

RUNNING HEAD: LINKS BETWEEN ADULTS, FECUNDITY AND RECRUITS

Eszter Z. Hidas¹, Kenneth G. Russell², David J. Ayre¹ and Todd E. Minchinton^{1*}

¹Institute for Conservation Biology and Environmental Management, School of
Biological Sciences, University of Wollongong, NSW, Australia, 2522

²School of Computing & Mathematics, Charles Sturt University, Locked Bag 588,
Wagga Wagga, NSW, Australia, 2678

*Corresponding author: E-mail: tminch@uow.edu.au

ABSTRACT

For benthic marine invertebrates with planktonic larvae, adult abundance at the species' range limit may depend on local fecundity, and recruitment from central, source populations. We test the importance of these processes using a hierarchical design at multiple spatial scales (from tens to hundreds of kilometres) to assess relationships between patterns of adult abundance, fecundity, and recruitment for the rocky intertidal barnacle, *Tesseropora rosea*, approaching its geographical range limit on the southeast coast of Australia. Across the 600 km region, adult abundance, recruitment and fecundity all declined towards the range limit, but only adult abundance and recruitment were positively correlated, suggesting that recruitment is negatively impacted at the range limit and, that recruitment plays a significant role in setting the geographical patterns of abundance of *T. rosea*. In contrast, at smaller spatial scales within this region, adult abundance and recruitment varied by up to 15- and 100-fold respectively, among some locations only tens of kilometres apart, and no obvious patterns in adult-recruit relationships were revealed towards the range limit. Local patterns of abundance and life-cycle relationships do not, therefore, translate to biogeographic scales, but investigation at this multitude of spatial scales help to identify the defining processes that account for spatial variation and determine population structure towards the range limits of benthic marine invertebrates.

KEYWORDS larval supply, adult abundance, intertidal barnacles, supply side ecology, recruitment

INTRODUCTION

Ecologists have long sought to understand the factors and processes that determine the abundance distribution and range limits of organisms (Grinnell 1917, Hutchins 1947). In recent years, several studies (Sagarin & Gaines 2002b, Gilman 2005, Hidas et al. 2010) have tested the applicability of the abundant-centre hypothesis, which predicts that the abundance of a species is greatest at the centre of its geographical range, to benthic marine invertebrates. The little empirical evidence supporting this type of distribution in marine habitats has highlighted the need to understand how the different life-cycle stages of benthic marine invertebrates interact at various spatial scales to determine adult abundance across the geographical range and at the range limit (Sagarin & Gaines 2002a, Sagarin et al. 2006).

For many benthic marine invertebrates that have a planktonic larval stage, adult abundances at a local scale can be determined by patterns of larval recruitment (Connell 1985, Gaines & Roughgarden 1985, Minchinton & Scheibling 1991). Because planktonic larvae disperse mostly passively with oceanic currents (Pechenik 1999, Kinlan & Gaines 2003), dominant surface currents are expected to cause the downstream accumulation of larvae originating from productive source populations within the middle of the range. If there were no environmental factors influencing post-settlement survival of recruits or adults at smaller spatial scales approaching the range limit, then this should lead to an increase in adult abundance towards the downstream range limit (Gaylord & Gaines

2000). Empirical studies assessing large-scale patterns of recruitment for benthic marine invertebrates approaching their range limits have shown a decline in recruit density gradually towards, or abruptly at, the range limit (Broitman et al. 2008, Herbert et al. 2009); decreased recruitment frequency at the range limit (Zacherl et al. 2003, Lima et al. 2006, Hidas et al. 2010); or no changes in recruitment patterns towards the range limit (Broitman et al. 2008, Lathlean et al. 2010). Assessing these variations in recruitment, and their relationship to adult abundance and fecundity at smaller spatial scales approaching the range limit, is necessary to understand the variable outcomes in patterns of abundance at geographical scales (Fowler-Walker et al. 2005).

Aside from the effects of oceanic currents, recruitment across the ranges of benthic marine invertebrates also partially depends on the intensity and success of larval production (i.e. fecundity) (Hughes 1990). Fecundity, however, may decline towards the range limit due to changes in air or sea surface temperature (Hughes et al. 2000, Ramirez Llorda 2002) or food availability (Ramirez Llorda 2002), particularly along the latitudinal distribution of many intertidal species (Lewis 1986), and thus may lead to a decrease in recruitment and adult abundance towards the range limit for species whose larvae are locally retained. The only previous large-scale study that has explored adult-fecundity-recruitment relationships of benthic marine invertebrates focused on broadcast spawning corals on the Great Barrier Reef in Australia (Hughes et al. 2000). For this species, adult abundance and recruitment were not correlated at either small or large spatial scales, but 72% of the variation in recruitment was explained by the fecundity of the adults at large spatial scales. This result is likely to reflect the highly limited dispersal and low degree of population connectivity of at least some coral species (Ayre & Hughes 2000).

Here we examine adult-fecundity-recruitment relationships at a hierarchy of spatial scales (from tens to hundreds of kilometres) across the southern third of the geographical range of the intertidal rocky shore barnacle, *Tesseropora rosea*, on the southeast coast of Australia (Edgar 1997, Hidas et al. 2007). The adult abundance of this species declines steeply from Sydney, New South Wales, to its southern range limit at Cape Conran in eastern Victoria (Fig. 1, Hidas et al. 2010), its larvae spend up to five weeks in the plankton (Wisely & Blick 1964, Caffey 1985), and the recruit stage of the life-cycle is easily recognisable. During the larval phase, its distribution is likely to be influenced by the warm temperate East Australian Current (EAC), which flows predominantly southwards along much of the eastern coast of Australia (Ridgway & Dunn 2003). The current partially deflects away from the coast at approximately the latitude of Sydney, and partially continues southwards as a weaker current, forming occasional coastal eddies as it reaches the southeast corner of Australia (Fig. 1; Ridgway & Dunn 2003). West of this point, sea surface temperatures change dramatically by up to five degrees compared with conditions to the north (Knox 1963, Ridgway & Dunn 2003), and for 300 km west of Cape Conran, rocky shore habitat is mostly unavailable, except for one isolated rocky patch (Fig. 1, Hidas et al. 2007). As a consequence of these complex hydrological, climatic and topographic conditions and the high dispersal potential of the species, we expect significant small-scale (over tens of kilometres) variation in demographic processes across the southern third of the species' range, which may also be reflected by a disparity in small-scale adult and recruit relationships. At large scales (over hundreds of kilometres), however, we expect to see a decrease in fecundity resulting from decreasing air or sea surface temperatures, a decline in recruit abundance resulting from the potential re-circulation of larvae by coastal eddies upstream, and as a result, despite the high

dispersal potential of this species, a significant link between adult abundance, fecundity, and recruitment.

MATERIALS AND METHODS

We examined the adult abundance, fecundity, sexual maturity and recruitment of the barnacle *Tesseropora rosea* along 600 km of coastline, from Garie Beach near Sydney, (34⁰10' S, 151⁰04' E) to Cape Conran in Victoria (37⁰08' S, 148⁰07' E) (Fig. 1), which constitutes approximately the southern third of the geographical range of the species (Edgar 1997, Hidas et al. 2007). *Tesseropora rosea* is found on exposed emergent rock platforms in the mid shore of the intertidal zone (Caffey 1985, Edgar 1997). This species is hermaphroditic, adults cross-fertilise with their nearest neighbours, and multiple broods are likely produced per breeding season as mature adults almost always contain distinct batches of embryos at different developmental stages (Justin Lathlean, pers. comm.). Mature larvae are released into the water column predominantly from January to April (Wisely & Blick 1964), but also sporadically throughout the year (Caffey 1985, Egan & Anderson 1988).

To assess spatial changes in adult abundance, recruitment, sexual maturity and fecundity from small to large spatial scales (i.e. from tens to hundreds of kilometres) towards the range limit, we surveyed at three different spatial scales using a hierarchical sampling design, with four, fixed sectors each covering 70 to 110 km of coastline, four locations within each sector, and three sites at each location (Fig. 1). To ensure that temporal variation did not influence spatial variation, we sampled four times during a 12-month period: May and September 2005, and January and May 2006. Sectors were separated by

30-100 km, locations by 10-50 km, and sites by 0-50m. Sites were approximately 30 m alongshore and of variable intertidal extent (from 6 to 50 m). Sectors 1 to 4 represented regions furthest from and closest to the range limit respectively, although sectors 1, 2 and 3 were all considered to be within the interior of the range of the species, while sector 4 was at the range limit (Fig. 1). Locations and sites accessible from land were randomly selected, although spaced apart at minimum distances indicated above to cover the geographical range along the coast, and chosen to have the same physical characteristics. Locations were positioned on moderately exposed headlands facing the open ocean, with a platform slope of no more than 30 degrees, but most were gradually sloping rock platforms. Rock type varied naturally between siltstone/sandstone (n = 13), granite (n = 2) and basalt (n = 1), but this variation was not systematic towards the range limit and therefore results were not spatially confounded in this way.

To estimate spatial variation in the abundance of adults and recruits we counted adults (≥ 3 mm aperture length) and recruits (< 3 mm aperture length) (Wisely and Blick 1964, Lathlean et al. 2010) within 20 haphazardly placed 0.15 x 0.15 m photoquadrats/site. To quantify levels of sexual maturity and fecundity of *Tesseropora rosea* at each location, we haphazardly collected samples of 20 to 60 (depending on availability) adults (i.e. individuals with ≥ 3 mm aperture length) and determined the proportion of these displaying gonads (i.e. sexually mature) and ready-to-release mature larvae (i.e. fecund, as indicated by orange or brown embryos with obvious eyed nauplii) (Wisely & Blick 1964), respectively.

We used Restricted Maximum Likelihood (REML; Patterson and Thompson, 1971) analysis (SPSS 20; IBM Corp. 2011) to test for differences in the mean densities of adults

and recruits among sectors (entered as a fixed factor), and among locations within sectors (entered as a random factor) along the southern distributional range of *Tesseropora rosea* for the time period between May 2005 and May 2006. Adult and recruit density data for each site at each location for each of the four sampling times (when available) was obtained and entered into the analysis by taking the mean of the 20 photoquadrats. Missing data for some sites for some sampling times, corresponding to days when sampling could not be carried out, rendered our analysis unbalanced and therefore, conventional mixed model ANOVA was not appropriate. Time was added into the analysis as a fixed factor blocking term to ensure that spatial patterns were not confounded by temporal variation. Data were log transformed as a Kolmogorov-Smirnov test indicated that they did not satisfy the assumptions of a normal distribution. Pairwise comparisons between each pair of sectors, and each pair of sampling times were carried out through Bonferroni post-hoc test. Restricted Maximum Likelihood analyses generate two test statistics: a standard F ratio for the fixed factor, and a Wald-Z value for the nested random factor; and denominator variances in the F-tests do not necessarily have a whole number for the degrees of freedom, because they are estimated by approximate methods.

To test for differences in the proportions of sexually mature and fecund individuals of *Tesseropora rosea* among sectors (entered as a fixed effect) and among locations within sectors (entered as a random effect) at the four different sampling times (entered as a fixed effect), we used generalized linear mixed models in GenStat 14.1 (VSN International 2011), using a binomial distribution with a logit link. Data were entered into the analysis as the numbers of individuals with and without gonads per location for

sexual maturity, and the numbers of individuals with and without mature embryos per location for fecundity, for each of the four sampling times.

Finally, we used simple correlation analyses (SPSS 20; IBM Corp. 2011) to test for relationships between recruit and adult densities of *Tesseropora rosea* across the 600 km study region (based on mean values at locations), and across sectors (based on mean values at sites). Additionally, using the same methodology we assessed the relationships between the proportion of adults with larvae (fecundity) and recruit densities, and between the proportion of adults with larvae (fecundity) and adult densities of *T. rosea*, based on mean values for each location across the 600 km study region. Recruit density data were initially transformed, but this did not result in a better fit for any of the relationships tested, and hence results are presented as untransformed data. To allow for the time lag between larval release and settlement, relationships between densities of adults and recruits, and between the proportions of adults with larvae and densities of recruits were examined by correlating the means of the May and September 2005, and January 2006 adult densities and their proportions of larvae at each location, with the means of the September 2005, and January and May 2006 recruit densities at each location.

RESULTS

Patterns of Adult Abundance, Recruitment and Fecundity

The mean densities of adult *Tesseropora rosea*, pooled over the four sampling times from May 2005 to May 2006, were 7- to 15-fold higher within sectors 1, 2 and 3 in the middle of the range, with values ranging from 14 to 30 adults per quadrat, than within sector 4 at

the species' range limit where on average only 3 adults were present per quadrat (Fig. 2a). There appeared to be a step-wise decline in adult density towards the range limit (Fig. 2a), with sectors 2 and 3 having considerably lower densities than sector 1. However, adult density only differed significantly between sector 1 and the range limit sector 4 (Fig. 2a, Table 1). There was significant temporal variation in adult densities between September 2005 and January 2006 (Table 1), and significant spatial variation among locations within sectors (Table 1), particularly in sectors 1 and 2 where densities varied between 4- and 100-fold among some locations respectively (Fig. 2e).

The mean densities of recruits, pooled over the four sampling times, mirrored the variation in adult densities across the four sectors (Fig. 2b). Mean density displayed a non-significant decline across sectors 1, 2 and 3, from 119 to 73 recruits per quadrat, and was significantly lowest at the range limit within sector 4, where we detected only a mean of 11 recruits per quadrat (Fig. 2b, Table 1). Similar to adult density, recruit density varied significantly among locations within sectors (Table 1), particularly in sector 2 where densities varied 15-fold between some locations (Fig. 2f), as well as between sampling times - namely between January and May 2006 (Table 1).

Interestingly, we found little evidence of variation in sexual maturity across sectors. The mean proportions of adults with gonads were consistently high (95-100 %) within all four sectors (Fig. 2d), and there was little variation among locations within sectors (Fig. 2h). Indeed, results of the generalised linear mixed model analysis showed no significant changes in sexual maturity across space (Table 2). There were, however, changes in sexual maturity across time (Table 2; Wald test, $p < 0.001$). Fecundity, in contrast, showed a decreasing trend towards the range limit, with a mean of only 6 % of individuals

containing mature embryos in sector 4, compared to 17 %, 20 % and 27 % within sectors 1, 2 and 3 respectively (Fig. 2c & 2g). These proportions, however, varied significantly among sampling times (Table 3), and there was thus a significant interaction between sector and time (Wald test, $p < 0.001$). There were no significant differences in fecundity among sectors in May 2005 and January 2006, whereas in September 2005 the proportion of fecund individuals was significantly greater in sector 3 than in other sectors, and in May 2006 the proportions of fecund individuals were significantly greater in sector 3 than sector 1 and significantly lower in sector 4 than all other sectors (Table 3).

Adult-Fecundity-Recruit Relationships

Across the entire 600 km region, adult and recruit densities were positively correlated, with 38 % of the variance in adult density explained by variation in recruit density (Fig. 3a). In contrast, fecundity was not significantly correlated with either the densities of adults (Fig. 3b) or recruits (Fig. 3c). Indeed, several locations within the middle of the range with low proportions of fecund adults still had high densities of adults and recruits (Fig. 3b&c). Within sectors, densities of recruits and adults were only significantly correlated within sector 2 (Fig. 3e), while in sectors 1, 3 and the range limit sector 4 (Figs 3d,f&g respectively) these relationships showed no significant patterns.

DISCUSSION

Our hierarchical survey of populations of the intertidal barnacle *Tesseropora rosea* revealed a dramatic decline in adult and recruit densities at locations towards the southern range limit on the southeast coast of Australia. Across the 600 km region, patterns of recruitment were positively correlated with variations in the number of adults, but they

were not significantly correlated with variations in the proportions of sexually mature or fecund adults, at each location. This contrasts with the outcome of the study by Hughes et al. (2000) who examined these relationships for corals at the centre of their geographical ranges on Australia's Great Barrier Reef, in which fecundity and recruitment were strongly positively associated at large spatial scales. The difference between our study and that of Hughes et al. (2000) may be due to the more widespread dispersal of *T. rosea* compared to coral species (Ayre et al. 2009), which may increase the scale at which fecundity and recruitment are related. Alternatively, in the Hughes et al. (2000) study, coral recruits were only identified to family or genus level, which may have confounded species-specific relationships. Our results suggest that either dispersal is limited, or some biotic or abiotic factor impacts negatively on the recruitment stage of the life-cycle of *T. rosea* as it approaches its southern range limit on the southeast coast of Australia.

In contrast to the positive relationship between adult and recruit densities of *Tesseropora rosea* across our 600 km study area, smaller-scale adult-recruit relationships were only evident within one of our four sectors. This disparity between adult and recruit densities at smaller spatial scales is not surprising, given the long larval stage of the species (Wisely & Blick 1964, Caffey 1985) and the potential for widespread dispersal provided by the predominantly southward flowing East Australian Current (Ridgway & Dunn 2003). Indeed, we were surprised to find a positive relationship between adults and recruits within one of the sectors positioned in the middle of our study area. This result suggests that adult-recruit relationships are not influenced by position within the geographical range but rather, by more local processes. This was also evident from the large differences in adult and recruit densities among some locations positioned only tens

of kilometres apart. Small-scale natural variations in abiotic and biotic conditions at each location, such as the density of already existing populations and their impacts on food and habitat availability (Connell 1985, Minchinton & Scheibling 1991), temperature (Helmuth et al. 2002), predation and competition (Denley & Underwood 1979) as well as the effects of currents and wave action specific to platforms (Connell 1985, Gaines & Roughgarden 1985) have been shown to influence adult abundance, and small-scale adult-recruit relationships of marine intertidal invertebrates. While every effort was made to choose study locations with comparable physical attributes, conditions inevitably varied among platforms independent of latitude and proximity to the range limit, thus influencing populations at a local scale.

An abrupt decline in recruit abundance at range limit locations of *Tesseropora rosea* was observed, which may have several possible explanations. If local populations of *T. rosea* at the range limit are dependent on the supply of larvae from source populations upstream, which is possible considering the high degree of genetic connectivity among populations of *T. rosea* and other taxa along the coasts of southern New South Wales and eastern Victoria (Ayre et al. 2009), then the weakening of the EAC and the occasional formation of coastal eddies on the southeast corner of Australia may fail to transport larvae in large enough numbers downstream, or may re-circulate larvae back to, or close to, their shores of origin (Murray-Jones & Ayre 1997, Ridgway & Dunn 2003). Indeed, analyses of size-frequency distributions of *T. rosea* along the same stretch of coast appeared to point towards less frequent or temporally variable recruitment at the range limit (Hidas et al. 2010). The influence of less frequent recruitment events at range limit locations on the southward distribution of the species may also be exacerbated by the

predominantly unsuitable rocky shore habitat for 300 km westwards of Cape Conran (Hidas et al. 2007), which may limit the further southward extension of the species (Lewis et al. 1986). An investigation into the latitudinal changes in larval abundance of *T. rosea* as it approaches its southern range limit would help to confirm these hypotheses.

The declining abundance of recruits at the range limit could also potentially reflect a decline in fecundity and larval development, particularly if local populations of *T. rosea* were mostly dependent on local larval recruitment. However, patterns of recruitment and fecundity of *Tesseropora rosea* were not significantly correlated across the 600km study. Fecundity varied considerably over space and time, was interestingly greatest in sector 3 during two of the four sampling times, and was only significantly lowest at the range limit during one of the four sampling times. There is faint evidence to suggest, therefore, that fecundity is negatively impacted at the range limit, but these results should be interpreted with some caution. Larval maturation may simply be delayed, larvae may fail to mature, or the number of broods of embryos produced or frequency of larval release may be reduced in populations approaching the range limit. Although almost all adults were sexually mature in all locations, fertilization success may also be reduced at lower densities because a decreased adult density may reduce the probability of encountering a near neighbour (Kent et al. 2003). Furthermore, *T. rosea* is considered to reproduce predominantly from January to April (Wisely & Blick 1964), however the timing and duration of the reproductive season has been observed to vary among years (Caffey 1985), and may vary with sea surface or air temperature (Knox 1963, Hughes et al. 2000, Ramirez Llorda 2002). For example, intertidal marine invertebrates on the European Atlantic coast have shorter breeding seasons at their northern range limits (Lewis 1986,

Burrows et al. 1992). A more detailed study, with a finer measurement of fecundity, such as the weight of egg lamellae or brooded larvae, is required to reveal more subtle effects of the range limit on fecundity.

Finally, recruit abundance may decrease at the range limit because the settlement or survival of larvae may be impacted onshore, at range limit locations. A myriad of factors may be influencing larval settlement and survival of *T. rosea* at its range limit locations, most notably the decline in air and sea temperature (Knox 1963, Ridgway & Dunn 2003, Lathlean et al. 2011), or the presence of potentially competitive or predatory species that do not inhabit shores further to the north (Hidas et al. 2007, Coulson et al. 2011).

Lathlean et al. (2010) did not find any significant differences in percent settlement, early post-settlement mortality, recruitment, post-recruitment mortality or adult mortality among sites between Garie Beach and Mallacoota over two years of rigorous sampling, but their sampling did not extend as far south as the range limit at Cape Conran (Fig. 1).

Our hierarchical sampling collectively suggests that: (1) there is considerable variation in the demography, and relationships among life-cycle stages of *Tesseropora rosea* at locations towards its range limit, (2) nevertheless, a declining trend in adult abundance, and a positive relationship between adult abundance and recruitment is evident towards the range limit and therefore, (3) the decline in adult abundance is likely due to one or a combination of range limit factors impacting on the recruitment stage of the life cycle. In the case of *T. rosea*, large-scale processes therefore appear to override the complexities of smaller-scale variation in recruitment towards the range limit (Fowler-Walker et al 2005). These conclusions are, however, based on a relatively short time-series of data, and recruitment for this species can vary greatly among years (e.g. Caffey 1985), indicating

that strong recruitment years could result in different outcomes. The system therefore deserves comparisons of pattern data over a multitude of years, or even decades (Lewis et al. 1986, Mieszkowska 2007) before it can be well understood, particularly at a time when the impacts of climate change on species' range shifts are increasingly evident (Sagarin et al. 1999, Helmuth et al. 2006, Hawkins et al. 2008, Wethey et al. 2011). Nevertheless, our results clearly highlight the benefits of investigating processes accounting for spatial variation across a range of scales towards the range limit, to gain greater insight into the factors that set range limits. Genetic data, or the tracking of larvae, may provide the key to identifying the source of recruits and the strength of larval connections between populations within the interior of the range and the range edge. Furthermore, careful assessment of the relationship between the abundance of settlers and consequent recruits (Connell 1985, Minchinton & Scheibling 1991), may reveal important information about how post-settlement factors and reproductive potential are affected towards the range limit.

ACKNOWLEDGEMENTS

Funding was provided via an Australian Research Council Discovery grant to DJA and TEM, the University of Wollongong's Institute for Conservation Biology and an Australian Postgraduate Award to EZH. We thank several students and volunteers from the University of Wollongong, particularly Justin Lathlean, for assisting with field- and lab-work. Research was carried out under permits from NSW DPI and Parks Victoria.

LITERATURE CITED

- Ayre DJ, Hughes TP (2000) Genotypic diversity and gene flow in brooding and spawning corals along the Great Barrier Reef, Australia. *Evolution* 54:1590-1605
- Ayre DJ, Minchinton TE, Perrin C (2009) Does life history predict past and current connectivity for rocky intertidal invertebrates across a marine biogeographic barrier? *Mol Ecol* 18:1887-1903
- Broitman BR, Blanchette CA, Menge BA, Lubchenco J, Krenz C, Foley M, Raimondi PT, Lohse D, Gaines SD (2008) Spatial and temporal patterns of invertebrate recruitment along the west coast of the United States. *Ecol Monogr* 78:403-421
- Burrows MT, Hawkins SJ, Southward AJ (1992) A comparison of reproduction in co-occurring chthamalid barnacles, *Chthamalus stellatus* (Poli) and *Chthamalus montagui* Southward. *J Exp Mar Biol Ecol* 160:229-249
- Caffey HM (1985) Spatial and temporal variation in settlement and recruitment of intertidal barnacles. *Ecol Monogr* 55:313-332
- Connell JH (1985) The consequence of variation in initial settlement vs. post-settlement mortality in rocky intertidal communities. *J Exp Mar Biol Ecol* 93:11-45
- Coulson LA, Perrin C, Roberts DG, Minchinton TE, Ayre DJ (2011) Can limited dispersal or biotic interaction explain the declining abundance of the whelk, *Morula marginalba*, at the edge of its range? *Biol J Linn Soc* 103:849-862
- Denley EJ, Underwood AJ (1979) Experiments on factors influencing settlement, survival and growth of two species of barnacles in New South Wales. *J Exp Mar Biol Ecol* 36:269-293

- Edgar GJ (1997) Australian marine life: The plants and animals of temperate waters.
Reed New Holland, Sydney, Australia
- Egan EA, Anderson DT (1988) Larval development of the coronuloid barnacles
Austrabalanus imperator (Darwin), *Tetraclitella purpureascens* (Wood), and
Tesseropora rosea (Krauss) (Cirripedia, Tetraclitidae). J Nat Hist 22:1379-1406
- Fowler-Walker MJ, Connell SD, Gillanders BM (2005) Variation at local scales need not
impede tests for broader scale patterns. Mar Biol 147:823-831
- Gaines S, Roughgarden J (1985) Larval settlement rate: A leading determinant of
structure in an ecological community of the marine intertidal zone. Proc Natl Acad
Sci U.S.A. 82:3707-3711
- Gaylord B, Gaines SD (2000) Temperature or transport? Range limits in marine species
mediated solely by flow. Am Nat 155:769-789
- Gilman SE (2005) A test of Brown's principle in the intertidal limpet *Colisella scabra*
(Gould, 1846). J Biogeogr 32:1583-1589
- Grinnell J (1917) Field tests of theories concerning distribution control. Am Nat 51:115-
128
- Hawkins SJ, Moore PJ, Burrows MT, Poloczanska E, Mieszkowska N, Herbert RJH,
Jenkins SR, Thompson RC, Genner MJ, Southward AJ (2008) Complex
interactions in a rapidly changing world: responses of rocky shore communities to
recent climate change. Clim Res 37:123-133

Helmuth B, Harley CDG, Halpin PM, O'Donnell M, Hofmann GE, Blanchette CA (2002)

Climate change and latitudinal patterns of intertidal thermal stress. *Science*
298:1015-1017

Helmuth B, Mieszkowska N, Moore P, Hawkins SJ (2006) Living on the edge of two
changing worlds: forecasting the responses of rocky intertidal ecosystems to
climate change. *Annu Rev Ecol Evol Syst* 37:373-404

Herbert RJH, Southward AJ, Clarke RT, Sheader M, Hawkins SJ (2009) Persistent
border: an analysis of the geographic boundary of an intertidal species. *Mar Ecol
Prog Ser* 379:135-150.

Hidas EZ, Costa TL, Ayre DJ, Minchinton TE (2007) Is the species composition of rocky
intertidal invertebrates across a biogeographic barrier in southeast Australia related
to their potential for dispersal? *J Mar Fresh Res* 58:835-842

Hidas EZ, Ayre DJ, Minchinton TE (2010) Patterns of demography for rocky-shore,
intertidal invertebrates approaching their range limits: tests of the abundant-
hypothesis in south-eastern Australia. *J Mar Fresh Res* 61:1243-1251.

Hughes TP (1990) Recruitment limitation, mortality, and population regulation in open
systems: a case study. *Ecology* 71:12-20

Hughes TP, Baird AH, Dinsdale EA, Moltschaniwskyj NA, Pratchett MS, Tanner JE,
Williams BL (2000) Supply-side ecology works both ways: the link between
benthic adults, fecundity, and larval recruits. *Ecology* 81:2241-2249

- Hutchins LW (1947) The bases for temperature zonation in geographical distribution.
Ecol Monogr 17:325-335
- IBM Corp. (2011) IBM SPSS Statistics for Windows, Version 20.0. Armonk, New York
- Kent A, Hawkins SJ, Doncaster CP (2003) Population consequences of mutual attraction between settling and adult barnacles. J Anim Ecol 72:941-952
- Kinlan BP, Gaines SD (2003) Propagule dispersal in marine and terrestrial environments: a community perspective. Ecology 84:2007-2020
- Knox GA (1963) The biogeography and intertidal ecology of the Australasian coasts.
Oceanogr Mar Biol Annu Rev 1:341-404
- Lathlean JA, Ayre DJ, Minchinton TE (2010) Supply-side biogeography: geographic patterns of settlement and early-mortality for a barnacle approaching its range limit.
Mar Ecol Prog Ser 412:141-150
- Lathlean JA, Ayre DJ, Minchinton TE (2011) Rocky intertidal temperature variability along the southeast coast of Australia and comparisons with satellite-derived SST and terrestrial weather stations. Mar Ecol Prog Ser 439: 83–95
- Lewis JR (1986) Latitudinal trends in reproduction, recruitment and population characteristics of some littoral molluscs and cirripedes. Hydrobiologia 142:1-13
- Lima FP, Queiroz N, Ribeiro PA, Hawkins SJ, Santos AM (2006) Recent changes in the distribution of a marine gastropod, *Patella rustica* Linnaeus, 1758, and their relationship to unusual climatic events. J Biogeogr 33:812-822

- Mieszkowska N, Hawkins SJ, Burrows MT, Kendall MA (2007) Long term changes in the geographic distribution and population structures of *Osilinus lineatus* (Gastropoda: Trochidae) in Britain and Ireland. *J Mar Biol Ass. U.K.* 87:537-545
- Minchinton TE, Scheibling RE (1991) The influence of larval supply and settlement on the population structure of barnacles. *Ecology* 72:1867-1879
- Murray-Jones SE, Ayre DJ (1997) High levels of gene flow in the surf bivalve *Donax deltoides* (Bivalvia: Donacidae) on the east coast of Australia. *Mar Biol* 128:83-89
- Patterson HD, Thompson R (1971) Recovery of inter-block information when block sizes are unequal. *Biometrika* 58:545-554
- Pechenik JA (1999) On the advantages and disadvantages of larval stages in benthic marine invertebrate life cycles. *Mar Ecol Prog Ser* 177:269-297
- Ramirez-Llorda E (2002) Fecundity and life-history strategies in marine invertebrates. *Adv Mar Biol* 43:87-170
- Ridgway KR, Dunn JR (2003) Mesoscale structure of the East Australian Current system and its relationship with topography. *Prog Oceanogr* 56:189-222
- Sagarin RD, Barry JP, Gilman SE, Baxter CH (1999) Climate related change in an intertidal community over short and long time scales. *Ecol Monogr* 69:465-490
- Sagarin RD, Gaines SD (2002a) The 'abundant centre' distribution: to what extent is it a biogeographical rule? *Ecol Lett* 5:137-147

- Sagarin RD, Gaines SD (2002b) Geographical abundance distributions of coastal invertebrates: using one-dimensional ranges to test biogeographic hypotheses. *J Biogeogr* 29:985-997
- Sagarin RD, Gaines SD, Gaylord B (2006) Moving beyond assumptions to understand abundance distributions across the ranges of species. *Trends Ecol Evol* 21:524-530
- VSN International (2011) *GenStat for Windows* 14th Edition. Hemel Hempstead, UK
- Wethey DS, Woodin SA, Hilbish TJ, Jones SJ, Lima FP, Brannock PM (2011) Responses of intertidal populations to climate: Effects of extreme events versus long term change. *J Exp Mar Biol Ecol* 400:132-144
- Wisely B, Blick RAP (1964) Seasonal abundance of first stage nauplii in 10 species of barnacles at Sydney. *J Mar Fresh Res* 15:162-171
- Zacherl D, Gaines SD, Lonhart SI (2003) The limits to biogeographical distributions: insights from the northern range extension of the marine snail, *Kelletia kelletii* (Forbes, 1852). *J Biogeogr* 30:913-924

TABLES

Table 1 Restricted Maximum Likelihood (REML) analysis of the adult and recruit densities of *Tesseropora rosea* among four fixed sectors and 16 nested, random locations on the southeast coast of Australia. REML is preferred to ANOVA for a mixed model analysis with unbalanced data (i.e. some missing data). Time has been added as a fixed blocking term. The denominator variances in the F-tests do not necessarily have a whole number for the degrees of freedom, because they have been estimated by approximate methods. The significance of the nested random factor is indicated by the *P*-value associated with the Wald Z.

VARIABLE	FIXED EFFECTS					COVARIANCES OF RANDOM EFFECTS				
	Source	Num. d.f.	Den. d.f.	F	P	Parameter	Estimate	Std. Error	Wald Z	P
ADULT DENSITY	Sector	3	11.684	5.474	0.014	Location(Sector)	0.628	0.282	2.228	0.026
	Time	3	113.062	2.945	0.036	Site(Location(Sector))	0.037	0.047	0.782	0.434
						Residual	0.406	0.054	7.464	<0.001
RECRUIT DENSITY	Sector	3	11.611	4.216	0.031	Location(Sector)	1.101	0.502	2.194	0.028
	Time	3	124.794	4.203	0.007	Site(Location(Sector))	0.061	0.076	0.799	0.425
						Residual	0.872	0.111	7.850	<0.001

Table 2 Predicted probabilities from a generalised linear mixed model analysis, for encountering sexually mature adult *Tesseropora rosea* (i.e. possessing gonads) at each of four sectors (1=furthest from range limit, 4=nearest to the range limit) for each of four sampling times (1=May 2005, 2=September 2005, 3=January 2006, 4=May 2006), along the southern third of the species' distributional range on the southeast coast of Australia.

		Sector			
		1	2	3	4
Time	1	0.9704	0.9273	0.9301	0.9825
	2	1.0000	1.0000	0.9935	0.9479
	3	0.9821	0.9405	0.8767	0.9066
	4	0.8974	0.8605	0.8812	0.9052

Table 3 Predicted probabilities from a generalised linear mixed model analysis, for encountering fecund adult *Tesseropora rosea* (i.e. possessing mature larvae) at each of four sectors (1=furthest from range limit, 4=nearest to the range limit) for each of four sampling times (1=May 2005, 2=September 2005, 3=January 2006, 4=May 2006), along the southern third of the species' distributional range on the southeast coast of Australia.

		Sector			
		1	2	3	4
Time	1	0.0966	0.1683	0.0641	0.0595
	2	0.0507	0.0272	0.3908	0.0540
	3	0.2262	0.2044	0.0975	0.1539
	4	0.2433	0.3038	0.4964	0.0108

FIGURE LEGENDS

Fig. 1 Map of the southeast coast of Australia showing the four sectors and corresponding 16 locations used to assess the adult abundance, fecundity, sexual maturity, and recruitment of *Tesseropora rosea* along the southern third of its distributional range, from Garie Beach near Sydney, New South Wales to Cape Conran, Victoria. Inset map shows Australia with its major states and territories (QLD = Queensland, NSW = New South Wales, VIC = Victoria, TAS = Tasmania, SA = South Australia, WA = Western Australia, NT = Northern Territory), and the bracket identifies the location of the larger map of this figure. Flow patterns of the southward flowing East Australian Current (EAC) and eastward flowing Zeehan Current (ZC) are also shown.

Fig. 2 Mean (+SE) adult (a & e) and recruit densities (b & f), and percentages of adults with gonads (c & g) and mature larvae (d & h) of *Tesseropora rosea* across four sectors and corresponding 16 locations, along the southern third of its distributional range from Garie Beach near Sydney, New South Wales to Cape Conran, Victoria, on the southeast coast of Australia, between May 2005 and May 2006. Within each graph, sectors marked by different letters are significantly different from one another.

Fig. 3 The relationships between (a) mean adult and recruit densities, (b) mean adult densities and the mean proportions of adults with mature larvae, and (c) the mean proportions of adults with mature larvae and mean recruit densities of *Tesseropora rosea* at 14-16 locations (depending on data availability) along 600 km of coastline constituting the southern third of the distributional range of this species on the southeast coast of Australia; and the relationships between adult and recruit densities at 12 sites within each

of the sectors 1-4 (d-g respectively) encompassing 70-100 km within this same coastline, and representing areas furthest to closest to the range limit, respectively. Note the different variables and scales represented on the x- and y-axes. A linear line of best fit highlights significant relationships.

FIGURES




Fig 1.




Fig 2.




Fig. 3