

2011

A systematic review of the validity and reliability of sedentary behaviour measures used with children and adolescents

D R. Lubans
University of Newcastle

K Hesketh
Deakin University

D P. Cliff
University of Wollongong, dylanc@uow.edu.au

L M. Barnett
Deakin University

J Salmon
Deakin University

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/edupapers>



Part of the [Education Commons](#)

Recommended Citation

Lubans, D R.; Hesketh, K; Cliff, D P.; Barnett, L M.; Salmon, J; Dollman, J; Morgan, P J.; Hills, A P.; and Hardy, L L.: A systematic review of the validity and reliability of sedentary behaviour measures used with children and adolescents 2011.
<https://ro.uow.edu.au/edupapers/1206>

Authors

D R. Lubans, K Hesketh, D P. Cliff, L M. Barnett, J Salmon, J Dollman, P J. Morgan, A P. Hills, and L L. Hardy

1 1. Title: A Systematic Review of the Validity and Reliability of Sedentary Behaviour
2 Measures used with Children and Adolescents

3
4
5
6

7 2. Authors: David R. Lubans¹, Kylie Hesketh², Dylan P. Cliff³, Lisa M. Barnett⁴, Jo
8 Salmon², Jim Dollman⁵, Philip J. Morgan¹, Andrew P. Hills⁶, and Louise L. Hardy⁷

9
10 3. Authors' addresses

11
12

13 ¹School of Education, University of Newcastle, Australia;

14 ²Centre for Physical Activity and Nutrition Research, Deakin University, Australia;

15 ³Faculty of Education, University of Wollongong, Australia;

16 ⁴School of Health & Social Development, Deakin University, Australia;

17 ⁵School of Health Sciences, University of South Australia, Australia;

18 ⁶School of Human Movement Studies, Queensland University of Technology, Australia;

19 ⁷NSW Centre for Overweight and Obesity, University of Sydney, Australia.

20

21 4. Key words: Measurement; validity; reliability

22

23 5. Running title: Measurement of Sedentary Behaviour in Youth

24

25 6. Acknowledgements: Funding for this review was provided by the Australasian Child
26 and Adolescent Obesity Research Network. Dylan Cliff is funded by a National
27 Heart Foundation of Australia- Macquarie Postdoctoral Research Fellowship.

28

29 7. *Corresponding author

30 David Lubans

31 University of Newcastle

32 School of Education

33 Callaghan Campus

34 NSW 2308

35 Australia

36 Email: David.Lubans@newcastle.edu.au

37 Telephone: +61 2 49212049

38 Fax: +61 2 49217407

39

40 8. Conflict of interest: The authors have no conflicts of interest to declare.

1 **Abstract**

2 The aim of this review was to evaluate the reliability and validity of methods used to
3 assess the multiple components of sedentary behaviour (i.e. screen time, sitting, not
4 moving and existing at low energy expenditure) in children and adolescents. Twenty-six
5 studies met our inclusion criteria and were reviewed. Thirteen studies reported the
6 reliability of self- and proxy-report measures of sedentary behaviour and seven of these
7 were found to have acceptable test-retest reliability. Evidence for the criterion validity of
8 self- and proxy-report measures was examined in three studies with mixed results. Seven
9 studies examined the reliability and/or validity of **direct** observation and the findings
10 were generally positive. Five studies demonstrated the utility of accelerometers to
11 accurately classify sedentary behaviour. Self-report measures provide reliable estimates
12 of screen-time, yet their validity remains largely untested. While accelerometers can
13 accurately classify participants' behaviour as sedentary, they do not provide information
14 about type of sedentary behaviour or context. Studies utilising measures of sedentary
15 behaviour need to more adequately report on the validity and reliability of the measures
16 used. We recommend the use of objective measures of sedentary behaviour such as
17 accelerometers, in conjunction with subjective measures (e.g. self-report) to assess type
18 and context of behaviour.

19

20

1 **Introduction**

2 The prevalence of paediatric obesity has become a major public health issue (1).
3 In addition to poor dietary patterns, reductions in physical activity and increased time
4 spent sedentary have been highlighted as the major contributors to the epidemic (2).
5 While much of the focus of obesity prevention and treatment has centred on the
6 promotion of physical activity, interventions targeting time spent in sedentary behaviour,
7 screen time in particular have demonstrated promise (3-5). The term sedentary behaviour
8 may be defined as minimal energy expenditure (1 to 1.5 metabolic equivalent multiples
9 of rest) that typically involves sitting or lying down (6). Time spent in sedentary
10 behaviour is distinct from lack of physical activity as these are considered unique
11 behavioural constructs that have independent relationships to various health outcomes
12 (7). Although time spent watching television has typically been the focus of sedentary
13 behaviour studies (8), other domain-specific sitting behaviours such as using the
14 computer, playing electronic games, reading, talking on the telephone and travelling by
15 bus, car, or train also contribute to young people's sedentary time. Notably, national
16 guidelines in many countries have included recommendations to minimise sedentary
17 behaviour including limiting the amount of time spent using screen-based recreation
18 pursuits to less than two hours per day (9, 10).

19 Time spent in sedentary behaviour among children and adolescents has been
20 linked positively to overweight and obesity and other adverse health outcomes in both
21 cross-sectional (11-13) and longitudinal studies (14, 15). In a recent large-scale 4-year
22 longitudinal study, higher levels of baseline self-reported TV viewing were positively
23 associated with a steeper body mass index (BMI) trajectory among U.S. adolescent girls

1 (15). A cross-sectional study of Portuguese children who participated in the European
2 Youth Heart Study found even after adjusting for sex, birth weight, pubertal status, and
3 total or central fat mass, there were positive associations between objectively-assessed
4 time spent sedentary (defined as <500 accelerometer counts per minute) and insulin
5 resistance (16). A further cross-sectional study that included more than 5,000 12-year old
6 children in the UK used accelerometers to assess sedentary time (defined as <200 counts
7 per minute) and found that for every hour spent sedentary per day, after adjusting for sex,
8 social factors, sleep, television viewing time and pubertal status, children were 32% more
9 likely to be obese (17). However, this association was attenuated when physical activity
10 was included in the model. Inconsistencies in study findings may be attributed to varying
11 definitions of sedentary time from accelerometry data. Given the increasing evidence
12 base on the adverse health consequences of time spent in sedentary behaviour, the valid
13 and reliable assessment of sedentary behaviour is an important public health priority and
14 a key issue for future research. Quality instruments for assessing sedentary behaviour
15 with known measurement psychometric properties are vital for understanding dose-
16 response relationships between sedentary behaviour and health and developmental
17 outcomes, for population health monitoring, for determining the correlates and predictors
18 of sedentary behaviour, and for determining the impact of health interventions targeting
19 reductions in sedentary time. While methodological issues relating to the assessment of
20 physical activity among children and adolescents have been explored in numerous
21 reviews (18-21), issues pertaining to the assessment of sedentary behaviour have received
22 little attention. Bryant and colleagues (22) published a systematic review of studies that
23 had included a measure of television exposure in children and adolescents. The authors

1 found a large number of studies that had used self-report measures and noted that the
2 validity and reliability of commonly used measures were rarely provided. While their
3 review provided important recommendations for assessing television viewing exposure in
4 youth, it did not explore the multiple components of sedentary behaviour (e.g. time spent
5 playing electronic games and computers, sitting time) and it did not include objective
6 measures of sedentary behaviour (e.g. accelerometers and direct observation). The
7 importance of assessing the multiple components of sedentary behaviour has been
8 highlighted in the literature (8). No previous review has evaluated the reliability and
9 validity of objective and subjective methods used to assess the multiple components of
10 sedentary behaviour. The primary aim of this review was to evaluate the reliability and
11 validity of methods used to assess the multiple components of sedentary behaviour in
12 children and adolescents (i.e. screen time, sitting, not moving and existing at low energy
13 expenditure) by systematically reviewing the existing literature.

14 **Methods**

15 *Identification of studies*

16 A systematic review of studies reporting validity and/or reliability of methods
17 used to assess the multiple components of sedentary behaviour (i.e. screen time, sitting,
18 not moving and existing at low energy expenditure) in youth was guided by the Preferred
19 Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) statement (23)
20 and was conducted in four phases. Firstly, we conducted a systematic search of published
21 literature using electronic databases (described in detail below). In the second phase we
22 conducted an internet-based search and search of authors' personal collections for
23 published literature examining measures of sedentary behaviour among children and

1 adolescents (aged 3-18 years). Articles were then hand-searched to identify key
2 researchers and programs of work examining sedentary behaviour in the target age group.
3 The third phase involved contacting key authors or research groups to identify measures
4 of sedentary behaviour they had used, or were aware of, with this age group. The fourth
5 phase was to identify any further articles from reference lists of retrieved articles.

6 Databases were searched from 1985 until the most recent published articles
7 (including in-press articles) as at May 2010. The databases searched included: Academic
8 Search Premier; CINAHL; Cochrane Central Register of Controlled Trials; Cochrane
9 Database of Systematic Reviews; Global Health; Health Source: Nursing / Academic;
10 MedLINE (PubMed); Psycharticles; Psychology and Behavioural Sciences Collection;
11 PsychInfo; SportsDiscus. Individualised search strategies for the different databases
12 focused on 1) behaviours, including combinations of the following key words:
13 ‘sedentar*’, ‘sitting’, ‘*screen’, ‘television’, ‘computer’, ‘electronic games’, ‘video’,
14 ‘DVD’, ‘video games’ and ‘electronic media’, in conjunction with 2) measurement
15 related words including: ‘instrument’, ‘survey’, ‘log’, ‘diary’, ‘questionnaire’, ‘self-
16 report’, ‘proxy report’, ‘accelerom*’, ‘inclinom*’, ‘actigraph’, ‘motion sensor’, ‘heart
17 rate’, ‘measure*’ or ‘assess*’ or ‘observ*’. When the database did not allow age limiters to be
18 set, words related to childhood and adolescence (i.e. ‘child*’, ‘adolescent’, ‘young
19 people’, ‘youth’) were also included. The keyword search was limited to words appearing
20 in the title and abstract.

21 *Criteria for inclusion/exclusion*

22 Two of the authors (JD and AH) independently assessed the eligibility of the
23 studies for inclusion according to the following criteria: i) child and adolescent

1 participants (aged 3-18 years); ii) direct observation (including video); self- or proxy-
2 report, or objective measure of sedentary behaviour; iii) validity and/or reliability of a
3 sedentary behaviour measurement tool reported; iv) published or in-press in a peer-
4 reviewed journal; and vi) published in English. Articles were only included if the
5 reliability and/or validity of the instruments' sedentary behaviour component were
6 analysed and reported separately. Reviews, positions statements, case studies, abstracts
7 and editorials were not included in the review. Articles that only included children or
8 adolescents with disabilities or developmental delays that may impact their ability to
9 accurately recall sedentary behaviour were excluded.

10 *Reliability of sedentary behaviour measures*

11 Reliability refers to the consistency of a response either across multiple tests
12 within a single assessment, generally called internal consistency, or across multiple
13 assessments, known as test-retest or stability reliability (18). In addition, inter-rater
14 reliability refers to the stability of observations between two or more testers measuring
15 the same behaviour (agreement between raters), while intra-rater reliability refers to the
16 consistency of observations made by the same observer on different days. Two authors
17 (LB and DRL) independently assessed the reported reliability of the sedentary behaviour
18 measures using a modified version of the checklist developed for assessing the qualitative
19 attributes and measurement properties of physical activity questionnaires (QAPAQ) (21,
20 24). Reliability was rated as acceptable, borderline, unacceptable, or indeterminate if it
21 was not possible to assess using the criteria provided. Intraclass correlation coefficient
22 (ICC) is the preferred method for estimating test-retest reliability (19) or Kappa for
23 dichotomous data or weighted Kappa for ordinal data (24). An ICC or Kappa of above

1 0.70 is considered acceptable (Pearson's correlation or Spearman's rank of > 0.80 was
2 also considered to be acceptable) (25). Borderline was reserved for an ICC or Kappa
3 between .60 - .69 (Pearson correlation or Spearman's rank > 0.70 was also considered to
4 be borderline). Intra-rater and inter-rater reliability of direct observation can be assessed
5 using ICC or Kappa and values above 0.70 were considered acceptable.

6 *Validity of sedentary behaviour measures*

7 Validity is the extent to which a method measures what it claims to measure (21).
8 There are numerous types of validity (i.e., criterion, concurrent, and content) relevant to
9 sedentary behaviour measurement. Criterion validity refers to the relationship between
10 results of the measure being assessed and the recognised measure or 'gold standard'(21).
11 Studies assessing the validity of physical activity questionnaires often use accelerometers
12 and direct observation as their criterion measures. Similarly, for the current review,
13 accelerometers and direct observation were considered to provide evidence of criterion
14 validity. Concurrent validity is the extent to which results are associated with those of
15 other existing measures (e.g. comparing results from a new sedentary behaviour
16 questionnaire to those from an existing measure). While comparing one method of
17 unknown validity against another method of unknown validity does not provide evidence
18 of criterion validity, agreement between measures indicates concurrent validity. Content
19 validity refers to the degree to which the content of an instrument adequately reflects all
20 aspects of the outcome of interest. As there is a lack of consensus on how high
21 correlations should be to demonstrate adequate criterion or concurrent validity (26),
22 classifications for direct observation, self- and proxy-report measures were not provided.

23 As reported in the Introduction section, accelerometry has been used to

1 objectively assess free-living sedentary behaviour among children and adolescents. To
2 utilise accelerometry for this purpose requires consistent cut-point definitions to be
3 applied to the data to categorise the counts accumulated by the device each epoch into
4 either sedentary behaviour or physical activity. Validity data for published cut-points
5 corresponding to different accelerometer models were reviewed. Results for sensitivity
6 (true positive rate), specificity (false positive rate), and area under the receiver operating
7 characteristic (ROC) curve (false-positive rate (1 – specificity) versus true-positive rate)
8 were extracted and reported. For area under the ROC curve analysis, an area of 1
9 indicates perfect classification accuracy, while an area of 0.5 represents a complete
10 absence of classification accuracy. Values of > 0.90 were rated as excellent, 0.80–0.90
11 good, 0.70–0.80 fair, and < 0.70 poor (27).

12 **Results**

13 *Study selection*

14 The initial search of 11 databases located 2862 potential articles. Of these studies
15 2813 were excluded based on titles and abstracts and 49 full-text articles were retrieved.
16 Further studies were located in the reference lists of these articles and additional studies
17 known to the authors were considered for inclusion. A review of the full content of the
18 papers reduced the number of studies to 26 that met the inclusion criteria (Figure 1).

19 *Sedentary behaviour measures and method of measurement*

20 Thirteen studies reported the reliability of self-report or proxy-report measures of
21 sedentary behaviour in children and adolescents (28-40). Time spent watching TV was
22 the most frequently measured sedentary behaviour. However, more recent self-report
23 measures of sedentary behaviour often included computer use and time playing electronic

1 games. Proxy-report measures of sedentary behaviour were used in four studies with
2 younger children (29, 30, 33, 36), while self-report measures were used in all of the
3 adolescent studies (> 12 years). Of both the proxy- and self-report measures, three (28,
4 35, 41) required participants to report their time in sedentary behaviour from the
5 previous week, but the majority of measures required participants or parents to report
6 usual weekday and weekend sedentary behaviour .

7 Three (29, 42, 43) and five (43-47) studies reported the inter-observer reliability
8 and concurrent validity of direct observation techniques, respectively. Five measures of
9 direct observation were designed to provide an assessment of time in physical activity,
10 but also reported the validity or reliability for time in sedentary behaviour (42, 44-46,
11 48). One study reported the inter-observer reliability for home observations using time-
12 lapse cameras (29). Five studies examined the validity of accelerometers for measuring
13 sedentary behaviour (49-53).

14 *Reliability of self- and proxy-report measures*

15 The reliability of self- and proxy-report measures of sedentary behaviour are
16 outlined in Table 1. Reliability was assessed using ICCs (28, 33, 34, 36-39) in most
17 studies, but bivariate correlation (29, 30), Kappa (31, 32), percent agreement (32) and
18 Spearman rank order correlations (32, 34, 40) were also used. Periods between test and
19 retest were generally one to two weeks. However, Anderson and colleagues evaluated the
20 1-month test-retest reliability for a 10-day TV viewing diary. Seven measures were found
21 to have acceptable test-retest reliability for specific components of sedentary behaviour
22 (28, 30, 34, 36, 37, 39, 40) and two measures were classified as borderline (29, 33). In
23 general, reliability was better for TV viewing than it was for computer use and playing

1 electronic games. The reliability of weekday sedentary behaviour was generally higher
2 than weekend sedentary behaviour. There were no obvious differences in the reliability of
3 sedentary behaviour measures for boys and girls.

4 *Criterion validity of self- and proxy-report measures*

5 Three studies examined the criterion validity of a self- or proxy-report measure of
6 sedentary behaviour by comparing the results to direct observation (29) or accelerometry
7 (41, 54) (Table 2). Hardy et al. (41), reported the mean weekly difference between self-
8 reported sedentary behaviour using the Adolescent Sedentary Activity Questionnaire
9 (ASAQ) and accelerometer estimates of sedentary behaviour. While this method of
10 assessing validity was not addressed in our predetermined criteria, the results indicate
11 that this measure has acceptable validity (less than 5% of data outside the limits of
12 agreement). Similarly, Wen and colleagues (54) examined the relationship between
13 sedentary behaviour using an accelerometer and proxy-reported sedentary behaviour and
14 found a positive correlation.

15 *Concurrent validity of self- and proxy-report measures*

16 Three studies compared the results from self-report measures with diary entries in
17 children and adolescents (28, 32, 34). One study examined the relationship between
18 child- and parent-reported sedentary behaviour (33) and another study reported a content
19 validity index. Four self-report measures were found to have correlation coefficients \geq
20 0.30 (28, 32-34). Liou and colleagues (38) reported a content validity index of 0.99, but
21 did not explain how this score was achieved and we were therefore, unable to classify the
22 validity of their self-report measure. Salmon and colleagues examined the concurrent

1 validity of self-report and proxy-report sedentary behaviour in youth (33) and found the
2 strongest association for TV viewing.

3 *Reliability and validity of direct observation*

4 Seven studies examined the psychometric properties of direct observation tools
5 for assessing sedentary behaviour at home (29, 42), in community settings (43, 44),
6 during physical education lessons (45, 48), or during breaks at school (46). Six studies
7 reported reliability results (29, 43, 45, 46, 48) and four studies provided validity data (43-
8 45, 48) (Table 4). Anderson and colleagues (29) used video-recordings to observe
9 children's time spent in the room with TV and their time spent directing their visual
10 attention towards the TV. Inter-observer reliabilities between ratings by two assessors
11 were 0.98 and 0.90 for presence in the viewing room and visual attention towards the TV,
12 respectively. DuRant et al.(42), used direct observation to assess children's time spent
13 television viewing by coding each minute throughout the day and also reported high
14 inter-observer reliability (96% agreement). The validity and inter-observer reliability of
15 the Children's Activity Rating Scale (CARS) was examined among young children (43) .
16 Percent VO₂ max and heart rate were found to differ between CARS category 1,
17 representing sedentary behaviour (stationary – no movement, e.g. lying and sitting), and
18 category 2 (stationary – with movement, e.g., standing and colouring). Inter-observer
19 agreement from 389 paired observation periods by 11 observers over 12 months was
20 84.1%. McKenzie et al (44) tested the validity of the Behaviours of Eating and Activity
21 for Children's Health Evaluation Systems (BEACHES) instrument using heart rate
22 monitoring. The authors found a linear relationship between heart rate and intensity of

1 activity, with the lowest average heart rate associated with lying down (99 beats/minute)
2 and the highest heart rate associated with 'very active' time (153 beats/minute).

3 Rowe and colleagues (45) tested the validity and reliability of the System for
4 Observing Fitness Instruction Time (SOFIT) categories (lying, sitting, standing, walking,
5 running) among students in 1st through to 8th grade during a structured activity protocol in
6 their physical education classes using heart rate monitoring. Heart rates during sedentary
7 behaviours (sitting and lying) differed from standing and walking, and heart rates during
8 sedentary behaviours had high internal consistency reliabilities ($r > 0.99$). Among 9th to
9 12th grade students, Rowe et al. (47), tested the validity and reliability of the SOFIT
10 categories against both heart rate and energy expenditure measured by indirect
11 calorimetry. Although heart rates differed for sedentary behaviours (sitting and lying)
12 compared with standing and walking, energy expenditure did not differ between lying,
13 sitting and standing, but did differ between those categories and walking. Internal
14 consistency reliabilities for sedentary categories were higher for heart rate ($r \geq 0.98$)
15 compared with energy expenditure ($r = 0.78-0.82$).

16 *Objective measures of sedentary behaviours*

17 Five studies examined the criterion validity of accelerometers for measuring
18 sedentary behaviour in youth by comparing accelerometer cut-points with direct
19 observation (49, 52), metabolic units (50, 53), calorimeter and heart rate telemetry (51).
20 Of the four studies examining cut-points for the Actigraph, three reported excellent
21 validity (50, 52, 53). Reilly and colleagues (49) developed and validated a sedentary
22 behaviour cut-point for the Actigraph against direct observation among 3- to 4-year-olds.
23 They found that a definition of <1100 counts/min provided optimal sensitivity (83%) and

1 specificity (82%) for young children's sedentary time. Similarly, Sirard et al. (52),
2 developed age-specific sedentary behaviour cut-points for the Actigraph using direct
3 observation. Sensitivity and specificity were high for all ages (92%-100%) and optimised
4 at <1204, <1452, and <1592 counts/min for 3-, 4-, and 5-year-olds, respectively. Treuth
5 and colleagues (50) developed cut-point definitions for the Actigraph among 13- to 14-
6 year-old adolescent girls using VO^2 measured by a portable indirect calorimetry system.
7 For sedentary behaviour (<1.5 METs), sensitivity (100%) and specificity (100%) were
8 optimised at <100 counts/min. Evenson and colleagues (53) also found that this cut-point
9 optimised sensitivity (95%) and specificity (93%) among 5- to 8-year-olds, where
10 portable indirect calorimetry was used to measure oxygen consumption.

11 Sedentary behaviour cut-points for the Actical and Actiwatch accelerometers have
12 been validated among children and adolescents, with one study reporting excellent
13 classification accuracy among children (53) and another reporting good classification
14 accuracy among children and adolescents (51). Evenson and colleagues (53) found that
15 sensitivity (97%) and specificity (98%) were optimised at < 44 counts/min for the Actical
16 among 5- to 8-year-olds. Puyau and colleagues (51) used calorimetry to determine cut-
17 points for sedentary behaviour (activity energy expenditure <0.01 kcal/kg/min), and
18 found that <100 and <50 counts/min provided good classification accuracy among 7- to
19 18-year-olds for the Actical and Actiwatch, respectively (area under ROC curve: Actical
20 = 0.85, Actiwatch = 0.85).

21 **Discussion**

22 This systematic review identified studies that reported on the reliability and/or
23 validity of measures of sedentary behaviour used in children and adolescents 0-18 years

1 of age. Despite the wide use of sedentary behaviour measurement tools in studies
2 involving children and adolescents, few studies report the reliability and validity of the
3 measures used. Further, the methods of assessing reliability and validity varied between
4 studies, making cross-study comparisons difficult. It is of additional concern that many
5 studies compared one method of unknown validity against another measure of unknown
6 validity to establish concurrent validity. While the varying utility of the measures
7 prohibits blanket recommendations for all study types, the results presented here provide
8 useful comparisons for researchers designing new studies and selecting measurement
9 tools.

10 Despite only being assessed in five studies, accelerometers appear to provide a
11 valid measure of sedentary behaviour. When assessed against direct observation,
12 metabolic monitoring and energy expenditure via calorimetry, accelerometers achieved
13 greater than 80% sensitivity and specificity. In two of the four studies, perfect (100%)
14 sensitivity and specificity were reported. Given the objective nature of accelerometry
15 measurement, it is perhaps not surprising that this method achieved such high validity
16 results. Where feasible, use of objective measures of sedentary behaviour is desirable to
17 provide accurate assessment of children and adolescents' sedentary behaviour that is not
18 marred by human error or bias. Accelerometers have the benefit of being able to assess
19 sedentary behaviour in free-living conditions, unlike other objective measures such as
20 calorimetry. However, the cost associated with the purchase of accelerometers, the
21 technical expertise required to transform the raw data into useable data and the additional
22 costs associated with retrieving the monitors from study participants may prohibit use of
23 accelerometers in many studies. In addition, accelerometers cannot differentiate sitting

1 from standing upright with minimal movement, nor can they provide information on the
2 type of sedentary behaviours children are engaging in and therefore would not be
3 appropriate for use in studies interested in investigating specific types of sedentary
4 behaviour. Despite the positive findings in this review, there is considerable variation in
5 the Actigraph cut-points used for sedentary behaviours especially among preschoolers.
6 This difference is possibly due to use of different criterion methods (direct observation vs
7 indirect calorimetry). There is a need for the cross-validation of cut-points in a single
8 study.

9 Seven studies reported reliability or validity of direct observation measures of
10 sedentary behaviour. This semi-objective measure performed well with inter-observer
11 reliability exceeding 90% (29, 42, 55) and validity assessed against heart rate monitoring
12 (44, 45, 47) and energy expenditure (indirect calorimetry) (47) was also high. Such
13 methods may provide a useful alternative to objective measurement, with less potential
14 for bias than self- or proxy-report measures. Direct observation has the added benefit of
15 allowing more comprehensive assessment including type and duration of sedentary
16 behaviour, as well as contextual factors associated with engagement in sedentary
17 behaviour (e.g. presence of other people). However, use of such measures can be costly
18 as it involves a large investment of time by research staff to collect and analyse the
19 observational data, which may be prohibitive for studies with large sample sizes. Because
20 of the time required to train observers, the length of the observation period, and the
21 tedious data-coding requirements, it is highly labour intensive and expensive (56).
22 Subject reactivity to observers is also a legitimate concern, but this problem can be
23 minimized by performing repeat observations. Another limitation of direct observation is

1 that it cannot feasibly be used to assess total habitual sedentary time, and it can only
2 assess sedentary behaviour in specific predefined settings such as the home, school class,
3 playground, parks, etc.

4 The reliability and validity of self- and proxy-report measures of children's and
5 adolescents' sedentary behaviour were most commonly reported. This is likely to be a
6 reflection of the popularity of these types of measures. Thirteen studies reported on the
7 reliability and/or validity of such measures but there was much less consistency in the
8 findings than for accelerometry or direct observation. A number of studies attempted to
9 establish the concurrent validity of self- and proxy-report measures by comparing the
10 results to other forms of self- or proxy-report (e.g. log book or activity diary). However,
11 this is problematic as it involves comparing one method of unknown validity against
12 another measure of unknown validity. The two studies (41, 54) which used an objective
13 criterion measure, accelerometry, reported lower levels of validity. Due to the lack of a
14 'gold standard', future studies examining the validity of sedentary behaviour measures
15 should consider adjusting correlation coefficients upward to attenuate for the weakening
16 effect of measurement error.

17 Reliability results for self- and proxy-report measures of children's and
18 adolescents' sedentary behaviour were mixed. It is difficult to draw conclusions from
19 these results as the measures varied substantially; in the type and aspect of sedentary
20 behaviour they assessed, the period of recall required, the method of administration, the
21 time lapse between assessments, and method of analyses. So while the inconsistent
22 results suggest that self- and proxy-report measures are less reliable than other methods
23 of assessing sedentary behaviour in children and adolescents, it is likely that some of

1 these measures are of higher quality and more comprehensive than others. While much
2 more susceptible to recall and reporting biases than more objective measures, self- and
3 proxy-report measures of sedentary behaviour have the advantage that they are relatively
4 low cost, easy to administer and thus can be easily applied in large scale studies. They are
5 also able to assess all aspects of sedentary behaviour including type, duration and
6 context.

7 While there are clear advantages and disadvantages to the use of the different
8 types of measures of sedentary behaviour in children and adolescents, it appears that
9 objective measures provide the most valid and reliable assessment. Decisions on the
10 choice of measures to use in a study will undoubtedly be largely driven by the study type
11 and resources available. Nonetheless, where possible we recommend the use of objective
12 measures of sedentary behaviour such as accelerometers, in conjunction with more
13 subjective measures (direct observation or self- or proxy-report) to assess aspects of
14 sedentary behaviour that are not captured by accelerometry such as type and context of
15 behaviour. In choosing self- or proxy-report instruments, we recommend researchers
16 select those instruments which have previously been shown to have acceptable reliability
17 and validity. We strongly recommend that studies utilising measures of sedentary
18 behaviour report on the validity and reliability of the measures used, particularly where
19 they have modified existing instruments.

20 **References**

21 1 Reilly J, Methven E, McDowell Z, Hacking B, Alexander D, Stewart L, Kelnar C.
22 Health consequences of obesity. *Arch Dis Child*. 2003 **88**: 748-52.

1 2 Reilly JJ, Dorosty AR. Epidemic of obesity in UK children. *Lancet*. 1999; **354**:
2 1874-75.

3 3 Salmon J, Ball K, Hume C, Booth M, Crawford D. Outcomes of a group-
4 randomized trial to prevent excess weight gain, reduce screen behaviors and
5 promote physical activity in 10-year-old children: Switch-Play. *Int J Obes*. 2008;
6 **32**: 601-12.

7 4 Robinson TN. Reducing children's television viewing to prevent obesity: A
8 randomized controlled trial. *JAMA*. 1999; **282**: 1561-67.

9 5 Simon C, Schweitzer B, Oujaa M, Wagner A, Arveiler D, Tribby E, Copin N,
10 Blanc S, Platat C. Successful overweight prevention in adolescents by increasing
11 physical activity: a 4-year randomized controlled intervention. *Int J Obes*. 2008;
12 **32**: 1489-98.

13 6 Pate RR, O'Neill JR, Lobelo F. The evolving definition of sedentary. *Exerc Sport*
14 *Sci Rev*. 2008; **36**: 173-78.

15 7 Dunstan DW, Barr EL, Healy GN, Salmon J, Shaw JE, Balkau B, Magliano DJ,
16 Cameron AJ, Zimmet PZ, Owen N. Television viewing time and mortality: the
17 Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Circulation*. 2010;
18 **26**: 384-91.

19 8 Marshall SJ, Biddle SJH, Gorely T, Cameron N, Murdey I. Relationships between
20 media use, body fatness and physical activity in children and youth: A meta-
21 analysis. *Int J Obes*. 2004; **28**: 1238-46.

22 9 American Academy of Pediatrics. Children, adolescents and television. *Pediatr*.
23 2001; **107**: 423-26.

- 1 10 Department of Health & Aging. Australia's physical activity recommendations for
2 5-12 year olds. Department of Health & Aging: Canberra 2004.
- 3 11 Mendoza JA, Zimmerman FJ, Christakis DA. Television viewing, computer use,
4 obesity, and adiposity in US preschool children. *Int J Behav Nutr Phys Act.* 2007;
5 **25**: doi:10.1186/479-5868-4-44.
- 6 12 Steele RM, van Sluijs EM, Cassidy A, Griffin SJ, Ekelund U. Targeting sedentary
7 time or moderate- and vigorous-intensity activity: independent relations with
8 adiposity in a population-based sample of 10-y-old British children. *Am J Clin*
9 *Nutr.* 2009; **90**: 1185-92.
- 10 13 Hardy LL, Dobbins TA, Denney-Wilson EA, Okely AD, Booth ML.
11 Sedentariness, small-screen recreation, and fitness in youth. *Am J Prev Med.*
12 2009; **36**: 120-25.
- 13 14 Barnett TA, O'Loughlin J, Sabiston CM, Karp I, Bélanger M, Van Hulst A,
14 Lambert M. Teens and screens: the influence of screen time on adiposity in
15 adolescents. *Am J Epidemiol.* 2010; **172**: 255-62.
- 16 15 Henderson VR. Longitudinal associations between television viewing and body
17 mass index among white and black girls. *J Adolesc Health.* 2007; **41**: 544-50.
- 18 16 Sardinha LB, Andersen LB, Anderssen SA, Quitério AL, Ornelas R, Froberg K,
19 Riddoch CJ, Ekelund U. Objectively measured time spent sedentary is associated
20 with insulin resistance independent of overall and central body fat in 9- to 10-
21 year-old Portuguese children. *Diabetes Care.* 2008; **31**: 569-75.

- 1 17 Mitchell JA, Mattocks C, Ness AR, Leary SD, Pate RR, Dowda M, Blair SN,
2 Riddoch C. Sedentary behavior and obesity in a large cohort of children. *Obes.*
3 2009; **17**: 1596-602.
- 4 18 Patterson P. Reliability, validity and methodological response to the assessment of
5 physical activity via self-report. *Res Q Exerc Sport.* 2000; **71**: 15-20.
- 6 19 Sallis JF, Saelens BE. Assessment of physical activity by self-report: Status,
7 limitations, and future directions. *Res Q Exerc Sport.* 2000; **71**: 1-14.
- 8 20 Sirard JR, Pate RR. Physical activity assessment in children and adolescents.
9 *Sports Med.* 2001; **31**: 439-54.
- 10 21 Terwee CB, Mokkink LB, van Poppel MN, Chinapaw MJ, van Mechelen W, de
11 Vet H. Qualitative attributes and measurement properties of physical activity
12 questionnaires: a checklist. *Sports Med.* 2010; **40**: 525-37.
- 13 22 Bryant MJ, Lucove JC, Evenson KR, Marshall S. Measurement of television
14 viewing in children and adolescents: a systematic review. *Obes Rev.* 2007; **8**: 197-
15 209.
- 16 23 Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for
17 systematic reviews and meta-analyses- the PRISMA statement. *BMJ.* 2009; **339**:
18 332-36.
- 19 24 Chinapaw MJM, Mokkink LB, van Poppel MNM, van Mechelen W, Terwee CB.
20 Physical activity questionnaires for youth: A systematic review of measurement
21 properties. *Sports Med.* 2010; **40**: 539-63.

- 1 25 Scientific Advisory Committee of the Medical Outcomes Trust. Assessing health
2 status and quality-of-life instruments: attributes and review criteria. *Qual Life Res.*
3 2002; **11**: 193-205.
- 4 26 Pols MA, Peeters PH, Kemper HC, Grobbee DE. Methodological aspects of
5 physical activity assessment in epidemiological studies. *Eur J Epidemiol.* 1998;
6 **14**: 63-70.
- 7 27 Zwieg MH, Campbell G. Receiver-operating characteristic (ROC) plots: a
8 fundamental evaluation tool in clinical medicine. *Clin Chem.* 1993; **39**: 561-77.
- 9 28 He M, Harris S, Piche L, Beynon C. Understanding screen-related sedentary
10 behavior and its contributing factors among school-aged children: a social-
11 ecologic exploration. *Am J Health Prom.* 2009; **23**: 299-308.
- 12 29 Anderson DR, Field DE, Collins PA, Lorch EP, Nathan JG. Estimates of young
13 children's time with television: a methodological comparison of parent reports
14 with time-lapse video home observation. *Child Dev.* 1985; **56**: 1345-57.
- 15 30 Taras HL, Sallis JF, Patterson TL, Nader PR, Nelson JA. Television's influence on
16 children's diet and physical activity. *J Dev Behav Pediatr.* 1989; **10**: 176-80.
- 17 31 Brener ND, Kann L, McManus T, Kinchen SA, Sundberg EC, Ross JG.
18 Reliability of the 1999 youth risk behavior survey questionnaire. *J Adolesc*
19 *Health.* 2002; **31**: 336-42.
- 20 32 Schmitz KH, Harnack L, Jacobs Jr DR, Gao S, Lytle LA, Coevering PV, Fulton
21 JE. Reliability and validity of a brief questionnaire to assess television viewing
22 and computer use. *J School Health.* 2004; **74**: 370-77.

- 1 33 Salmon J, Timperio A, Cleland V, Venn A. Trends in children's physical activity
2 and weight status in high and low socio-economic status areas of Melbourne,
3 Victoria, 1985-2001. *Aust NZ J Pub Health*. 2005; **29**: 337-42.
- 4 34 Vereecken CA, Todd J, Roberts C, Mulvihill C, Mae L. Television viewing
5 behaviour and associations with food habits in different countries. *Pub Health*
6 *Nutr*. 2006; **9**: 244-50.
- 7 35 Koezuka N, Koo M, Allison KR, Adlaf EM, Dwyer JJM, Faulkner G, Goodman J.
8 The relationship between sedentary activities and physical inactivity among
9 adolescents: results from the Canadian Community Health Survey. *J Adoles*
10 *Health*. 2006; **39**: 515-22.
- 11 36 Salmon J, Campbell KJ, Crawford DA. Television viewing habits associated with
12 obesity risk factors: A survey of Melbourne schoolchildren. *MJA*. 2006; **184**: 64-
13 67.
- 14 37 Hardy LL, Booth ML, Okely AD. The reliability of the Adolescent Sedentary
15 Activity Questionnaire (ASAQ). *Prev Med*. 2007; **45**: 71-74.
- 16 38 Liou YM, Liou TH, Chang LC. Obesity among adolescents: sedentary leisure
17 time and sleeping as determinants. *J Adv Nurs*. 2010; **66**: 1246-56.
- 18 39 Liu Y, Wang M, Tynjälä J, Lv Y, Villberg J, Zhang Z, Kannas L. Test-retest
19 reliability of selected items of Health Behaviour in School-aged Children (HBSC)
20 survey questionnaire in Beijing, China. *BMC Med Res Methodol*. 2010; **10**: doi:
21 10.1186/471-2288-10-73.
- 22 40 Rey-Lopez JP, Vicente-Rodriguez Gn, Ortega FB, Ruiz JR, Martinez-GÃ³mez D,
23 De Henauw S, Manios Y, Molnar D, Polito A, Verloigne M, Castillo MJ,

- 1 Sjostrom M, De Bourdeaudhuij I, Moreno LA. Sedentary patterns and media
2 availability in European adolescents: The HELENA study. *Prev Med.* 2010; **51**:
3 50-55.
- 4 41 Hardy LL, Bass SL, Booth ML. Changes in sedentary behavior among adolescent
5 girls : a 2.5-year prospective cohort study. *J Adoles Health.* 2007; **40**: 158-65.
- 6 42 DuRant RH, Baranowski T, Johnson M, Thompson WO. The relationship among
7 television watching, physical activity, and body composition of young children.
8 *Pediatr.* 1994; **94**: 449-55.
- 9 43 Puhl J, Greaves K, Hoyt M, Baranowski T. Children's Activity Rating Scale
10 (CARS): description and calibration. *Res Q Exerc Sport.* 1990; **61**: 26-36.
- 11 44 McKenzie TL, Sallis JF, Nader PR, Patterson TL, Elder JP, Berry CC, Rupp JW,
12 Atkins CJ, Buono MJ, Nelson JA. BEACHES: an observational system for
13 assessing children's eating and physical activity behaviors and associated events. *J*
14 *Appl Behav Anal.* 1991; **24**: 141-51.
- 15 45 Rowe PJ, Schuldheisz JM, van der Mars H. Validation of SOFIT for measuring
16 physical activity of first-to eighth-grade students. *Pediatr Exerc Sci.* 1997; **9**: 136-
17 49.
- 18 46 McKenzie T, Marshall S, Sallis J, Conway TL. Leisure-time physical activity in
19 school environments: an observational study using SOPLAY. *Prev Med.* 2000;
20 **30**: 70-77.
- 21 47 Rowe DA, Mahar MT, Raedeke TD, Lore J. Measuring physical activity in
22 children with pedometers: Reliability, reactivity, and replacement of missing data.
23 *Pediatr Exerc Sci.* 2004; **16**: 343-54.

- 1 48 Rowe PJ, van der Mars H, Schuldheisz JM, Fox S. Measuring students' physical
2 activity levels: validating SOFIT for use with high-school students. *J Teach Phys*
3 *Educ.* 2004; **23**: 235-51.
- 4 49 Reilly JJ, Coyle J, Kelly L, Burke G, Grant S, Paton JY. An objective method for
5 measurement of sedentary behavior in 3- to 4-year olds. *Obes Res.* 2003; **11**:
6 1155-58.
- 7 50 Treuth MS, Schmitz K, Catellier DJ, McMurray RG, Murray DM, Almeida MJ,
8 Going S, Norman JE, Pate R. Defining accelerometer thresholds for activity
9 intensities in adolescent girls. *Med Sci Sports Exerc.* 2004; **36**: 1259-66.
- 10 51 Puyau MR, Adolph AL, Vohra FA, Zakeri I, Butte NF. Prediction of activity
11 energy expenditure using accelerometers in children. *Med Sci Sports Exerc.* 2004;
12 **36**: 1625-31.
- 13 52 Sirard JR, Trost SG, Pfeiffer KA, Dowda M, Pate RR. Calibration and evaluation
14 of an objective measure of physical activity in preschool children. *J Phys Act*
15 *Health.* 2005; **3**: 345-57.
- 16 53 Evenson KR, Cattellier D, Gill K, Ondrak K, McMurray RG. Calibration of two
17 objective measures of physical activity for children. *J Sports Sci.* 2008; **26**: 1557-
18 65.
- 19 54 Wen LM, Van der Ploeg HP, Kite J, Cashmore A, Rissel C. A validation study of
20 assessing physical activity and sedentary behavior in children aged 3 to 5 years.
21 *Pediatr Exerc Sci.* 2010; **22**: 408-20.

1 55 McKenzie TL, Marshall SJ, Sallis JF, Conway TL. Leisure-time physical activity
2 in school environments: an observational study using SOPLAY. *Prev Med.* 2000;
3 **30**: 70-7.

4 56 Trost SG. State of the art reviews: measurement of physical activity in children
5 and adolescents. *Am J Lifestyle Med.* 2007; **1**: 299-314.

6

7

Table 1: Reliability of self- and proxy-report measures of sedentary behaviour in children and adolescents

Study	Study sample	Behaviour assessed	Methods	Results	Rating
Anderson et al. (29)	N = 334 families with 5 year old children United States	<i>Proxy-report home TV viewing 10-day diary-</i> parents reported the time the TV was turned on and whether or not child was in the room.	1-month test-retest for 10-day viewing diary using bivariate correlation.	$r = 0.72$	Borderline
Taras et al. (30)	N = 66 mothers of children aged 3-8 years United States	<i>Interviewer administered proxy-report of TV viewing-</i> parents reported their children's time spent watching TV during and between meals for a typical weekday, a typical Saturday and a typical Sunday.	14-21 day test-retest reliability using PC.	$r = 0.80$	Acceptable
Brener et al (31)	N = 4619 children 13-18 years United States	<i>Self-report measure of TV viewing-</i> as part of the Youth Risk Behavior Survey Questionnaire participants report ≤ 2 hours watching TV on an average school day.	2-week test-retest reliability using Kappa.	$k = 0.47\%$	Unacceptable
Schmitz et al. (32)	N = 245 children 11-15 years United States	<i>Self-report measure of TV viewing and computer use-</i> participants report their weekday and weekend time watching TV and using the computer. <i>Self-report measure of TV viewing-</i> the TV viewing question from the 1999 Youth Risk Behaviour Questionnaire (YRBS).	1-week test-retest reliability. Reliability assessments included % agreement, Kappa and SROC.	% agreement ranged from 35% (weekend TV Summer) to 50% (computer use).	Unacceptable
				SROC for TV viewing and computer use ranged from $\rho = 0.55$ (weekend TV summer) to $\rho = 0.68$ (weekday TV school year).	Unacceptable
				Kappa ranged from 0.42 (weekend TV Summer) to 0.55 (weekday school year)	Unacceptable
				YRBS weekday TV item % agreement (48%), SROC (ρ	Unacceptable

= 0.68) and Kappa (0.55).

Salmon et al. (33)	N = 156 parents 40.0 ± 5.2 years	<i>Proxy-report sedentary behaviour measure-</i> parents reported time their child usually spent watching TV, playing electronic games and using the computer in a typical week and on a typical weekend.	2-week test-retest reliability using ICC	Proxy-report ICC (based on mean minutes per day) ranged from 0.6 to 0.8.	Borderline
	N = 147 children 10-12 years Australia	<i>Self-report sedentary behaviour measure-</i> children reported time their child usually spent watching TV, playing electronic games and using the computer in a typical weekday and on a typical weekend.	1-week test-retest reliability using ICC.	Self-report ICC not reported.	Indeterminate
Vereecken et al. (34)	N = 112 children 11-15 years Germany	<i>Self-report measure of TV viewing-</i> participants reported their usual hours of TV viewing (including videos) in free time on weekdays and weekend days.	7-day test -retest reliability using ICC.	ICC (average TV viewing per day) for boys = 0.76 ICC for girls = 0.81	Acceptable
Koezuka et al. (35)	N = 7982 children 12-19 years Canada	<i>Self-report measure of sedentary behaviour</i> - participants reported the number of hours per week (categorical) spent during leisure time using computers, playing video games, watching TV and reading.	Internal consistency of the measure assessed using SROC among the 4 sedentary behaviours.	SROC among the 4 sedentary behaviours ranged from $\rho = 0.04$ to $\rho = 0.13$.	Indeterminate
Salmon et al. (36)	N = 133 parents of children aged 5-12 Australia	<i>Proxy-report of TV viewing-</i> parents reported TV viewing by children on a usual weekday and weekend day.	7-14 day test-retest reliability using ICC.	ICC of usual daily TV = 0.78	Acceptable
Hardy et al. (37)	N = 250 11-15 years Australia	<i>Self-report measure of sedentary behaviour</i> - participants completed the Adolescent Sedentary Activity Questionnaire (ASAQ) which requires participants to report their time spent using small screen recreation devices	2-week test-retest reliability using ICC.	ICC for total time spent in sedentary behaviour was ≥ 0.70 (except for Grade 6 boys = 0.57). ICC values were high for all	Acceptable

		(e.g. watching TV/DVDs), doing homework (with/without computer and tutoring), traveling (motorized), in cultural activities (e.g. hobbies, playing a musical instrument), and socializing (e.g. sitting with friends, using the telephone) and travel.		students for small screen recreation, education and cultural sedentary behaviour with only one or two borderline exceptions. In addition, ICC was unacceptable for 'education' weekdays for Grade 6 boys and girls, ICC was also unacceptable for 'cultural' weekend for Grade 8 boys, There were a range of ICC unacceptable values for 'social' and travel. ICC values for weekend days were lower than for weekdays.	
He et al. (28)	Sample not reported Canada	<i>Self-report measure of sedentary behaviour</i> - participants completed a modified version of the Child Sedentary Activity Questionnaire (CSAQ). The CSAQ requires participants to recall the hours spent each day of the previous week watching TV/videos and playing computer and video games outside school hours.	2-week test-retest reliability using ICC	ICC = 0.98	Acceptable
Liou et al. (38)	Sample not reported China	<i>Self-report measure of sedentary behaviour</i> - participants report average number of hours weekend and weekday spent watching TV, using computers (not for school), reading, traveling in a vehicle and studying/completing homework.	ICC used for test-retest reliability but period between assessments is not described.	ICC = 0.84.	Indeterminate

Liu et al. (39)	N = 95 11-15 years China	<i>Self-report measure of sedentary behaviour</i> - participants completed 8 items related to sedentary behaviour from the Health Behaviour in School-aged Children survey translated into Chinese. Items related to time spent on weekdays and weekend days for the following items: time watching TV/DVDs, using the computer, playing computer and console games, and doing homework.	3-week test-retest reliability using ICC.	ICC for TV and Homework (both weekday and weekend) were acceptable ICC for PC and console games was borderline for weekend and unacceptable for weekdays ICC for using computer was unacceptable	Acceptable Unacceptable Unacceptable
Rey-Lopez et al. (40)	N = 183 13-18 years Europe	<i>Self-report measure of sedentary behaviour</i> - participants reported the hours TV watching (categorical), playing computer games and console games, surfing the internet for non-study reasons, surfing the internet for study reasons and studying both on weekends and weekdays.	1-week test-retest reliability using Kappa.	Weekday TV viewing $k = 0.71$ Computer games $k = 0.82$ Console games $k = 0.82$ Internet no study $k = 0.86$ Internet study $k = 0.46$ Studying $k = 0.73$ Weekend TV viewing $k = 0.68$ Computer games $k = 0.79$ Console games $k = 0.81$ Internet no study $k = 0.71$ Internet study $k = 0.33$ Studying $k = 0.82$	Acceptable Acceptable Acceptable Acceptable Unacceptable Acceptable Acceptable Acceptable Acceptable Acceptable Unacceptable Acceptable

Note. The studies are provided in chronological order, then alphabetical order. PA = physical activity; ICC = intraclass correlation, SROC = Spearman rank order correlation; ρ = Spearman coefficient; k = Kappa coefficient, r = Pearson correlation coefficient, PC = Pearson correlation, NR = not reported; PAL = physical activity level.

Table 2: Criterion validity of self-report and proxy-report measures of sedentary behaviour in children and adolescents

Study	Study sample	Description of measure	Methods	Criterion	Results
Anderson et al. (29)	334 families with 5 year old children United States	<p><i>Proxy-report questionnaire-</i> parents reported child's number of hours of TV viewing time each day of the week during the morning, afternoon and evening. Parents also completed a daily activity chart in which the parents indicated child's daily schedule including TV viewing.</p> <p><i>Home TV viewing diary-</i> parents reported the time the TV was turned on and whether or not child was in the room.</p> <p><i>Home-observations-</i> Using time-lapse video cameras placed in the homes of children which filmed TV and room area. Video equipment began recording when the TV was turned on and stopped recording when the TV was turned off.</p>	<p>The study involved 4 phases:</p> <ol style="list-style-type: none"> 1. Home visit by researcher to observe child's TV viewing and parents completed TV questionnaire 2. Parents provided with 10-day viewing diary 3. 1 month later, a second 10-day viewing diary was issued with experimental group having recording equipment installed 4. Post-test and debrief <p>Bivariate correlation used to examine the relationship between proxy-report estimates and actual TV viewing time.</p>	<p>Video observation of actual behaviour</p>	<p>Correlation between proxy-report of TV viewing and home TV viewing diary was $r = 0.62$</p> <p>Correlation between time lapse videos and diary estimates was $r = 0.84$ when any uncertainty was treated as the child not present</p> <p>Correlation between diary 1 and daily activity chart was $r = 0.48$</p> <p>Correlation between diary 1 and direct estimate of hours watching TV $r = 0.60$</p>
Hardy et al. (41)	N = 172 girls 12-15 years Australia	<p><i>Self-report measure of sedentary behaviour -</i> Participants report time in the following sedentary behaviour before and after school, on a weekend or weekday: watching TV or videos, playing video games, using a</p>	<p>Construct validity of the sedentary behaviour measure self – report questionnaire was determined by accelerometry. At each data collection subjects wore an MTI accelerometer for 7</p>	<p>Accelerometer</p>	<p>Mean weekly difference between self report and accelerometer based measures was -3.2 hours/week.</p> <p>Less than 5% of data points were outside the limits of agreement ($\pm 2SD$) -26.5 to 20.1</p>

		computer for fun or study, doing homework / study, reading, talking on the phone, sitting with friends, doing hobbies or crafts, music/practice, traveling in a car, bus, ferry or train and going to the cinema.	consecutive days (except whilst sleeping or in water). The mean weekly difference between self-report and accelerometer-based sedentary behaviour and limits of agreement were calculated.		h/week).
Wen et al. (54)	N = 34 parents and their children 3-5 years Australia	<i>Proxy-report of child's physical activity and sedentary behaviour</i> - parents reported their child's behaviour in terms of number of times, hrs and mins. in a 7-day diary. A number of items were used to assess the amount of time in sedentary behaviour including watching TV, videos, DVD, computer or computer games inc. Playstation, playing indoors in a stationary way, reading, napping /sleeping, eating and sitting in a pram.	Children wore accelerometers for 7-days. SROC was used to examine the relationship between the diary entries and accelerometer activity counts.	Accelerometer	Time spent in sedentary behaviour recorded by the diary was positively correlated with sedentary behaviour time assessed by the accelerometer ($\rho = 0.24$). Time in screen time was ($\rho = 0.08$).

Note. PA = physical activity; ICC = intraclass correlation, SROC = Spearman rank order correlation; ρ = Spearman coefficient; k = Kappa coefficient, r = Pearson correlation coefficient, PC = Pearson correlation, NR = not reported; PAL = physical activity level.

Table 3: Concurrent validity of self-report and proxy-report measures of sedentary behaviour in children and adolescents

Study	Study sample	Description of measure	Methods	Criterion	Results
Schmitz et al. (32)	N = 245 children 11-15 years United States	<i>Self-report measure of TV viewing and computer use-</i> participants report their weekday and weekend time watching TV and using the computer in a diary.	To assess concurrent validity, participants completed TV and computer logs for 7 days. Validity assessments included SROC and mean difference using t-tests.	Self-reported TV and computer diary.	SROC ranged from 0.37 (weekend TV) to 0.47 (average week TV) Mean difference in hours ranged from -0.09 (average week TY) to 0.68 (computer only). SROC for YRBS item (weekday TV) was $\rho = 0.46$. Mean difference in hours for YRBS item was -0.04 hours.
Salmon et al. (33)	N = 156 parents Mean age = 40.0 ± 5.2 years N = 147 children Mean age = 11.8 ± 0.8 years Australia	<i>Proxy-report sedentary behaviour measure-</i> parents reported time their child usually spent watching TV, playing electronic games and using the computer in a typical week and on a typical weekend. <i>Self-report sedentary behaviour measure-</i> children reported time their child usually spent watching TV, playing electronic games and using the computer in a typical weekday and on a typical weekend.	Convergent validity between parents' proxy-report data and children self-report was tested using correlation. NR	Parent report of child's TV viewing, electronic game and computer usage.	TV viewing ($\rho = 0.61$), computer use ($\rho = 0.47$) and playing electronic games ($\rho = 0.44$). NR

Vereecken et al. (34)	N = 112 children 11-15 years Germany	<i>Self-report measure of TV viewing-</i> participants reported their usual hours of TV viewing (including videos) in free time on weekdays and weekend days.	ICC was used to assess convergent validity of the self-report questions and a 7-day TV diary completed by participants.	Self-reported TV diary.	Boys ICC = 0.36 Girls ICC = 0.54
He et al. (28)	Sample not reported Canada	<i>Self-report measure of sedentary behaviour -</i> participants completed a modified version of the Child Sedentary Activity Questionnaire (CSAQ). The CSAQ requires participants to recall the hours spent each day of the previous week watching TV/videos and playing computer and video games outside school hours.	Criterion for validity was assessed using an activity diary. ICC values provided.	Self-reported activity diary.	ICC ranged from 0.5-0.8.
Liou et al (38)	Sample not reported China	<i>Self-report measure of sedentary behaviour-</i> items related to time spent on weekdays and weekend days for the following items: time watching TV/DVDs, using the computer, playing computer and console games, and doing homework.	Methods for assessing validity are not described.	Not reported.	Content validity index = 0.99

Note. PA = physical activity; ICC = intraclass correlation, SROC = Spearman rank order correlation; ρ = Spearman coefficient; k = Kappa coefficient, r = Pearson correlation coefficient, PC = Pearson correlation, NR = not reported; PAL = physical activity level.

Table 4: Reliability and validity of direct observation of sedentary behaviour in children and adolescents

Study	Study sample	Description of measure	Methods	Criterion	Results
Anderson et al. (29)	N = 334 families with a 5 year old child United States	<i>Home-observations-</i> Using time-lapse video cameras placed in the homes of families.	TV viewing (time spent in the TV room and time spent directing visual attention towards the TV) was observed using video-taped recording equipment in the families' homes. Inter-observer reliability for presence in the viewing room and visual attention was tested using bivariate correlation. Reliability based on 14 viewers rated by 2 observers.	N/A	Inter-observer reliability for presence in the viewing room and visual attention were $r = 0.98$ and $r = 0.90$, respectively.
Puhl et al. (43)	Validation: N = 25 children 6 years Inter-observer Reliability: N = 192 children 3-4 years United States	Children's Activity Rating Scale (CARS)	CARS categories (1. stationary – no movement, 2. stationary – with movement, 3. translocation – slow/easy, 4. translocation – medium/moderate. 5. translocation - translocation – medium/moderate. 5. Translocation –fast, very fast/strenuous) were validated against HR monitoring and indirect calorimetry during a 50-minute protocol. HR and VO ₂ were collected continuously. Reliability based on 389 paired observation periods by 11 observers over 12 months.	HR monitoring and indirect calorimetry	Mean %Max VO ₂ differed for Category 1 - lying (14.6) and sitting (14.5) vs Category 2 - standing/colouring (21.2) and standing/ball activity (23.0). Mean HR (bpm) differed for Category 1 vs Category 2. Mean HR (bpm) differed for lying (89) vs sitting vs standing/ colouring (116) vs standing/ball activity (112). Inter-observer percent agreement = 84.1%.
McKenzie et al. (44)	N = 19 children 4-9 years United States	<i>Direct observation of children's behaviour—</i> The Behaviours of Eating and Activity for Children's Health Evaluation Systems (BEACHES) is a direct observation of	HR was measured using a UNIQ Heart watch while children participated in specific activities	HR monitoring	HR increased across activities HR lying = 99 BPM HR sitting = 107 BPM HR standing = 130 BPM HR walking = 130 BPM HR very active = 153 BPM

		children's physical activity and eating behaviours and related environmental events using momentary time sampling.			
DuRant et al. (42)	N = 191 children 3-4 years United States	<i>Direct observation of children's TV viewing-</i> The Children's Activity Rating Scale (CARS) was used to determine when a child was in the room with a TV on and attending to the program.	Children were observed using CARS from 6-12 hours across 4 days in one year. Every minute the child watched TV was recorded	N/A	TV viewing inter-observer agreement was 96%.
Rowe et al. (45)	N = 173 adolescents Mean age = 10.6 years United States	<i>Direct observation of children's activity levels in PE lessons-</i> System for Observing Fitness Instruction Time (SOFIT)	SOFIT categories (lying, sitting, standing, walking, running) were validated against HR monitoring collected at 5 sec intervals during a 36-minute protocol. Internal consistency reliability for HR during each category was examined. Test-retest reliability was examined among 47 students.	HR monitoring	Mean HR (BPM) differed for: i) lying (87) and sitting (91) vs ii) standing (103) vs iii) walking (121). Internal consistency reliabilities were – HR: lying = 0.99, sitting = 0.99. Intraclass correlations were – HR: lying = 0.88, sitting = 0.88.
McKenzie et al. (46)	N = 24 middle schools Children in grades 6-8 United States	<i>Direct observation of children's activity levels during breaks-</i> The System for Observing Play and Leisure Activity in Youth (SOPLAY) is based on momentary time sampling and is used to determine the number of participants and their physical	Independent inter-observer reliability on 14 observations using ICC.	N/A	ICC for sedentary time = 0.98

		activity levels during play.			
Rowe et al. (48)	N = 35 adolescents Mean age = 15.7 years United States	<i>Direct observation of children's activity levels in PE lessons-System for Observing Fitness Instruction Time (SOFIT)</i>	SOFIT categories (lying, sitting, standing, walking, running) were validated against HR monitoring and indirect calorimetry during a 42-minute protocol. HR and VO ₂ were collected at 5 sec and 20 sec intervals, respectively. Internal consistency reliability for HR and EE during each category was examined.	HR monitoring	Mean HR (BPM) differed for: i) lying (71.6) and sitting (77.1) vs ii) standing (85.7) vs iii) walking (106.3). Mean EE (O ₂ /kg ¹ /min ¹ differed for: i) lying (3.9), sitting (4.2), and standing (4.2) vs ii) walking (15.2). Internal consistency reliabilities were – HR: lying = 0.98, sitting = 0.98; EE: lying = 0.82, sitting = 0.78.

Note. HE = heart rate, EE = energy expenditure, BPM = beats per minute.

Table 5: Validity of objective measures of sedentary behaviour in children and adolescents

Study	Study sample	Measure	Methods	Criterion	Equation/cut-points	Results	Rating*
Reilly et al. (49)	Development: N = 30 children, 3-4 years Cross-validation: N = 50 children, 3-4 years Scotland	Actigraph 7164 accelerometer	Activity monitors were compared against a validated direct observation technique.	Direct observation	Sedentary behaviour <1100 counts/min	Sensitivity: 83% Specificity: 82%	Good
Treuth et al. (50)	N = 24 adolescent girls 13-14 years United States	Actigraph 7164 accelerometer	Participants wore accelerometers and Cosmed metabolic units to determine oxygen consumption and heart rate. Participants completed a range of sedentary activities including lying on a bed, sitting in a chair watching a movie and sitting in a chair playing a computer game. Sedentary behaviour <1.5 METs.	Metabolic measurement system	METs = 2.01 + 0.000856 counts/min Sedentary behaviour <100 counts/min	Sensitivity: 100% Specificity: 100% Area under ROC curve: 1.00.	Excellent
Puyau et al. (51)	N = 32 7-18 years United States	Actical accelerometer	Activity monitors were validated and calibrated against continuous 4-hour measurements of EE by respiration room calorimetry and heart rate by telemetry. While they were in the calorimeter, the children adhered to a structured protocol of physical activities. Sedentary activities included playing Nintendo, and working on a computer. Correlation used to compare accelerometer counts with EE. Sedentary behaviour- AEE <0.01 kcal·kg ⁻¹ ·min ⁻¹ .	Calorimeter and heart rate telemetry	Actical AEE = 0.00423 + 0.00031 counts ^{0.653} Sedentary behaviour <100 counts/min	Actical Area under ROC curve: 0.85.	Good
		Actiwatch accelerometer			Actiwatch AEE = 0.00441 + 0.00032 counts ^{0.724} Sedentary behaviour <50 counts/min	Actiwatch Area under ROC curve: 0.85.	Good

Sirard et al. (52)	Development: N = 16 children, 3-5 years Validation: N = 269 children, 3-5 years United States	Actigraph 7164 accelerometer	Participants completed 5 structured activities while wearing the accelerometers. Sedentary activities included sitting and talking and sitting and playing. Activity monitors were compared against a validated direct observation technique.	Direct observation	3y: <1204 counts/min 4y: <1452 counts/min 5y: <1592 counts/min	3y	Excellent
						Sensitivity: 100%	
						Specificity: 100%	
						Area under ROC curve: 1.00	Excellent
						4y	
						Sensitivity: 100%	Excellent
						Specificity: 100%	
						Area under ROC curve: 1.00	
						5y	
						Sensitivity: 94%	
						Specificity: 92%	
						Area under ROC curve: 0.97	
Evenson et al. (53)	N = 33 children 5-8 years United States	Actical accelerometer	Participants wore accelerometers and Cosmed metabolic system to determine oxygen consumption. Sedentary activities included sitting in a chair watching a movie and sitting in a chair colouring.	Metabolic measurement system	Actical Sedentary behaviour <44 counts/min	Actical	Excellent
		Actigraph 7164 accelerometer				Sensitivity: 97%	
						Specificity: 98%	
						Area under ROC curve: 0.99	
					Actigraph Sedentary behaviour <100 counts/min	Actigraph	Excellent
						Sensitivity: 95%	
						Specificity: 93%	
						Area under ROC curve: 0.98	

Note. AEE = activity energy expenditure; EE = energy expenditure; HR = heart rate; ROC = receiver operating characteristic

*According to area under ROC curve results: > 0.90 = excellent, 0.80–0.90 = good, 0.70–0.80 = fair, and < 0.70 = poor (27).