



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers

Faculty of Science, Medicine and Health

2014

Late Pleistocene glaciation of the Kodar Mountains, south-central Siberia, constrained by Be-10 exposure dating

Martin Margold

Stockholm University, martin.margold@natgeo.su.se

John Jansen

Stockholm University, jjansen@uow.edu.au

Artem L. Gurinov

Lomonosov Moscow State University

Natalya V. Reznichenko

Durham University

Alexandru T. Codilean

University of Wollongong, codilean@uow.edu.au

See next page for additional authors

Publication Details

Margold, M., Jansen, J. D., Gurinov, A. L., Reznichenko, N. V., Codilean, A. T. & Fink, D. (2014). Late Pleistocene glaciation of the Kodar Mountains, south-central Siberia, constrained by Be-10 exposure dating. *Geophysical Research Abstracts of EGU General Assembly 2014* (pp. 1-1). UK: ADS.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Late Pleistocene glaciation of the Kodar Mountains, south-central Siberia, constrained by Be-10 exposure dating

Abstract

Geophysical Research Abstracts of EGU General Assembly 2014, held 27 April - 2 May, 2014 in Vienna, Austria.

Keywords

GeoQuest

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Margold, M., Jansen, J. D., Gurinov, A. L., Reznichenko, N. V., Codilean, A. T. & Fink, D. (2014). Late Pleistocene glaciation of the Kodar Mountains, south-central Siberia, constrained by Be-10 exposure dating. Geophysical Research Abstracts of EGU General Assembly 2014 (pp. 1-1). UK: ADS.

Authors

Martin Margold, John Jansen, Artem L. Gurinov, Natalya V. Reznichenko, Alexandru T. Codilean, and David Fink



Late Pleistocene glaciation of the Kodar Mountains, south-central Siberia, constrained by Be-10 exposure dating

Martin Margold (1), John D. Jansen (2), Artem L. Gurinov (3), Natalya V. Reznichenko (1), Alexandru T. Codilean (4), and David Fink (5)

(1) Durham University, Department of Geography, Durham, United Kingdom (martin.margold@durham.ac.uk), (2) Stockholm University, Department of Physical Geography and Quaternary Geology, Stockholm, Sweden, (3) Department of Geomorphology and Paleogeography, Lomonosov Moscow State University, Moscow, Russia, (4) University of Wollongong, School of Earth and Environmental Sciences, Wollongong, Australia, (5) Australian Nuclear Science and Technology Organisation, Institute for Environmental Research, Lucas Heights, Australia

The glacial history of the mountainous Transbaikalia region NE of Lake Baikal, Siberia, has so far received little attention. The Kodar Mountains exceed 3000 m in elevation and small cirque glaciers currently occur in the central parts of the range. Yet, greatly expanded glaciation in the past is evidenced by massive moraine complexes at the mouth of glacial valleys in the Chara Depression (part of the Baikal rift system), and along the Vitim River valley to the NW. The moraines document the existence of large valley glaciers that reached to over 120 kilometres in length. We applied Be-10 exposure dating to determine the timing of glacier advances that formed the major moraines: 19 samples from boulders on 8 moraine ridges in 5 moraine complexes. Our results indicate extensive glaciation in the Kodar Mountains at the time of the Last Glacial Maximum, with our ages spanning from about 20 ka to the Late Glacial. We attribute the age spread on some of the moraines to boulder exhumation linked to permafrost dynamics.