

University of Wollongong

Research Online

Faculty of Health and Behavioural Sciences -
Papers (Archive)

Faculty of Science, Medicine and Health

1-1-2012

Seven out of ten most popular vegetables remain top choice during first 3 months of a 12 month weight loss dietary intervention trial

Jane O'Shea

University of Wollongong, janeo@uow.edu.au

Yasmine Probst

University of Wollongong, yasmine@uow.edu.au

Rebecca Thorne

University of Wollongong, beck@uow.edu.au

Sara Grafenauer

University of Wollongong, sarag@uow.edu.au

Linda Tapsell

University of Wollongong, ltapsell@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/hbspapers>



Part of the [Arts and Humanities Commons](#), [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

O'Shea, Jane; Probst, Yasmine; Thorne, Rebecca; Grafenauer, Sara; and Tapsell, Linda: Seven out of ten most popular vegetables remain top choice during first 3 months of a 12 month weight loss dietary intervention trial 2012, 528.

<https://ro.uow.edu.au/hbspapers/3060>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Seven out of ten most popular vegetables remain top choice during first 3 months of a 12 month weight loss dietary intervention trial

Abstract

Increasing vegetable consumption is part of dietary advice for weight loss, but how this converts to vegetable choices is not known. In this context, our aim was to identify the main vegetable groups reported by the study sample in the Healthy Eating and Lifestyle Trial [ACTRN1260000784011] at baseline (Oct 2010-Feb 2011) and after 3mo (Feb - May 2011). Data from diet histories (DH) were analyzed using the AUSNUT 2007 database in Foodworks (Xyris, version 6.0.2562) to determine the average daily amount (g) of vegetables consumed by the study sample. Baseline DH (n=113 subjects; 85 female) exposed 32 vegetable categories. The top 10 : tomato (69.9g), potato (58.2g), cucumber (27.5g), carrot (25.7g), lettuce (19.4g), broccoli(17.0g), mixed vegetables (15.3g), leafy greens(12.2g), onion(12.0g), avocado (12.0g) contributed to 74% of the total (365g/day). After 3mo (n=109 subjects), 7 remained in the top 10 (contributing 72% of average consumption, 505g/day), with legumes (35.0g), capsicum (22.8g), pumpkin (15.4g) replacing leafy greens, onion, avocado . Tomato remained top ranked but potato dropped from 2nd to 5th rank (58.2g vs36.5g), and mixed vegetables shifted from 7th to 4th (15.3 g vs 39.1g). Baseline intakes were similar to National Nutrition Survey 1995 levels (240g -300g). Consumption appeared to increase for the group, above the Australian Guide to Healthy Eating recommendation of ~ 375g/day. Advice to increase vegetables may influence vegetables choices, with tomatoes a mainstay and frozen vegetables a feasible option. More detailed analysis may be informative for strategies to increase vegetable consumption in this context.

Keywords

month, 12, first, trial, popular, intervention, months, 3, during, choice, top, remain, vegetables, most, ten, out, seven, dietary, loss, weight

Disciplines

Arts and Humanities | Life Sciences | Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

O'Shea, J., Probst, Y., Thorne, R., Grafenauer, S. & Tapsell, L. (2012). Seven out of ten most popular vegetables remain top choice during first 3 months of a 12 month weight loss dietary intervention trial. 16th International Congress of Dietetics (p. 528).

Seven out of ten most popular vegetables remain top choice during first 3 months of a 12 month weight loss dietary intervention trial

Jane O'Shea¹, Yasmine Probst¹, Rebecca Thorne¹, Sara Grafenauer¹, Linda Tapsell¹
¹*Smart Foods Centre, University of Wollongong, Wollongong, NSW 2522, Australia*

Increasing vegetable consumption is part of dietary advice for weight loss, but how this converts to vegetable choices is not known. In this context, our aim was to identify the main vegetable groups reported by the study sample in the Healthy Eating and Lifestyle Trial [ACTRN1260000784011] at baseline (Oct 2010-Feb 2011) and after 3mo (Feb – May 2011). Data from diet histories (DH) were analyzed using the AUSNUT 2007 database in Foodworks (Xyris, version 6.0.2562) to determine the average daily amount (g) of vegetables consumed by the study sample. Baseline DH (n=113 subjects; 85 female) exposed 32 vegetable categories. The top 10 : tomato (69.9g), potato (58.2g), cucumber (27.5g), carrot (25.7g), lettuce (19.4g), broccoli(17.0g), mixed vegetables (15.3g), leafy greens(12.2g), onion(12.0g), avocado (12.0g) contributed to 74% of the total (365g/day). After 3mo (n=109 subjects), 7 remained in the top 10 (contributing 72% of average consumption, 505g/day), with legumes (35.0g), capsicum (22.8g), pumpkin (15.4g) replacing leafy greens, onion, avocado . Tomato remained top ranked but potato dropped from 2nd to 5th rank (58.2g vs36.5g), and mixed vegetables shifted from 7th to 4th (15.3 g vs 39.1g). Baseline intakes were similar to National Nutrition Survey 1995 levels (240g -300g). Consumption appeared to increase for the group, above the Australian Guide to Healthy Eating recommendation of ~375g/day. Advice to increase vegetables may influence vegetables choices, with tomatoes a mainstay and frozen vegetables a feasible option. More detailed analysis may be informative for strategies to increase vegetable consumption in this context.

Funding support for this research has been provided by Horticulture Australia Limited
Contact author – janeo@uow.edu.au