



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science - Papers (Archive)

Faculty of Science, Medicine and Health

2012

Antimalarial alkaloids from a Bhutanese traditional medicinal plant *Corydalis dubia*

Phurpa Wangchuk

University of Wollongong, pw54@uow.edu.au

Paul A. Keller

University of Wollongong, keller@uow.edu.au

Stephen G. Pyne

University of Wollongong, spyne@uow.edu.au

Anthony C. Willis

Australian National University

Sumalee Kamchonwongpaisan

BIOTEC, Pathum Thani 12120, Thailand

Publication Details

Wangchuk, P., Keller, P. A., Pyne, S. G., Willis, A. C., Kamchonwongpaisan, S. (2012). Antimalarial alkaloids from a Bhutanese traditional medicinal plant *Corydalis dubia*. *Journal of Ethnopharmacology*, 143 (1), 310-313.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Antimalarial alkaloids from a Bhutanese traditional medicinal plant *Corydalis dubia*

Abstract

Ethnopharmacological relevance: *Corydalis dubia* is used in Bhutanese traditional medicine as a febrifuge and for treating infections in the blood, liver and bile which correlate to the signs and symptoms of malarial and microbial infections.

Aim of the study: To validate the ethnopharmacological uses of the plant and to discover potential new therapeutic drug leads. Materials and methods *C. dubia* was collected from Bhutan and the alkaloids were obtained using acid–base fractionation and separation by repeated column and preparative plate chromatography. The alkaloids were identified from analysis of their physiochemical and spectroscopic data and were tested for antiplasmodial, antimicrobial and cytotoxicity activities.

Results: A systematic extraction and isolation protocol yielded one new natural product, dubiamine, and seven known isoquinoline alkaloids, scoulerine, cheilanthifoline, protopine, capnoidine, bicuculline, corydecumbine and hydrastine. Among the four alkaloids tested, scoulerine showed the best antiplasmodial activity with IC₅₀ values of 5.4 μ M and 3.1 μ M against the antifolate sensitive and the multidrug resistant *P. falciparum* strains: TM4/8.2 and K1CB1, respectively. None of the alkaloids tested showed significant antimicrobial or cytotoxicity activities.

Conclusions: The antiplasmodial test results, of the isolated alkaloid components, are commensurated with the ethnopharmacological uses of this plant.

Keywords

alkaloids, bhutanese, traditional, antimalarial, medicinal, dubia, plant, corydalis, CMMB

Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

Wangchuk, P., Keller, P. A., Pyne, S. G., Willis, A. C., Kamchonwongpaisan, S. (2012). Antimalarial alkaloids from a Bhutanese traditional medicinal plant *Corydalis dubia*. *Journal of Ethnopharmacology*, 143 (1), 310-313.

Antimalarial alkaloids from a Bhutanese traditional medicinal plant *Corydalis dubia*

Phurpa Wangchuk^{a,b}, Paul A. Keller^{a*}, Stephen G. Pyne^{a,*}, Anthony C. Willis^c, Sumalee Kamchonwongpaisan^d

^a*School of Chemistry, University of Wollongong, Wollongong, NSW, 2522, Australia*

^b*Pharmaceutical and Research Unit, Ministry of Health, Thimphu, Bhutan*

^c*Research School of Chemistry, The Australian National University, Canberra, A.C.T. 0200, Australia*

^d*Medical Molecular Biology Research Unit, National Center for Genetic Engineering and Biotechnology, National Science and Technology Development Agency, Thailand Science Park, Pathumthani, 12120, Thailand.*

*Corresponding author: School of Chemistry, University of Wollongong, NSW, 2522, Australia. Tel.: +61 2 4221 4388; Fax: +61 2 4221 4287; E-mail address: spyne@uow.edu.au

ABSTRACT

Ethnopharmacological relevance: *Corydalis dubia* is used in Bhutanese traditional medicine as a febrifuge and for treating infections in the blood, liver and bile which correlate to the signs and symptoms of malarial and microbial infections.

Aim of the study: To validate the ethnopharmacological uses of the plant and to discover potential new therapeutic drug leads.

Materials and methods: *C. dubia* was collected from Bhutan and the alkaloids were obtained using acid-base fractionation and separation by repeated column and preparative plate chromatography. The

alkaloids were identified from analysis of their physiochemical and spectroscopic data and were tested for antiplasmodial, antimicrobial and cytotoxicity activities.

Results: A systematic extraction and isolation protocol yielded one new natural product, dubiamine, and seven known isoquinoline alkaloids, scoulerine, cheilanthifoline, protopine, capnoidine, bicuculline, corydecumbine and hydrastine. Among the four alkaloids tested, scoulerine showed the best antiplasmodial activity with IC₅₀ values of 5.4 µM and 3.1 µM against the antifolate sensitive and the multidrug resistant *P. falciparum* strains: TM4/8.2 and K1CB1, respectively. None of the alkaloids tested showed significant antimicrobial or cytotoxicity activities.

Conclusions: The antiplasmodial test results, of the isolated alkaloid components, are commensurated with the ethnopharmacological uses of this plant and generated a new antiplasmodial drug lead.

Keywords

Bhutanese traditional medicine; medicinal plant, *Corydalis dubia*; antimalarial; scoulerine

1. Introduction

Corydalis dubia Prain (Fumariaceae), which is locally known as *Re-skon*, is an annual yellow-flowering glabrous herb of lower stature arising from tuberous rootstock. This plant is endemic to Bhutan, Sikkim (India) and Chumbi Valley of Tibet (Grierson and Long, 1984). In Bhutan, it occurs in the wild in localized parts of Lingzhi and Bumthang region at an altitude range of 4294-5039 meters above sea level (Wangchuk et al., 2008; Anonymous, 2008). It grows sparsely with recorded density of 0.6 to 3.6 plants/m² (Anonymous, 2008). Therapeutic uses of this plant were reported for detoxifying impure blood, treating fever arising from the infections of liver, heart, lung, pancreas and kidney and alleviating neuralgia and complicated disorders that have resulted from a combination of defective air, bile and phlegm (Wangchuk et al., 2011). These are symptoms which correlate to both malarial and microbial infections. A preliminary phytochemical screening demonstrated the presence of alkaloids

and that the crude MeOH and CHCl₃ alkaloid extracts showed significant antiplasmodial activity against chloroquine and antifolate resistant *Plasmodium falciparum* strains: TM4/8.2 and K1CB1 (Wangchuk et al., 2011). This prompted us to investigate the alkaloid constituents of *C. dubia* with a view to generate potential new therapeutic agents and at the same time validate the use of this plant in Bhutanese traditional medicine (BTM) for the treatment of malarial or microbial infections.

A systematic extraction and isolation protocol yielded one new natural product which we named as dubiamine (**1**) drawn from the species name of the plant. Seven known isoquinoline alkaloids were also isolated which were identified as scoulerine (**2**), cheilanthifoline (**3**), protopine (**4**), capnoidine (**5**), bicuculline (**6**), corydecumbine (**7**) and hydrastine (**8**). Compounds **1**, **2**, **5** and **6** were investigated for their antiplasmodial, antimicrobial and cytotoxicity activities. Compounds **2** showed significant antiplasmodial activity which is reported here for the first time. The antiplasmodial activities of compounds **3** and **4** were reported earlier by us and are included in Table 1.

2. Materials and methods

2.1. Plant material

Wild *C. dubia* (whole part) was collected from Thruenchela (Altitude: 4651 m; Latitude: 27°56' 00.1"; Longitude: 89° 26' 11.6"; global positioning system (GPS) point number: 167; Site number: P167; Slope: 30°; Aspect: North-West), under the Lingzhi region in Bhutan in August 2009. The herbarium voucher specimen with accession number 78 was authenticated by Samten, a pharmacognosist at the Pharmaceutical and Research Unit (PRU) and was deposited at the herbarium unit of PRU, Thimphu, Bhutan.

2.2. Extraction, isolation and identification of alkaloids

The extraction and separation of the alkaloids from the crude methanol extract (65.12 g obtained from 2 kg dry plant material) was undertaken via an acid/base extraction procedure using the analytical chemicals and tools as reported in Wangchuk et al. (2012). This protocol furnished the CH₂Cl₂ extract (17.87 g) which contained a mixture of acidic and neutral compounds and the CHCl₃ extract (1.13 g) which contained the basic alkaloids. Focusing on the alkaloids, the CHCl₃ extract (1.2 g) was chromatographed on a silica gel (120 g, 200-300 mesh) column, eluting with a gradient solvent system of MeOH-CHCl₃ (total combined volume of 200 mL in a v/v ratio of 0:100, 0.5:99.5, 1.5:98.5, 3:97, 4:96, 5:95, 7:93, 10:90, 15:85, 20:80, 30:70, 40:60) to obtain fractions F1-F11. Fraction F4, upon crystallization from CHCl₃/MeOH (1:1), yielded compound **5** (80 mg). Its mother liquor was purified by column chromatography eluting with a gradient solvent system of MeOH-CHCl₃ to obtain five sub-fractions F4.1-F4.5. The separation of these sub-fractions F4.2, F4.3, F4.4 and F4.5 using preparative TLC plates (mobile phase; CHCl₃/ethyl acetate/NH₄OH) (55:45:4 drops) yielded compounds **3** (15 mg), **6** (18 mg), **7** (5 mg), and **8** (0.7 mg). Fractions 5 and 6 were combined (F56) and then separated by the column chromatography with a gradient solvent system of ethyl acetate-CHCl₃ (50 mL, v/v ratio of 0:100, 20:80, 40:60, 50:50, 60:40, 80:20, 100:0) to obtain sub-fractions F56.1-F56.6. Further separation of sub-fraction F56.2 on preparative TLC plates (mobile phase; CHCl₃/ethyl acetate (55:45)) gave compound **2** (9 mg). Crystallization of the sub-fraction F56.3 from CHCl₃/MeOH (1:1) furnished compound **1** (7 mg). Fraction F7 was crystallized from the ethyl acetate/hexane (1:1) to give compound **4** (160 mg), the major alkaloid of the plant.

Compound **1** was identified as a new natural product which we named as dubiamine after the name of the plant species. It was obtained as colorless needle-like crystals with mp 192.8-194.6 °C; ESI-MS (*m/z*) 300 [M + H⁺]; EI-MS (*m/z*): 299 [M⁺], 298, 192, 176, 164, 148 (100%), 134, 121, 107; and HR-ESI-MS (*m/z*) 300.1597 [M + H]⁺ (300.1600 calc. for C₁₈H₂₂NO₃). ¹H-NMR (500 MHz, CDCl₃): δ 2.73 (2H, *t*, *J* = 5.8 Hz, H-3), 2.81 (2H, *t*, *J* = 5.8 Hz, H-4), 3.53 (2H, *s*, H-1), 3.60 (2H, *s*,

H-2), 3.80 (3H, *s*, OMe on C-7), 3.83 (3H, *s*, OMe on C-6), 6.48 (1H, *s*, H-8), 6.58 (1H, *s*, H-5), 6.75 (2H, *d*, $J = 8.0$ Hz, H-3' and H-5'), 7.22 (2H, *d*, $J = 7.5$ Hz, H-2' and H-6'). ^{13}C -NMR (125 MHz, CDCl_3): δ 28.7 (C-4), 50.7 (C-3), 55.6 (C-1), 56.0 (OMe), 56.1 (OMe), 62.2 (C-2), 109.7 (C-8), 111.6 (C-5), 115.3 (C-3' and C-5'), 126.3 (C-8a), 126.7 (C-4a), 130.1 (C-1'), 130.7 (C-2' and C-6'), 147.6 (C-6 and C-7).




Fig. 1. Alkaloids isolated from *Corydalis dubia*.




Fig. 2. Single crystal structure of dubiamine (**1**) with CCDC 872799.

The structure of **1** (Fig. 1) was confirmed from the analysis of its single crystal X-ray crystallographic data (Fig. 2): (C₁₈H₂₁NO₃), M_r 299.37. Monoclinic, P2₁/n, a = 6.4140 (1) Å, b = 13.0324 (3) Å, c = 19.2120 (4) Å, β = 98.8791 (11)°, V = 1586.68 (6) Å³, Z = 4, F(000) = 640, D_x = 1.253 Mg m⁻³, MoKα radiation, λ = 0.71073 Å, cell parameters from 26826 reflections, θ = 26-27.5°, μ = 0.09 mm⁻¹, specimen = 0.42 × 0.35 × 0.22 mm (colorless block). 46844 reflections were measured to 2θ_{max} = 55° and merged to 3625 unique reflections. Final R(3174 reflns with F² > 2σ(F²)) = 0.037, wR (all data) = 0.101, S = 0.99.

The crystal structure has been deposited at the Cambridge Crystallographic Data Centre and allocated the deposition number CCDC 872799.

While **1** has not been reported as a natural product, it has been synthesized *in vitro* from the enzymatic oxidation of *d,l*-N-norarmepavine which was assigned the IUPAC name as 4-((6,7-dimethoxy-3,4-dihydroisoquinolin-2(1*H*)-yl)methyl)phenol (Inubushi et al., 1969). This alkaloid was closely related to sendaverine and its derivatives (Pandey and Tiwari, 1980; Kametani and Ohkubo, 1965; Suau et al., 1994). Other known compounds **2-8** were identified through MS library matching techniques (NIST08s) and subsequent comparison of their melting point, optical rotation, mass spectra and NMR spectra (¹H and ¹³C) with those reported. Compound **2** was identified as scoulerine (Brochman-Hanssen and Nielson 1966; Cheng et al., 2008), compound **3** as cheilanthifoline (Okamoto

et al., 1971; Haisova and Slavik, 1973; Yeola and Mali, 1984; Rucker et al., 1994; Cheng et al., 2008), compound **4** as protopine (Abou-Donia et al., 1980; Takahashi et al., 1985; Seger et al., 2004), compound **5** as capnoidine (Blasko et al., 1982; Ribar and Meszaros, 1991; Pilar et al., 2010) whose structure was also confirmed from a X-ray crystal structure (not shown here), compound **6** as bicuculline (Edwards and Handa, 1961; Abou-Donia et al., 1980; Blasko et al., 1982; Elango, et al., 1982; Basha et al., 2002), compound **7** as corydecumbine (Basnet et al., 1993) and compound **8** as hydrastine (Blasko et al., 1982; Elango, et al., 1982).

2.3. *Bioassay methods*

Antiplasmodial, antimicrobial and cytotoxicity bioassays were carried out using the standard protocols reported by Wangchuk et al., (2011). The test strains of *Plasmodium falciparum* used for the antiplasmodial assay were K1CB1, a multidrug resistant strain; and TM4/8.2, a wild type chloroquine and antifolate sensitive strain. Chloroquine, cycloguanil and pyrimethamine were used as reference drugs or positive controls in the antiplasmodial assays. DMSO (0.1%) and distilled water were used as controls to rule out the solvent effects on the bioassay results of the test samples. All the experiments were performed three times in duplicate (3x2).

3. **Results and Discussions**

Given the traditional use of *C. dubia* to treat fever and various disorders bearing relevance to malaria and microbial infections and the evidence that the crude alkaloid extracts exhibited significant antiplasmodial activity (Wangchuk et al., 2011), we investigated the bioactive alkaloid constituents of this plant. Out of eight alkaloids (compounds **1-8**, Fig. 1) isolated from this plant, we examined compounds **1**, **2**, **5** and **6** for their antiplasmodial, cytotoxicity and antimicrobial and activities (Table 1).

Table 1

Antiplasmodial activity (IC₅₀ in µg/mL) of the pure compounds **1**, **2**, **5** and **6** isolated from *C. dubia*.

Samples	Antiplasmodial		Cytotoxicity		a original activity taken from Wangchuk et al. (2011). b original activity taken from Wangchuk et al. (2010). c original activity taken from Wangchuk et al. (2012). d Reference
	TM4/8.2	K1CB1	Vero cells	KB cells	
MeOH extract	6.38 ± 0.62 ^a	9.50 ± 0.84 ^a	>10 ^a	>10 ^a	
CHCl ₃ alkaloid extract	2.21 ± 0.43 ^a	2.84 ± 0.32 ^a	>10 ^a	>10 ^a	
Dubiamine (1)	>20	>20	>20	>20	
Scoulerine (2)	1.78 ± 0.72	1.04 ± 0.08	>20	>20	
Cheilanthifoline (3)	0.90 ± 0.27 ^b	1.22 ± 0.84 ^b	Not tested	Not tested	
Protopine (4)	1.45 ± 0.53 ^c	1.38 ± 0.31 ^c	>3.5 ^c	>3.5 ^c	
Capnoidine (5)	>20	>20	>20	>20	
Bicuculline (6)	>10	>10	>10	>10	
Chloroquine ^d	0.010	0.089			
Cycloguanil ^d	0.009	0.810			
Pyrimethamine ^d	0.020	7.700			
Ellipticine ^e			0.093		
Doxorubicin ^e				0.56	

drugs for antiplasmodial activity.

^e Reference drugs for cytotoxicity activity.

Interestingly, compound **2** demonstrated significant *in vitro* antiplasmodial activity against the *P. falciparum* strains: TM4/8.2 (a wild type chloroquine and antifolate sensitive strain) and K1CB1 (multidrug resistant strain) with IC₅₀ values of 1.78 µg/mL (5.47 µM) and 1.04 µg/mL (3.1 µM), respectively. Compounds **3** and **4** were isolated from different plant species and their antiplasmodial activities were determined earlier by us (Wangchuk et al., 2010; Wangchuk et al., 2012) and are included in this study (Table 1). Compounds **2**, **3**, and **4** exhibited similar antiplasmodial activity. However, the activities of compound **2** were found to be almost two fold better than the CHCl₃ crude alkaloid extract (2.21 and 2.84 µg/mL against TM4/8.2 and K1CB1 strains, respectively); seven fold

better than the crude MeOH extract (6.38 µg/mL for TM4/8.2 and 9.50 µg/mL for K1CB1 strains); and eight fold better than the conventional antimalarial drug pyrimethamine (reference drug) against the K1CB1 strain (Table 1). Although the antiplasmodial activity of scoulerine (**2**) does not meet the criteria of the medicines for malaria venture (MMV), it represents a potential new scaffold based on which the improved antimalarial lead compounds can be drawn, particularly as it did not show mammalian cell line (KB and Vero cells) toxicity (Table 1) and has not been previously reported. Scoulerine has been reported to exhibit other activities such as: anti-topoisomerase I (Cheng et al., 2008), GABA_A receptor agonist (Eisenreich et al., 2003), and antiemetic, antitussive and antibacterial activities (Taylor and Francis, 2011).

Compounds **1**, **5** and **6** did not show antiplasmodial activity even at the highest tested concentration of 10-20 µM. Due to the smaller quantities of compounds **7** and **8**, their biological activity studies were not carried out.

Since compounds **2**, **3** and **4** have shown significant antiplasmodial activities and that protopine (**4**) was the major alkaloid present, it may be deduced that these three alkaloids (alone or synergistically) may be responsible for the significant activity of the CHCl₃ crude alkaloid extract of this plant. The antimicrobial assay of the pure alkaloids (**1-6**) did not identify any compounds with significant activity against *Escherichia coli*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *MRSA*, *Bacillus subtilis*, *Vibrio cholera* and *Candida albicans* with all showing MIC values greater than 250 µg/mL.

The significant antiplasmodial activities of the crude extracts and the isolated three pure compounds **2**, **3** and **4** provide support for the use of *C. dubia* in BTM for treating fever associated with malaria and other infections of the blood, liver and bile. Thus, our study verified the ethnomalarial uses of the plant.

4. Conclusions and future directions

In summary, our study found the following: a) a new natural product (**1**) was isolated which we named as dubiamine and its structure was confirmed from its single crystal X-ray structure b) compound **2** showed significant *in vitro* antiplasmodial activity without mammalian cell toxicity and was identified as a potential antimalarial hit scaffold c) the *in vitro* antiplasmodial activity of the crude extracts and the pure compounds **2**, **3** and **4** is commensurated with the ethnopharmacological uses of this plant and thus substantiated its usage in BTM for the aforementioned disorders. However, it may be premature to claim that the plant extract formulated and administered according to traditional protocols would be safe and effective for treating humans. Therefore, *in vivo* animal model of malaria using methods of traditional preparation and administration would more confidently support the use of the plants in treating humans. As resistance to the front line antimalarial drug, artesunate, appears to be emerging (Dondorp et al., 2009), the above *in vivo* experiments on crude extracts and also on compound **2**, **3** and **4** which showed significant antiplasmodial activity against antifolate sensitive and multidrug resistant strains are essential and will be possibly carried out in future.

Acknowledgements

Financial support has been provided by the government of Australia in a form of an Endeavour Postgraduate Award to PW. Plant materials have been obtained from the Pharmaceutical and Research Unit, Ministry of Health, Thimphu, Bhutan. We are grateful to Dr. Malai Taweechotipatr, Srinakharinwirot University, Bangkok, Thailand for the antimicrobial testing. SK is an international Research Scholar of the Howard Hughes Medical Institute, USA.

References

- Abou-Donia, A.H., El-Masry, S., Saleh, M.R.I., Phillipson, J.D., 1980. Alkaloids from *Fumaria judaica*. *Planta Medica* 40, 295-298.
- Anonymous., 2008. Guidelines for Identification and Collection of Medicinal Plants in Bhutan. Department of Agriculture, Ministry of Agriculture, Thimphu, Bhutan, pp.25-28.
- Basha, S.A., Mishra, R.K., Jha, R.N., Pandey, V.B., Singh, U.P., 2002. Effect of berberine and (±)-bicuculline isolated from *Corydalis chaerophylla* on spore germination of some fungi. *Folia Microbiologica* 47, 161-165.
- Basnet, P., Kadota, S., Li, J.X., Wu, C.Z., Namba, T., 1993. (-)-Corydecumbine, a novel phthalideisoquinoline alkaloid from *Corydalis decumbens*. *Heterocycles* 36, 2205-2208.
- Blasko, G., Gula, D.J., Shamma, M., 1982. The phthalideisoquinoline alkaloids. *Journal of Natural Products* 45, 105-122.
- Brochmann-Hanssen, E., Nielson, B., 1966. The isolation of (-)-scoulerine from opium. *Tetrahedron Letters* 20, 2261-2263.
- Cheng, X., Wang, D., Jiang, L., Yang, D., 2008. DNA Topoisomerase I inhibitory alkaloids from *Corydalis saxicola*. *Chemistry and Biodiversity* 5, 1335-1344.
- Dondorp, A.M., Nosten, F., Yi, P., Das, D., Phyto, A.P., Tarning, J., Lwin, K.M., Ariey, F., Hanpithakpong, W., Lee, S.J., Ringwald, P., Silamut, K., Imwong, M., Chotivanich, K., Lim, P., Herdman, T., An, S.S., Yeung, S., Singhasivanon, P., Day, N.P.J., Lindergardh, N., Socheat, D., White, N.J., 2009. Artemisinin resistance in *Plasmodium falciparum* malaria. *The New England Journal of Medicine* 361, 455-467.
- Edwards, O.E., Handa, K.L., 1961. The alkaloids of *Corydalis govaniiana*. *Canadian Journal of Chemistry* 39, 1801-1804.
- Eisenreich, W.J., Hofner, G., Bracher, F., 2003. Alkaloids from *Croton flavans* L. and their affinities to GABA-receptors. *Natural Product Research* 17, 437-440.

- Elango, V., Freyer, A.J., Blasko, G., Shamma, M., 1982. The NMR spectra and conformations of the phthalideisoquinolines. *Journal of Natural Products* 45, 517-522.
- Grierson, A.J.C., Long, D.G., 1984. *Flora of Bhutan: Including a Record of Plants from Sikkim*, vol. 1, part 2. Royal Botanic Garden, Edinburgh.
- Haisova, K., Slavik, J., 1973. Alkaloids of the Papaveraceae. II. On alkaloids from *Argemone ochroleuca* Sweet. *Collection of Czechoslovakia Chemical Communication* 38, 2307-2312.
- Inubushi, Y., Aoyagi, Y., Matsuo, M., 1969. The enzymatic oxidation of isoquinoline alkaloids. *Tetrahedron Letters* 28, 2363-2366.
- Kametani, T., Ohkubo, K., 1965. The structure of sendaverine and its total synthesis. *Tetrahedron Letters* 48, 4317-4326.
- Okamoto, Y., Tanaka, S., Kitayama, K., Isomoto, M., Masaishi, M., Yanagawa, H., Kunitomo, J., 1971. Studies of the alkaloids of Menispermaceous plants. CCLXI. Alkaloids of *Menispermum dauricum* DC. *Yakugaku Zasshi* 91, 684-687.
- Pandey, G.D., Tiwari, K.P., A new total synthesis of sendaverine. *Indian Journal of Chemistry* 19, 160-161.
- Pilar, M.D., Soriano, C., Shankaraiah, N., Santos, L.S., 2010. Short synthesis of noscapine, biculline, egenine, capnoidine, and corytensine alkaloids through the addition of 1-siloxy-isobenzofurans to imines. *Tetrahedron Letters* 51, 1770-1773.
- Ribar, B., Meszaros, C., 1991. Structure of capnoidine, a phthalideisoquinoline alkaloid. *Acta Crystallographica C* 47, 2191-2193.
- Rucker, G., Breitmaier, E., Zhang, G.L., Mayer, R., 1994. Alkaloids from *Dactylicapnos torulosa*. *Phytochemistry* 36, 519-523.

- Seger, C., Sturm, S., Strasser, E.M., Ellmerer, E., Stuppner, H., 2004. ^1H and ^{13}C NMR signal assignment of benzyloquinoline alkaloids from *Fumaria officinalis* L. (Papaveraceae). *Magnetic Resonance in Chemistry* 42, 882-886.
- Suau, R., Silva, M.V., Ruiz, I., Valpuesta, M., 1994. N-Benzyloquinoline alkaloids from *Ceratocarpus heterocarpus*. *Phytochemistry* 36, 241-243.
- Takahashi, H., Iguchi, M., Onda, M., 1985. Utilization of protopine and related alkaloids. XVII. Spectroscopic studies on the ten-membered ring conformations of protopine and α -Allocryptopine. *Chemistry and Pharmaceutical Bulletin* 33, 4775-4782.
- Taylor and Francis, 2011. *Dictionary of natural products*. CHEMnetBase, DNP 19.2.
Chemical Database Online at: <http://www.chemnetbase.com.ezproxy.uow.edu.au/>.
- Wangchuk, P., Bremner, J.B., Samten, Rattanajak, R., Kamchongpaisan., 2010. Anti-plasmodial agents from the Bhutanese medicinal plant *Corydalis calliantha*. *Phytotherapy Research* 24, 481-485.
- Wangchuk, P., Keller, P.A., Pyne, S.G., Taweechotipatr, M., Tonsomboon, A., Rattanajak, R., Kamchonwongpaisan, S., 2011. Evaluation of an ethnopharmacologically selected Bhutanese medicinal plants for their major classes of phytochemicals and biological activities. *Journal of Ethnopharmacology* 137, 730-742.
- Wangchuk, P., Keller, P.A., Pyne, S.G., Sastraruji, T., Taweechotipatr, M., Tonsomboon, A., Rattanajak, R., Kamchonwongpaisan, S., 2012. Phytochemical and biological activity studies of the Bhutanese medicinal plant *Corydalis crispa*. *Natural Product Communications* (in press).
- Wangchuk, P., Samten., Ugyen, Thinley, J., Afaq, S.H., 2008. High altitude medicinal plants used in Bhutanese traditional medicine (*g.so-ba-rig-pa*). *Ethnobotany* 20, 54-64.
- Yeola, S.N., Mali, R.S., 1984. Convenient total synthesis of (\pm)-tetrahydrogroenlandicine. *Indian Journal of Chemistry* 23, 818-820.