



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers

Faculty of Science, Medicine and Health

2013

Anthropogenic biogeochemical impacts on coral reefs in the Pacific Islands-An overview

R.J. Morrison

University of Wollongong, johnm@uow.edu.au

G.R.W. Denton

University of Guam

U bale Tamata

University of the South Pacific, ubt16@uow.edu.au

J Grignon

Griffith University

Publication Details

Morrison, R. J., Denton, G., Tamata, U. Bale. & Grignon, J. (2013). Anthropogenic biogeochemical impacts on coral reefs in the Pacific Islands-An overview. *Deep-Sea Research Part 2: Topical Studies in Oceanography*, 96 5-12.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Anthropogenic biogeochemical impacts on coral reefs in the Pacific Islands-An overview

Abstract

Coral reefs dominate the coastal environment in many Pacific Islands, being present as atolls, coral platforms, barrier and fringing reefs. With ever increasing populations and migration of people to the coast, the anthropogenic impacts on these reefs have increased dramatically in the last 30 years. While research on these impacts has been limited, some important progress has been made. This paper reviews some of the completed studies, with outcomes from American Samoa, Fiji, French Polynesia, Guam, Saipan, New Caledonia and Tonga presented. These studies indicate that the most significant impacts have been found in locations close to major urban centres or industrial and mining activities. The extent of impact varies from place to place with minimal impacts in the more isolated and less industrialised communities. Common anthropogenic impacts are contamination caused by inadequate sewage treatment, erosion from adjacent agricultural and urban expansion activities, poor waste management, eutrophication, inefficient and/or inappropriate pesticide use and hydrocarbons use, storage and management. The outcomes include contaminated sediments (trace metals, pesticides, PCBs, hydrocarbons) with some impacts on resident biota. In some instances, the edible quality of local fisheries resources has been significantly compromised. Even in locations with small populations, increasing populations and poor economic conditions have resulted in noticeable effects on the adjacent fringing reefs, including dramatic algal proliferation and declines in fish numbers resulting from increasing nutrient discharges and increased herbivore fish catches. Recovery measures including fishing bans and alternative fishing practices have been implemented to address these issues in some areas.

Keywords

pacific, islands, reefs, overview, coral, anthropogenic, biogeochemical, impacts

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Morrison, R. J., Denton, G., Tamata, U. Bale. & Grignon, J. (2013). Anthropogenic biogeochemical impacts on coral reefs in the Pacific Islands-An overview. *Deep-Sea Research Part 2: Topical Studies in Oceanography*, 96 5-12.

Anthropogenic Biogeochemical Impacts on Coral Reefs in the Pacific Islands – An Overview

¹R. J. Morrison, ²G.R.W. Denton, ³U. Bale Tamata and ⁴J. Grignon

¹GeoQuEST Research Centre, University of Wollongong, NSW2522, AUSTRALIA

²Water and Environment Research Institute, University of Guam, GUAM

³Institute of Applied Sciences, University of the South Pacific, Suva, FIJI

⁴International Ecotourism Research Centre, Griffith University, Gold Coast, AUSTRALIA

ABSTRACT

Coral reefs dominate the coastal environment in many Pacific Islands, being present as atolls, coral platforms, barrier and fringing reefs. With ever increasing populations and migration of people to the coast, the anthropogenic impacts on these reefs have increased dramatically in the last 30 years. While research on these impacts has been limited, some important progress has been made. This paper will review some of the completed studies, with outcomes from American Samoa, Fiji, French Polynesia, Guam, Saipan, New Caledonia and Tonga presented.

These studies indicate that the most significant impacts have been found in locations close to major urban centres or industrial and mining activities. The extent of impact varies from place to place with minimal impacts in the more isolated and less industrialised communities. Common anthropogenic impacts are contamination caused by inadequate sewage treatment, erosion from adjacent agricultural and urban expansion activities, poor waste management, eutrophication, inefficient and/or inappropriate pesticide use and hydrocarbons use, storage and management. The outcomes include contaminated sediments (trace metals, pesticides, PCBs, hydrocarbons) with some impacts on resident biota. In some instances, the edible quality of local fisheries resources has been significantly compromised.

Even in locations with small populations, increasing populations and poor economic conditions have resulted in noticeable effects on the adjacent fringing reefs, including dramatic algal proliferation and declines in fish numbers resulting from increasing nutrient discharges and increased herbivore fish catches. Recovery measures including fishing bans and alternative fishing practices have been implemented to address these issues in some areas.

INTRODUCTION

South Pacific Region/Coral Reefs

Coral reefs are a dominant feature of the Pacific Islands. They are unique to tropical and certain subtropical oceans since the reef-building organisms require water temperatures in excess of 22°C. Coral structures include atolls (essentially reefs of variable thickness built up by corals (and other organisms) resting on a volcanic base) and reef platforms having an elevation generally less than 5 m above mean sea level. In the central and south Pacific, some countries consist entirely of low elevation coral structures, e.g., Tuvalu, others contain atoll groups, e.g., the Cook Islands, and some

countries consist of mainly volcanic islands with a few isolated atolls, e.g., Ontong Java in the Solomon Islands. Fringing and barrier reefs are common around 'high' islands (those having a volcanic or other non-calcareous features rising to elevations above 50 m (Cumberland, 1956).

Most major population centres in the Pacific Islands have evolved in coastal locations that afforded some form of protection against storms where shipping is the main mode of transport. Many of these centres are located in bays protected by barrier reefs, by islands or by riverine deltas. However, since these geomorphological features also have the effect of limiting, to some extent, mixing of near-shore and open ocean waters (Viles and Spencer, 1995), they tend also to facilitate contaminant accumulation in the near-shore waters with which most people commonly interact.

This paper reviews various studies that have been carried out on investigation of the impact of anthropogenic activities on coral reefs and related water bodies in the Pacific Islands (Figure 1). It is not meant to be comprehensive, rather it uses case studies to illustrate the main issues that are currently of greatest concern. The sites referred to are, for the most part, major urban centres (Saipan, Guam, Suva, Fanga'uta), although one site (Coral Coast, Fiji) examines impacts of a growing population in a rural area, while another (Tutuila) compares urban and isolated rural areas around one island. Some important work completed in the French territories in the Pacific is also discussed.

CASE STUDIES

Western Coast, Saipan, CNMI

Saipan (15°12'N, 145°43'E)(Figure 1) is the major population centre of the Commonwealth of the Northern Mariana Islands (CNMI). The resident population is nearing 60,000, and there is a substantial influx of tourists, mainly from SE Asia and Japan. Saipan has been a shipping centre in Micronesia for about 400 years, but major increases in shipping activity have taken place since the 1940s during and following World War II. Tanapag Lagoon is a typical high-island barrier reef lagoon bordering the western shore of central Saipan, and adjacent to the main port area on that island. It is about 9 km long and 3 km at its widest point, and covers an area of about 13 km². Large expanses of patch reef, interspersed among sand and rubble, provide for a diversity of shallow water habitats and harbour rich assemblages of flora and fauna (Doty and Marsh 1977, Amesbury *et al.* 1979). In addition to its ecological significance, the Lagoon supports a variety of recreational activities, and local people traditionally harvest many fisheries resources for food.

Over the last quarter century, the southern, nearshore fringing reef section of Tanapag Lagoon has become heavily impacted by human activities. Primary sources of anthropogenic disturbances between Muchot Point and Flores Point, a distance of approximately 3 km, include a commercial port (Saipan Harbour) and bulk fuel facility, a sewer outfall, a municipal waste dump, and two small-boat marinas. The area is also heavily inundated by stormwater runoff during prolonged periods of wet weather. Several studies have examined the impacts on these human activities on the sediments and biota in the Lagoon (e.g., Denton et al, 2006a, 2009).

Sediments have been sampled adjacent to the shoreline and further offshore, up to 2250 m. Analysis for a range of trace metals showed that offshore sediments were not significantly contaminated, but there was evidence of accumulation of anthropogenic metals, especially Cu, Pb, Zn, and to a lesser degree Cd, Ni, Cr, Hg, in nearshore waters around the old municipal dump and adjacent to boat marinas. In general, metal concentrations decreased on moving seaward and/or away from identified sources of metal contamination. PCB analyses showed all offshore sites were relatively free of contamination (concentrations < 1 ng/g); relatively high values were found adjacent to the old dump (16.6 ng/g) and in the area around the port (8-11 ng/g). Detailed studies of individual congener profiles indicated that Aroclor 1260 and possibly 1254 were the main PCB sources in Tanapag Lagoon, probably leaked from electrical transformers. PAH concentrations were also very low in offshore sites (<0.5 µg/g), but nearshore sites showed higher values (e.g., 2.4 µg/g near the port, 3.2 µg/g near the old dump). The PAH profiles were dominated by higher molar mass compounds, but the actual source (e.g., combustion or aged petrochemical spills) could not be confirmed. Overall, the sediments in Tanapag Lagoon were considered relatively clean apart from problems around the main port, the old waste dump site and the small boat marina (Denton et al, 2006a).

In two related studies, dominant biotic representatives were sampled from various sites within the Lagoon (Denton et al, 2009, 2010) and analysed for trace metals. Preference was given to species traditionally harvested as food by local residents or considered to have bioindicator potential. Many organisms (e.g., algae, seagrass, bivalves) were found to be enriched in some trace metals (Cr, Cu, Pb, Zn) at one or more contaminated sites identified at the southern end of the lagoon (dry dock, old dump, boat marina). From a human health perspective only lead and copper were identified as elements of concern and only in bivalves from the old dump area (Denton et al, 2009). Mercury levels in axial muscle of over 300 popular table fish (65 species) from the Lagoon proper were all well below the current U.S. Food and Drug Administration guideline of 1.0 µg/g wet weight (US FDA 1998) with 84% of the overall catch yielding values below 0.01 µg/g wet weight (Denton et al, 2010). While Hg concentrations were somewhat elevated in representatives from around the port area and municipal dump, the highest levels encountered were generally found in specimens captured some distance from obvious sources of pollution known at the time.

A follow-up investigation traced the mercury back to an old incinerator site at the Commonwealth Health Center, Saipan's only public hospital (Denton et al, 2011). The incinerator was used for the destruction of medical wastes from the hospital and other medical clinics on island. It was eventually shut down for multiple violations of the Clean Air Act in 2006 (US EPA, 2005), just 18 months after the fish survey had been completed. Mercury levels in certain species from the impacted area were measured again in 2007 (Denton et al, 2011) and found to be substantially lower than those recorded during the earlier survey

Pago Gay, Guam

Guam (13°28'N, 144°45'E) has been dramatically impacted by human activities since World War II and has undergone considerable economic growth and urban expansion over the last 30 years. It also has a thriving tourist industry that currently attracts over

one million visitors annually. Today, the island is poised on the brink of another economic boom associated with the relocation of 18,000 U.S. marines and their dependants to Guam from Okinawa, Japan, in 2014. The population, which presently stands at around 185,000, is expected to approach 250,000 by 2020 as a direct result of this military build-up. Environmental problems currently experienced on the island are largely associated with solid and hazardous waste disposal, storm water runoff, and the treatment and disposal of domestic and industrial wastewaters.

Guam has been the major shipping centre in Micronesia for about 400 years. Trace metals, PCBs and PAHs in sediments and biota have been investigated in the island's major port and smaller boat harbours (Denton et al. 2005, 2006b-e). Relatively few studies, however, have examined the impacts of Guam's growth and development on the coastal environment. This is particularly so on the western side of central Guam where the greatest intensity of commercial and industrial activities exists. Pago Bay on the eastern side of the island has been subjected to far less direct human intervention, although the possible impact on its fisheries of a poorly managed landfill, in the nearby village of Ordot, has been of long-standing concern. The landfill is located in the catchment of the Longfit/Pago River system that drains into the southern half of Pago Bay, and has been releasing heavy metal enriched leachate into the watershed for over 60 years

Recent studies have identified mild to moderate heavy metal contamination in Lonfit River surface waters and sediments immediately adjacent to the landfill but no measurable contamination in either water or bottom sediment matrices further downstream (Denton et al, 2007, 2008). Likewise, there were no abnormal metal levels in aquatic organisms anywhere within the watershed itself (Denton et al. 2007), or in the Bay (Denton and Morrison 2009). It was concluded that soluble metals released from the landfill during dry weather (low flow conditions) are rapidly sequestered by aluminium and iron oxyhydroxides present in streams and subsequently deposited in the beds of tributaries and at the confluence of the Longfit River. In subsequent heavy rains, the contaminant laden sediments are scoured and carried away from the river to offshore deepwater beyond the reef front, via the Pago Bay channel, a flooded river valley. This natural cleansing process prevents metals discharged from the landfill from accumulating in the watershed and in the Bay. Any residual contamination from the landfill that accumulates in and around the river mouth is periodically flushed from the system by major storms (typhoons) that frequently impact the eastern side of the island. While these studies clearly demonstrated that the Lonfit/Pago River system and Pago Bay are not permanent sinks for metal contaminants leached from the landfill, they did reveal some mild Pb enrichment in sediments and bivalves near an old military firing range at the southern end of the Bay. Marginally elevated levels of Zn and Hg were also identified in both media near sources of ground water intrusion, septic-tank leachate and urban runoff at the northern end of the fringing reef (Denton and Morrison, 2009). Nevertheless, the impacts of these metal sources on the edible quality of aquatic resources in these areas were considered inconsequential.

Suva Lagoon, Fiji

Suva, the capital of the Fiji Islands, located on the southeast corner of Viti Levu at 18°06'S, 178°30'E (Figure 1), is the major commercial, shipping and industrial centre

in the South Pacific islands. The city has rapidly expanded in the last 50 years with significant population growth (50,000 to >200,000) in the catchment of the adjacent lagoon over the last 30 years. Suva Lagoon is located within a well developed barrier reef system and is made up of two parts – Laucala Bay on the east and Suva Harbour on the west. Significant small industry development around the shoreline has occurred in recent years including ship-building and repair, metal manufacturing, food processing and packaging, oil storage and handling, in addition to activities associated with the operations of Suva port. Studies on pollution in Suva lagoon have been undertaken by various researchers over the past 30 years, and the major findings are summarised in Morrison et al (2006).

Nutrient data for Suva Lagoon span a period of about 25 years. The data sets are not consistent in terms of methods used and the parameters measured (total element versus reactive components). In some instances quality control procedures have also been questioned. Nevertheless, some valid conclusions can be drawn from the available data. For example, dissolved inorganic nitrogen (DIN) and dissolved inorganic phosphorus (DIP) concentrations are often high compared to open ocean values, with reported values ranging from 50-450 $\mu\text{mol/L}$ for DIN and 8-12 $\mu\text{mol/L}$ for DIP. Values for both parameters vary with tide, season, amount of runoff (wet weather) and source. Singh et al (2009) considered sewage discharges to be a major source of nutrients in dry periods, while rivers were considered to be the dominant source during wet weather when high flows carry sewage material from small treatment plants and septic overflows as well as diffuse land-based source materials. At sites near the barrier reef, nutrient concentrations are about DIN < 1 $\mu\text{mol/L}$, DIP ~ 0.1 $\mu\text{mol/L}$, values close to those in the open ocean. There is evidence, however, that despite these low average concentrations near the barrier reef, algal growth has been increasing in recent years (including Sargassum that was not found in the 1970s and 1980s) (L.F. Zann, personal communication).

Sediment trace metal studies in Suva Lagoon have identified 2 major sources - riverine non-carbonate materials and reef derived carbonate materials (Morrison et al, 2001, Fernandez et al, 2006) with the carbonate influence increasing on moving towards the barrier reef. On top of these variations, concentrations of metals of concern for health are generally higher in nearshore waters (Morrison et al, 2001), especially around 'hot spot areas', such as, the Lami dump, closed in 2006 (Naidu and Morrison, 1994), Lami industrial area (Gangaiya et al, 2001), and the Walu Bay industrial area (Pb, Sb and Zn: Naidu and Morrison, 1994). Suva Lagoon sediments have been reported to have significant TBT contamination, including the highest reported TBT values in the world (92-360 $\mu\text{g/g}$, Stewart and De Mora, 1992, Maata and Koshy, 2001). However, monobutyl tin (MBT)/dibutyl tin (DBT) ratios are generally below 0.1, indicating that little or no additions of TBT in recent years (Maata and Koshy, 2001).

Some data exist for organic contaminant in Suva lagoon, with emphasis organochlorines and PCBs. The highest concentrations found in Suva lagoon sediments (up to 14 ng/g DDT and metabolites, 8 ng/g dieldrin and 30 ng/g total PCBs) are relatively low compared with other parts of the world (Morrison et al, 1996). Organochlorine pesticides were used until 1980, for mosquito control in the city and for agricultural purposes in adjacent farming areas. PCBs were components of transformers used in the old electricity generating facility adjacent to Nabukalou Creek

in the centre of Suva city. There is no data for any organochlorinated compounds in shellfish or other biota from Suva lagoon.

The disposal and untreated sewage into Suva Lagoon presents a constant challenge to environmental managers from a human health perspective. Typically, the hygienic quality of recreational waters is monitored using faecal indicator bacteria. In Fiji, it is recommended that faecal coliform (FC) counts should be less than 400 colony forming units (CFUs) per 100 mL water and less than 2 FC/g flesh wet weight for shellfish consumption. Data for Suva lagoon show faecal coliform (FC) densities ranging from 0 to >10,000 CFUs/100 mL with sites around Kinoya, Nubukalou Creek and Lami showing the highest values and minimal change over the period 1978-2005 (Morrison et al, 2006). FC contents in mangrove oysters (*Crassostrea mordax*) taken from Suva Lagoon range from 0.7-24,000 FC/g (Morrison et al, 2006). The potential health risks associated with eating raw shellfish from Suva lagoon can be high, although less so if shellfish are depurated, marinated or cooked appropriately.

Overall, the contaminant data from Suva lagoon is indicative of relatively low levels of contamination by trace metals, persistent organic pollutants, apart from a few localised hotspots. Identifying problem areas is unfortunately hampered by the absence of any management plan for the Lagoon or its catchment, due to the fact that while legislative/administrative powers for pollution control are distributed among several agencies, interagency interactions are often limited.

Coral Coast, Fiji

The Coral Coast region is located on the south coast of Viti Levu (Figure 1), the main island of the Fiji group. It has a stretch of about 60 km of coastline where fringing reefs lie adjacent to a high island. Traditionally, its population consisted of a number of coastal villages with small populations and subsistence lifestyles. Since the 1960s, this has been one of the fastest developing regions in Fiji, mainly as a result of tourism activities. This has led to increases in village populations with more intensive fishing and reef gleaning and more intensive root crop farming, chicken and pig production. These changes have contributed to increased nutrient concentrations (up to >10 $\mu\text{mol/L}$ average DIN, > 1 $\mu\text{mol/L}$ average DIP) in coastal waters (Tamata, 2007) sourced from sewage and animal sources particularly during wet weather (confirmed by stable isotope studies). For example, at a Valase site undergoing development, dry weather concentrations averaged relatively low values (nitrate 0.07 $\mu\text{mol/L}$, phosphate 0.10 $\mu\text{mol/L}$). while average wet weather concentrations were much higher (nitrate 6.06 $\mu\text{mol/L}$ and phosphate 0.37 $\mu\text{mol/L}$). The herbivore fish population was decreased due to overfishing and this resulted in extensive growth of Sargassum and some filamentous algal species that led to corals in some parts of the reef system being overwhelmed (Tamata, 2007).

Alarmed by these changes in their reef systems in terms of food supply and tourism, local communities solicited assistance from local scientists. The scientists carried out investigations, including grazing exclusion trials that clearly revealed the impact of herbivory in Sargassum control. This finding led to the establishment of locally managed marine protected areas under local community control with *tabu* (bans) on fishing, in certain areas, especially for herbivorous species. The local community has also developed strategies for improved management of nutrient sources, including

improved solid waste management practices, replanting of buffer strips of vegetation along river banks, relocating piggeries away from river banks, and the construction of wetlands for treatment of wastewater (Tamata, 2007)

Another important outcome is the active participation of local community members in monitoring fisheries stocks through the catch per unit effort (CPUE) information recording, which is then passed on to the scientists for analysis. Community-coordinated re-afforestation of catchment areas, and mangrove replanting along the coastline have emerged along the Coral Coast. In these initiatives scientists play a facilitation and advisory role to the communities.

Fanga'uta Lagoon, Tonga

Fanga'uta Lagoon is the major coastal water body of Tongatapu (Figure 1), the main island of the Tonga group located at 21°12'S, 175°12'W. The island of Tongatapu is a raised coral platform covered with soils derived from volcanic ash and corals. Fanga'uta Lagoon's catchment has a resident population of about 40,000 people, including a significant portion of the capital, Nuku'alofa. The lagoon has suffered many anthropogenic impacts including substantial coastal modification, vegetation and mangrove removal, surface runoff from roads and farmland, sedimentation, polluted groundwater inputs, overfishing, waste dumping, and a general lack of appreciation of ecological benefits of the system (Kaly and Morrison, 2005).

Anthropogenic impact investigations were carried out in 1983-4, 1988-9, 1992, and 1998-2001. Bottom sediments were mixtures of coralline and volcanic-soil derived materials (Morrison and Brown, 2003). There was no evidence of any trace metal contamination of sediments or shellfish (Morrison and Brown, 2003). Low levels of organochlorine pesticides and PCBs were found in 1992 in lagoon sediments (Harrison et al, 1996) close to a known illegal dumping site. In a complementary study in 2000, low concentrations of chlorfluazuron and flusilazole found in sediments, probably originating from use in nearby agricultural fields (Chen et al, 2000).

In the late 1990s, a significant decline in seagrass cover, with increased epiphyte growth relative to the 1980s was observed (e.g., see Kaly, 1998). In 1992, the lagoon water was transparent with coral and seagrass communities visible to depths of at least 2 m. In 1993, after a period of occasional algal blooms, the lagoon suddenly and permanently turned green, and has stayed that way until the present time. *Caulerpa* and *Halimeda* algal species increased in abundance and increased turbidity was measured. Fish and shellfish catches, in terms of both total mass and sizes of individuals declined, with the continuing (often illegal) removal of mangroves for housing believed to be contributing to major habitat loss for juvenile fish. Increasing urbanisation, some of which occurred in cleared mangrove areas, along with poor building standards control were considered contributing problems, e.g., inappropriate septic tank location was contributing to nutrient runoff and eutrophication in the lagoon.

Based on the outcomes of scientific studies and community consultations, a full management plan for the lagoon was prepared (Prescott et al, 2001), but this has not yet been implemented to any extent due to financial constraints.

Tutuila, American Samoa

Tutuila, located at 14°17'S, 170°41'W (Figure 1), is the main island of the American Samoa group and is the site of Pago Pago, the capital and main port. Tutuila is globally a remote location, distanced from any major global sources of pollution, unless pollutants have been transported atmospherically over long distances (Peshut, 2009). At this remote location, trace elements in sediments would be expected to reflect naturally occurring background conditions for volcanic high island sites not influenced by anthropogenic impacts. Tutuila is a rugged volcanic island dominated by olivine basalts (Stearns, 1944; Staudigel et al., 2006). The coastline (200 km long) is extremely irregular with numerous small open bays, which lack bars or barrier reefs, and are subject to the sea conditions of exposed coasts. Pago Pago Harbour is one of the few protected areas and maintains most of its natural shoreline, except for some filled areas near the main port, which have been developed for commercial shipping activities and two fish canneries. Tutuila has no estuaries and most of the coastline is rocky with abrupt elevation changes immediately above breaking waves. Surface waters are limited to a few dozen perennial streams, most of which have short, steep reaches, and typically low base flows. Coral reefs are the dominant marine habitat for Tutuila near-shore waters, with 60% of the coastline occupied by narrow fringing reefs.

As part of a larger ecological investigation (Peshut, 2009), levels of sedimentary trace metals (As, Cd, Cu, Fe, Hg, Mn, Ni, P, Sr, Zn) were examined at six study sites on Tutuila. Five of these sites were located in near pristine coastal areas while one (Loa) was located in Pago Pago Harbour, the most human-impacted zone in Tutuila. Sediment samples from all sites were dominated with carbonates (aragonite and calcite) with minor contributions of quartz, kaolinite, illite and chlorite). There was no evidence of trace element contamination at any sites (Morrison et al, 2010). Indeed, the observed variations could all be explained by assuming the sediments were predominantly of coralline origin, mixed with varying amounts of basalt-derived material. Strong correlations (coefficients >0.8) were found between the main basaltic elements (Al, Fe, Mn, K, Ti, Si, P) all of which showed strong negative correlations with Ca. Two notable features among the basalt derived elements were: (1) Cu showed effectively no strong correlations with any other element examined (not even Zn, as has been found at other similar Pacific locations - Morrison et al., 1997, Denton et al., 2005b); (2) As showed strong correlations with 9 other elements (positive with Al, Fe, K, Mn, Ti, Zn, P, Si and negative with Ca) which again is different from some other Pacific sites, where As usually shows few correlations.

Surprisingly, the elemental data for the Loa samples taken in Pago Pago Harbour showed no clear differences from the other sites, with the exception of Hg (total) which was 3-9 times higher than elsewhere in the study area. Inter-site differences for MeHg were smaller, however, with variation factors of 1-3 (Peshut, 2009). While there are limited sources of pollution in Tutuila, some activities in the harbour are potential sources of contamination (ship repair and maintenance, cargo and fuel spillage, fish cannery effluents, illegal dumping). The absence of any metal pollution indication (at Loa) is inconsistent with results from Suva, Fiji (Naidu and Morrison, 1994), Apra Harbour, Guam (Denton et al, 2005) and Tanapag Lagoon, Saipan, CNMI (Denton et al, 2009), where significant metal pollution of harbour sediments has been observed. The lack of metal contamination at Loa may simply relate to the lack of industrial

activity, as compared to other small Pacific Island harbour sites where economic development is significantly greater relative to Tutuila.

French Territories in the South Pacific

The French Pacific territories exhibit a wide diversity of island types including continental islands (New Caledonia), archipelagos (French Polynesia) and isolated atolls (Gardes and Salvat, 2008). The majority of reefs in these territories are considered healthy (Salvat *et al.*, 2008), but increasing anthropogenic activity, e.g., mining, fisheries, tourism, is leading to significant impacts in some areas, especially near urban centres. Nuclear experiments carried out in French Polynesia from 1966-1996 are also of interest in terms of environmental impacts.

a. New Caledonia

Nickel mining and the related metal-processing industry constitute both a threat to coastal and marine biodiversity in New Caledonia (22°15'S, 166°26'E), and the driving force supporting the economy of the country (Pascal *et al.*, 2007, David *et al.*, 2010). Economic development and environmental preservation remain two contradictory processes and significant efforts have been made to identify the unique diversity present in New Caledonia for the addition of a large part of its reefs to the UNESCO World Heritage List (Andrefouet and Wantiez, 2010). This is particularly important on the eve of major anthropogenic perturbation linked to the development of a new nickel mine on the island's north-western coast (Chabanet *et al.*, 2010).

In New Caledonia, the major studies on anthropogenic biogeochemical impacts have recently emerged from a large interdisciplinary group funded by the 'Programme National Environnement Cotier' (PNEC) from 2000 to 2008. This program aimed to continue the study of the impact of anthropogenic activities on the functioning of the southwest lagoon of New Caledonia initiated by the 'ECOTROPE' program in 1996. Recent studies have extended the preliminary research on the indicators of human-induced physical disturbance (Chabanet *et al.*, 2005) and particle inputs on coral reefs (Fichez *et al.*, 2005) by developing a framework and the tools for anthropogenic biogeochemical impact studies. Physical processes studied include coastal erosion (Dumas *et al.*, 2010), the circulation and transport of suspended sediment (Ouillon *et al.*, 2010), and the biogeochemical typology and temporal variability of the lagoon waters of New Caledonia (Fichez *et al.*, 2010).

Trace metal contamination and bioaccumulation mechanisms have been studied for the brown alga *Lobophora variegata* (Metian *et al.*, 2008a), the tropical scallop *Comptopallium radula* (Metian *et al.*, 2008b), the tropical oysters *Isognomon isognomon* and *Malleus regula* (Hedouin *et al.*, 2010a,b), the clam *Gafrarium tumidum* (Hedouin *et al.*, 2010b), and the butterfly fish *Chaetodon speculum* (Labonne *et al.*, 2008). With the exception of the butterfly fish, all species were useful bioindicator species for surveying metal contamination. Transplantation studies revealed, however, that due to the very efficient metal retention capacity of the shellfish, particular attention to the origin of bioindicators was required if used in low contamination areas (Hedouin *et al.*, 2011).

A study of coral assemblages from fringing reefs to oceanic barrier reefs in the southwestern lagoon of New Caledonia addressed the specific relationships between the spatial patterns and recruitment processes of coral and water and sediment quality, in particular, the presence of nutrients and trace metals (Adjeroud *et al.*, 2010). Surprisingly, results revealed no clear cross-shelf gradient in richness, abundance, and percent cover for all coral genera from most to least disturbed areas. Instead, the composition and abundance of coral assemblages appeared correlated to the substrate type rather than water quality or the metal concentrations in sediments. The relationship between the distribution of juveniles and adults predicted relatively limited replenishment capacities in these reefs. Consequently, the results of this study illustrated the importance of management and conservation efforts along fringing reefs within bays, where anthropogenic activities and subsequent pollution are important.

The southwest lagoon of New Caledonia was also the site of a study of the spatial and temporal distribution of zooplanktonic prey of fish larvae, i.e., small crustaceans and small copepods, in relation to environmental conditions. Results indicated that the volume and total density of zooplankton were correlated to increasing chlorophyll *a* and particulate organic matter (POM) concentration along a gradient from the lagoon to the bays (Carassou *et al.*, 2010). Differences in assemblage composition between the lagoon and the bays were influenced by wind speed, surface temperature, chlorophyll *a* and POM. The high abundance of zooplanktonic prey in bays suggested that sheltered bays, most influenced by terrigenous inputs, were likely to provide the best feeding conditions and therefore should receive more attention in management and conservation efforts.

b. French Polynesia

Studies on pesticide impacts on the coral reefs of Tahiti (see Figure 1) and Moorea have been summarized by Roche *et al.* (2011). As pesticide contamination has been identified as one of the main stressors of coral reefs and is a potential major threat to coral reef ecosystems (Fabricius *et al.*, 2005; Ramade & Roche, 2006), the French section of the International Coral Reef Initiative (ICRI), the IFRECOR, launched a survey on pesticide residues in key organisms from the coral reef trophic webs in several French Territories, including Tahiti and Moorea.

On each island, surveys were carried out on two inshore coral reef sites using control (traditional agriculture) versus impacted (intensive agriculture) sites in the wet and dry seasons. Key species included the green coralline alga *Halimeda crassata*, the clam *Tridacna maxima*, scleractinian coral species *Fungia spp.*, the holothurian *Halodeima atra*, and fishes *Chlorurus sordidus* and *Epinephelus merra*. Organochlorine pesticides including chlordecone and herbicides from chloracetic acid, triazine and substituted urea families were studied.

The results first demonstrated the presence of chlordecone in all key organisms at all sites, despite this material never having been officially used in French Polynesia (Roche *et al.*, 2011). Additionally, chlordecone average concentrations were higher in organisms from reef sites located off the rural traditional agriculture areas than the ones from intensive agriculture areas, suggesting unregulated widespread use of pesticides in traditional subsistence agriculture (Roche *et al.*, 2011). Secondly, herbicides such as atrazine, simazine, alachlor, metolachlore, and terbutylazine were

also detected in reef organisms from most sites. In some instances, recorded concentrations were sufficient to induce significant photosynthesis inhibition in *Symbodium* (Jones and Kerswell, 2003). Finally, while average organochlorine insecticide levels were generally low, a few samples showed Σ DDT concentrations as high as 1080 ng/g in the liver of grouper and parrot-fish from Tahiti and Σ DDT 393 ng/g in holothuria from Moorea (Roche *et al.*, 2011).

Overall, the contamination of coral reef communities by organochlorine pesticides, insecticides and some herbicides, although widespread, did not uncorrelate with the agriculture intensity in the cultivated coastal areas of Tahiti and Moorea. Some insecticide concentrations observed were among the highest recorded in coral reef communities to date (Deichmann *et al.*, 1972; Glynn *et al.*, 1995; Von Westerhagen & Klumpp, 1995). One of the major results of this study was the unexpected ubiquitous detection of chlordecone in all sites and all organisms sampled. Due to the extreme persistence of this pesticide, the contamination of coral reef communities raises concern for public health as some of the organisms sampled are currently used for consumption by local populations and the risks of long-term exposures are unknown.

c. Mururoa Atoll

Between 1966 and 1996, France conducted 193 nuclear experiments on Mururoa (21°53'S, 138°52'W) and Fangataufa (22°16'S, 138°47'W) atolls in French Polynesia, including 41 in the atmosphere, 137 underground and 15 security tests without nuclear detonation. Since 1966, comprehensive monitoring of radiological conditions has been carried out in order to assess environmental impacts and radiation exposure rates of local populations in the five archipelagos of French Polynesia. This was completed in 1996 by an independent study by the International Atomic Energy Agency (IAEA) following the end of the nuclear experiments. The radiation levels detected in 1996 are now used as a benchmark for the radioactive contamination of the local environment.

Large landslides occurred in Mururoa atoll as a result of nuclear tests in 1979 (Flouzat 2011). While limited to the south-west of Mururoa atoll, these events were responsible for generating large swells that submerged part of the neighbouring island of Tureia, located 105 km northeast of Mururoa at 20°50'S, 138°32'W. As a result of concerns raised by Tureia residents, a numeric model was used in 2011 to determine the hydrologic effects of collapses of portions of the coral structure (volume <50 to 670 million m³) of the northern part of Mururoa atoll (Flouzat, 2011). Simulation results indicated that the predicted hydrologic effects from the collapse of small portions of the coral structure would remain localized to Mururoa atoll without any consequences for neighboring islands. The collapse of a large portion of the northern part of Mururoa atoll, however, was predicted to generate a wave 20 m high within a 500 m radius, traveling at 600 km/hr and with a swell length of several kilometers. At Mururoa, the sea water level would reach 2-5 m in populated areas of the island. Furthermore, the wave induced by the collapse of Mururoa atoll would be expected to reach the south of Tureia atoll in approximately 10 minutes and the north of that island in 13 minutes (Flouzat, 2011). The south and western parts of Tureia atoll would be expected to be submerged and face destruction equivalent to cyclonic waves. In the northern part of the Tureia, areas less than 3 m above sea level would be partially to totally submerged. The majority of the population lives more than 3 m above sea level in the northern part

of the atoll, but for people living less than 3 m above sea level, this potential event would pose a severe threat.

To put the risks from atoll collapse into perspective, coral reefs in French Polynesia, as in most places in Pacific, are subjected to major natural disturbances responsible for damaging the physical and biological structure of coral reefs and their communities. Between 1980 and 2005, French Polynesia has been affected by 15 cyclones, seven bleaching events, several *Acanthaster planci* outbreaks and dystrophic crises (Salvat *et al*, 2008). Out of the 13 islands monitored between 1992 and 2002, only two were unaffected by these natural disturbances and the impact of coral bleaching and cyclones varied greatly at local and regional scales (Adjeroud *et al*, 2005). Cyclones are usually the most destructive and were directly responsible for the destruction of 80% and 40% of coral reefs on the outer-slope of Tikehau and Takapoto islands, respectively (Harmelin-Vivien and Laboute, 1983; Laboute, 1985; Salvat and Wilkinson, 2011). Similarly, historical records and seismic monitoring activities show that French Polynesia is also frequently affected by tsunamis of various amplitudes, the Marquesas Islands and Rurutu Island being the most at risk (Schindele *et al*, 2006; Sladen *et al*, 2007). The 1946 Aleutian and 1960 Chilean events were the last two ocean-wide tsunamis that generated significant damage in most archipelagos and no earthquake of magnitude higher than 8.4 occurred since 1965. Only one tsunami was generated by a local source, in 1999, due to a cliff landslide on Fatu Hiva Island and caused serious damage in adjacent Omoa Bay (Hebert *et al*, 2002).

CONCLUSIONS

The studies described above illustrate that anthropogenic biogeochemical impacts in the Pacific Islands are significant, but most of the problems occur near major population centres or industrial zones. Other issues have been identified but not discussed here – microbial contamination, sedimentation, dynamite fishing and reef blasting for boat passages, cyanide and other toxic chemical fishing. While a large proportion of the early investigations were completed by outsiders/visitors, there is now a body of well-trained local scientists to carry on this work. Unfortunately, many of these professionals remain limited by inadequacies of funding and facilities. The development of sound coastal management and resource protection policies lags behind the more developed countries of the world despite improved interactions between scientists and community leaders, regulators and environmental managers.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the assistance of Dominique Ponton, Bernard Salvat and Rene Galzin with the provision of recent materials on the French Territories in the Pacific.

REFERENCES

Adjeroud, M., Chancerelle, Y., Schrimm, M., Perez, T., Lecchini, D., Galzin, R. and Salvat, B. 2005. Detecting the effects of natural disturbances on coral assemblages in French Polynesia: A decade survey at multiple scales. *Aquatic Living Resources* 18, 111-123.

- Adjeroud, M., Fernandez, J.M., Carroll, A.G., Harrison, P.L. and Penin, L., 2010. Spatial patterns and recruitment processes of coral assemblages among contrasting environmental conditions in the southwestern lagoon of New Caledonia. *Mar. Pollut. Bull.* 61, 375-386.
- Amesbury, S.S., Lassuy, D.R., Myers, R.F. and Vaughan Tyndzik, 1979. *A Survey of the Fish Resources of Saipan Lagoon*. University of Guam Marine Laboratory Technical Report No. 52., Mangilao, Guam, 58 pp.
- Andrefouet, S., and Wantiez, L. 2010. Characterizing the diversity of coral reef habitats and fish communities found in a UNESCO World Heritage Site: The strategy developed for lagoons of New Caledonia. *Mar. Pollut. Bull.* 61, 612–620.
- Carassou, L., Le Borgne, R., Rolland, E. and Ponton, D., 2010. Spatial and temporal distribution of zooplankton related to the environmental conditions in the coral reef lagoon of New Caledonia, Southwest Pacific. *Mar. Pollut. Bull.* 61, 367-374.
- Chabanet, P., Adjeroud, M., Andrefouet, S., Bozec, Y-M., Ferraris, J., Garcia-Charton, J-A. and Schrimm, M., 2005. Human-induced physical disturbances and their indicators on coral reef habitats: A multi-scale approach. *Aquat. Living Resour.* 18, 215–230.
- Chabanet, P., Guillemot, N., Kulbicki, M., Vigliola, L. and Sarra megna, S., 2010. Baseline study of the spatio-temporal patterns of reef fish assemblages prior to a major mining project in New Caledonia (South Pacific). *Mar. Pollut. Bull.* 61, 598-611.
- Chen, Z, Kookana, R., Naidu, R. and Morrison, R.J. 2000. *Pesticide residues in sediment and soil samples from the Kingdom of Tonga: A preliminary study*. Report from CSIRO Land and Water, Glen Osmond, South Australia, to the Department of Environment, Nuku'alofa, Tonga.
- Cumberland, K.B. 1956. *Southwest Pacific: a geography of Australia, New Zealand and their Pacific Island Neighbours*. McGraw-Hill, New York, pp 8-12.
- David, G., Leopold, M., Dumas, P.S., Ferraris, J., Herrenschmidt, J.B. and Fontenelle, G., 2010. Integrated coastal zone management perspectives to ensure the sustainability of coral reefs in New Caledonia. *Mar. Pollut. Bull.* 61, 323-334.
- Deichmann, W.B., Cubit, D.A., MacDonald, W.E., Beasley, A.G., 1972. Organochlorine pesticides in the tissue of the Great barracuda (*Sphyrna barracuda*). *Arch. Toxikol.* 29, 287-309.
- Denton, G.R.W., Concepcion, L.P., Wood, H.R. and Morrison R.J. 2005. Trace Metals in Sediments of Four Harbours in Guam. *Marine Pollution Bulletin*, 50: 1133-1141.
- Denton, G.R.W., B.G. Bearden, Lucrina P. Concepcion, H.R. Wood and R.J. Morrison, 2006a. Contaminant assessment of surface sediments from Tanapag Lagoon, Saipan, Commonwealth of the Northern Marianas Islands. *Marine Pollution Bulletin*, **52**, 703-710.

Denton, G.R.W., Concepcion, L.P., Wood, H.R. and Morrison R.J. 2006b. Polychlorinated Biphenyls (PCBs) in Marine Organisms from Four Harbours in Guam. *Marine Pollution Bulletin* 52: 214-225

Denton, G.R.W., Concepcion, L.P., Wood, H.R. and Morrison R.J. 2006c. Polychlorinated Biphenyls (PCBs) in Sediments from Four Harbours in Guam. *Marine Pollution Bulletin*, 52: 710-718.

Denton, G.R.W., Concepcion, L.P., Wood, H.R. and Morrison R.J. 2006d. Polycyclic Aromatic Hydrocarbons (PAHs) in Small Island Coastal Environments: A Case Study from Harbours in Guam, Micronesia. *Marine Pollution Bulletin*, 52: 1090-1117

Denton, G.R.W., Concepcion, L.P., Wood, H.R. and Morrison R.J. 2006e. Trace Metals in Marine Organisms from Four Harbours in Guam. *Marine Pollution Bulletin*, 52: 1786-1804

Denton, G.R.W., M.C. Olsen and Y. Wen 2007. Solid Waste Disposal on Guam: Impact of an Unsanitary Landfill on the Heavy Metal Status of Adjacent Community Representatives. In: Wang, Y., Li, S., Huang, P., Yuzhong, Y., Ying, A., Sun, X. (Eds.) *Progress in Environmental Science and Technology, Volume 1*. pp. 1169-1176, Proceedings of the 2007 International Symposium on Envir. Science and Technology, Beijing, China, November 13-16, 2007, Science Press USA. Available at: <http://www.weriguam.org/pdf/pollution-monitoring-and-assessment-program/recent-publications/8.pdf>

Denton, G.R.W., Golabi, M.H., Iyekar, C., Wood, H.R., Concepcion, L.P., and Wen, Y. 2008. Impact of Ordot Dump on Water Quality of the Lonfit River Basin in Central Guam. 2. Aqueous Chemical and Biological Contaminants. *Micronesica* 40: 149-167. Available at: <http://www.weriguam.org/pdf/pollution-monitoring-and-assessment-program/recent-publications/7.pdf>

Denton, G.R.W., and R.J. Morrison, 2009. The impact of a landfill on the trace metal status of Pago Bay, Guam. *Marine Pollution Bulletin*, **58**, 150-162.

Denton, G.R.W., R.J. Morrison, B.G. Bearden, P. Houk, J.A. Starmer and H.R. Wood, 2009. Impact of a coastal landfill in a tropical lagoon on trace metal concentrations in surrounding marine biota: A case study from Saipan, Commonwealth of the Northern Mariana Islands (CNMI). *Marine Pollution Bulletin*, **58**, 424-431.

Denton, G.R.W., M.S. Trianni and M.C. Tenorio, 2010. Impact of Land-Based Sources of Pollution on Coastal Water Quality of Saipan, Commonwealth of the Northern Mariana Islands (CNMI): Arsenic, Mercury and PCBs in Popular Table Fish from Saipan Lagoon. Technical Report No. 130. Water and Environmental Research Institute, University of Guam. 98 pp. Available at: <http://www.weriguam.org/reports/item/impact-of-land-based-sources-of-pollution-on-coastal-water-quality-of-saipan-commonwealth-of-the-northern-mariana>

Denton, G.R.W., M.S. Trianni, B.G. Bearden, P.C. Houk, J.A. Starmer, 2011. Impact of a medical waste incinerator on mercury levels in lagoon fish from a small tropical island in the Western Pacific. *J. Toxicology & Environmental Health, Part A*, **74**, 1-5.

Doty, J.E. and Marsh, J.A. Jr. 1977. Marine Survey of Tanapag, Saipan: The Power Barge "Impedance". University of Guam Marine Laboratory Technical Report No 33, Mangilao, Guam, 147 pp.

Dumas, P., Printemps, J., Mangeas, M. and Luneau, G., 2010. Developing erosion models for integrated coastal zone management: A case study of The New Caledonia west coast. *Mar. Pollut. Bull.* **61**, 519–529.

Fabricius, K., De'ath, G., McCook, L., Turak, E. and Williams, D.L., 2005. Changes in algal, coral and fish assemblages along water quality gradients on the inshore Great Barrier Reef. *Mar. Pollut. Bull.* **51**, 384-398.

Fernandez, J-M., Cadiou, G., Moreton, B., Legendre, R., Fischez, R. and Badie, C. (2006). Overview of the metal distribution in surface deposits and sedimentary records in Suva Lagoon. In: R.J. Morrison and W.G.L. Aalbersberg (Editors) *At the Crossroads - Science and Management of Suva Lagoon*, University of the South Pacific, Suva, Fiji, pp 106-119.

Fischez, R., Adjeroud, M., Bozec, Y-M., Breau, L., Chancerelle, Y., Chevillon, C., Douillet, P., Fernandez, J-M., Frouin, P., Kulbicki, M., Moreton, B., Ouillon, S., Payri, C., Perez, T., Sasal, P. and Thébault, J., 2005. A review of selected indicators of particle, nutrient and metal inputs in coral reef lagoon systems. *Aquat. Living Resour.* **18**, 125–147.

Fischez, R., Chifflet, S., Douillet, P., Gerard, P., Gutierrez, F., Jouon, A., Ouillon, S. and Grenz, C., 2010. Biogeochemical typology and temporal variability of lagoon waters in a coral reef ecosystem subject to terrigenous and anthropogenic inputs (New Caledonia). *Mar. Pollut. Bull.* **61**, 309-322.

Flouzat, M., 2011. Surveillance geomechanique de l'atoll de Mururoa: conséquences pour l'atoll de Tureia d'un glissement de terrain de grande ampleur a Mururoa. Rapport du Commissariat a l'Énergie Atomique et aux Energies Alternatives, Arpajon cedex, 13 p.

Gangaiya, P., Tabudravu, J., South, R. and Sotheeswaran, S. (2001). Heavy metal contamination of the Lami coastal environment, Fiji. *S. Pac. J. Natural Sci.*, **19**, 24-29.

Gardes, L. and Salvat B. 2008. Coral reefs in French Overseas Territories: a retrospective study of changes in health conditions of these diversified and vulnerable ecosystems recorded by monitoring networks. *Rev. Ecol. (Terre Vie)* **63**: 23-32.

Glynn, P.W., Rumbold, D.G. and Snedaker, S.C., 1995. Organochlorine pesticides residues in marine sediments and biota from the northern Florida reef tract. *Mar. Pollut. Bull.* **30**, 384-398.

Harmelin-Vivien, M. and Laboute, P. 1983. Preliminary data on underwater effects of cyclones on the outer slopes of Tikehau Island (Tuamotu, French Polynesia) and its fish fauna. *Inter. Soc. For Reef Studies, Nice, Abstracts*: 27.

- Harrison, N.L., P. Gangaiya and R.J. Morrison, 1996. Organochlorines in the coastal marine environment of Vanuatu and Tonga. *Marine Pollution Bulletin*, **32**, 575-579.
- Hebert, H., Piatanesi, A., Heinrich, P., Schindele, F. and Okal EA. 2002. Numerical modeling of the September 13, 1999 landslide and tsunami on Fatu Hiva Island (French Polynesia). *Geophys. Res. Lett.* **29**(10):1484 DOI10.1029/2001GL013774
- Hedoin, L., Metian, M., Teyssie, J.L., Fichez, R. and Warnau, M., 2010a. Delineation of heavy metal contamination pathways (seawater, food and sediment) in tropical oysters from New Caledonia using radiotracer techniques. *Mar. Pollut. Bull.* **61**, 542-553.
- Hedoin, L., Metian, M., Lacoue-Labarthe, T., Fichez, R., Teyssie, J.L., Bustamante, P. and Warnau, M., 2010b. Influence of food on the assimilation of selected metals in tropical bivalves from the New Caledonia lagoon: Qualitative and quantitative aspects. *Mar. Pollut. Bull.* **61**, 568-575.
- Hedoin, L., Pringault, O., Bustamante, P., Fichez, R. and Warnau, M., 2011. Validation of two tropical marine bivalves as bioindicators of mining contamination in the New Caledonia lagoon: Field transplantation experiments. *Water. Res.* **45**, 483-496.
- Jones, R.J. and Kerswell, A.P., 2003. Phytotoxicity of Photosystem II (PSII) herbicides to coral. *Mar. Ecol. Prog. Ser.* **261**, 149-151.
- Kaly, U.L. 1998. Monitoring Training and Lagoon Baseline Survey using the Fanga'uta Lagoon as a Case Study. Working Paper 16. Tonga Environmental Management and Planning Project, Department of Environment, Tonga, 77 p.
- Kaly, U. and R.J. Morrison, 2005. Four years in the life of Fanga'uta Lagoon, Tonga. In: Kay, R. and Alder, J., *Coastal Planning and Management*, 2nd Edition, Taylor & Francis Group, Oxford, pp 317-318.
- Labonne, M., Morize, E., Kulbicki, M., Ponton, D. and Marec, L., 2008. Otolith chemical signature and growth of *Chaetodon speculum* in coastal areas of New Caledonia. *Est. Coast. Shelf Sci.* **78**, 493-504.
- Laboute, P. 1985. Evaluation of damage done by cyclones of 1982-1983 to the outer slopes of the Tikehau and Takapoto atolls (Tuamotu Archipelago). *Proc. Fifth Inter. Coral Reef Congr., Tahiti*, **3** : 323-329.
- Maata, M. and Koshy, K. (2001). A study on tributyltin contamination of marine sediments in the major ports of Fiji. *S. Pacific J. Natural Sci.*, **19**, 1-4.
- Metian, M., Giron, E., Borne, V., Hedouin, L., Teyssie, J.L. and Warnau, M., 2008a. The brown alga *Lobophora variegata*, a bioindicator species for surveying metal contamination in tropical marine environments. *J. Exp. Mar. Biol. Eco.* **362**, 49-54.
- Metian, M., Bustamante, P., Hedouin, L. and Warnau, M., 2008b. Accumulation of nine metals and one metalloid in the tropical scallop *Comptopallium radula* from coral reefs in New Caledonia. *Env. Pollut.* **152**, 543-552.

Morrison, R.J., Harrison, N.L. and Gangaiya, P. (1996). Organochlorines in the estuarine and coastal marine environment of the Fiji Islands. *Environmental Pollution*, **93**, 159-167.

Morrison, R.J., S.P. Narayan and P. Gangaiya. 2001. Trace metal studies in Laucala Bay, Suva, Fiji. *Marine Pollution Bulletin*, **42**, 397-404.

Morrison, R.J. and P.L. Brown, 2003. Trace metals in Fanga'uta Lagoon, Kingdom of Tonga. *Marine Pollution Bulletin*, **46**, 139-152.

Morrison, R.J., P. Gangaiya, S. Garimella, S. K. Singh, M. Maata and A. Chandra, 2006. Contamination of Suva Lagoon. In: R.J. Morrison and W.G.L. Aalbersberg (Editors) *At the Crossroads - Science and Management of Suva Lagoon*, University of the South Pacific, Suva, Fiji, pp 146-155.

Morrison, R.J., Peshut, P.J. and Lasorsa, B.K. 2010. Elemental composition and mineralogical characteristics of coastal marine sediments of Tutuila, American Samoa. *Marine Pollution Bulletin*, **60**, 925-30.

Naidu, S.D. and Morrison, R.J. 1994. Contamination of Suva Harbour, Fiji. *Mar. Poll. Bull.*, **29**, 126-130.

Ouillon, S., Douillet, P., Lefebvre, J.P., Le Gendre, R., Jouon, A., Bonneton, P., Fernandez, J-M., Chevillon, C., Magand, O., Lefèvre, J., Le Hir, P., Laganier, R., Dumas, F., Marchesiello, P., Bel Madani, A., Andréfouët, S., Panché, J-Y. and Fichez, R., 2010. Circulation and suspended sediment transport in a coral reef lagoon: The south-west lagoon of New Caledonia. *Mar. Pollut. Bull.* **61**, 269-296.

Pascal, M., Richer de Forges, B., Le Guyader, H. and, Simberloff, D., 2007. Mining and other threats to the New Caledonia biodiversity hotspot. *Cons. Biol.* **22**, 498-499.

Peshut, P.J. 2009. Mercury Investigations in Remote Oceania: Wet Deposition and Bio-accumulation on Tutuila Island. PhD Thesis, School of Earth and Environmental Sciences, University of Wollongong, Wollongong, Australia, 246 p.

Prescott, N., Kaly, U.L., Taufu, P., Matoto, L., Lepa, S.T., Niu, L., Faletau, T., Hibberd, J.K. and Palaki, A. 2001. Environmental Management Plan for the Fanga'uta Lagoon System. Department of Environment, Tonga, 50 p.

Ramade, F. and Roche, H., 2006. Effets des polluants sur les écosystèmes récifaux. *Rev. Ecol. (Terre Vie)*. **61**, 3-33.

Roche, H., Salvat, B. and Ramade, F., 2011. Assessment of the pesticides pollution of coral reefs communities from French Polynesia. *Rev. Ecol. (Terre Vie)*. **66**, 3-10.

Salvat, B. and Wilkinson, C. 2011. Cyclones and climate change in the South Pacific. *Rev. Ecol. (Terre Vie)*, **66**: 105-115.

Salvat, B., Aubanel, A., Adjeroud, M., Bouisset, P., Calmet, D., Chancerelle, Y., Cochenne, N., Davies, N., Fougere, A., Galzin, R., Lagouy, E., Lo, C., Monier, C., Ponsonnet, C., Remoissenet, G., Schneider, D., Stein, A., Tatarata, M. and Villiers, L.. 2008. Le suivi de l'état des récifs coralliens de Polynésie Française et leur récente évolution. *Rev. Ecol. (Terre Vie)*. **63**, 103-135.

Schindele, F., Hebert, H., Raymond, D. and Sladen, A. 2006. L'aléa tsunami en Polynésie française : synthèse des observations et des mesures. *C. R. Geoscience*. **338**, 1133-1140.

Singh, S., Aalbersberg, W.G.L. and Morrison, R.J. 2009. Nutrient pollution in Laucala Bay, Fiji. *Water Air and Soil Pollution*, **204**, 363-372.

Sladen, A., Hebert, H., Schindele, F. and Raymond, D. 2007. L'aléa tsunami en Polynésie française : apports de la simulation numérique. *C. R. Geoscience*. **339**, 303-316.

Staudigel, H., Hart, S.R., Pile, A., Bailey, B.E., Baker, E.T., Brooke, S., Connelly, D.P., Haucke, L., German, C.R., Hudson, I., Jones, D., Koppers, A.A.P., Konter, J., Lee, R., Pietsh, T.W., Tebo, B.M., Templeton, A.S., Zierenberg, R., Young, C.M. 2006. Vailulu'u seamount, Samoa: Life and death on an active submarine volcano. *Proc. National Academy of Sciences of the United States of America*, **103**, 6448-6453.

Stearns, H.T. 1944. Geology of the Samoan Islands. *Bulletin of the Geological Society of America*, **55**, 1279-1332.

Stewart, C. and de Mora, S.J. (1992). Elevated tri(n-butyl)tin concentrations in shellfish and sediments from Suva Harbour, Fiji. *Appl. Organometallic Chem.*, **6**, 507-512.

Tamata, U.B. 2007. *Studies of nutrient variability and the consequences for benthic communities on the Coral Coast fringing reefs, Fiji*. PhD Thesis, University of Wollongong, Wollongong, Australia, 277 p + appendices.

US EPA (2005). *EPA Orders Medical Waste Incinerator in Saipan to Close*. U.S. Environmental Protection Agency. Available at:
<http://yosemite.epa.gov/opa/admpress.nsf/665323ad33bb55ee852572a000657b63/78d76fafe47b8823852570d8005e178d!OpenDocument>

USFDA (1998). Appendix 5: FDA and EPA Guidance Levels. In: *Fish and Fisheries Products Hazards and Controls Guide, Chapter 9: Environmental Chemical Contaminants and Pesticides (A Chemical Hazard)*. U.S. Food & Drug Administration, Center for Food Safety and Applied Nutrition, Washington, D.C.

Viles, H. and Spenser, T. (1995). *Coastal Problems*. Edward Arnold, London, 350 p.

Von Westerhagen, H. and Klumpp, D.W., 1995. Xenobiotics in fish from Australian tropical coastal waters, including Great Barrier Reef. *Mar. Pollut. Bull.* **51**, 876-881.

Legend for Figure

Figure 1. The South Pacific Region

