

University of Wollongong

Research Online

Faculty of Engineering and Information
Sciences - Papers: Part A

Faculty of Engineering and Information
Sciences

1-1-2015

A note on the boundary behavior for a modified Green function in the upper-half space

Y Zhang

Henan Institute of Education

Valery Piskarev

University of Wollongong

Follow this and additional works at: <https://ro.uow.edu.au/eispapers>



Part of the [Engineering Commons](#), and the [Science and Technology Studies Commons](#)

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

A note on the boundary behavior for a modified Green function in the upper-half space

Abstract

Motivated by (Xu et al. in Bound. Value Probl. 2013:262, 2013) and (Yang and Ren in Proc. Indian Acad. Sci. Math. Sci. 124(2):175-178, 2014), in this paper we aim to construct a modified Green function in the upper-half space of the n -dimensional Euclidean space, which generalizes the boundary property of general Green potential.

Keywords

function, green, modified, behavior, space, half, boundary, upper, note

Disciplines

Engineering | Science and Technology Studies

Publication Details

Zhang, Y. & Piskarev, V. (2015). A note on the boundary behavior for a modified Green function in the upper-half space. Boundary Value Problems, 114 1-7.

RESEARCH

Open Access



A note on the boundary behavior for a modified Green function in the upper-half space

Yulian Zhang¹ and Valery Piskarev^{2*}

*Correspondence:

v.piskarev@outlook.com

²Faculty of Science and Technology,
University of Wollongong,
Wollongong, NSW 2522, Australia
Full list of author information is
available at the end of the article

Abstract

Motivated by (Xu *et al.* in *Bound. Value Probl.* 2013:262, 2013) and (Yang and Ren in *Proc. Indian Acad. Sci. Math. Sci.* 124(2):175-178, 2014), in this paper we aim to construct a modified Green function in the upper-half space of the n -dimensional Euclidean space, which generalizes the boundary property of general Green potential.

Keywords: modified Green function; capacity; upper-half space

1 Introduction and main results

Let \mathbf{R}^n ($n \geq 2$) denote the n -dimensional Euclidean space. The upper half-space H is the set $H = \{x = (x_1, x_2, \dots, x_n) \in \mathbf{R}^n : x_n > 0\}$, whose boundary and closure are ∂H and \bar{H} respectively.

For $x \in \mathbf{R}^n$ and $r > 0$, let $B(x, r)$ denote the open ball with center at x and radius r .

Set

$$E_\alpha(x) = \begin{cases} -\log|x| & \text{if } \alpha = n = 2, \\ |x|^{\alpha-n} & \text{if } 0 < \alpha < n. \end{cases}$$

Let G_α be the Green function of order α for H , that is,

$$G_\alpha(x, y) = E_\alpha(x - y) - E_\alpha(x - y^*), \quad x, y \in \bar{H}, x \neq y, 0 < \alpha \leq n,$$

where $*$ denotes reflection in the boundary plane ∂H just as $y^* = (y_1, y_2, \dots, -y_n)$.

In case $\alpha = n = 2$, we consider the modified kernel function, which is defined by

$$E_{n,m}(x - y) = \begin{cases} E_n(x - y) & \text{if } |y| < 1, \\ E_n(x - y) + \Re(\log y - \sum_{k=1}^{m-1} (\frac{x^k}{ky^k})) & \text{if } |y| \geq 1. \end{cases}$$

In case $0 < \alpha < n$, we define

$$E_{\alpha,m}(x - y) = \begin{cases} E_\alpha(x - y) & \text{if } |y| < 1, \\ E_\alpha(x - y) - \sum_{k=0}^{m-1} \frac{|x|^k}{|y|^{n-\alpha+k}} C_k^{\frac{n-\alpha}{2}}(\frac{x \cdot y}{|x||y|}) & \text{if } |y| \geq 1, \end{cases}$$

where m is a non-negative integer, $C_k^\omega(t)$ ($\omega = \frac{n-\alpha}{2}$) is the ultraspherical (or Gegenbauer) polynomial (see [1]). The expression arises from the generating function for Gegenbauer polynomials

$$(1 - 2tr + r^2)^{-\omega} = \sum_{k=0}^{\infty} C_k^\omega(t) r^k, \quad (1.1)$$

where $|r| < 1$, $|t| \leq 1$ and $\omega > 0$. The coefficient $C_k^\omega(t)$ is called the ultraspherical (or Gegenbauer) polynomial of degree k associated with ω , the function $C_k^\omega(t)$ is a polynomial of degree k in t .

Then we define the modified Green function $G_{\alpha,m}(x, y)$ by

$$G_{\alpha,m}(x, y) = \begin{cases} E_{n,m+1}(x - y) - E_{n,m+1}(x - y^*) & \text{if } \alpha = n = 2, \\ E_{\alpha,m+1}(x - y) - E_{\alpha,m+1}(x - y^*) & \text{if } 0 < \alpha < n, \end{cases}$$

where $x, y \in \overline{H}$ and $x \neq y$. We remark that this modified Green function is also used to give unique solutions of the Neumann and Dirichlet problem in the upper-half space [2–4].

Write

$$G_{\alpha,m}(x, \mu) = \int_H G_{\alpha,m}(x, y) d\mu(y),$$

where μ is a non-negative measure on H . Here note that $G_{2,0}(x, \mu)$ is nothing but the general Green potential.

Let k be a non-negative Borel measurable function on $\mathbf{R}^n \times \mathbf{R}^n$, and set

$$k(y, \mu) = \int_E k(y, x) d\mu(x) \quad \text{and} \quad k(\mu, x) = \int_E k(y, x) d\mu(y)$$

for a non-negative measure μ on a Borel set $E \subset \mathbf{R}^n$. We define a capacity C_k by

$$C_k(E) = \sup \mu(\mathbf{R}^n), \quad E \subset H,$$

where the supremum is taken over all non-negative measures μ such that S_μ (the support of μ) is contained in E and $k(y, \mu) \leq 1$ for every $y \in H$.

For $\beta \leq 0$, $\delta \leq 0$ and $\beta \leq \delta$, we consider the kernel function

$$k_{\alpha,\beta,\delta}(y, x) = x_n^{-\beta} y_n^{-\delta} G_\alpha(x, y).$$

Now we prove the following result. For related results in a smooth cone and tube, we refer the reader to the papers by Qiao (see [5, 6]) and Liao-Su (see [7]), respectively. The readers may also find some related interesting results with respect to the Schrödinger operator in the papers by Su (see [8]), by Polidoro and Ragusa (see [9]) and the references therein.

Theorem *Let $n + m - \alpha + \delta + 2 \geq 0$. If μ is a non-negative measure on H satisfying*

$$\int_H \frac{y_n^{\delta+1}}{(1 + |y|)^{n+m-\alpha+\delta+2}} d\mu(y) < \infty, \quad (1.2)$$

then there exists a Borel set $E \subset H$ with properties:

$$(1) \quad \lim_{x_n \rightarrow 0, x \in H-E} \frac{x_n^{n-\alpha-\beta+\delta+1}}{(1+|x|)^{n+m-\alpha+\delta+2}} G_{\alpha,m}(x, \mu) = 0;$$

$$(2) \quad \sum_{i=1}^{\infty} 2^{i(n-\alpha+\beta+\delta)} C_{k_{\alpha,\beta,\delta}}(E_i) < \infty,$$

where $E_i = \{x \in E : 2^{-i} \leq x_n < 2^{-i+1}\}$.

Remark By using Lemma 4 below, condition (2) in Theorem with $\alpha = 2$, $\beta = 0$, $\delta = 0$ means that E is 2-thin at ∂H in the sense of [10].

2 Some lemmas

Throughout this paper, let M denote various constants independent of the variables in questions, which may be different from line to line.

Lemma 1 *There exists a positive constant M such that $G_{\alpha}(x, y) \leq M \frac{x_n y_n}{|x-y|^{n-\alpha+2}}$, where $0 < \alpha \leq n$, $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$ in H .*

This can be proved by a simple calculation.

Lemma 2 *Gegenbauer polynomials have the following properties:*

$$(1) \quad |C_k^{\omega}(t)| \leq C_k^{\omega}(1) = \frac{\Gamma(2\omega+k)}{\Gamma(2\omega)\Gamma(k+1)}, \quad |t| \leq 1;$$

$$(2) \quad \frac{d}{dt} C_k^{\omega}(t) = 2\omega C_{k-1}^{\omega+1}(t), \quad k \geq 1;$$

$$(3) \quad \sum_{k=0}^{\infty} C_k^{\omega}(1) r^k = (1-r)^{-2\omega};$$

$$(4) \quad |C_k^{\frac{n-\alpha}{2}}(t) - C_k^{\frac{n-\alpha}{2}}(t^*)| \leq (n-\alpha) C_{k-1}^{\frac{n-\alpha}{2}}(1) |t - t^*|, \quad |t| \leq 1, |t^*| \leq 1.$$

Proof (1) and (2) can be derived from [1], p.232. Equality (3) follows from expression (1.1) by taking $t = 1$; property (4) is an easy consequence of the mean value theorem, (1) and also (2). \square

Lemma 3 *For $x, y \in \mathbb{R}^n$ ($\alpha = n = 2$), we have the following properties:*

$$(1) \quad |\Im \sum_{k=0}^m \frac{x^k}{y^{k+1}}| \leq \sum_{k=0}^{m-1} \frac{2^k x_n |x|^k}{|y|^{k+2}};$$

$$(2) \quad |\Im \sum_{k=0}^{\infty} \frac{x^{k+m+1}}{y^k}| \leq 2^{m+1} x_n |x|^m;$$

$$(3) \quad |G_{n,m}(x, y) - G_n(x, y)| \leq M \sum_{k=1}^m \frac{k x_n y_n |x|^{k-1}}{|y|^{k+1}};$$

$$(4) \quad |G_{n,m}(x, y)| \leq M \sum_{k=m+1}^{\infty} \frac{k x_n y_n |x|^{k-1}}{|y|^{k+1}}.$$

The following lemma can be proved by using Fuglede (see [11], Théorème 7.8).

Lemma 4 *For any Borel set E in H , we have $C_{k_{\alpha}}(E) = \hat{C}_{k_{\alpha}}(E)$, where $\hat{C}_{k_{\alpha}}(E) = \inf \lambda(H)$, $k_{\alpha} = k_{\alpha,0,0}$, the infimum being taken over all non-negative measures λ on H such that $k_{\alpha}(\lambda, x) \geq 1$ for every $x \in E$.*

Following [10], we say that a set $E \subset H$ is α -thin at the boundary ∂H if

$$\sum_{i=1}^{\infty} 2^{i(n-\alpha)} C_{k_{\alpha}}(E_i) < \infty,$$

where $E_i = \{x \in E : 2^{-i} \leq x_n < 2^{-i+1}\}$.

3 Proof of Theorem

We write

$$\begin{aligned} G_{\alpha,m}(x, \mu) &= \int_{G_1} G_{\alpha}(x, y) d\mu(y) + \int_{G_2} G_{\alpha}(x, y) d\mu(y) + \int_{G_3} [G_{\alpha,m}(x, y) - G_{\alpha}(x, y)] d\mu(y) \\ &\quad + \int_{G_4} G_{\alpha,m}(x, y) d\mu(y) + \int_{G_5} G_{\alpha,m}(x, y) d\mu(y) \\ &= U_1(x) + U_2(x) + U_3(x) + U_4(x) + U_5(x), \end{aligned}$$

where

$$\begin{aligned} G_1 &= \left\{ y \in H : |x - y| \leq \frac{x_n}{2} \right\}, & G_2 &= \left\{ y \in H : |y| \geq 1, \frac{x_n}{2} < |x - y| \leq 3|x| \right\}, \\ G_3 &= \{ y \in H : |y| \geq 1, |x - y| \leq 3|x| \}, & G_4 &= \{ y \in H : |y| \geq 1, |x - y| > 3|x| \}, \\ G_5 &= \left\{ y \in H : |y| < 1, |x - y| > \frac{x_n}{2} \right\}. \end{aligned}$$

We distinguish the following two cases.

Case 1. $0 < \alpha < n$.

By assumption (1.2) we can find a sequence $\{a_i\}$ of positive numbers such that $\lim_{i \rightarrow \infty} a_i = \infty$ and $\sum_{i=1}^{\infty} a_i b_i < \infty$, where

$$b_i = \int_{\{y \in H : 2^{-i-1} < y_n < 2^{-i+2}\}} \frac{y_n^{\delta+1}}{(1 + |y|)^{n+m-\alpha+\delta+2}} d\mu(y).$$

Consider the sets

$$E_i = \left\{ x \in H : 2^{-i} \leq x_n < 2^{-i+1}, \frac{x_n^{n-\alpha-\beta+\delta+1}}{(1 + |x|)^{n+m-\alpha+\delta+2}} U_1(x) \geq a_i^{-1} 2^{(i-1)\beta} \right\}$$

for $i = 1, 2, \dots$. Set

$$G = \bigcup_{x \in E_i} B\left(x, \frac{x_n}{2}\right).$$

Then $G \subset \{y \in H : 2^{-i-1} < y_n < 2^{-i+2}\}$. Let ν be a non-negative measure on H such that $S_{\nu} \subset E_i$, where S_{ν} is the support of ν . Then we have $k_{\alpha,\beta,\delta}(y, \nu) \leq 1$ for $y \in H$ and

$$\begin{aligned} \int_H d\nu &\leq a_i 2^{(-i+1)\beta} \int_H \frac{x_n^{n-\alpha-\beta+\delta+1}}{(1 + |x|)^{n+m-\alpha+\delta+2}} U_1(x) d\nu(x) \\ &\leq M a_i 2^{(-i+1)\beta} 2^{(-i+1)(n-\alpha+\delta+1)} \int_G k_{\alpha,\beta,\delta}(y, \nu) \frac{y_n^{\delta}}{(1 + |y|)^{n+m-\alpha+\delta+2}} d\mu(y) \\ &\leq M a_i 2^{(-i+1)\beta} 2^{(-i+1)(n-\alpha+\delta+1)} 2^{i+1} \int_{\{y \in H : 2^{-i-1} < y_n < 2^{-i+2}\}} \frac{y_n^{\delta+1}}{(1 + |y|)^{n+m-\alpha+\delta+2}} d\mu(y) \\ &\leq M 2^{n-\alpha+\beta+\delta+2} 2^{-i(n-\alpha+\beta+\delta)} a_i b_i. \end{aligned}$$

So that

$$C_{k_{\alpha,\beta,\delta}}(E_i) \leq M2^{-i(n-\alpha+\beta+\delta)} a_i b_i,$$

which yields

$$\sum_{i=1}^{\infty} 2^{i(n-\alpha+\beta+\delta)} C_{k_{\alpha,\beta,\delta}}(E_i) < \infty.$$

Setting $E = \bigcup_{i=1}^{\infty} E_i$, we see that (2) in Theorem is satisfied and

$$\lim_{x_n \rightarrow 0, x \in H-E} \frac{x_n^{n-\alpha-\beta+\delta+1}}{(1+|x|)^{n+m-\alpha+\delta+2}} U_1(x) = 0. \quad (3.1)$$

For $U_2(x)$, by Lemma 1 we have

$$\begin{aligned} |U_2(x)| &\leq Mx_n \int_{G_2} \frac{y_n}{|x-y|^{n-\alpha+2}} d\mu(y) \\ &\leq Mx_n^{\alpha-n-1} |x|^{n+m-\alpha+\delta+2} \int_{G_2} \frac{1}{y_n^{\delta}} \frac{y_n^{\delta+1}}{(1+|y|)^{n+m-\alpha+\delta+2}} d\mu(y) \\ &\leq Mx_n^{\alpha-n-1} |x|^{n+m-\alpha+2} \int_{G_2} \frac{y_n^{\delta+1}}{(1+|y|)^{n+m-\alpha+\delta+2}} d\mu(y). \end{aligned} \quad (3.2)$$

Note that $C_0^{\omega}(t) \equiv 1$. By (3) and (4) in Lemma 2, we take $t = \frac{x \cdot y}{|x||y|}$, $t^* = \frac{x \cdot y^*}{|x||y^*|}$ in Lemma 2(4) and obtain

$$\begin{aligned} |U_3(x)| &\leq \int_{G_3} \sum_{k=1}^m \frac{|x|^k}{|y|^{n-\alpha+k}} 2(n-\alpha) C_{k-1}^{\frac{n-\alpha+2}{2}}(1) \frac{x_n y_n}{|x||y|} \frac{2|y|^{n+m-\alpha+\delta+2}}{(1+|y|)^{n+m-\alpha+\delta+2}} d\mu(y) \\ &\leq Mx_n |x|^m \sum_{k=1}^m \frac{1}{4^{k-1}} C_{k-1}^{\frac{n-\alpha+2}{2}}(1) \int_{G_3} \frac{y_n^{\delta+1}}{(1+|y|)^{n+m-\alpha+\delta+2}} d\mu(y) \\ &\leq Mx_n |x|^m. \end{aligned} \quad (3.3)$$

Similarly, we have by (3) and (4) in Lemma 2

$$\begin{aligned} |U_4(x)| &\leq \int_{G_4} \sum_{k=m+1}^{\infty} \frac{|x|^k}{|y|^{n-\alpha+k}} 2(n-\alpha) C_{k-1}^{\frac{n-\alpha+2}{2}}(1) \frac{x_n y_n}{|x||y|} \frac{2|y|^{n+m-\alpha+\delta+2}}{(1+|y|)^{n+m-\alpha+\delta+2}} d\mu(y) \\ &\leq Mx_n |x|^m \sum_{k=m+1}^{\infty} \frac{1}{2^{k-1}} C_{k-1}^{\frac{n-\alpha+2}{2}}(1) \int_{G_4} \frac{y_n^{\delta+1}}{(1+|y|)^{n+m-\alpha+\delta+2}} d\mu(y) \\ &\leq Mx_n |x|^m. \end{aligned} \quad (3.4)$$

Finally, by Lemma 1, we have

$$|U_5(x)| \leq Mx_n^{\alpha-n-1} \int_{G_5} \frac{y_n^{\delta+1}}{(1+|y|)^{n+m-\alpha+\delta+2}} d\mu(y). \quad (3.5)$$

Combining (3.1), (3.2), (3.3), (3.4) and (3.5), by Lebesgue's dominated convergence theorem, we prove Case 1.

Case 2. $\alpha = n = 2$.

In this case, $U_1(x)$, $U_2(x)$ and $U_5(x)$ can be proved similarly as in Case 1. Here we omit the details and state the following facts:

$$\lim_{x_n \rightarrow 0, x_n \in H-E} \frac{x_n^{\delta-\beta+1}}{(1+|x|)^{m+\delta+2}} U_1(x) = 0, \quad (3.6)$$

where $E = \bigcup_{i=1}^{\infty} E_i$ and $\sum_{i=1}^{\infty} 2^{i(\beta+\delta)} C_{k_{\alpha,\beta,\delta}}(E_i) < \infty$,

$$\lim_{x_n \rightarrow 0, x_n \in H} \frac{x_n^{\delta-\beta+1}}{(1+|x|)^{m+\delta+2}} [U_2(x) + U_5(x)] = 0. \quad (3.7)$$

By Lemma 3(3), we obtain

$$\begin{aligned} |U_3(x)| &\leq \int_{G_3} \sum_{k=1}^m \frac{k x_n y_n |x|^{k-1}}{|y|^{k+1}} \frac{2|y|^{m+\delta+2}}{y_n^{\delta+1}} \frac{y_n^{\delta+1}}{(1+|y|)^{m+\delta+2}} d\mu(y) \\ &\leq M x_n |x|^m \sum_{k=1}^m \frac{k}{4^{k-1}} \int_{G_3} \frac{y_n^{\delta+1}}{(1+|y|)^{m+\delta+2}} d\mu(y) \\ &\leq M x_n |x|^m. \end{aligned} \quad (3.8)$$

By Lemma 3(4), we have

$$\begin{aligned} |U_4(x)| &\leq \int_{G_4} \sum_{k=m+1}^{\infty} \frac{k x_n y_n |x|^{k-1}}{|y|^{k+1}} \frac{2|y|^{m+\delta+2}}{y_n^{\delta+1}} \frac{y_n^{\delta+1}}{(1+|y|)^{m+\delta+2}} d\mu(y) \\ &\leq M x_n |x|^m \sum_{k=m+1}^{\infty} \frac{k}{2^{k-1}} \int_{G_4} \frac{y_n^{\delta+1}}{(1+|y|)^{m+\delta+2}} d\mu(y) \\ &\leq M x_n |x|^m. \end{aligned} \quad (3.9)$$

Combining (3.6), (3.7), (3.8) and (3.9), we prove Case 2.

Hence the proof of the theorem is completed.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Author details

¹College of Mathematics and Statistics, Henan Institute of Education, Zhengzhou, 450046, China. ²Faculty of Science and Technology, University of Wollongong, Wollongong, NSW 2522, Australia.

Acknowledgements

The authors are highly grateful for the referees' careful reading and comments on this paper. This work was completed while the authors were visiting the Department of Mathematical Sciences at the University of Wollongong, and they are grateful for the kind hospitality of the Department.

Received: 13 April 2015 Accepted: 28 May 2015 Published online: 07 July 2015

References

1. Szegő, G: Orthogonal Polynomials. American Mathematical Society Colloquium Publications, vol. 23. Am. Math. Soc., Providence (1975)
2. Ren, YD, Yang, P: Growth estimates for modified Neumann integrals in a half space. *J. Inequal. Appl.* **2013**, 572 (2013)
3. Xu, G, Yang, P, Zhao, T: Dirichlet problems of harmonic functions. *Bound. Value Probl.* **2013**, 262 (2013)
4. Yang, DW, Ren, YD: Dirichlet problem on the upper half space. *Proc. Indian Acad. Sci. Math. Sci.* **124**(2), 175-178 (2014)
5. Qiao, L: Integral representations for harmonic functions of infinite order in a cone. *Results Math.* **61**, 62-74 (2012)
6. Qiao, L, Pan, GS: Generalization of the Phragmén-Lindelöf theorems for subfunctions. *Int. J. Math.* **24**(8), 1350062 (2013)
7. Liao, Y, Su, BY: Solutions of the Dirichlet problem in a tube domain. *Acta Math. Sin.* **57**(6), 1209-1220 (2014)
8. Su, BY: Dirichlet problem for the Schrödinger operator in a half space. *Abstr. Appl. Anal.* **2012**, Article ID 578197 (2012)
9. Polidoro, S, Ragusa, MA: Harnack inequality for hypoelliptic ultraparabolic equations with a singular lower order term. *Rev. Mat. Iberoam.* **24**(3), 1011-1046 (2008)
10. Armitage, H: Tangential behavior of Green potentials and contractive properties of L^p -potentials. *Tokyo J. Math.* **9**, 223-245 (1986)
11. Fuglede, B: Le théorème du minimax et la théorie fine du potentiel. *Ann. Inst. Fourier* **15**, 65-88 (1965)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com